



## In situ disintegration of protein gels (egg white) by pepsin based on time-lapse confocal microscopy

Juliane Flouri, Geeshani Somaratne, Didier Dupont, Francoise Nau

### ► To cite this version:

Juliane Flouri, Geeshani Somaratne, Didier Dupont, Francoise Nau. In situ disintegration of protein gels (egg white) by pepsin based on time-lapse confocal microscopy. Première réunion plénière du GDR SLAMM (Solliciter LA Matière Molle), Institut National de Recherche Agronomique (INRA). UMR Laboratoire d'Ingénierie des Systèmes Biologiques et des Procédés (0792)., Nov 2018, Hyères, France. hal-01929031

HAL Id: hal-01929031

<https://hal.science/hal-01929031>

Submitted on 5 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.




# *In situ* disintegration of protein gels (egg white) by pepsin based on time-lapse confocal microscopy

G. SOMARATNE<sup>1</sup>, F. NAU<sup>2</sup>, D. DUPONT<sup>2</sup>, J. FLOURY<sup>2</sup>

<sup>1</sup>Ridder Institute, Massey University, Palmerston North, New-Zealand

<sup>2</sup>STLO, Agrocampus Ouest, INRA, Rennes, France

*juliane.floury@agrocampus-ouest.fr*



Première réunion plénière du GDR  
SLAMM  
Mon 12 – Thu 15 Nov 2018  
Hyères, France

# Context



# **FOOD** **NUTRITION**



# HEALTH



# How **food characteristics** can modulate the digestion and **nutrient utilization** by the human body?

# FOOD MATRIX STRUCTURE

# PROTEIN DIGESTION

# Previous results



The extent of ovalbumin *in vitro* digestion and the nature of generated peptides are modulated by the morphology of protein aggregates

Kéra Nyemb <sup>a,b</sup>, Catherine Guérin-Dubiard <sup>a,b</sup>, Didier Dupont <sup>a,b</sup>, Julien Jardin <sup>a,b,\*</sup>  
Françoise Nau <sup>a,b,\*</sup>



## Investigating the ovalbumin digestion multivariate data

Kéra Nyemb<sup>a,b</sup>, Julien J. Luton<sup>a,b</sup>, David Causeur<sup>c</sup>, Catherine Guérin-Dubiard<sup>a,b</sup>, Didier Dupont<sup>a,b</sup>, Shane M. Rutherford<sup>d</sup>, Françoise Nau<sup>a,b,\*</sup>



Investigating the impact of egg white gel structure on peptide kinetics profile during *in vitro* digestion


Kéra Nyemb-Diop<sup>a,b</sup>, David Causeur<sup>c</sup>, Julien Jardin<sup>a,b</sup>, Valérie Briard-Bion<sup>a,b</sup>, Catherine Guérin-Dubiard<sup>a,b</sup>, Shane M. Rutherford<sup>d</sup>, Didier Dupont<sup>a,b</sup>, Françoise Nau<sup>a,b,\*</sup>

**Food matrix structure**  
a deciding parameter for  
**food disintegration**  
during digestion

The gastric phase is a key step in the digestion of semi-solid to solid foods, including protein gels





How do gastric acid secretion and pepsin reach their substrates in the stomach?



# Strategy

Two egg white gels (EWG) [1]

- Identical composition : 10w/w% protein
- **Different microstructures :**



- **100 µL EW** in an IBIDI µ-Slide system
- **Fast Green** added as a protein marker ( $\lambda_{ex}$  633 nm)
- Heat gelation at 80°C/5min

[1] Nyemb et al. (2016) Food Hydrocolloids, 54B, 315-327

# Strategy

EWG disintegration during gastric digestion was observed using the confocal microscope **ZEISS LSM880** equipped with the new **Airy scan detector**

→ high spatial & temporal resolution




Addition of 150 µL SGF\*  
+ pepsin (8000 U/ml)



\*Simulated Gastric Fluid

## Airy scan detector



- ✓ Each of the 32 detector elements acts as its own small pinhole
- ✓ 1.7× **higher resolution & increases signal-to-noise ratio** by 4–8× compared with conventional CSLM using an optimized 1-AU pinhole (Airy Unit)



Tile scan set up to cover **1 mm field of view from the gel surface:**  
→ 15 images captured with the **x63 oil objective**  
→ every 15 min over a **120 min digestion period (37°C)**

Treatment & controls	EWG pH 5 & 9
SGF* pH2 + pepsin	3 rep
SGF* pH2 (no pepsin)	1 rep
Water only	1 rep
Photo-bleaching	1 rep

# Image analysis of the gel microstructures



Black – Particles ; White - Background

## White top hat filter

→ to enhance protein structures

## Smooth

→ to remove acquisition noise


## Otsu threshold

→ to get binary image

## Quantification of microstructural parameters using specific plugins:


- *MorphoLibJ*
- *Granulometry*
- *Geodesics*

- ✓ Area fraction of particles
- ✓ Perimeter density of particle aggregates
- ✓ Pore size distribution
- ✓ Particle size distribution
- ✓ Tortuosity



Silva et al., Food Hydro. 2015

## Results. Initial microstructure of the gels




Area fraction of particle	Perimeter density of particle ( $\mu\text{m}/\mu\text{m}^2$ )	Tortuosity
---------------------------	---	------------

$0.31 \pm 0.08$


$0.92 \pm 0.20$

$1.14 \pm 0.14^*$

Particle size ( $\mu\text{m}$ )



Pore size ( $\mu\text{m}$ )





$0.32 \pm 0.02$

$0.29 \pm 0.02$

$1.93 \pm 0.11^*$

$1.07 \pm 0.01$




Protein particles in green, scale bar 10μm

\*  $p$ -value < 0.05

**EWG5**

**Digestion time (min)**

Distance from gel surface ( $\mu\text{m}$ )





**EWG9**

Digestion time (min)

Distance from gel surface ( $\mu\text{m}$ )


tile	$\mu\text{m}$
15	1019
14	951
13	883
12	815
11	747
10	679
9	611
8	544
7	475
6	408
5	340
4	272
3	204
2	136
1	68



**EWG5**Particle area  
fraction (%)**Digestion time (min)**Distance from gel surface ( $\mu\text{m}$ )

10

EWG9



## Discussion


Diffusion coefficient of FITC-pepsin measured by FRAP (20°C)

Medium	$D_{\text{eff}}$ ( $\mu\text{m}^2 \text{s}^{-1}$ )
Water	$104.5 \pm 10.7^{\text{a}}$
pH5-granular EWG	$52.5 \pm 5.3^{\text{b}}$
pH9-smooth EWG	$44.2 \pm 6.1^{\text{c}}$

It is not possible to conclude if proteolysis is limited by :

- the **diffusion of pepsin inside the gels**
- the **local pH**, which is of **paramount importance on pepsin activity**

Activity of pepsin on ovalbumin in function of the pH



## Perspectives

These technique is valuable to get a better picture on how food characteristics can modulate digestion kinetics

BUT :

Improvement of the proposed methodology is needed to **decouple the acidification phase by SGF from the enzymatic action by pepsin.**

Moreover, more research is required **to produce different microstructures of protein gels with both identical composition and pH.**