



**HAL**  
open science

## Insecticide resistance to organophosphates in *Culex pipiens* complex from Lebanon

Mike Abboud Osta, Zeinab J Rizk, Pierrick Labbe, Mylène Weill, Khouzama Knio

► **To cite this version:**

Mike Abboud Osta, Zeinab J Rizk, Pierrick Labbe, Mylène Weill, Khouzama Knio. Insecticide resistance to organophosphates in *Culex pipiens* complex from Lebanon. *Parasites & Vectors*, 2012, 5 (1), pp.132. 10.1186/1756-3305-5-132 . hal-01938155

**HAL Id: hal-01938155**

**<https://hal.science/hal-01938155>**

Submitted on 28 Nov 2018

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

RESEARCH

Open Access

# Insecticide resistance to organophosphates in *Culex pipiens* complex from Lebanon

Mike A Osta<sup>1\*</sup>, Zeinab J Rizk<sup>1</sup>, Pierrick Labbé<sup>2</sup>, Mylène Weill<sup>2</sup> and Khouzama Knio<sup>1</sup>

## Abstract

**Background:** Analysis of *Culex pipiens* mosquitoes collected from a single site in Lebanon in 2005, revealed an alarming frequency of *ace-1* alleles conferring resistance to organophosphate insecticides. Following this, in 2006 the majority of municipalities switched to pyrethroids after a long history of organophosphate usage in the country; however, since then no studies have assessed the impact of changing insecticide class on the frequency of resistant *ace-1* alleles in *C. pipiens*.

**Methods:** *C. pipiens* mosquitoes were captured indoors from 25 villages across the country and subjected to established methods for the analysis of gene amplification at the *Ester* locus and target site mutations in *ace-1* gene that confer resistance to organophosphates.

**Results:** We conducted the first large-scale screen for resistance to organophosphates in *C. pipiens* mosquitoes collected from Lebanon. The frequency of carboxylesterase (*Ester*) and *ace-1* alleles conferring resistance to organophosphates were assessed among *C. pipiens* mosquitoes collected from 25 different villages across the country between December 2008 and December 2009. Established enzymatic assay and PCR-based molecular tests, both diagnostic of the major target site mutations in *ace-1* revealed the absence of the F290V mutation among sampled mosquitoes and significant reduction in the frequency of G119S mutation compared to that previously reported for mosquitoes collected from Beirut in 2005. We also identified a new duplicated *ace-1* allele, named *ace-1<sup>D13</sup>*, exhibiting a resistant phenotype by associating a susceptible and a resistant copy of *ace-1* in a mosquito line sampled from Beirut in 2005. Fisher's exact test on *ace-1* frequencies in the new sample sites, showed that some populations exhibited a significant excess of heterozygotes, suggesting that the duplicated allele is still present. Starch gel electrophoresis indicated that resistance at the *Ester* locus was mainly attributed to the *Ester<sup>2</sup>* allele, which exhibits a broad geographical distribution.

**Conclusions:** Our analysis suggests that the frequency of resistant *ace-1* alleles in mosquito populations can be downshifted, and in certain cases (F290V mutation) even eliminated, by switching to a different class of insecticides, possibly because of the fitness cost associated with these alleles.

**Keywords:** Insecticide resistance, Acetylcholine esterase, *Culex pipiens*

## Background

An important global strategy to contain mosquito-borne diseases is vector control using chemical insecticides. However, the strong dependence on insecticides for mosquito control worldwide and the use of such chemicals in agriculture has led to the physiological resistance of important mosquito vectors in recent years, including

*Anopheles gambiae* [1,2], *Aedes aegypti* [3-5], and *Culex pipiens* [6,7]. Hence, monitoring insecticide resistance in mosquito populations is crucial in order to ensure the sustainability of vector control programs [8].

Lebanon is a temperate country where two potentially important mosquito vectors of disease are prevalent, *C. pipiens*, which transmits filarial worms, West Nile (WNV) and several encephalitis viruses [9,10], and *Aedes albopictus* the vector for Chikungunya (CHIKV) [11] and dengue viruses (DENV) [12]. *Ae. aegypti*, the primary vector of DENV is not present in the country.

\* Correspondence: mo07@aub.edu.lb

<sup>1</sup>Department of Biology, American University of Beirut, Bliss Street, Beirut 11072020, Lebanon

Full list of author information is available at the end of the article

Despite the prevalence of potential vectors of disease, arboviral diseases are absent from Lebanon with the exception of some cases of WNV infections [13], while WNV has been responsible for numerous and iterative outbreaks in Israel [14], a country at the southern border of Lebanon. In the past, however, both DENV [15] and WNV [16] were highly prevalent in the country and a Dengue epidemic affected thousands of individuals in Beirut between the years 1945 and 1946 [15].


Mosquito control in Lebanon depended heavily on organophosphate (OPs) usage before the year 2006. The most commonly used OPs included, dichlorvos, malathion, diazinon and chlorpyrifos. However, after that date the use of OPs dwindled; dichlorvos, malathion and diazinon were eventually discontinued, while chlorpyrifos has remained in use in a limited number of villages. On the other hand, there has been a significant shift towards the use of pyrethroids in most villages, according to the feedback obtained from several municipalities and major insecticide distributors across the country. The most commonly used pyrethroids are alpha-cypermethrin, deltamethrin and tetramethrin. In Lebanon, insecticides are used almost exclusively to control adult mosquito populations by spraying along the roads in villages and around houses, while no strategies exist to identify and treat larval habitats. Resistance to OPs can be metabolic or due to target site modifications. The former is characterized by the amplification of esterases A and B that sequester these insecticides [17], preventing them from reaching their target, the *ace-1* gene-encoded acetylcholinesterase. Target-site modifications are due to three distinct mutations in *ace-1*, resulting in three substitutions, G119S found in several mosquito species, F290V found only in *Culex pipiens*, and F331W found only in *Culex tritaeniorhynchus*, (numbered according to *Torpedo californica ace* [18]), which independently render the enzyme less sensitive to OP insecticides.

Data from *C. pipiens* mosquitoes sampled in 2005 from Beirut indicated a high frequency of both G119S and F209V mutations [19]. Here, we conducted a large-scale one-year survey, between December 2008 and December 2009, to measure the impact of switching to pyrethroids on the residual OP resistance in *C. pipiens* mosquito populations. The study involved analysis of gene amplification at the *Ester* locus and target site mutations in *ace-1* gene in mosquitoes captured indoors across the country.

## Methods

### Collection sites and mosquito strains

*C. pipiens* mosquitoes were collected indoors from 25 villages across Lebanon (Figure 1) in a one year survey between December 2008 and December 2009. *C. pipiens* reference strains included: SLAB, the susceptible strain



lacking overproduced esterases [20]; SA2, a resistant strain homozygous for *Ester*<sup>2</sup>, characterized by overproduction of esterases A2-B2 [21], and the resistant SR strain homozygous for the G119S mutation [22]. Mosquitoes were reared at room temperature. Larval stages were kept in plastic trays (30 cm x 20 cm) and fed ground fish food (PRODAC), while adult mosquitoes were kept on 10% sucrose solution.

### Starch gel electrophoresis

The frequency of *Ester*<sup>2</sup> (A2-B2) locus among the sampled mosquitoes was determined by Starch gel electrophoresis using TEM 7.4 buffer systems and revealed according to Pasteur *et al.* [23]. The reference strain SA2 [21] was used as a control for esterase amplification at *Ester*<sup>2</sup>.

### Detection of the G119S and F290V mutations in *ace-1*

The G119S mutation was detected using a diagnostic PCR test followed by RFLP, as previously described [24]. Briefly, legs of individual mosquitoes were ground in extraction buffer (0.1 M Tris-HCl, pH 8.0, 0.01 M EDTA, 1.4 M NaCl, 2% cetyltrimethyl ammonium bromide), DNA was then extracted with chloroform, precipitated in isopropanol, and resuspended in sterile water. A 374 bp amplicon was amplified from exon 3 of *ace-1* gene using the primers CpEx3dir, 5'-CGACTCGGACCCACTCGT-3'; and CpEx3-rev, 5'-GACTTGCGACACGGTACTGCA-3'; and the

following PCR conditions: 30 cycles, 95 °C for 5 min, 95 °C for 40 sec, 60 °C for 1 min and 72 °C for 50 sec. Since the G119S mutation in exon 3 creates an AluI restriction site [24], the PCR product was digested with AluI (this generates two fragments of 272 and 102 bp) and products were analyzed on 1.5% agarose gel. Detection of the F290V mutation was performed using the PASA diagnostic test as previously described [25] except that the PCR conditions were modified as follows: 30 cycles, 95 °C for 5 min, 95 °C for 40s, 61 °C for 1 min and 72 °C for 50s.


#### Témoin-dichlorvos-propoxur- (TDP) test

The TDP test was performed exactly as described by Alout *et al.* [19] to identify all possible phenotypes at the *ace-1* locus: phenotype [V], mosquitoes containing only the F290V mutated enzyme; [RR], mosquitoes containing only the G119S mutated enzyme, [SS] those containing susceptible enzyme; [VS], [VR], [RS] and [VRS], mosquitoes containing two or three (VRS) enzyme forms.

#### Detection of *ace-1* gene duplications

The genetic test developed by Labbé *et al.* [6] was used to detect the presence of duplicated alleles. Briefly, females from field *C. pipiens* populations were crossed with males from the susceptible reference SLAB strain. After blood feeding, females were isolated individually, allowed to produce egg rafts and then phenotyped using the TDP enzymatic test [25]. The progeny of each female appearing as heterozygotes [R/S] were reared separately and second instar larvae selected with 1 mg/l propoxur, a dose that kills all susceptible [S/S] individuals. Only the progenies of females carrying a duplicated allele displayed no mortality following the exposure to this insecticide. Real heterozygous [R/S] non-duplicated individuals generate [S/S] individuals in their progenies after crossing with susceptible males (for more explanations, see Figure 2 in [6]). DNA was extracted from frozen females, a fragment of the *ace-1* gene encompassing intron 2 and exon 3 was PCR-amplified, and the products were cloned (to separate the different copies), as previously described [6]. Sequences were then analyzed using the Mega 5 software (<http://www.megasoftware.net/> [26]): Beirut duplicated allele sequences were aligned with known duplicated alleles using ClustalW and a similarity tree was built using Neighbor-Joining with default parameters.

Due to the similarity of duplicated allele sequences with non-duplicated *ace-1* alleles, it is not possible to directly detect the presence of the duplicated allele in natural populations [6,27]. However, an excess of heterozygote phenotypes at the *ace-1* locus have been shown to suggest robustly the presence of a duplicated allele in a population [28,29]. Thus, excess of heterozygotes (as



indicated by a negative  $F_{is}$ , i.e. a significant departure from the expected frequencies under Hardy-Weinberg assumptions) was tested for each sampled population using the Genepop software (Fisher's exact tests, [30]).

## Results and discussion

### Prevalence of *Ester*<sup>2</sup> in sampled mosquitoes

There are many *Ester* alleles associated with resistance to OPs, the most common type is *Ester*<sup>2</sup>, which exhibits a worldwide distribution [31,32]. We used starch gel electrophoresis to identify amplifications at the *Ester* locus. The reference strain SA2 [21] was used as a control for the presence of the *Ester*<sup>2</sup> allele, since preliminary data indicated a high frequency of this allele in a sample collected from Beirut in 2005 (M. Weill, unpublished data). The 2008–2009 survey indicated that approximately 40.8% of sampled mosquitoes carried the *Ester*<sup>2</sup> (A2-B2) resistance allele (Table 1), while the rest showed no other resistance allele at that locus. This,

**Table 1 Frequency of *Ester*<sup>2</sup> phenotype in the mosquito populations sampled**

Location	N	None	[ <i>Ester</i> <sup>2</sup> ]
Beirut	64	0.67 (43)	0.33 (21)
Mount Lebanon	90	0.60 (54)	0.40 (36)
South Lebanon	183	0.51 (93)	0.49 (90)
North Lebanon	153	0.61 (94)	0.39 (59)
Beqaa	134	0.63 (85)	0.37 (49)
<b>Total number</b>	<b>624</b>	<b>369</b>	<b>255</b>

Indicated are the frequencies of the normal (None) and *Ester*<sup>2</sup> phenotypes together with the corresponding number of individuals (in brackets) from the different geographical locations. N is the number of mosquitoes analyzed.

apparently high prevalence, probably reflects the continuous, though limited use of chlorpyrifos in some villages. Moreover, cypermethrin, a pyrethroid used in many municipalities in Lebanon, is known in other organisms to select for resistance due to elevated esterase production [33].

#### Geographical distribution of AChE1 G119S substitution and evidence for *ace-1* duplication

The analysis of insecticide resistance in a small sample of *C. pipiens* mosquitoes collected from Beirut in 2005 revealed an alarming frequency of the G119S resistant AChE1 distributed as follows [19]: 41.4% [RS], 37.9% [RR], 8.6% [VS] and 12.1% [VRS]. None of the mosquitoes sampled exhibited the [SS] phenotype (susceptible enzyme), suggesting that in 2005 there was indeed a strong selective pressure on the *ace-1* locus due to the past heavy usage of OPs. Here, we screened a mosquito line, that was selected with OP for several generations from this 2005 Beirut sample, for the presence of a duplicated *ace-1* allele as previously reported [6]. The duplicated *ace-1* allele (or haplotype) originated from duplication of the *ace-1* gene (termed *ace-1*<sup>D</sup> or D) which associates a resistant and a susceptible copy (termed D(R) and D(S), respectively), resulting in "permanent heterozygotes" [6]. This screen revealed the presence of a new duplicated allele associating the D<sub>7</sub>S susceptible copy already found in Algeria [34] and a new DR resistant copy (Figure 2), hence exhibiting an overall resistant phenotype. This new allele (as of the complete haplotype) should thereafter be named *ace-1*<sup>D13</sup>, according to previous nomenclature [34] (*ace-1*<sup>D13</sup> susceptible and resistant copies are available in GenBank under references JX007790 and JX007791, respectively. All other duplicated alleles can be found in GenBank under references JX007766 to JX007789). Also, considering only the G119S mutation, the *ace-1* allele frequencies in the 2005 sample reported above [19] show a significant departure from the Hardy-Weinberg equilibrium (Genepop, [30], Fisher's exact-test for an excess of heterozygotes;  $F_{is} = -0.34$ ,  $p$ -value = 0.018). This provides

additional support to the presence of at least one *ace-1* duplicated allele in the 2005 Beirut sample, as previously reported in other locations [27,35].

In our more recent samples collected between December 2008 and December 2009, we identified only two genotypes associated with the *ace-1* locus using PCR analysis; mosquitoes carrying only the susceptible *ace-1* allele [SS] and mosquitoes carrying one susceptible and one resistant *ace-1* alleles [RS]. The frequency of [SS] ranged from 0.74 to 0.82 across the five geographical regions, while [RS] ranged between 0.17 and 0.25 (Table 2). The TDP assay gave similar results: the frequency of [SS] and [RS] were 0.76 and 0.24, respectively, in the pool of sampled mosquitoes. Our data suggest that switching to pyrethroids resulted in a dramatic reduction in frequency of resistant mosquitoes, possibly because of the fitness cost associated with this mutation in the absence of a selecting insecticide [36,37]; Indeed, [RR] mosquitoes, supposedly the ones with the highest fitness cost, disappeared. The G119S mutation is known to significantly reduce the enzymatic activity of AChE1 in cholinergic synapses affecting the behavior of mosquitoes, which is probably the cause of this fitness cost [38]. We did not perform genetic crosses to determine whether the [RS] phenotype represented a heterozygous state or was due to the presence of the duplicated allele, since no mosquito lines were maintained from the 2008–2009 collection sites. However, statistical analyses on the *ace-1* frequencies showed that all populations exhibited an excess of heterozygotes, which was significant in the Mount and South Lebanon populations (Table 2), suggesting that the duplicated allele is still present [27,35]. This type of duplicated allele is thought to be less costly than the single R allele, although it is

**Table 2 Frequency of *ace-1* alleles based on diagnostic PCR analysis**

Location	N	[SS]	[RS]	[RR]	[V]	$F_{is}$	$p$ -value
Beirut	76	0.763 (58)	0.236 (18)	0	0	-0.13	0.32
Mount Lebanon	182	0.73 (133)	0.27 (49)	0	0	-0.15	<b>0.02</b>
South Lebanon	159	0.742 (118)	0.258 (41)	0	0	-0.14	<b>0.05</b>
North Lebanon	120	0.80 (96)	0.20 (24)	0	0	-0.11	0.28
Beqaa	107	0.822 (88)	0.178 (19)	0	0	-0.09	0.41
<b>Total number</b>	<b>644</b>	<b>493</b>	<b>151</b>	<b>0</b>	<b>0</b>	<b>-</b>	<b>-</b>

Indicated are the frequencies for each genotype together with the corresponding number of individuals in brackets. N is the number of mosquitoes analyzed. SS, susceptible mosquitoes; RS, mosquitoes heterozygous for G119S; RR, mosquitoes homozygous for G119S; V, mosquitoes containing the F290V mutation. The  $F_{is}$  value indicates whether there is an excess (negative value) or a deficit (positive value) of heterozygotes. The  $p$ -value (Genepop [30], Fisher's exact test) is also indicated. Significant values are in bold.

not always the case [6,35]. However, the observed reduction of [RS] indicates that the duplicated allele is undoubtedly less fit than the susceptible one in the absence of insecticides. We endeavored to collect information on the history of insecticide usage from the municipalities corresponding to the different collection sites; this task was difficult because Lebanon is a free market and there are no unified guidelines for insecticide usage in the country. The information gathered from 31 villages across the country revealed that 5-10% of municipalities are still using chlorpyrifos (an OP) either alone or in combination with pyrethroids, which probably explains why the G119S mutation is still present at low frequencies in *C. pipiens* populations.

Interestingly, the F290V substitution that was previously detected at low frequency only in five Mediterranean countries including Lebanon [19], was not detected in the present study using both the PASA (Table 2) and TDP tests, despite the large samples collected across the country. A plausible explanation for the disappearance of the F290V substitution from natural populations is the increased resort to pyrethroids, which do not select for this mutation. Moreover, even in municipalities where chlorpyrifos is still used, the resistance conferred by the F290V mutation to this insecticide is 150-fold weaker than that conferred by G119S, according to Alout *et al.* [25]. Thus, high fitness cost in addition to weak insecticide selection of the F290V AChE1 may explain the loss of this mutation from *C. pipiens* in Lebanon. The reason why F290V was prevalent at low frequencies in samples collected from Beirut in 2005 may be due to the insecticide dichlorvos which was commonly used at that time together with chlorpyrifos. F290V was shown to confer approximately 10 fold higher resistance to dichlorvos compared to the G119S mutation [25].

## Conclusions

In summary, this study shows that the frequencies of resistant *ace-1* alleles carrying the G119S and F290V substitutions were dramatically reduced in *C. pipiens* mosquitoes collected between 2008 and 2009 when compared to those recorded in 2005. This is probably due to increased dependence on pyrethroid insecticides in recent years that do not select for these mutations. We propose that using a rotation system, whereby the use of different classes of insecticides is alternated on a yearly basis, should maintain resistance *ace-1* alleles at a low frequency in *C. pipiens* mosquitoes. Nevertheless, cross-resistance between pyrethroids and OP through elevated esterases should be considered and evaluated.

## Competing interests

The authors declare that they have no competing interests.

## Acknowledgements

We thank all friends who helped in mosquito collections. We also thank the different municipalities, Boecker International Sal and Debbane Group for providing information regarding insecticide usage in Lebanon. MAO and KK were funded by AUB-URB and Lebanese CNRS, MW was funded by CNRS institutional grants. All sequence data were obtained on the Environmental Genomic Platform of the SFR Montpellier-Environnement-Biodiversité.

## Author details

<sup>1</sup>Department of Biology, American University of Beirut, Bliss Street, Beirut 11072020, Lebanon. <sup>2</sup>Team Genomics of Adaptation, CNRS UMR 5554, Institut des Science de l'Evolution, Université Montpellier 2, Place E. Bataillon, 34095 Montpellier cedex 05, France.

## Authors' contributions

MAO, KK and MW conceived the study. ZR performed the diagnostic PCR analysis and TDP enzymatic assays. Statistical analysis and *ace-1* duplication studies were performed by MW and PL. All authors were involved in data analysis and interpretation. MAO drafted the manuscript. KK, PL and MW critically revised the manuscript. All authors read and approved the final version.

Received: 28 March 2012 Accepted: 3 July 2012

Published: 3 July 2012

## References

1. Koekemoer LL, Spillings BL, Christian RN, Lo TC, Kaiser ML, Norton RA, Oliver SV, Choi KS, Brooke BD, Hunt RH, *et al*: Multiple insecticide resistance in *Anopheles gambiae* (Diptera: Culicidae) from pointe Noire, Republic of the Congo. *Vector Borne Zoonotic Dis* 2011, **11**:1193-1200.
2. Koffi AA, Alou LP, Adja MA, Kone M, Chandre F, N'Guessan R: Update on resistance status of *Anopheles gambiae* s.s. to conventional insecticides at a previous WHOPEs field site, "Yaokoffikro", 6 years after the political crisis in Cote d'Ivoire. *Parasit Vectors* 2012, **5**:68.
3. Dusfour I, Thalmensy V, Gaborit P, Issaly J, Carinci R, Girod R: Multiple insecticide resistance in *Aedes aegypti* (Diptera: Culicidae) populations compromises the effectiveness of dengue vector control in French Guiana. *Mem Inst Oswaldo Cruz* 2011, **106**:346-352.
4. Kamgang B, Marcombe S, Chandre F, Nchoutpouen E, Nwane P, Etang J, Corbel V, Paupy C: Insecticide susceptibility of *Aedes aegypti* and *Aedes albopictus* in Central Africa. *Parasit Vectors* 2011, **4**:79.
5. Lima EP, Paiva MH, de Araujo AP, da Silva EV, da Silva UM, de Oliveira LN, Santana AE, Barbosa CN, de Paiva Neto CC, Goulart MO, *et al*: Insecticide resistance in *Aedes aegypti* populations from Ceara Brazil. *Parasit Vectors* 2011, **4**:5.
6. Labbe P, Berthomieu A, Berticat C, Alout H, Raymond M, Lenormand T, Weill M: Independent duplications of the acetylcholinesterase gene conferring insecticide resistance in the mosquito *Culex pipiens*. *Mol Biol Evol* 2007, **24**:1056-1067.
7. Liu Y, Zhang H, Qiao C, Lu X, Cui F: Correlation between carboxylesterase alleles and insecticide resistance in *Culex pipiens* complex from China. *Parasit Vectors* 2011, **4**:236.
8. Ranson H, N'Guessan R, Lines J, Moiroux N, Nkuni Z, Corbel V: Pyrethroid resistance in African anopheline mosquitoes: what are the implications for malaria control? *Trends Parasitol* 2010, **27**:91-98.
9. Farajollahi A, Fonseca DM, Kramer LD, Marm Kilpatrick A: "Bird biting" mosquitoes and human disease: a review of the role of *Culex pipiens* complex mosquitoes in epidemiology. *Infect Genet Evol* 2011, **11**:1577-1585.
10. Kilpatrick AM: Globalization, land use, and the invasion of West Nile virus. *Science* 2011, **334**:323-327.
11. Thiboutot MM, Kawalekar OU, Shedlock DJ, Khan AS, Sarangan G, Srikanth P, Weiner DB, Muthumani K: Chikungunya: a potentially emerging epidemic? *PLoS Negl Trop Dis* 2010, **4**:e623.
12. Lambrechts L, Scott TW, Gubler DJ: Consequences of the expanding global distribution of *Aedes albopictus* for dengue virus transmission. *PLoS Negl Trop Dis* 2010, **4**:e646.
13. Gallian P, de Micco P, Ghorra P: Seroprevalence of West Nile virus in blood donors at Hotel Dieu de France, Beirut, Lebanon. *Transfusion* 2010, **50**:1156-1158.

14. Weinberger M, Pitlik SD, Gandacu D, Lang R, Nassar F, Ben David D, Rubinstein E, Izhaki A, Mishal J, Kitzes R, et al: **West Nile fever outbreak, Israel, 2000: epidemiologic aspects.** *Emerg Infect Dis* 2001, **7**:686–691.
15. Hitti J, Khairallah A: **A report of recent epidemic of Dengue in Beirut, Lebanon, and some of its complications.** *J Pales Arab Med Assoc* 1945, **1**:150.
16. Garabedian GA, Matossian RM, Musalli MN: **Serologic evidence of arbovirus infection in Lebanon.** *J Med Liban* 1971, **24**:339–350.
17. Vaughan A, Hawkes N, Hemingway J: **Co-amplification explains linkage disequilibrium of two mosquito esterase genes in insecticide-resistant *Culex quinquefasciatus*.** *Biochem J* 1997, **325**(Pt 2):359–365.
18. Duval N, Bon S, Silman I, Sussman J, Massoulie J: **Site-directed mutagenesis of active-site-related residues in Torpedo acetylcholinesterase. Presence of a glutamic acid in the catalytic triad.** *FEBS Lett* 1992, **309**:421–423.
19. Alout H, Labbe P, Berthomieu A, Pasteur N, Weill M: **Multiple duplications of the rare ace-1 mutation F290V in *Culex pipiens* natural populations.** *Insect Biochem Mol Biol* 2009, **39**:884–891.
20. Georghiou GP, Metcalf RL, Gidden FE: **Carbamate-resistance in mosquitos. Selection of *Culex pipiens fatigans* Wiedemann (=C. quinquefasciatus Say) for resistance to Baygon.** *Bull World Health Organ* 1966, **35**:691–708.
21. Berticat C, Rousset F, Raymond M, Berthomieu A, Weill M: **High Wolbachia density in insecticide-resistant mosquitoes.** *Proc Biol Sci* 2002, **269**:1413–1416.
22. Berticat C, Boquien G, Raymond M, Chevillon C: **Insecticide resistance genes induce a mating competition cost in *Culex pipiens* mosquitoes.** *Genet Res* 2002, **79**:41–47.
23. Pasteur N, Iseki A, Georghiou GP: **Genetic and biochemical studies of the highly active esterases A' and B associated with organophosphate resistance in mosquitoes of the *Culex pipiens* complex.** *Biochem Genet* 1981, **19**:909–919.
24. Weill M, Malcolm C, Chandre F, Mogensen K, Berthomieu A, Marquine M, Raymond M: **The unique mutation in ace-1 giving high insecticide resistance is easily detectable in mosquito vectors.** *Insect Mol Biol* 2004, **13**:1–7.
25. Alout H, Berthomieu A, Hadjivassilis A, Weill M: **A new amino-acid substitution in acetylcholinesterase 1 confers insecticide resistance to *Culex pipiens* mosquitoes from Cyprus.** *Insect Biochem Mol Biol* 2007, **37**:41–47.
26. Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S: **MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods.** *Mol Biol Evol* 2011, **28**:2731–2739.
27. Djogbenou L, Labbe P, Chandre F, Pasteur N, Weill M: **Ace-1 duplication in *Anopheles gambiae*: a challenge for malaria control.** *Malaria J* 2009, **8**:70.
28. Djogbenou L, Noel V, Agnew P: **Costs of insensitive acetylcholinesterase insecticide resistance for the malaria vector *Anopheles gambiae* homozygous for the G119S mutation.** *Malaria J* 2010, **9**:12.
29. Lenormand T, Guillemaud T, Bourguet D, Raymond M: **Appearance and sweep of a gene duplication: adaptive response and potential for new functions in the mosquito *Culex pipiens*.** *Evolution* 1998, **52**:1705–1712.
30. Raymond M, Rousset F: **Genepop (Version 1.2): population genetics software for exact tests and ecumenicism.** *J Hered* 1995, **86**:248–249.
31. Labbe P, Lenormand T, Raymond M: **On the worldwide spread of an insecticide resistance gene: a role for local selection.** *J Evol Biol* 2005, **18**:1471–1484.
32. Raymond M, Berticat C, Weill M, Pasteur N, Chevillon C: **Insecticide resistance in the mosquito *Culex pipiens*: what have we learned about adaptation?** *Genetica* 2001, **112**–113:287–296.
33. Zhang L, Gao X, Liang P: **Beta-cypermethrin resistance associated with high carboxylesterase activities in a strain of house fly, *Musca domestica* (Diptera: Muscidae).** *Pestic Biochem Physiol* 2007, **89**:65–72.
34. Alout H, Labbe P, Pasteur N, Weill M: **High incidence of ace-1 duplicated haplotypes in resistant *Culex pipiens* mosquitoes from Algeria.** *Insect Biochem Mol Biol* 2010, **41**:29–35.
35. Labbe P, Berticat C, Berthomieu A, Unal S, Bernard C, Weill M, Lenormand T: **Forty years of erratic insecticide resistance evolution in the mosquito *Culex pipiens*.** *PLoS Genet* 2007, **3**:e205.
36. Berticat C, Bonnet J, Duchon S, Agnew P, Weill M, Corbel V: **Costs and benefits of multiple resistance to insecticides for *Culex quinquefasciatus* mosquitoes.** *BMC Evol Biol* 2008, **8**:104.
37. Rivero A, Magaud A, Nicot A, Vezilier J: **Energetic cost of insecticide resistance in *Culex pipiens* mosquitoes.** *J Med Entomol* 2011, **48**:694–700.
38. Bourguet D, Guillemaud T, Chevillon C, Raymond M: **Fitness costs of insecticide resistance in natural breeding sites of the mosquito *Culex pipiens*.** *Evolution* 2004, **58**:128–135.

doi:10.1186/1756-3305-5-132

**Cite this article as:** Osta et al.: Insecticide resistance to organophosphates in *Culex pipiens* complex from Lebanon. *Parasites & Vectors* 2012 **5**:132.

**Submit your next manuscript to BioMed Central and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
www.biomedcentral.com/submit

