



Morphological observations of fluorescence in different tick species (Family: Ixodidae)

Dakota A. Shade, Anna R. Oller, Stephen W. Wilson

► To cite this version:

Dakota A. Shade, Anna R. Oller, Stephen W. Wilson. Morphological observations of fluorescence in different tick species (Family: Ixodidae). *Acarologia*, 2018, 58 (4), pp.989-994. 10.24349/acarologia/20184305 . hal-01940211v2

HAL Id: hal-01940211

<https://hal.science/hal-01940211v2>

Submitted on 16 Apr 2024


HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Morphological observations of fluorescence in different tick species (Family: Ixodidae)

Dakota A. Shade ^{a,b}, Anna R. Oller ^a, Stephen W. Wilson^a

^a Department of Biology and Earth Science, University of Central Missouri, WCM 306, Warrensburg, MO 64093, USA.

^b Independent Researcher, Kansas City, MO, USA.

Corrected version 16 April 2024

ABSTRACT


Ticks are important ectoparasites due to their ability to transmit harmful pathogens and their study is thus significant for medical and veterinary practices around the world. Fluorescence has been found in numerous biological organisms, including ticks. However, the overall function, evolutionary significance, and distribution of fluorescence in ticks are unknown. This study examined tick fluorescence in purchased, unfed adult *Amblyomma americanum* (Linnaeus), *A. maculatum* Koch, *Dermacentor variabilis* (Say), *D. andersoni* (Stiles), *Ixodes scapularis* Say, and *Rhipicephalus sanguineus* Latreille ticks. An Olympus SZX7 microscope and NightSea SFA system with ultraviolet (UV) and royal blue sets were used for observing tick morphology. Ticks were shown to have widespread fluorescence in all species observed. The results corroborated previous studies, but with the addition of more fluorescent structures. Tick fluorescence could potentially play roles in communication or predator avoidance. Studies on tick behavior in conjunction with fluorescence are now called for to elucidate potential roles.

Keywords fluorescence; ticks; morphology; ixodid; ultraviolet

Introduction

Ticks are important ectoparasites to the medical and veterinary world due to their capability of transmitting pathogenic organisms (Ghosh *et al.*, 2007). Fluorescence is a phenomenon that can be observed in numerous biological systems and has been previously found in ticks (Dillinger and Kesel, 2002). It occurs due to the excitation of an electron into a higher energy orbital. The return of the electron to the ground state results in the emission of a photon of lower energy (Lakowicz, 2006). The phenomenon can be due to different compounds, microscopic structures and a mixture of both. Little knowledge exists as to the overall function, distribution, and evolution of fluorescence in biological systems (Lagorio *et al.*, 2015). Fluorescence has been found in multiple arthropods including crustaceans (Giurginca *et al.*, 2015), insects (Phillips, 1959; Vukusic and Hooper, 2005; Jiang *et al.*, 2012), and arachnids (Volschenk, 2005; Gaffin *et al.*, 2012; Andrews *et al.*, 2007). Andrews *et al.* (2007) showed that every spider they examined had several fluorescent structures. As the presence of external fluorescence varied among spiders, they concluded it must have an adaptive role in some species (Andrews *et al.*, 2007). Jumping spiders that preyed on orb weaving spiders were shown to approach UV reflective webs more often than non-UV reflective webs, suggesting UV may possibly play roles in communication and interactions between arthropods (Li and Lim, 2005). Dillinger and Kesel (2002) were the first to observe fluorescence in *Ixodes ricinus* Linnaeus (Acari: Ixodidae) ticks. They described fluorescence in the alloscutum, leg joints, between the idiosoma and capitulum, setae, and various structures of the mouthparts (Dillinger and Kesel, 2002). The

Received 22 January 2018
Accepted 21 November 2018
Published 30 November 2018

Corresponding author
Dakota A. Shade 
dakota.a.shade@gmail.com

Academic editor
McCoy, Karen

DOI
10.24349/acarologia/20184305

 Copyright
Shade D. A. *et al.*

Licensed under
Creative Commons CC-BY 4.0

presence and role of fluorescence in other ticks is unknown. The scope of this study is thus to perform morphological observations of fluorescence in different tick species and discuss its potential adaptive significance.

Materials and Methods

Six individuals of each of six species of unfed adults (three males and three females) were purchased from the Oklahoma State University tick lab: *Amblyomma americanum* (Linnaeus), *A. maculatum* Koch, *Dermacentor variabilis* (Say), *D. andersoni* (Stiles), *Ixodes scapularis* Say, and *Rhipicephalus sanguineus* Latreille (Acari: Ixodidae). Ticks were sacrificed and stored in 70% ethanol 24 hours prior to imaging. Images were captured using an Olympus SZX7 stereo microscope (Olympus Corporation, Tokyo, Japan) with an attached Zeiss axiocam 105 camera (Carl Zeiss, Oberkochen, Germany). The excitation sources were a NightSea SFA-UV (360-380 nm) with a 412 nm longpass filter and NightSea SFA-Royal Blue (RB) (440-460 nm) with a 500 nm longpass filter (NightSea, Lexington Massachusetts). ZEN imaging software was used for imaging and ImageJ for image enhancements.

Results

Under UV excitation (360-380 nm), the fluorescence appeared blue in structures. Under RB excitation (440-460 nm), the fluorescence was green/yellow in color. All species observed in this study showed similar fluorescence with few exceptions (Figures 1-3).

Table 1 shows the positive and negative results for fluorescence in both sexes. The following structures showed fluorescence in all species and sexes: anal groove, genital pore, spiricals, setae, ventral idiosoma, hypostome, palpal articles, cheliceral sheath, leg segments, arthrodial membrane, and the apotele. All species with eyes and ornamentation present showed fluorescence in those structures (Figure 1). Fovae and lamellae were intentionally observed



Figure 1 Micrographs showing a dorsal view of *Dermacentor variabilis* (male) to compare each light source. White light (left), royal blue (RB, 440-460 nm) (middle) and ultraviolet (UV, 360-380 nm) (right). Variation in ornamentation illuminated by the different light sources can also be noted.

only in *A. americanum* females due to limited access to a fluorescent microscope, but these structures are presumed to be fluorescent in females of the other species. Portions of the basis capitulum are not fluorescent in *I. scapularis* contrary to the other species.

Table 1 Positive (+) and negative (-) fluorescence in body structures among tick species and sexes (M for Male and F for Female). Multiple plus signs indicate a general increase in visual fluorescence compared to other structures. This is due to some structures appearing brighter in one sex compared to the other.

Fluorescent Structure	<i>A. americanum</i>	<i>A. maculatum</i>	<i>D. andersoni</i>	<i>D. variabilis</i>	<i>I. scapularis</i>	<i>R. sanguineus</i>
Idiosoma						
Anal Groove	+++ (F) / + (M)	+(F) / +(M)	+(F)	+(F)	++ (F) / + (M)	+
Genital Pore	++ (F) / + (M)	++ (F) / + (M)	+(F) / ++ (M)	+	++	+
Spirical	+	+	+	+	++ (F) / + (M)	+
Scutum	+ (Patchy)	-	-	-	-	-
Ornamentation	++	++	++	++	-	-
Eyes	+++	+	+++ (F) / + (M)	++	-	++
Setae	+	+	+	+	++	+
Ventral Idiosoma	+	+(F) / ++ (M)	+(F) / ++ (M)	+(F) / ++ (M)	++ (F) / + (M)	+
Fovae	+					
Lamellae	+(F)					
Dorso-Posterior Groove					++ (M)	
Capitulum						
Hypostome	+	+	+	+	++	++
Articles	+	+	+	+	+	+
Basis Capitulum	+	+	+	+	-	+
Cheliceral Sheath	+	+	+	+	+	+
Interior Palps					+	
Legs						
Leg Segments	+	+	+	+	+	+
Arthroidal Membrane	+	+	+	+	+	+
Apotele	+	+	+	+	+	+
Syncoxae					+	

Discussion

Fluorescence appears to be widespread in the six species studied and is likely the case for all species of ixodid ticks. Much of the observed fluorescence appears to be related to flexible cuticle and may be a by-product of the compounds used to make this layer. Dillinger and Kesel (2002) hypothesized fluorescence found in *I. ricinus* was due to resilin, but the study did not test for the protein. Andersen and Roepstorff (2005) concluded the alloscutum of *I. ricinus* contains 8% chitin, meaning most proteins had a Rebers-Riddiford consensus and are bound to other proteins by cross-links with other compounds. The components of resilin, dityrosine and trityrosine, were found to be present, however, the amount was much less compared to insect resilin. 3-monochlorotyrosine and 3,5-dichlorotyrosine were isolated in the distended alloscutum, but were concluded to play a role in cross-linking or were products of oxidative stress (Andersen and Roepstorff, 2005). The nature of these compounds and their role in fluorescence is unknown.

The anal groove of females in almost all species observed appears to have brighter fluorescence compared to their male counterparts (Figure 2). The fluorescence of the female anal groove could be due to pre-emptive preparation for expansion while feeding. As the tick feeds, the alloscutum extends to a considerable size, but the anal groove does not expand in size during this process. In contrast, ixodid males feed very little in the adult stage and do not increase significantly in size, suggesting a link between feeding morphology and fluorescence.




Figure 2 Fluorescent (RB, 440-460 nm) ventral micrographs of *Amblyomma americanum* male (left) and female (right). The anal groove is found in the lower-center of each image.

I found that *I. scapularis* has fluorescent syncoxae on the underside of the coxa of legs 1 and 2 (Figure 3). The purpose of these structures are unknown, but they do not expand during engorgement. Fluorescence in such structures may thus be by-products of shared compounds used in flexible cuticle e.g., leg segments. In other observed fluorescent structures that do not appear to be involved in expansion, the molecular make-up may be different. For example, tick ornamentation does not have a structural purpose, but also shows fluorescence that differs in comparison to other fluorescent structures. Interestingly, most ornate tick species have eyes. Ornamentation fluorescence may therefore serve one or more functions in ticks. Indeed, Kaltenrieder *et al.* (1989) showed that the spectral acuity of tick eyes has features of both spider eyes and insect ocelli. The use of visual cues may aid in aggregation and mate choice, ensuring viable copulations occur on or off the host. Ornamentation could also potentially serve a function in disruptive/cryptic coloration. Crypsis is a trait that reduces an animal's risk of being detected (Stevens and Merilaita, 2009). Ticks may utilize fluorescent ornamentation as a form of coloration to break up their overall shape. Indeed, the ornamentation fluorescence contrasts well with the dark non-fluorescent portions of the tick and this contrast can be deceptive to a visual predator. Crypsis/disruptive coloration could potentially benefit those tick species that quest in more open habitat compared to those that quest in leaf litter and/or dense wooded areas. All the species examined here tend to search for their hosts in open habitats. Intraspecific and interspecific behavioral studies will now be needed to ascertain whether this is a viable avenue to explore further.

Conclusion

This study showed that ticks exhibit wide range fluorescence like other related arthropods. New fluorescent structures have been identified and the number is likely to increase as more tick species are viewed under UV light. Fluorescence may play a role in intraspecific communication in tick species or act in crypsis to avoid detection by potential predators. Chemical and behavior




Figure 3 Fluorescent (UV, 360-380 nm) micrograph of the ventral view of an *Ixodes scapularis* male.

studies on ticks is now called for to better understand the true significance of fluorescence in these arthropods. The future potential for exploring fluorescence in ticks appears to be very fruitful, as it may uncover more unknown biological characteristics of ticks.

Acknowledgements

I wish to thank my daughter Molly for all her help in the lab. “Don’t touch that!”

ORCID

Dakota A. Shade  <https://orcid.org/0000-0003-4753-5232>

Anna R. Oller  <https://orcid.org/0000-0003-1671-6540>

References

- Andersen, S. O., Roepstorff, P., 2005. The extensible alloscutal cuticle of the tick, *Ixodes ricinus*. *Insect Biochemistry and Molecular Biology*. 35(10): 1181-1188. [doi:10.1016/j.ibmb.2005.05.009](https://doi.org/10.1016/j.ibmb.2005.05.009)
- Andrews, K., Reed, S. M., Masta, S. E., 2007. Spiders fluoresce variably across many taxa. *Biology Letters*, 3(3): 265-267. [doi:10.1098/rsbl.2007.0016](https://doi.org/10.1098/rsbl.2007.0016)
- Dillinger, S. C. G., Kesel, A. B., 2002. Changes in the structure of the cuticle of *Ixodes ricinus* L. 1758 (Acari, Ixodidae) during feeding. *Arthropod Structure and Development*. 31(2): 95-101. [doi:10.1016/S1467-8039\(02\)00042-7](https://doi.org/10.1016/S1467-8039(02)00042-7)
- Gaffin, D. D., Bumm, L. A., Taylor, M. S., Popokina, N. V., Mann, S., 2012. Scorpion fluorescence and reaction to light. *Animal Behaviour*. 83(2): 429-436. [doi:10.1016/j.anbehav.2011.11.014](https://doi.org/10.1016/j.anbehav.2011.11.014)
- Ghosh, S., Azhahianambi, P., Yadav, M. P., 2007. Upcoming and future strategies of tick Control: a review. *Journal of Vector Borne Diseases*. 44(2): 79-89.
- Giurginca, A., Šustr, V., Tajovský, K., Giurginca, M., Matei, I., 2015. Spectroscopic parameters of the cuticle and ethanol extracts of the fluorescent cave isopod *Mesoniscus graniger* (Isopoda, Oniscidea). *ZooKeys*. (515): [doi:10.3897/zookeys.515.9395](https://doi.org/10.3897/zookeys.515.9395)

- Jiang, L., Dong, B., Liu, X., Liu, F., Zi, J., 2012. Structural origin of sexual dichromatic coloration and luster in the beetle *Goliathus cacicus*. Chinese Science Bulletin. 57(24): 3211-3217. doi: [10.1007/s11434-012-5343-4](https://doi.org/10.1007/s11434-012-5343-4)
- Kaltenrieder, M., Labhart, T., Hess, E., 1989. Spectral sensitivity, absolute threshold, and visual field of two tick species, *Hyalomma dromedarii* and *Amblyomma variegatum*. Journal of Comparative Physiology A. 165(2): 155-164. doi:[10.1007/BF00619190](https://doi.org/10.1007/BF00619190)
- Lagorio, M. G., Cordon, G. B., Iriel, A., 2015. Reviewing the relevance of fluorescence in Biological systems. Photochemical and Photobiological Sciences. 14(9): 1538-1559. doi:[10.1039/C5PP00122F](https://doi.org/10.1039/C5PP00122F)
- Lakowicz, J. R. 2006. Principles of fluorescence spectroscopy, 3rd Edn. Springer. New York, NY. pp. 63-69. doi:[10.1007/978-0-387-46312-4_3](https://doi.org/10.1007/978-0-387-46312-4_3)
- Li, D., Lim, M. L., 2005 Ultraviolet cues affect the foraging behaviour of jumping spiders. Animal Behaviour. 70(4): 771-776. doi:[10.1016/j.anbehav.2004.12.021](https://doi.org/10.1016/j.anbehav.2004.12.021)
- Phillips, L. S., 1959. Fluorescence in the colors of certain Lepidoptera observed under ultraviolet light. Journal of the Lepidopterists' Society. 13(2): 73-77.
- Stevens, M., Merilaita, S., 2009. Animal camouflage: current issues and new perspectives. Philosophical Transactions of the Royal Society B: Biological Sciences. 364(1516): 423-427. doi:[10.1098/rstb.2008.0217](https://doi.org/10.1098/rstb.2008.0217)
- Volschenk, E. S., 2005. A new technique for examining surface morphosculpture of scorpions. Journal of Arachnology. 33(3): 820-825. doi:[10.1636/S03-047.1](https://doi.org/10.1636/S03-047.1)
- Vukusic, P., Hooper, I., 2005. Directionally controlled fluorescence emission in butterflies. Science. 310(5751): 1151. doi:[10.1126/science.1116612](https://doi.org/10.1126/science.1116612)