



Comment on: 'Homozygous knockout of the piezo1 gene in the zebrafish is not associated with anemia'

Adèle Faucherre, K. Kissa, Joël Nargeot, Matteo E. Mangoni, Chris Jopling

► To cite this version:

Adèle Faucherre, K. Kissa, Joël Nargeot, Matteo E. Mangoni, Chris Jopling. Comment on: 'Homozygous knockout of the piezo1 gene in the zebrafish is not associated with anemia'. *Haematologica*, 2016, 101 (1), pp.e38. 10.3324/haematol.2015.137398 . hal-01940741

HAL Id: hal-01940741

<https://hal.science/hal-01940741>

Submitted on 30 Nov 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Comment on: 'Homozygous knockout of the *piezo1* gene in the zebrafish is not associated with anemia'

We read with interest the recent letter by Shmuckler et al. 'Homozygous knockout of the *piezo1* gene in the zebrafish is not associated with anemia' published in *Haematologica*.¹ In their letter the authors describe the characterization of a zebrafish *piezo1* knockout (KO) line generated using zinc finger nucleases. Having ensured that the *piezo1* mRNA had been truncated the authors then compared their *piezo1* KO zebrafish with our previous results on *piezo1* morphants (generated using antisense morpholinos to block mRNA translation or splicing) published as an article in *Haematologica* 'Piezo1 plays a role in erythrocyte volume homeostasis'.² In our article we reported that knockdown of *piezo1* in zebrafish embryos resulted in fragile, overhydrated, spherocytic erythrocytes leading to hemolysis and anemia. In contrast, Shmuckler et al. report that they did not observe any of these defects in erythrocytes obtained from either embryonic or adult *piezo1* KO zebrafish save for a slight reduction in volume as opposed to the increase in volume we observed in *piezo1* morphants. The authors subsequently discuss why there is a discrepancy between our two reports and while they highlight the possibility that the *piezo1* morpholinos are non-specific (although 2 different morpholinos that have the same effect specifically on erythrocytes would be highly unusual) they also acknowledge that the problem may also lie with their *piezo1* KO line. Lastly, the authors compare their findings with a recent article published in *eLife* by Cahalan et al. entitled 'Piezo1 links mechanical forces to red blood cell volume' in which *piezo1* was specifically knocked out in mouse erythrocytes.³ They conclude that the findings of Cahalan et al. are consistent with their own observations that knocking out *piezo1* has little effect on erythrocytes.

However, having carefully studied the letter by Shmuckler et al. we believe an oversight may have been made by the authors which has subsequently led to a possible misinterpretation of the data from Cahalan et al. First, we would like to offer some background information on the current morpholino/knockout controversy within the zebrafish community. At the beginning of 2015 Kok et al. published an article in *Developmental Cell* comparing morphant phenotypes with KO phenotypes in zebrafish.⁴ The conclusion of this article was that around 80% of the published morphant phenotypes did not occur when the same genes were knocked out. They concluded that these discrepancies were most likely due to non-specific effects of morpholinos.⁴ This was obviously very worrisome, with some members of the zebrafish community suggesting that all morpholino-based phenotypes must be verified by KO of the same gene. However, an article was published later this year in *Nature* by Rossi et al. which has added a new angle to this debate.⁵ In their article the authors describe how knockdown of a particular gene using morpholinos leads to a specific vascular phenotype in zebrafish embryos. However, when they generated the KO line they did not observe this phenotype. By comparing the KO and morphant proteomes and transcriptomes they were able to determine that a compensatory response had been activated in the KO line that was absent from the morphants. In particular they found a number of genes were upregulated in the KO line which effectively rescued the phenotype observed in morphants. This argues that just because a KO zebrafish line fails to produce the same phenotype as a

morphant it cannot be immediately concluded that the morphant is at fault.

This phenomenon may also have led to an oversight by Shmuckler et al. In their letter the authors conclude that the lack of any observable erythrocyte phenotype in their *piezo1* KO zebrafish line is in agreement with the mouse erythrocyte specific *piezo1* KO phenotype described by Cahalan et al. However, the *piezo1* KO zebrafish line is a global KO of *piezo1* and should first be compared to the mouse global KO of *piezo1* not the erythrocyte-specific KO line. Complete KO of *piezo1* in mice, as Shmuckler et al. indicate, results in early embryonic lethality associated with defective vasculogenesis. This phenotype is very different from the zebrafish *piezo1* KO line which appears to all intents and purposes to have no observable phenotype. The authors do not explain why there appears to be such a large discrepancy between these phenotypes. We suggest at least two possibilities, firstly, *piezo1* could be completely dispensable in zebrafish or a compensatory mechanism has been triggered in the zebrafish *piezo1* KO line, similar to the observations made by Rossi et al. Furthermore, we believe that the lack of any phenotype in the *piezo1* KO zebrafish line has possibly led the authors to misinterpret the data from Cahalan et al. Cahalan et al. showed that erythrocyte-specific KO of *piezo1* in mice leads to overhydrated, fragile erythrocytes which undergo intravascular hemolysis. This would appear to be more in line with our own observations of *piezo1* morphant zebrafish. In fact Shmuckler et al. described that the only measurable phenotype in the zebrafish *piezo1* KO line is a slight reduction in erythrocyte volume which appears to be at odds with the data from Cahalan et al. Although we agree the anemic phenotype we observe in *piezo1* morphants is more pronounced than that of the mouse phenotype it would appear that the actual mechanism by which *piezo1* regulates erythrocyte volume is conserved.

Adèle Faucherre,^{1,2,3} Karima Kissa,⁴ Joël Nargeot,^{1,2,3} Matteo E. Mangoni,^{1,2,3} and Chris Jopling^{1,2,3}

¹CNRS, UMR-5203, Institut de Génomique Fonctionnelle, Département de Physiologie, Labex ICST, Montpellier; ²INSERM, U 1191, Montpellier; ³Universités de Montpellier, UMR-5203; and ⁴CNRS, UMR-5235, Dynamique des Interactions Membranaires Normales et Pathologiques, Université Montpellier, F-34094, France

Correspondence: chris.jopling@igf.cnrs.fr
doi:10.3324/haematol.2015.137398

Key words: *piezo 1*, zebrafish, anemia, erythrocytes.

Information on authorship, contributions, and financial & other disclosures was provided by the authors and is available with the online version of this article at www.haematologica.org.

References

- Shmuckler BE, Huston NC, Thon JN, et al. Homozygous knockout of the *piezo1* gene in the zebrafish is not associated with anemia. *Haematologica*. 2015 Aug 20. [Epub ahead of print]
- Faucherre A, Kissa K, Nargeot J, Mangoni ME, Jopling C. Piezo1 plays a role in erythrocyte volume homeostasis. *Haematologica*. 2014;99(1):70-75.
- Cahalan SM, Lukacs V, Ranade SS, Chien S, Bandell M, Patapoutian A. Piezo1 links mechanical forces to red blood cell volume. *Elife*. 2015;4.
- Kok FO, Shin M, Ni CW, et al. Reverse genetic screening reveals poor correlation between morpholino-induced and mutant phenotypes in zebrafish. *Dev Cell*. 2015;32(1):97-108.
- Rossi A, Kontarakis Z, Gerri C, et al. Genetic compensation induced by deleterious mutations but not gene knockdowns. *Nature*. 2015;524(7564):230-233.