



HAL
open science

Comparison of local spherical characters and the Ichino-Ikeda conjecture for unitary groups

Raphaël Beuzart-Plessis

► **To cite this version:**

Raphaël Beuzart-Plessis. Comparison of local spherical characters and the Ichino-Ikeda conjecture for unitary groups. *Journal de l'Institut de Mathématiques de Jussieu*, 2021, 20 (6), pp.1803-1854. 10.1017/S1474748019000707 . hal-01941343

HAL Id: hal-01941343

<https://hal.science/hal-01941343>

Submitted on 30 Nov 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Comparison of local spherical characters and the Ichino-Ikeda conjecture for unitary groups

Raphaël Beuzart-Plessis *

November 30, 2018

Abstract

In this paper, we prove a conjecture of Wei Zhang on comparison of certain local spherical characters from which we draw some consequences for the Ichino-Ikeda conjecture for unitary groups.

Contents

1	Introduction	2
2	Preliminaries	6
2.1	General notations and conventions	6
2.2	Analytic families of distributions	7
2.3	Base Change for unitary groups	8
2.4	The local Gan-Gross-Prasad conjecture for unitary groups	11
2.5	Measures	12
3	Spherical characters, the Ichino-Ikeda conjecture and Zhang’s conjecture	14
3.1	Global spherical characters	14
3.2	Local spherical characters	15
3.3	Orbital integrals	17
3.4	Correspondence of orbits and transfer	20
3.5	Transfer of spherical characters, Zhang’s conjecture and Ichino-Ikeda conjecture	23
3.6	A globalization result	27

*Université d’Aix-Marseille, I2M-CNRS(UMR 7373), Campus de Luminy, 13288 Marseille Cédex 9, France
rbeuzart@gmail.com

MSC 2010 classification: 22E50, 22E55 (primary), 11F70 (secondary)

4	Proof of Zhang’s conjecture	28
4.1	A result of Zhang on truncated local expansion of the spherical character I_{Π}	29
4.2	Weak comparison of local spherical characters	31
4.3	A local trace formula	32
4.4	Proof of Theorem 4.0.1	33
4.5	A first corollary	36
4.6	Proof of Theorem 3.5.8	37
A	Compaision of relative trace formulae	37
A.1	Norms on adelic varieties	37
A.2	Relative trace formulae	52
A.3	Proof of Theorem 3.5.1	56

1 Introduction

Let E/F be a quadratic extension of number fields. Let V be a $(n+1)$ -dimensional hermitian space over E and let $W \subset V$ be a nondegenerate hyperplane. Set $G = U(W) \times U(V)$ and $H = U(W)$. We view H as a subgroup of G via the natural diagonal embedding. Let π be a cuspidal automorphic representation of $G(\mathbb{A})$. Define the H -period of π to be the linear form $\mathcal{P}_H : \pi \rightarrow \mathbb{C}$ given by

$$\mathcal{P}_H(\phi) = \int_{H(F)\backslash H(\mathbb{A})} \phi(h)dh, \quad \phi \in \pi$$

where dh stands for the Tamagawa Haar measure on $H(\mathbb{A})$ (the integral is absolutely convergent by cuspidality of π). Let $BC(\pi)$ be the base change of π to $GL_n(\mathbb{A}_E) \times GL_{n+1}(\mathbb{A}_E)$ (known to exist thanks to the recent work of Mok [Mok] and Kaletha, Mínguez, Shin and White [KMSW]). We may decompose $\pi = \pi_n \boxtimes \pi_{n+1}$ with π_n, π_{n+1} cuspidal automorphic representations of $U(W)$ and $U(V)$ respectively. We have a similar decomposition $BC(\pi) = BC(\pi_n) \boxtimes BC(\pi_{n+1})$ with $BC(\pi_n), BC(\pi_{n+1})$ two automorphic representations of $GL_{n,E}$ and $GL_{n+1,E}$ respectively. Let $L(s, BC(\pi))$ denote the L -function of pair $L(s, BC(\pi_n) \times BC(\pi_{n+1}))$ defined by Jacquet, Piatetskii-Shapiro and Shalika. If π is tempered everywhere (meaning that for all place v the local representation π_v is tempered), a famous conjecture of Gan, Gross and Prasad links the nonvanishing of the period \mathcal{P}_H to the nonvanishing of the central value $L(1/2, BC(\pi))$ (see [GGP,conjecture 24.1] for a precise statement). In the influential paper [II], Ichino and Ikeda have proposed a refinement of this conjecture for orthogonal groups in the form of an exact formula relating these two invariants. This conjecture has been suitably extended to unitary groups by N. Harris in his Ph.D. thesis ([Ha]). These formulas are modeled on the celebrated work of Waldspurger ([Wald3]) on toric periods for GL_2 .

In two recent papers ([Zh1], [Zh2]), W.Zhang has proved both the Gan-Gross-Prasad and the Ichino-Ikeda conjectures for unitary groups under some local assumptions on π . More precisely, Zhang proves the Gan-Gross-Prasad conjecture under some mild local assumptions

(mainly that π is supercuspidal at one place of F which splits in E , see [Zh1, Theorem 1.1]) but he only gets the Ichino-Ikeda conjecture under far more stringent assumptions (see [Zh2, Theorem 1.2]). This discrepancy is due to some local difficulties that we shall discuss shortly. In [Zh2], Zhang makes a series of conjectures (one for every place of F) which if true would allow to considerably weaken the assumptions of [Zh2, Theorem 1.2]. The goal of this paper is to prove this conjecture at all nonarchimedean place of F . Thus, it will allow us to derive new cases of the Ichino-Ikeda conjecture.

Let us now formulate the Ichino-Ikeda conjecture in a form suitable to our purpose. We assume from now on that π is everywhere tempered. Set

$$\mathcal{L}(s, \pi) := \Delta_{n+1} \frac{L(s, BC(\pi))}{L(s + \frac{1}{2}, \pi, Ad)}$$

where Δ_{n+1} is the following product of special values of Hecke L -functions

$$\Delta_{n+1} := \prod_{i=1}^{n+1} L(i, \eta_{E/F}^i)$$

$\eta_{E/F}$ being the idele class character associated to the extension E/F and where the adjoint L -function of π is defined by

$$L(s, \pi, Ad) := L(s, BC(\pi_n), As^{(-1)^n}) L(s, BC(\pi_{n+1}), As^{(-1)^{n+1}})$$

(see [GGP, §7] for the definition of the Asai L -functions). For all place v of F , we will denote by $\mathcal{L}(s, \pi_v)$ the corresponding quotient of local L -functions. To the period \mathcal{P}_H we associate a global spherical character J_π . It is a distribution on the Schwartz space $\mathcal{S}(G(\mathbb{A}))$ of $G(\mathbb{A})$ given by

$$J_\pi(f) = \sum_{\phi \in \mathcal{B}_\pi} \mathcal{P}_H(\pi(f)\phi) \overline{\mathcal{P}_H(\phi)}$$

for all $f \in \mathcal{S}(G(\mathbb{A}))$ and where \mathcal{B}_π is a (suitable) orthonormal basis of π for the Petersson inner product

$$(\phi, \phi')_{Pet} = \int_{G(F) \backslash G(\mathbb{A})} \phi(g) \overline{\phi'(g)} dg$$

(where dg is the Tamagawa Haar measure on $G(\mathbb{A})$). We also define local spherical characters as follows. Fix factorizations $dg = \prod_v dg_v$ and $dh = \prod_v dh_v$ of the Tamagawa Haar measures on $G(\mathbb{A})$ and $H(\mathbb{A})$ respectively. For all place v of F , we define a local spherical character $J_{\pi_v} : \mathcal{S}(G(F_v)) \rightarrow \mathbb{C}$ (where $\mathcal{S}(G(F_v))$ denotes the Schwartz space of $G(F_v)$) by

$$J_{\pi_v}(f_v) = \int_{H(F_v)} \text{Trace}(\pi_v(h)\pi_v(f_v)) dh_v, \quad f_v \in \mathcal{S}(G(F_v))$$

(the integral is absolutely convergent by temperedness of π_v). For almost all place v of F , if f_v is the characteristic function of $G(\mathcal{O}_v)$ we have

$$J_{\pi_v}(f_v) = \mathcal{L}\left(\frac{1}{2}, \pi_v\right) \text{vol}(H(\mathcal{O}_v)) \text{vol}(G(\mathcal{O}_v))$$

We define a normalized spherical character $J_{\pi_v}^{\natural}$ by

$$J_{\pi_v}^{\natural}(f_v) = \frac{J_{\pi_v}(f_v)}{\mathcal{L}\left(\frac{1}{2}, \pi_v\right)}$$

Finally, we will write S_{π} for the component group associated to the L -parameter of π . It is a 2-abelian group and if $BC(\pi)$ is cuspidal we have $S_{\pi} \simeq (\mathbf{Z}/2\mathbf{Z})^2$. We can now state the Ichino-Ikeda conjecture as follows:

Conjecture 1.0.1 (Ichino-Ikeda) *Assume that π is everywhere tempered. Then, for all factorizable test function $f = \prod_v f_v \in \mathcal{S}(G(\mathbb{A}))$ we have*

$$J_{\pi}(f) = |S_{\pi}|^{-1} \mathcal{L}\left(\frac{1}{2}, \pi\right) \prod_v J_{\pi_v}^{\natural}(f_v)$$

Note that the Ichino-Ikeda conjecture is not usually stated this way but rather in a form involving directly the (square of the absolute value of the) period \mathcal{P}_H and some local periods (see [II,conjecture 1.5] and [Ha,conjecture 1.2]), see however [Zh2, lemma 1.7] for the equivalence between the two formulations.

The main tool used by Zhang to attack conjecture 1.0.1 is a comparison of certain (simple) relative trace formulae that have been proposed by Jacquet and Rallis ([JR]). To carry this comparison, we need a fundamental lemma and the existence of smooth matching. The fundamental lemma for the case at hand has been proved by Yun ([Yu]) in positive characteristic and extended by J. Gordon to characteristic 0 in the appendix to [Yu]. The existence of smooth matching at nonarchimedean places is one of the main achievements of Zhang in [Zh1]. It has been recently extended in a weak form by Xue ([Xue]) to archimedean places. The comparison between the two trace formulae has been done by Zhang in [Zh1]. The output is an identity relating the spherical J_{π} (under some mild local assumptions on π) to certain periods on the base-change of π . More precisely, there is a certain spherical character $I_{BC(\pi)}$ attached to these periods and we get an equality between $J_{\pi}(f)$ and $I_{BC(\pi)}(f')$ up to an explicit factor for *nice* matching functions f and f' (see [Zh2, Theorem 4.3] and Theorem 3.5.1 below). Thanks to the work of Jacquet, Piatetskii-Shapiro and Shalika on Rankin-Selberg convolutions we know an explicit factorization for $I_{BC(\pi)}$ in terms of local (normalized) spherical characters $I_{BC(\pi_v)}^{\natural}$ (see [Zh2, Proposition 3.6]). As a consequence, we also get an explicit factorization of J_{π} . However, this factorization is still in terms of the local spherical characters $I_{BC(\pi_v)}^{\natural}$ which are living on (products of) general linear groups. In order to get the Ichino-Ikeda conjecture we need to compare them with our original local spherical characters $J_{\pi_v}^{\natural}$. It is precisely the content of the following conjecture of Zhang (see [Zh2, conjecture 4.4] and conjecture 3.5.5 for precise statements):

Conjecture 1.0.2 (Zhang) *Let v be a place of F . Then for all matching functions $f_v \in \mathcal{S}(G(F_v))$ and $f'_v \in \mathcal{S}(G'(F_v))$ we have*

$$I_{BC(\pi)_v}^\natural(f'_v) = C(\pi_v) J_{\pi_v}^\natural(f_v)$$

where $C(\pi_v)$ is some explicit constant.

Together with the above-mentioned comparison of relative trace formulae, this conjecture implies the Ichino-Ikeda conjecture under mild local assumptions (see [Zh2, Proposition 4.5]). Zhang was able to verify his conjecture in certain particular cases. More precisely, in [Zh2] the above conjecture is proved for split places or when the representation π_v is unramified (and the residual characteristic is sufficiently large) or supercuspidal (see Theorem 4.6 of *loc.cit*). This explains the very strong conditions that are imposed on π in [Zh2, Theorem 1.2]. The main purpose of this paper is to prove conjecture 1.0.2 at every nonarchimedean place. Our main result thus reads as follows (see Theorem 3.5.7):

Theorem 1.0.3 *For every nonarchimedean place v of F , conjecture 1.0.2 holds at v .*

As a consequence of this theorem we obtain the following result towards conjecture 1.0.1 (see Theorem 3.5.8):

Theorem 1.0.4 *Let π be cuspidal automorphic representation of $G(\mathbb{A})$ which is everywhere tempered. Assume that all the archimedean places of F split in E and that there exists a nonarchimedean place v_0 of F such that $BC(\pi_{v_0})$ is supercuspidal. Then conjecture 1.0.1 holds for π .*

The main new ingredient in the proof of Theorem 1.0.3 is a group analog of the local relative trace formula for Lie algebras developed by Zhang in [Zh1, §4.1]. Actually, this local trace formula can be derived directly from results contained in [Zh1] and [Beu] so that the proof of it is rather brief (see §4.3). We then deduce Theorem 1.0.3 from a combination of this local trace formula with certain results of Zhang on truncated local expansion of spherical characters (see [Zh2, §8] and §4.1).

We now briefly describe the content of each section. In section 1, we set up the notations, fix the measures and recall a number of results (in particular concerning global and local base-change for unitary groups and the local Gan-Gross-Prasad conjecture) that will be needed in the sequel. In section 2 we mainly recall the work of Zhang on comparison of global relative trace formulae, we state precisely conjecture 1.0.2 as well as the main results (Theorem 3.5.7 and Theorem 3.5.8). Section 3 is devoted to the proofs of Theorem 1.0.3 and Theorem 1.0.4. In section 4, we explain how we can remove the temperedness assumption in Theorem 3.5.8. Finally, we have included an appendix to prove that the simple Jacquet-Rallis trace formulae are still absolutely convergent for test functions which are not necessarily compactly supported (but nevertheless rapidly decreasing). For this, we define certain norms on the automorphic quotient $[G] := G(F) \backslash G(\mathbb{A})$ and establish their basic properties. This material

is certainly classical but the author hasn't be able to find a convenient reference, hence we provide complete proofs. It has however interesting consequences e.g. for H a closed subgroup of G we can give a criterion under which every cuspidal form on $[G]$ is integrable on $[H]$ (see Proposition A.1.1 (ix), the criterion simply being that the variety $H \backslash G$ is quasi-affine).

Acknowledgement: I thank Volker Heiermann, Wee-Teck Gan and Hang Xue for useful comments on an earlier draft of this paper. This work has been done while the author was a Senior Research Fellow at the National University of Singapore and the author would like to thank this institution fo its warm hospitality.

2 Preliminaries

2.1 General notations and conventions

In this paper E/F will always be a quadratic extension of number fields or of local fields of characteristic zero. We will always denote by $Tr_{E/F}$ the corresponding trace and by $x \mapsto \bar{x}$ the nontrivial F -automorphism of E . Moreover, we will fix a nonzero element $\tau \in E$ such that $Tr_{E/F}(\tau) = 0$. The notation $R_{E/F}$ will stand for the Weil restriction of scalars from E to F . For every finite dimensional hermitian space V over E we will denote by $U(V)$ the corresponding unitary group and we will write $\mathfrak{u}(V)$ for its Lie algebra. The standard maximal unipotent subgroup of GL_n will be denoted by N_n . For all connected reductive group G over F we will write Z_G for the center of G . For all $n \geq 1$ we define a variety S_n over F by

$$S_n := \{s \in R_{E/F}GL_n; s\bar{s} = 1\}$$

and its "Lie algebra" \mathfrak{s}_n by

$$\mathfrak{s}_n := \{X \in R_{E/F}M_n; X + \bar{X} = 0\}$$

We have a surjective map $\nu : R_{E/F}GL_n/GL_n \rightarrow S_n$ given by $\nu(g) = g\bar{g}^{-1}$ which, by Hilbert 90, is surjective at the level of k -points for any field k . We will denote by \mathfrak{c} the Cayley map $\mathfrak{c} : X \mapsto (X + 1)(X - 1)^{-1}$ which realizes a birational isomorphism between \mathfrak{s}_n and S_n and also between $\mathfrak{u}(V)$ and $U(V)$ for all finite dimensional hermitian space V over E .

Assume that the fields E and F are local. We will then denote by $|\cdot|_F$ the normalized absolute value on F (and similarly for E) and by $\eta_{E/F}$ the quadratic character of F^\times associated to the extension E/F . We will also fix an extension η' of $\eta_{E/F}$ to E^\times and a nontrivial additive character $\psi : F \rightarrow \mathbb{C}^\times$. We will set $\psi_E(z) = \psi(\frac{1}{2}Tr_{E/F}(z))$ for all $z \in E$. Let G be a reductive connected group over F . By a representation of $G(F)$ we will always mean a smooth representation if F is p -adic and an admissible smooth Fréchet representation of moderate growth if F is archimedean (see [BK], [Ca] and [Wall, section 11]). We will denote by $Irr(G)$, $Irr_{unit}(G)$ and $Temp(G)$ the set of isomorphism classes of irreducible, irreducible unitary, irreducible tempered representations of $G(F)$ respectively. We will endow these sets

with the Fell topology (see [Tad]). For any parabolic subgroup $P = MU$ of G (U denoting the unipotent radical of P and M a Levi factor) and for any irreducible representation σ of $M(F)$ we will denote by $i_P^G(\sigma)$ the normalized parabolic induction of σ . The notation $\Psi_{unit}(G)$ will stand for the group of unitary unramified characters of $G(F)$. The space of Schwartz functions $\mathcal{S}(G(F))$ consists of locally constant compactly supported functions if F is p -adic and functions rapidly decreasing with all their derivatives if F is archimedean (see [Beu, §1.4]). If F is p -adic and Ω is a finite union of Bernstein components of $G(F)$ (see [BD]), we will denote by $\mathcal{S}(G(F))_\Omega$ the corresponding summand of $\mathcal{S}(G(F))$ (for the action by left translation). Finally, if π is an irreducible generic representation of $GL_n(E)$ we will denote by $\mathcal{W}(\pi, \psi_E)$ the Whittaker model of π with respect to ψ_E . It is a space of smooth functions $W : G(F) \rightarrow \mathbb{C}$ satisfying the relation

$$W(ug) = \psi_E\left(\sum_{i=1}^{n-1} u_{i,i+1}\right)W(g)$$

for all $u \in N_n(E)$ and such that π is isomorphic to $\mathcal{W}(\pi, \psi_E)$ equipped with the $G(F)$ -action by right translation.

In the number field case, we will denote by \mathbb{A} and \mathbb{A}_E the adèle rings of F and E respectively and by $\eta_{E/F}$ the idele class character associated to the extension E/F . We will fix an extension η' of $\eta_{E/F}$ to \mathbb{A}_E^\times . For every place v of F we will denote by F_v the corresponding completion, $\mathcal{O}_v \subset F_v$ the ring of integers (if v is nonarchimedean) and we will set $E_v = E \otimes_F F_v$, $\mathcal{O}_{E,v} = \mathcal{O}_E \otimes_{\mathcal{O}_F} \mathcal{O}_v$ where $\mathcal{O}_F, \mathcal{O}_E$ denote the ring of integers in F and E respectively. If S is a finite set of places of F , we define $F_S = \prod_{v \in S} F_v$. If Σ is a (usually infinite) set of places of F , we will write \mathbb{A}_Σ for the restricted product of the F_v for $v \in \Sigma$. We will also fix a nontrivial additive character $\psi : \mathbb{A}/F \rightarrow \mathbb{C}^\times$ and we will set $\psi_E(z) = \psi(\frac{1}{2}Tr_{E/F}(z))$ for all $z \in \mathbb{A}_E$. For all place v of F , we will denote by $\psi_v, \psi_{E,v}$ and η'_v the local components at v of ψ, ψ_E and η' respectively. Let G be a connected reductive group over F . We will set $[G] = G(F) \backslash G(\mathbb{A})$ and for all place v of F we will denote by G_v the base-change of G to F_v . The Schwartz space $\mathcal{S}(G(\mathbb{A}))$ of $G(\mathbb{A})$ is by definition the restricted tensor product of the local Schwartz spaces $\mathcal{S}(G(F_v))$. We will denote by $\mathcal{U}(\mathfrak{g}_\infty)$ the enveloping algebra of the complexification of the Lie algebra \mathfrak{g}_∞ of $\prod_{v|\infty} G(F_v)$ and by $C_G \in \mathcal{U}(\mathfrak{g}_\infty)$ the Casimir element. If a maximal compact subgroup $K = \prod_v K_v$ of $G(\mathbb{A})$ has been fixed, we will also denote by $C_K \in \mathcal{U}(\mathfrak{g}_\infty)$ the Casimir element of $K_\infty := \prod_{v|\infty} K_v$. Finally if $\eta : \mathbb{A}^\times / F^\times \rightarrow \mathbb{C}^\times$ is an idele class character and $g \in GL_n(\mathbb{A})$ we will usually abbreviate $\eta(\det g)$ by $\eta(g)$.

2.2 Analytic families of distributions

Assume that F is a local field. Let G be a connected reductive group over F and let $\pi \mapsto L_\pi$ be a family of (continuous if F is archimedean) linear forms on $\mathcal{S}(G(F))$ indexed by the set $Temp(G(F))$ of all irreducible tempered representations of $G(F)$. Assume that the following condition is satisfied:

For all parabolic subgroup $P = MU$ of G and for all square-integrable

representation σ of $M(F)$ there is at most one irreducible subrepresentation π of $i_P^G(\sigma)$ such that $L_\pi \neq 0$.

This condition is for example automatically satisfied if $G = GL_n$ (as in this case the representation $i_P^G(\sigma)$ is always irreducible). If this condition is satisfied, we may extend the family of distributions $\pi \mapsto L_\pi$ to any induced representation $i_P^G(\sigma)$ as above by setting $L_{i_P^G(\sigma)} = L_\pi$ if π is the unique irreducible subrepresentation of $i_P^G(\sigma)$ such that $L_\pi \neq 0$ and $L_{i_P^G(\sigma)} = 0$ if no such subrepresentation exists. We then say that this family is *analytic* if for all $f \in \mathcal{S}(G(F))$, all parabolic subgroup $P = MU$ and all square-integrable representation σ of $M(F)$ the function

$$\chi \in \Psi_{unit}(M) \mapsto L_{i_P^G(\sigma \otimes \chi)}(f)$$

is analytic (recall that $\Psi_{unit}(M)$ being a compact real torus has a natural structure of analytic variety).

2.3 Base Change for unitary groups

Let E/F be a quadratic extension of local fields of characteristic zero (either archimedean or p -adic). Let V be a n -dimensional hermitian space over E . Recall that the set of Langlands parameters for $U(V)$ is in one-to-one correspondence with the set of $(-1)^{n+1}$ -conjugate dual continuous semisimple representations φ of the Langlands group \mathcal{L}_E of E (see [GGP, §3] for a definition of ϵ -conjugate dual representations). In what follows, by a Langlands parameter for $U(V)$ we shall mean a representation φ of this sort. By the recent results of Mok [Mok] and Kaletha-Minguez-Shin-White [KMSW] on the local Langlands correspondence for unitary groups together with the work of Langlands [Lan] for real groups, we know that there exists a canonical decomposition

$$Irr(U(V)) = \bigsqcup_{\varphi} \Pi^{U(V)}(\varphi)$$

indexed by the set of all Langlands parameters for $U(V)$. The sets $\Pi^{U(V)}(\varphi)$ are finite (some of them may be empty) and called L -packets. By the Langlands classification, the above decomposition boils down to an analog decomposition of the tempered dual

$$Temp(U(V)) = \bigsqcup_{\varphi} \Pi^{U(V)}(\varphi)$$

where the union is over the set of *tempered* Langlands parameters for $U(V)$ i.e. the parameters φ whose image is bounded. This last decomposition admits a characterization in terms of endoscopic relations (see [Mok, Theorem 3.2.1] and [KMSW, Theorem 1.6.1]) and of the (known) Langlands correspondence for $GL_d(E)$ ([He], [HT], [S]). By this Langlands correspondence, every parameter φ of $U(V)$ determines an irreducible representation $\pi(\varphi)$ of $GL_n(E)$. If π is in the L -packet corresponding to φ we will write $BC(\pi) := \pi(\varphi)$. If π is tempered then so is $BC(\pi)$ and conversely. However it might happen that π is supercuspidal

or square-integrable but $BC(\pi)$ is not. Aubert, Moussaoui and Solleveld [AMS] have recently proposed a very general conjecture on how to detect supercuspidal representations in L -packets. Moreover, Moussaoui [Mou] has been able to verify this conjecture for orthogonal and symplectic groups. Most probably his work will soon cover unitary groups too. We will need the following particular case of the Aubert-Moussaoui-Solleveld conjecture for which however we can give a direct proof.

Lemma 2.3.1 *Assume that F is p -adic. Let $\pi \in \text{Irr}(U(V))$ and assume that $BC(\pi)$ is supercuspidal. Then so is π .*

Proof: We will use the following characterization of supercuspidal representations

- (1) π is supercuspidal if and only if the Harish-Chandra character Θ_π of π is compactly supported modulo conjugation.

The necessity is an old result of Deligne ([De]). The sufficiency follows for example from Clozel's formula for the character ([Cl1, Proposition 1]).

Let φ be the Langlands parameter of π . Then by our assumption the L -packet $\Pi^{U(V)}(\varphi)$ is a singleton. Introduce the *twisted* group $\widetilde{GL}_n(E) = GL_n(E)\theta_n$ where $\theta_n(g) = {}^t\bar{g}^{-1}$. It is the set of F -points of the nonneutral connected component of the non-connected group $G^+ = R_{E/F}GL_n \rtimes \{1, \theta_n\}$. Since φ is a conjugate-dual representation of \mathcal{L}_E , it follows that $BC(\pi)$ may be extended to a representation $BC(\pi)^+$ of $G^+(F)$. Denote by $\widetilde{BC}(\pi)$ the restriction of $BC(\pi)^+$ to $\widetilde{GL}_n(E)$ and denote by $\Theta_{\widetilde{BC}(\pi)}$ the Harish-Chandra character of $\widetilde{BC}(\pi)$ (the Harish-Chandra theory of characters has been extended to twisted groups by Clozel [Cl2]). Since $BC(\pi)$ is supercuspidal, the character $\Theta_{\widetilde{BC}(\pi)}$ is compactly supported modulo conjugation (this follows for example from the equality up to a factor between $\Theta_{\widetilde{BC}(\pi)}$ and weighted orbital integrals of coefficients of $\widetilde{BC}(\pi)$ see [Wald2, théorème 7.1]). By the endoscopic characterization of the local Langlands correspondence for unitary groups, there is a relation between Θ_π and $\Theta_{\widetilde{BC}(\pi)}$. More precisely there is a correspondence between (stable) regular conjugacy classes in $U(V)(F)$ and $\widetilde{GL}_n(E)$ (see [Beu2, §3.2], in this particular case the correspondence takes the form of an injective map $U(V)_{reg}(F)/stab \hookrightarrow \widetilde{GL}_n(E)_{reg}/stab$) and for all regular elements $y \in U(V)(F)$, $\tilde{x} \in \widetilde{GL}_n(E)$ that correspond to each other we have (see [Mok, Theorem 3.2.1] and [KMSW, Theorem 1.6.1])

$$\Theta_{\widetilde{BC}(\pi)}(\tilde{x}) = \Delta(y, \tilde{x})\Theta_\pi(y)$$

where $\Delta(y, \tilde{x})$ is (up to a sign) a certain transfer factor. From this relation we easily infer that Θ_π is compactly supported modulo conjugation and hence that π is supercuspidal by (1). ■

We now move on to a global setting. Thus E/F is a quadratic extension of number fields and V is a n -dimensional hermitian space over E . If v is a place of F which splits in E then

we have isomorphisms $U(V)(F_v) \simeq GL_n(F_v)$ and $(R_{E/F}GL_n)(F_v) \simeq GL_n(F_v) \times GL_n(F_v)$ and we define a base change map $BC : Irr(U(V)_v) \rightarrow Irr((R_{E/F}GL_n)_v)$ by $\pi \mapsto \pi \boxtimes \pi^\vee$. By Theorem 2.5.2 of [Mok] and Theorem 1.7.1/Corollary 3.3.2 of [KMSW] we may associate to any cuspidal automorphic representation π of $U(V)$ an isobaric conjugate-dual automorphic representation $BC(\pi)$ of $GL_n(E)$, the base-change of π , satisfying the following properties:

(1) The Asai L -function

$$L(s, BC(\pi), As^{(-1)^{n+1}})$$

has a pole at $s = 1$ and moreover if $BC(\pi)$ is cuspidal this pole is simple (see [GGP, §7] for the definition of the Asai L -functions);

- (2) Let v be a place of F . Then, if $BC(\pi)$ is generic or v splits in E we have $BC(\pi_v) = BC(\pi)_v$;
- (3) If $BC(\pi)$ is generic then the multiplicity of π in $L^2([U(V)])$ is one (see Theorem 2.5.2/Remark 2.5.3 of [Mok] and Theorem 5.0.5, Theorem 1.7.1 and the discussion thereafter of [KMSW]).

Let v be a place of F and $\pi \in Irr(U(V)(F_v))$. Assume first that v is inert in E . By the Langlands classification there exist

- a parabolic subgroup $P = MN$ of $U(V)_v$ with

$$M \simeq R_{E_v/F_v}GL_{n_1} \times \dots \times R_{E_v/F_v}GL_{n_r} \times U(V')$$

where $V' \subset V_v$ is a nondegenerate subspace;

- tempered representations $\pi_i \in Temp(GL_{n_i}(E_v))$, $1 \leq i \leq r$, and $\pi' \in Temp(U(V'))$;
- real numbers $\lambda_1 > \dots > \lambda_r > 0$,

such that π is the unique irreducible quotient of

$$i_P^{U(V)_v} (|\det|_{E_v}^{\lambda_1} \pi_1 \boxtimes \dots \boxtimes |\det|_{E_v}^{\lambda_r} \pi_r \boxtimes \pi')$$

The r -uple $(\lambda_1, \dots, \lambda_r)$ only depends on π and we will set $\mathbf{c}(\pi) = \lambda_1$ if $r \geq 1$, $\mathbf{c}(\pi) = 0$ if $r = 0$ (i.e. if π is tempered). Assume now that v splits in E . Then, we have an isomorphism $U(V)_v \simeq GL_{n, F_v}$ and there exists a r -uple (n_1, \dots, n_r) of positive integers such that $n_1 + \dots + n_r = n$, tempered representations $\pi_i \in Temp(GL_{n_i}(F_v))$ $i = 1, \dots, r$ and real numbers $\lambda_1 > \lambda_2 > \dots > \lambda_r$ such that π is the unique irreducible quotient of

$$i_P^{GL_n} (|\det|_{F_v}^{\lambda_1} \pi_1 \boxtimes \dots \boxtimes |\det|_{F_v}^{\lambda_r} \pi_r)$$

where P denotes the standard parabolic subgroup of GL_n with Levi $GL_{n_1} \times \dots \times GL_{n_r}$. In this case, we set $\mathbf{c}(\pi) = \max(|\lambda_1|, |\lambda_r|)$. This depends only on π and in particular not on the choice of the isomorphism $U(V)_v \simeq GL_{n, F_v}$ (which is only defined up to an automorphism of GL_{n, F_v} since it involves the choice of a place of E above v).

In any case, for $c > 0$ we define $Irr_{\leq c}(U(V)_v)$ to be the set of irreducible representations $\pi \in Irr(U(V)_v)$ such that $\mathbf{c}(\pi) \leq c$. Combining the above global results of Mok and Kaletha-Minguez-Shin-White with the bounds toward the Ramanujan conjecture for GL_n of Luo-Rudnick-Sarnak [LRS] suitably extended to ramified places independently by Müller-Speh and Bergeron-Clozel ([MS], [BC]), we get the following:

Lemma 2.3.2 *Set $c = \frac{1}{2} - \frac{1}{n^2+1}$. Let π be a cuspidal automorphic representation of $U(V)(\mathbb{A})$ such that $BC(\pi)$ is generic. Then, for all place v of F we have*

$$\pi_v \in Irr_{\leq c}(U(V)_v)$$

2.4 The local Gan-Gross-Prasad conjecture for unitary groups

Let E/F be a quadratic extension of local fields of characteristic zero (either archimedean or p -adic). Let W be a n -dimensional hermitian space over E and define the hermitian space V by $V = W \oplus^\perp e$ where $(e, e) = 1$. Set $H = U(W)$ and $G = U(W) \times U(V)$. We view H as a subgroup of G via the diagonal embedding. We will say that an irreducible representation π of $G(F)$ is H -distinguished if the space $Hom_H(\pi, \mathbb{C})$ of $H(F)$ -invariant (continuous in the archimedean case) linear forms on π is nonzero. By multiplicity one results (see [AGRS], [JSZ]) we always have $\dim Hom_H(\pi, \mathbb{C}) \leq 1$. We will denote by $Irr_H(G)$ and $Temp_H(G)$ the subsets of H -distinguished representations in $Irr(G)$ and $Temp(G)$ respectively. Let φ be a generic Langlands parameter for G . We have the following conjecture of Gan, Gross and Prasad ([GGP, conjecture 17.1])

Conjecture 2.4.1 *The L -packet $\Pi^G(\varphi)$ contains at most one H -distinguished representation.*

By [Beu, Theorem 12.4.1] and [GI, Proposition 9.3], the following cases of this conjecture are known.

Theorem 2.4.2 (Beuzart-Plessis, Gan-Ichino) .

- (i) *Let φ be a tempered Langlands parameter for G . Then conjecture 2.4.1 holds for φ .*
- (ii) *Assume that F is p -adic. Then conjecture 2.4.1 holds for any generic Langlands parameter φ of G .*

2.5 Measures

We will use the same normalization of measures as in [Zh2, §2]. Let us recall these choices. We actually define two sets of Haar measures: the *normalized* and the *unnormalized*. We will use the normalized Haar measures apart in section 4 where we will use the unnormalized one. From now on and until section 4, where we will switch to a local setting, we fix a quadratic extension E/F of number fields. We will denote by $\eta_{E/F}$ the idele class character corresponding to this extension. We will also fix a nonzero character $\psi : \mathbb{A}/F \rightarrow \mathbb{C}^\times$ and a nonzero element $\tau \in E$ such that $\text{Tr}_{E/F}(\tau) = 0$. We will denote by ψ_E the character of \mathbb{A}_E given by $\psi_E(z) = \psi(\frac{1}{2}\text{Tr}_{E/F}(z))$.

Let v be a place of F . We endow F_v with the self-dual Haar measure for ψ_v . Similarly, we endow E_v with the self-dual Haar measure for $\psi_{E,v}$. On F_v^\times , we define a normalized measure

$$d^\times x_v = \zeta_{F_v}(1) \frac{dx_v}{|x_v|_{F_v}}$$

and an unnormalized one

$$d^* x_v = \frac{dx_v}{|x_v|_{F_v}}$$

More generally, for all $n \geq 1$, we equip $GL_n(F_v)$ with the following normalized Haar measure

$$dg_v = \zeta_{F_v}(1) \frac{\prod_{ij} dg_{v,ij}}{|\det g_v|_{F_v}^n}$$

as well as with the following unnormalized one

$$d^* g_v = \frac{\prod_{ij} dg_{v,ij}}{|\det g_v|_{F_v}^n}$$

and similarly for $GL_n(E_v)$. Recall that N_n denotes the standard maximal unipotent subgroup of GL_n . We will give $N_n(F_v)$ and $N_n(E_v)$ the Haar measures

$$du_v = \prod_{1 \leq i < j \leq n} du_{v,ij}$$

We equip \mathbb{A}^\times , $N_n(\mathbb{A})$, $N_n(\mathbb{A}_E)$, $GL_n(\mathbb{A})$ and $GL_n(\mathbb{A}_E)$ with the global Tamagawa Haar measures given by

$$d^\times x = \prod_v d^\times x_v, \quad du = \prod_v du_v, \quad dg = \prod_v dg_v$$

Recall that $S_n = \{s \in R_{E/F}GL_n; s\bar{s} = 1\}$ and its *Lie algebra* $\mathfrak{s}_n = \{X \in R_{E/F}M_n; X + \bar{X} = 0\}$. Let V be a n -dimensional hermitian space over E and denote by $\mathfrak{u}(V)$ the Lie algebra of $U(V)$. Choosing a basis of V we get an embedding $\mathfrak{u}(V) \hookrightarrow R_{E/F}M_n$. Let us denote by $\langle \cdot, \cdot \rangle$ the $GL_n(E_v)$ -invariant bilinear pairing on $M_n(E_v)$ given by

$$\langle X, Y \rangle := \text{Trace}(XY)$$

Note that the restrictions of $\langle \cdot, \cdot \rangle$ to $\mathfrak{sl}_n(F_v)$ and $\mathfrak{u}(V)(F_v)$ are F_v -valued and nondegenerate. We define a Haar measure dX on $\mathfrak{u}(V)(F_v)$ such that the Fourier transform

$$\widehat{\varphi}(Y) = \int_{\mathfrak{u}(V)(F_v)} \varphi(X) \psi_v(\langle X, Y \rangle) dX$$

and its dual

$$\check{\varphi}(X) = \int_{\mathfrak{u}(V)(F_v)} \varphi(Y) \psi_v(-\langle Y, X \rangle) dY$$

are inverse of each other. We define similarly a Haar measure and Fourier transforms $\varphi \mapsto \widehat{\varphi}$, $\varphi \mapsto \check{\varphi}$ on $\mathfrak{sl}_n(F_v)$.

The Cayley map $\mathfrak{c} : X \mapsto \mathfrak{c}(X) = (1 + X)(1 - X)^{-1}$ induces birational isomorphisms from \mathfrak{sl}_n to S_n and from $\mathfrak{u}(V)$ to $U(V)$. We define the *unnormalized* Haar measure d^*g_v on $U(V)(F_v)$ to be the unique Haar measure such that the Jacobian of \mathfrak{c} at the origin is 1. The normalized Haar measure on $U(V)(F_v)$ is defined by

$$dg_v = L(1, \eta_{E_v/F_v}) d^*g_v$$

Similarly, we endow $S_n(F_v)$ with an unnormalized measure d^*s_v which is the unique $GL_n(E_v)$ -invariant measure for which the Jacobian of the Cayley map \mathfrak{c} at the origin is 1. The corresponding normalized measure is given by

$$ds_v = L(1, \eta_{E_v/F_v}) d^*s_v$$

Note that d^*s_v (resp. ds_v) can also be identified with the quotient of the unnormalized (resp. normalized) Haar measures on $GL_n(E_v)$ and $GL_n(F_v)$ via the isomorphism $\nu : GL_n(E_v)/GL_n(F_v) \simeq S_n(F_v)$, $\nu(g) = g\bar{g}^{-1}$. Finally, we equip $U(V)(\mathbb{A})$ with the global Haar measure given by

$$dg = \prod_v dg_v$$

It is not the Tamagawa measure since there is a factor $L(1, \eta_{E/F})^{-1}$ missing. Note that the local normalized Haar measure dg_v can be identified with the quotient of the normalized Haar measures on E_v^\times and F_v^\times via the isomorphism $E_v^\times/F_v^\times \simeq U(1)(F_v)$, $x \mapsto x/\bar{x}$. Hence, as the Tamagawa number of $U(1)$ is 2, we have

$$(1) \quad \text{vol}(E^\times \mathbb{A}^\times \backslash \mathbb{A}_E^\times) = \text{vol}([U(1)]) = 2L(1, \eta_{E/F})$$

3 Spherical characters, the Ichino-Ikeda conjecture and Zhang's conjecture

In this section E/F will be a quadratic extension of number fields and we will use normalized Haar measures (see §2.5). Let W be an hermitian space of dimension n over E . We will set $V = W \oplus^\perp Ee$ where $(e, e) = 1$, $G = U(W) \times U(V)$ and $H = U(W)$. We view H as a subgroup of G via the diagonal embedding. We will fix a maximal compact subgroup $K = \prod_v K_v$ of $G(\mathbb{A})$. We will say that an irreducible representation $\pi = \otimes'_v \pi_v$ of $G(\mathbb{A})$ is *abstractly H -distinguished* if for all place v of F the representation π_v is H_v -distinguished i.e. if $\text{Hom}_{H_v}(\pi_v, \mathbb{C}) \neq 0$. Set $G' = R_{E/F}(GL_n \times GL_{n+1})$. We define two subgroups $H'_1 = R_{E/F}GL_n$ and $H'_2 = GL_n \times GL_{n+1}$ of G' (H'_1 is embedded diagonally). We also define a character η of $H'_2(\mathbb{A})$ by

$$\eta(g_1, g_2) = \eta_{E/F}(g_1)^{n+1} \eta_{E/F}(g_2)^n$$

for all $(g_1, g_2) \in H'_2(\mathbb{A}) = GL_n(\mathbb{A}) \times GL_{n+1}(\mathbb{A})$. We will also fix a maximal compact subgroup $K' = \prod_v K'_v$ of $G'(\mathbb{A})$ such that $K'_v = GL_n(\mathcal{O}_{E,v}) \times GL_{n+1}(\mathcal{O}_{E,v})$ for all nonarchimedean place v of F . Finally, if π and Π are cuspidal automorphic representations of $G(\mathbb{A})$ and $G'(\mathbb{A})$ respectively then we endow them with the following Petersson inner products

$$(\phi_1, \phi_2)_{Pet} := \int_{[G]} \phi_1(g) \overline{\phi_2(g)} dg, \quad \phi_1, \phi_2 \in \pi$$

$$(\phi'_1, \phi'_2)_{Pet} := \int_{[Z_{G'} \backslash G']} \phi'_1(g') \overline{\phi'_2(g')} dg', \quad \phi'_1, \phi'_2 \in \Pi$$

3.1 Global spherical characters

For any cuspidal automorphic representation π of $G(\mathbb{A})$ we define the H -period $\mathcal{P}_H : \pi \rightarrow \mathbb{C}$ by

$$\mathcal{P}_H(\phi) = \int_{[H]} \phi(h) dh, \quad \phi \in \pi$$

The integral is absolutely convergent (see Proposition A.1.1(ix)). We will say that the cuspidal automorphic representation π is *globally H -distinguished* if the period \mathcal{P}_H is not identically zero on π . We may associate to this period a (global) spherical character $J_\pi : \mathcal{S}(G(\mathbb{A})) \rightarrow \mathbb{C}$ defined as follows. Let $f \in \mathcal{S}(G(\mathbb{A}))$ and choose a compact-open subgroup $K_f \subset G(\mathbb{A}_f)$ by which f is biinvariant. Let $\mathcal{B}_\pi^{K_f}$ be an orthonormal basis for the Petersson inner product of π^{K_f} whose elements are C_G and C_K eigenvectors. Then we set

$$J_\pi(f) = \sum_{\phi \in \mathcal{B}_\pi^{K_f}} \mathcal{P}_H(\pi(f)\phi) \overline{\mathcal{P}_H(\phi)}$$

The sum is absolutely convergent and does not depend on the choice of the basis $\mathcal{B}_\pi^{K_f}$ (see Proposition A.1.2).

Let Π be a cuspidal automorphic representation of $G'(\mathbb{A})$ whose central character is trivial on $Z_{H'_2}(\mathbb{A}) = \mathbb{A}^\times \times \mathbb{A}^\times$. We define two periods $\lambda : \Pi \rightarrow \mathbb{C}$ and $\beta : \Pi \rightarrow \mathbb{C}$ by

$$\lambda(\phi) = \int_{[H'_1]} \phi(h_1) dh_1$$

$$\beta(\phi) = \int_{[Z_{H'_2} \backslash H'_2]} \phi(h_2) \eta(h_2) dh_2$$

for all $\phi \in \Pi$. The two above integrals are absolutely convergent (see Proposition A.1.1(ix)). We also define a (global) spherical character $I_\Pi : \mathcal{S}(G'(\mathbb{A})) \rightarrow \mathbb{C}$ as follows. Let $f' \in \mathcal{S}(G'(\mathbb{A}))$ and choose a compact-open subgroup $K_{f'} \subset G(\mathbb{A}_f)$ by which f' is biinvariant. Let $\mathcal{B}_\Pi^{K_{f'}}$ be an orthonormal basis for the Petersson inner product of $\Pi^{K_{f'}}$ whose elements are $C_{G'}$ and $C_{K'}$ eigenvectors. Then we set

$$I_\Pi(f') = \sum_{\phi \in \mathcal{B}_\Pi^{K_{f'}}} \lambda(\Pi(f')\phi) \overline{\beta(\phi)}$$

The sum is absolutely convergent and does not depend on the choice of the basis $\mathcal{B}_\Pi^{K_{f'}}$ (see Proposition A.1.2).

3.2 Local spherical characters

Let v be a place of F . Let π_v be a tempered representation of $G(F_v)$. We define a distribution $J_{\pi_v} : \mathcal{S}(G(F_v)) \rightarrow \mathbb{C}$ (the local spherical character associated to π_v) by

$$J_{\pi_v}(f_v) = \int_{H(F_v)} \text{Trace}(\pi_v(h)\pi_v(f_v)) dh_v, \quad f_v \in \mathcal{C}(G(F_v))$$

By [Beu, §8.2], the above integral is absolutely convergent. Choosing models for G and H over \mathcal{O}_F , for almost all v if $f_v = \mathbf{1}_{K_v}$, we have

$$J_{\pi_v}(f_v) = \mathcal{L}\left(\frac{1}{2}, \pi_v\right) \text{vol}(H(\mathcal{O}_v)) \text{vol}(G(\mathcal{O}_v))$$

(see the introduction for the definition of $\mathcal{L}(s, \pi_v)$). Hence, we define a *normalized* spherical character $J_{\pi_v}^\sharp$ by

$$J_{\pi_v}^\sharp = \frac{1}{\mathcal{L}\left(\frac{1}{2}, \pi_v\right)} J_{\pi_v}$$

By [Beu, Theorem 8.2.1] we have

- (1) π_v is H_v -distinguished if and only if $J_{\pi_v} \neq 0$.

Moreover, by [Beu, Corollary 8.6.1], for all parabolic subgroup $P = MU$ of G_v and for all square-integrable representation σ of M there is at most one irreducible subrepresentation $\pi \subset i_P^G(\sigma)$ such that $J_\pi \neq 0$. Thus, we are in the situation of §2.2 and the family of distributions $\pi_v \in \text{Temp}(G_v) \mapsto J_{\pi_v}$ is analytic.

Let Π_v be a generic unitary representation of $G'(F_v)$. We may write $\Pi_v = \Pi_{n,v} \boxtimes \Pi_{n+1,v}$ where $\Pi_{n,v}$ and $\Pi_{n+1,v}$ are generic and unitary representations of $GL_n(E_v)$ and $GL_{n+1}(E_v)$ respectively. Let $\mathcal{W}(\Pi_{n,v}, \overline{\psi}_E)$ and $\mathcal{W}(\Pi_{n+1,v}, \psi_E)$ be the Whittaker models of $\Pi_{n,v}$ and $\Pi_{n+1,v}$ corresponding to the characters $\overline{\psi}_E$ and ψ_E respectively. Set $\mathcal{W}(\Pi_v) = \mathcal{W}(\Pi_{n,v}, \overline{\psi}_E) \otimes \mathcal{W}(\Pi_{n+1,v}, \psi_E)$. We define linear forms (the local Flicker-Rallis periods)

$$\beta_{n,v} : \mathcal{W}(\Pi_{n,v}, \overline{\psi}_E) \rightarrow \mathbb{C}, \quad \beta_{n+1,v} : \mathcal{W}(\Pi_{n+1,v}, \psi_E) \rightarrow \mathbb{C}$$

and scalar products

$$\theta_{n,v} : \mathcal{W}(\Pi_{n,v}, \overline{\psi}_E) \times \mathcal{W}(\Pi_{n,v}, \overline{\psi}_E) \rightarrow \mathbb{C}, \quad \theta_{n+1,v} : \mathcal{W}(\Pi_{n+1,v}, \psi_E) \times \mathcal{W}(\Pi_{n+1,v}, \psi_E) \rightarrow \mathbb{C}$$

by

$$\beta_{k,v}(W_k) = \int_{N_{k-1}(F_v) \backslash GL_{k-1}(F_v)} W_k(\epsilon_k(\tau)g_{k-1}) \eta_{E_v/F_v}(\det g_{k-1})^{k-1} dg_{k-1}$$

$$\theta_{k,v}(W_k, W'_k) = \int_{N_{k-1}(E_v) \backslash GL_{k-1}(E_v)} W_k(g_{k-1}) \overline{W'_k(g_{k-1})} dg_{k-1}$$

for all $k = n, n+1$, all $W_n, W'_n \in \mathcal{W}(\Pi_{n,v}, \overline{\psi}_E)$ and all $W_{n+1}, W'_{n+1} \in \mathcal{W}(\Pi_{n+1,v}, \psi_E)$ where $\epsilon_k(\tau) = \text{diag}(\tau^{k-1}, \tau^{k-2}, \dots, \tau, 1)$ (recall that τ is a fixed nonzero element of E such that $\text{Tr}_{E/F}(\tau) = 0$). The above integrals are absolutely convergent (see [JS] Propositions 1.3 and 3.16 for the absolute convergence of $\theta_{k,v}$, the proof for $\beta_{k,v}$ is identical). Set $\beta_v = \beta_{n,v} \otimes \beta_{n+1,v}$ and $\theta_v = \theta_{n,v} \otimes \theta_{n+1,v}$. If E_v/F_v , Π_v , $\psi_{E,v}$ are unramified, τ is a unit in E_v and $W_v \in \mathcal{W}(\Pi_v)$ is the unique K'_v -invariant vector such that $W_v(1) = 1$, we have (see [JS, Proposition 2.3] and [Zh2, §3.2])

$$\beta_v(W_v) = \text{vol}(K'_v) L(1, \Pi_{n,v}, As^{(-1)^{n-1}}) L(1, \Pi_{n+1,v}, As^{(-1)^n})$$

and

$$\theta_v(W_v) = \text{vol}(K'_v) L(1, \Pi_{n,v} \times \Pi_{n,v}^\vee) L(1, \Pi_{n+1,v} \times \Pi_{n+1,v}^\vee)$$

Hence, we define normalized versions β_v^\natural and θ_v^\natural of β_v and θ_v by

$$\beta_v^\natural = \frac{\beta_v}{L(1, \Pi_{n,v}, As^{(-1)^{n-1}}) L(1, \Pi_{n+1,v}, As^{(-1)^n})}, \quad \theta_v^\natural = \frac{\theta_v}{L(1, \Pi_{n,v} \times \Pi_{n,v}^\vee) L(1, \Pi_{n+1,v} \times \Pi_{n+1,v}^\vee)}$$

For all $s \in \mathbb{C}$, we also have the local Rankin-Selberg period $\lambda_v(s, \cdot) : \mathcal{W}(\Pi_v) \rightarrow \mathbb{C}$ defined by

$$\lambda_v(s, W_n \otimes W_{n+1}) = \int_{N_n(E_v) \backslash GL_n(E_v)} W_n(g_n) W_{n+1}(g_n) |\det g_n|_{E_v}^s dg_n$$

for all $(W_n, W_{n+1}) \in \mathcal{W}(\Pi_{n,v}, \overline{\psi_E}) \times \mathcal{W}(\Pi_{n+1,v}, \psi_E)$, and its normalization $\lambda_v^\natural(s, \cdot)$ given by

$$\lambda_v^\natural(s, \cdot) = \frac{\lambda_v(s, \cdot)}{L(s + \frac{1}{2}, \Pi_{n,v} \times \Pi_{n+1,v})}$$

The integral defining $\lambda_v(s, \cdot)$ is absolutely convergent for $Re(s) \gg 0$ and $\lambda_v^\natural(s, \cdot)$ extends to an entire function on \mathbb{C} (see [JPSS] and [Jac] for the archimedean case). We will set $\lambda_v^\natural = \lambda_v^\natural(0, \cdot)$. Obviously λ_v^\natural defines a $H_1'(F_v)$ -invariant linear form on Π_v . Moreover by [JPSS] and [Jac], there exists $W \in \mathcal{W}(\Pi_v)$ such that $\lambda_v^\natural(W) = 1$. Hence λ_v^\natural defines a nonzero element in $Hom_{H_1'}(\Pi_v, \mathbb{C})$. If Π_v is tempered then $\lambda_v(s, \cdot)$ is absolutely convergent for $Re(s) > -1/2$ and we will set $\lambda_v = \lambda_v(0, \cdot)$.

We are now ready to define the (normalized) local spherical character $I_{\Pi_v}^\natural : \mathcal{S}(G'(F_v)) \rightarrow \mathbb{C}$ attached to Π_v . Let $f'_v \in \mathcal{S}(G'(F_v))$. If v is nonarchimedean then choose a compact-open subgroup $K_{f'_v}$ of $G'(F_v)$ by which f'_v is biinvariant and let \mathcal{B}_{Π_v} be an orthonormal basis of $\Pi_v^{K_{f'_v}}$ for the scalar product θ_v^\natural . If v is archimedean, we let \mathcal{B}_{Π_v} be any orthonormal basis of Π_v for the scalar product θ_v^\natural consisting of $C_{K'_v}$ -eigenvectors. Then we set

$$I_{\Pi_v}^\natural(f'_v) = \sum_{W \in \mathcal{B}_{\Pi_v}} \lambda_v^\natural(\Pi_v(f'_v)W) \overline{\beta_v^\natural(W)}$$

The sum is absolutely convergent and does not depend on the choice of \mathcal{B}_{Π_v} . If moreover Π_v is tempered then we define an *unnormalized* local spherical character $I_{\Pi_v} : \mathcal{S}(G'(F_v)) \rightarrow \mathbb{C}$ by using θ_v , β_v and λ_v instead of θ_v^\natural , β_v^\natural and λ_v^\natural . Finally, the proofs of [JS, Proposition 1.3] and [JPSS, Theorem 2.7] easily show that the family of distributions $\Pi_v \in Temp(G'_v) \mapsto I_{\Pi_v}$ is analytic in the sense of §2.2.

3.3 Orbital integrals

Consider the action of $H \times H$ on G by left and right translations. Then, an element $\delta \in G$ is said to be *regular semisimple* for this action if its orbit is closed and its stabilizer is trivial. Denote by G_{rs} the open subset of regular semisimple element in G . Let v be a place of F and $\delta \in G_{rs}(F_v)$ be regular semisimple. We define the (relative) orbital integral associated to δ as the distribution given by

$$O(\delta, f_v) = \int_{H(F_v) \times H(F_v)} f_v(h\delta h') dh dh', \quad f_v \in \mathcal{S}(G(F_v))$$

There is another way to see these orbital integrals. For all $f_v \in \mathcal{S}(G(F_v))$, we define a function $\tilde{f}_v \in \mathcal{S}(U(V)(F_v))$ by

$$\tilde{f}_v(x) = \int_{H(F_v)} f_v(h(1, x)) dh, \quad x \in U(V)(F_v)$$

This defines a surjective linear map $\mathcal{S}(G(F_v)) \rightarrow \mathcal{S}(U(V)(F_v))$. Let us say that an element $x \in U(V)$ is regular semisimple if it is so for the action of $U(W)$ by conjugation i.e. if the $U(W)$ -conjugacy class of x is closed and the stabilizer of x in $U(W)$ is trivial. Denote by $U(V)_{rs}$ the open subset of regular semisimple element in $U(V)$. For all $x \in U(V)_{rs}(F_v)$ we define the orbital integral associated to x as the distribution

$$O(x, \varphi) = \int_{U(W)(F_v)} \varphi(hxh^{-1})dh, \quad \varphi \in \mathcal{S}(U(V)(F_v))$$

For all $\delta = (\delta_W, \delta_V) \in G_{rs}$ the element $x = \delta_W^{-1}\delta_V$ is regular semisimple in $U(V)$ and this defines a surjection $G_{rs} \rightarrow U(V)_{rs}$. Moreover, for all $\delta \in G_{rs}(F_v)$ and all $f \in \mathcal{S}(G(F_v))$, we have the equality

$$O(\delta, f) = O(x, \tilde{f})$$

where $x = \delta_W^{-1}\delta_V$.

We can also define orbital integrals on the space $\mathcal{S}(\mathfrak{u}(V)(F_v))$. Call an element $X \in \mathfrak{u}(V)$ regular semisimple if it is so for the adjoint action of $U(W)$. Let us denote by $\mathfrak{u}(V)_{rs}$ the open subset of regular semisimple elements. Then, for all $X \in \mathfrak{u}(V)_{rs}(F_v)$ we can define an orbital integral by

$$O(X, \varphi) = \int_{U(W)(F_v)} \varphi(h^{-1}Xh)dh, \quad \varphi \in \mathcal{S}(U(V)(F_v))$$

The Cayley map $\mathfrak{c} : X \mapsto (1+X)(1-X)^{-1}$ realizes a $U(W)$ -equivariant isomorphism between the open subsets $\mathfrak{u}(V)^\circ = \{X \in \mathfrak{u}(V); \det(1-X) \neq 0\}$ and $U(V)^\circ = \{x \in U(V); \det(1+x) \neq 0\}$. Assume that v is nonarchimedean and let $\omega \subset \mathfrak{u}(V)^\circ(F_v)$ and $\Omega \subset U(V)^\circ(F_v)$ be open and closed $U(W)(F_v)$ -invariant neighborhoods of 0 and 1 respectively such that the Cayley map restricts to an analytic isomorphism between ω and Ω preserving measures. For all $\varphi \in \mathcal{S}(U(V)(F_v))$, we define a function $\varphi_{\mathfrak{h}}$ by

$$\varphi_{\mathfrak{h}}(X) = \begin{cases} \varphi(\mathfrak{c}(X)) & \text{if } X \in \omega \\ 0 & \text{otherwise} \end{cases}$$

Then for all $X \in \omega_{rs} = \omega \cap \mathfrak{u}(V)_{rs}(F_v)$ and all $\varphi \in \mathcal{S}(U(V)(F_v))$ we have

$$O(\mathfrak{c}(X), \varphi) = O(X, \varphi_{\mathfrak{h}})$$

Consider now the action of $H'_1 \times H'_2$ on G' by left and right translations. As before, an element $\gamma \in G'$ is said to be *regular semisimple* for this action if its orbit is closed and its stabilizer is trivial. Denote by G'_{rs} the open subset of regular semisimple element in G' . Let v be a place of F and $\gamma \in G'_{rs}(F_v)$ be regular semisimple. We define the (relative) orbital integral associated to γ as the distribution given by

$$O(\gamma, f'_v) = \int_{H'_1(F_v) \times H'_2(F_v)} f'_v(h_1^{-1}\gamma h_2)\eta(h_2)dh_1dh_2, \quad f'_v \in \mathcal{S}(G'(F_v))$$

There is another way to see these orbital integrals. Recall that $S_{n+1}(F_v) = \{s \in GL_{n+1}(E_v); s\bar{s} = 1\}$ and that we have a surjective map $\nu : GL_{n+1}(E_v) \rightarrow S_{n+1}(F_v)$, $\nu(g) = g\bar{g}^{-1}$. For all $f'_v \in \mathcal{S}(G'(F_v))$, we define a function $\tilde{f}'_v \in \mathcal{S}(S_{n+1}(F_v))$ by

$$\tilde{f}'_v(s) = \int_{H'_1(F_v)} \int_{GL_{n+1}(F_v)} f'_v(h_1(1, gh_2)) dh_2 dh_1, \quad g \in GL_{n+1}(E_v), s = \nu(g)$$

if n is even and

$$\tilde{f}'_v(s) = \int_{H'_1(F_v)} \int_{GL_{n+1}(F_v)} f'_v(h_1(1, gh_2)) \eta'_v(gh_2) dh_2 dh_1, \quad g \in GL_{n+1}(E_v), s = \nu(g)$$

if n is odd. In any case, this defines a surjective linear map $\mathcal{S}(G'(F_v)) \rightarrow \mathcal{S}(S_{n+1}(F_v))$. The group GL_n acts on S_{n+1} by conjugation and we shall say that an element $s \in S_{n+1}$ is regular semisimple if it is so for this action i.e. if the GL_n -conjugacy class of s is closed and the stabilizer of s in GL_n is trivial. We will denote by $S_{n+1,rs}$ the open subset of regular semisimple elements in S_{n+1} . For all $s \in S_{n+1,rs}(F_v)$ we define the orbital integral associated to s as the distribution

$$O(s, \varphi') = \int_{GL_n(F_v)} \varphi'(h^{-1}sh) \eta_{E_v/F_v}(h) dh, \quad \varphi' \in \mathcal{S}(S_{n+1}(F_v))$$

For $\gamma = (\gamma_1, \gamma_2) \in G'_{rs}$ the element $s = \nu(\gamma_1^{-1}\gamma_2) \in S_{n+1}$ is regular semisimple and this defines a surjection $G'_{rs} \rightarrow S_{n+1,rs}$. Moreover, for all $\gamma \in G'_{rs}(F_v)$ and all $f' \in \mathcal{S}(G'(F_v))$, we have the equality

$$O(\gamma, f') = \begin{cases} O(s, \tilde{f}') & \text{if } n \text{ is even,} \\ \eta'_v(\gamma_1^{-1}\gamma_2) O(s, \tilde{f}') & \text{if } n \text{ is odd.} \end{cases}$$

where $s = \nu(\gamma_1^{-1}\gamma_2)$.

We can also define orbital integrals on the space $\mathcal{S}(\mathfrak{s}_{n+1}(F_v))$. Call an element $X \in \mathfrak{s}_{n+1}$ regular semisimple if it is so for the adjoint action of GL_n . Let us denote by $\mathfrak{s}_{n+1,rs}$ the open subset of regular semisimple elements. Then, for all $X \in \mathfrak{s}_{n+1,rs}(F_v)$ we can define an orbital integral by

$$O(X, \varphi') = \int_{GL_n(F_v)} \varphi'(h^{-1}Xh) \eta_{E_v/F_v}(h) dh, \quad \varphi' \in \mathcal{S}(\mathfrak{s}_{n+1}(F_v))$$

The Cayley map $\mathbf{c} = \mathbf{c}_{n+1} : X \mapsto (1+X)(1-X)^{-1}$ realizes a GL_n -equivariant isomorphism between the open subsets $\mathfrak{s}_{n+1}^\circ = \{X \in \mathfrak{s}_{n+1}; \det(1-X) \neq 0\}$ and $S_{n+1}^\circ = \{s \in S_{n+1}; \det(1+s) \neq 0\}$. Let $\omega' \subset \mathfrak{s}_{n+1}^\circ(F_v)$ and $\Omega' \subset S_{n+1}^\circ(F_v)$ be open and closed $GL_n(F_v)$ -invariant neighborhoods of 0 and 1 respectively such that the Cayley map restricts to an analytic isomorphism between ω' and Ω' preserving measures. For all $\varphi' \in \mathcal{S}(S_{n+1}(F_v))$, we define a function $\varphi'_\natural \in \mathcal{S}(\mathfrak{s}_{n+1}(F_v))$ by

$$\varphi'_\mathfrak{h}(X) = \begin{cases} \varphi'(\mathfrak{c}(X)) & \text{if } X \in \omega' \\ 0 & \text{otherwise} \end{cases}$$

Then for all $X \in \omega'_{rs} = \omega' \cap \mathfrak{s}_{n+1,rs}(F_v)$ and all $\varphi' \in \mathcal{S}(S_{n+1}(F_v))$ we have

$$O(\mathfrak{c}(X), \varphi') = O(X, \varphi'_\mathfrak{h})$$

3.4 Correspondence of orbits and transfer

We now recall the correspondence between orbits following [Zh1, §2.4]. We will denote by $H'_1 \backslash G' / H'_2$, $H \backslash G / H$, S_{n+1} / GL_n and $U(V) / U(W)$ the geometric quotients of G' , G , S_{n+1} and $U(V)$ by $H'_1 \times H'_2$, $H \times H$, GL_n and $U(W)$ respectively (the last two actions being given by conjugation). We will also write $(H'_1 \backslash G' / H'_2)_{rs}$, $(H \backslash G / H)_{rs}$, $(S_{n+1} / GL_n)_{rs}$ and $(U(V) / U(W))_{rs}$ for the regular semisimple loci in these geometric quotients. These are the image of G'_{rs} , G_{rs} , $S_{n+1,rs}$ and $U(V)_{rs}$ by the natural projections. The maps $(\gamma_1, \gamma_2) \in G' \mapsto \nu(\gamma_1^{-1} \gamma_2)$ and $(\delta_W, \delta_V) \in G \mapsto \delta_W^{-1} \delta_V$ induce isomorphisms

$$H'_1 \backslash G' / H'_2 \simeq S_{n+1} / GL_n \text{ and } H \backslash G / H \simeq U(V) / U(W)$$

and similarly for the regular semisimple loci. Moreover, there is a natural isomorphism (see [Zh1, §3.1])

$$(1) \quad H'_1 \backslash G' / H'_2 \simeq H \backslash G / H$$

which preserves the regular semisimple loci. For all field extension k of F we have $(H'_1 \backslash G' / H'_2)_{rs}(k) = H'_1(k) \backslash G'_{rs}(k) / H'_2(k)$ and $H(k) \backslash G_{rs}(k) / H(k)$ is a subset of $(H \backslash G / H)_{rs}(k)$. The above isomorphism thus induces injections

$$(2) \quad H(k) \backslash G_{rs}(k) / H(k) \hookrightarrow H'_1(k) \backslash G'_{rs}(k) / H'_2(k)$$

and

$$(3) \quad U(V)_{rs}(k) / U(W)(k) \hookrightarrow S_{n+1,rs}(k) / GL_n(k)$$

This last map admits the following explicit description. Choosing a basis of V whose last element is e we get an embedding $U(V)(k) \hookrightarrow GL_{n+1}(k \otimes_F E)$. By [Zh3, lemma 2.3] any regular semisimple element $x \in U(V)_{rs}(k)$ is $GL_n(k \otimes_F E)$ -conjugated to a regular semisimple element of $S_{n+1}(k)$ which is unique up to $GL_n(k)$ -conjugation. The $GL_n(k)$ -conjugacy class of this element is exactly the image of x by the map (3).

We have a similar situation at the level of Lie algebras: we have a canonical isomorphism between geometric quotients

$$(4) \quad \mathfrak{s}_{n+1}/GL_n \simeq \mathfrak{u}(V)/U(W)$$

which preserves the regular semisimple loci $(\mathfrak{s}_{n+1}/GL_n)_{rs} = \mathfrak{s}_{n+1,rs}/GL_n$ and $(\mathfrak{u}(V)/U(W))_{rs} = \mathfrak{u}(V)_{rs}/U(W)$. For all field extension k of F this induces an injection

$$(5) \quad \mathfrak{u}(V)_{rs}(k)/U(W)(k) \hookrightarrow \mathfrak{s}_{n+1,rs}(k)/GL_n(k)$$

We now define, following [Zh2, §4.1], two families of *transfer factors* $\Omega_v : G'_{rs}(F_v) \rightarrow \mathbb{C}^\times$ and $\omega_v : \mathfrak{s}_{n+1,rs}(F_v) \rightarrow \mathbb{C}^\times$, v a place of F , satisfying the following conditions:

- For all v and all $\gamma \in G'_{rs}(F_v)$ (resp. all $X \in \mathfrak{s}_{n+1,rs}(F_v)$) we have $\Omega_v(h_1\gamma h_2) = \eta_v(h_2)\Omega_v(\gamma)$ (resp. $\omega_v(h^{-1}Xh) = \eta_{E_v/F_v}(h)\omega_v(X)$) for all $(h_1, h_2) \in H'_1(F_v) \times H'_2(F_v)$ (resp. for all $h \in GL_n(F_v)$);
- For all $\gamma \in G'_{rs}(F)$ (resp. all $X \in \mathfrak{s}_{n+1,rs}(F)$), we have the product formula $\prod_v \Omega_v(\gamma) = 1$ (resp. $\prod_v \omega_v(X) = 1$) where almost all terms in the product are equal to 1.

Let v be a place of F . For all $s \in S_{n+1,rs}(F_v)$ and all $X \in \mathfrak{s}_{n+1,rs}(F_v)$, we set

$$(6) \quad \Omega_v(s) = \eta'_v \left(\det(s)^{-[\frac{n+1}{2}]} \det(e_{n+1}, e_{n+1}s, \dots, e_{n+1}s^n) \right)$$

$$(7) \quad \omega_v(X) = \eta'_v (\det(e_{n+1}, e_{n+1}X, \dots, e_{n+1}X^n))$$

where $e_{n+1} = (0, \dots, 0, 1)$ and η'_v is the local component at v of the character $\eta' : \mathbb{A}_E^\times \rightarrow \mathbb{C}^\times$ extending $\eta_{E/F}$ that we fixed at the beginning. Note that (see the proof of [Zh1, lemma 3.5])

$$(8) \quad \Omega_v(\mathfrak{c}(X)) = \eta_v(2)^{n(n+1)/2} \omega_v(X)$$

for all $X \in \mathfrak{s}_{n+1,rs}(F_v)$ sufficiently close to 0. Finally for all $\gamma = (\gamma_1, \gamma_2) \in G'_{rs}(F_v)$, we set

$$\Omega_v(\gamma) = \begin{cases} \Omega_v(s) & \text{if } n \text{ is even,} \\ \eta'_v(\gamma_1^{-1}\gamma_2)\Omega_v(s) & \text{if } n \text{ is odd.} \end{cases}$$

where $s = \nu(\gamma_1^{-1}\gamma_2)$. For future reference, we record the following formula

$$(9) \quad \Omega_v(\gamma)O(\gamma, f') = \Omega_v(s)O(s, \tilde{f}')$$

for all $f' \in \mathcal{S}(G'(F_v))$, all $\gamma \in G'_{rs}(F_v)$ and where we have set $s = \nu(\gamma_1^{-1}\gamma_2)$.

Using the transfer factors we can define the notion of matching functions as follows. Let v be a place of F . We say that functions $f' \in \mathcal{S}(G'(F_v))$ and $f \in \mathcal{S}(G(F_v))$ *match* each other or that they are *smooth transfer* of each other if we have the equality

$$O(\delta, f) = \Omega_v(\gamma)O(\gamma, f')$$

for every $\delta \in G_{rs}(F_v)$ and $\gamma \in G'_{rs}(F_v)$ whose orbits correspond to each other via the embedding (2). Similarly, we say that functions $\varphi' \in \mathcal{S}(\mathfrak{s}_{n+1}(F_v))$ and $\varphi \in \mathcal{S}(\mathfrak{u}(V)(F_v))$ *match* each other or that they are *smooth transfer* of each other if we have the equality

$$O(X, \varphi_W) = \omega_v(Y)O(Y, \varphi')$$

for every $X \in \mathfrak{u}(V)_{rs}(F_v)$ and $Y \in \mathfrak{s}_{n+1,rs}(F_v)$ whose orbits correspond to each other via the embedding (5).

If the place v splits in E then the existence of smooth transfer is easy (see [Zh1, Proposition 2.5]). One of the main achievement of [Zh1] was to prove the existence of smooth transfer for nonarchimedean places. In other words, we have the following (see [Zh1, Theorem 2.6]):

Theorem 3.4.1 (Zhang) *Let v be a nonarchimedean place of F .*

- (i) *For every function $f' \in \mathcal{S}(G'(F_v))$ there exists a function $f \in \mathcal{S}(G(F_v))$, matching f' and conversely for every function $f \in \mathcal{S}(G(F_v))$ there exists a function $f' \in \mathcal{S}(G'(F_v))$ which matches f .*
- (ii) *For every function $\varphi' \in \mathcal{S}(\mathfrak{s}_{n+1}(F_v))$ there exists a function $\varphi \in \mathcal{S}(\mathfrak{u}(V)(F_v))$ matching φ' and conversely for every function $\varphi \in \mathcal{S}(\mathfrak{u}(V)(F_v))$ there exists a function $\varphi' \in \mathcal{S}(\mathfrak{s}_{n+1}(F_v))$ which matches φ .*

One of the main ingredient in the proof of Zhang was the following (see [Zh1, Theorem 4.17])

Theorem 3.4.2 (Zhang) *Let v be any place of F . If $\varphi \in \mathcal{S}(\mathfrak{u}(V)(F_v))$ and $\varphi' \in \mathcal{S}(\mathfrak{s}_{n+1}(F_v))$ match then so do $\epsilon(\frac{1}{2}, \eta_{E_v/F_v}, \psi)^{n(n+1)/2} \widehat{\varphi}$ and $\widehat{\varphi}'$.*

In a recent paper Xue ([Xue]) was able to extend Zhang results to obtain a weak version of smooth transfer at archimedean places (which however is sufficient for many global applications). In order to state Xue's result in the generality that we need, we have to vary the hermitian space W . Let us denote momentarily the groups G and H by G^W and H^W . To every hermitian space W' of rank n over E we can associate similar groups $G^{W'}$ and $H^{W'}$ and replacing W by W' everywhere in the previous paragraphs we have a notion of matching between test functions in $\mathcal{S}(G'(F_v))$ and test functions in $\mathcal{S}(G^{W'}(F_v))$, v a place of F . Then, Xue's result reads as follows:

Theorem 3.4.3 (Xue) *Let v be an archimedean place of F . Then, the space of functions $f' \in \mathcal{S}(G'(F_v))$ admitting a smooth transfer to $\mathcal{S}(G^{W'}(F_v))$ for all hermitian space W' of rank n over E is dense in $\mathcal{S}(G'(F_v))$. Similarly, the space of functions $f \in \mathcal{S}(G^W(F_v))$ such that*

there exists a function $f' \in \mathcal{S}(G'(F_v))$ matching f and with the property that for all hermitian space W' of rank n over E for which $W'_v \not\cong W_v$ the function f' match $0 \in \mathcal{S}(G^{W'}(F_v))$, is dense in $\mathcal{S}(G(F_v))$.

Parallel to the existence of smooth transfer there is also a fundamental lemma for the case at hand. This fundamental lemma has been proved by Yun in (sufficiently large) positive characteristic [Yu] and extended to the characteristic zero case by J.Gordon in the appendix to [Yu]. It can be stated as follows:

Theorem 3.4.4 (Yun-Gordon) *There exists a constant $c(n)$ depending only on n such that for every place v of F of residual characteristic greater than $c(n)$ the following holds: If W_v admits a self-dual lattice L_v then the function $f'_v = \mathbf{1}_{K'_v}$ match the function $f_v = \mathbf{1}_{G(\mathcal{O}_v)}$ where we have defined a model of G over \mathcal{O}_v using the self-dual lattice L_v , otherwise the function $f'_v = \mathbf{1}_{K'_v}$ match the function $f_v = 0$.*

3.5 Transfer of spherical characters, Zhang's conjecture and Ichino-Ikeda conjecture

We shall say of a function $f \in \mathcal{S}(G(\mathbb{A}))$ that it is *nice* if it satisfies the following conditions:

- f is factorizable: $f = \otimes_v f_v$;
- There exists a nonarchimedean place v_1 of F and a finite union Ω_1 of cuspidal Bernstein components of $G(F_{v_1})$ such that $f_{v_1} \in \mathcal{S}(G(F_{v_1}))_{\Omega_1}$;

We define the notion of *nice* function on $G'(\mathbb{A})$ similarly. To state the next theorem, we will need to consider more than one pair of hermitian spaces (W, V) . Recall that we have an orthogonal decomposition $V = W \oplus^\perp Ee$ where $(e, e) = 1$. To any (isomorphism class of) n -dimensional hermitian space W' over E we associate the pair (W', V') where $V' = W' \oplus^\perp Ee$. Using such a pair we may construct a new pair $(H^{W'}, G^{W'})$ of reductive groups over F where $H^{W'} = U(W')$ and $G^{W'} = U(W') \times U(V')$. Note that if $W' = W$ then $(H^{W'}, G^{W'}) = (H, G)$. The discussions of the previous paragraphs of course apply verbatim to $(H^{W'}, G^{W'})$. In particular we have a notion of matching between functions in $\mathcal{S}(G^{W'}(F_v))$ and $\mathcal{S}(G'(F_v))$, v a place of F , and a notion of nice function on $G^{W'}(\mathbb{A})$. We shall say that a nice function $f' \in \mathcal{S}(G'(\mathbb{A}))$ match a tuple of nice functions $(f^{W'})_{W'}$, $f^{W'} \in \mathcal{S}(G^{W'}(\mathbb{A}))$ and W' running over all isomorphism classes of n -dimensional hermitian spaces over E , if for all W' and all place v of F the functions f'_v and $f_v^{W'}$ match. Comparing two (simple) global relative trace formulae that have been proposed by Jacquet and Rallis ([JR]), Zhang proves the following (see [Zh1, Proposition 2.10] and [Zh2, Theorem 4.3]):

Theorem 3.5.1 (Zhang) *Let π be an abstractly H -distinguished cuspidal automorphic representation of $G(\mathbb{A})$ such that $BC(\pi)$ is cuspidal and for all non-split archimedean place v the representation π_v is tempered. Let $f \in \mathcal{S}(G(\mathbb{A}))$ and $f' \in \mathcal{S}(G'(\mathbb{A}))$ be nice functions and*

assume that there exists a tuple $(f^{W'})_{W'}$, $f^{W'} \in \mathcal{S}(G^{W'}(\mathbb{A}))$, of nice functions matching f' such that $f^W = f$. Then, we have

$$J_\pi(f) = 2^{-2}L(1, \eta_{E/F})^{-2}I_{BC(\pi)}(f')$$

Remark 3.5.2 *The above theorem differs slightly from [Zh1, Proposition 2.10] and [Zh2, Theorem 4.3] for the following reasons:*

- *First we are not assuming that our test functions are supported in the regular semi-simple locus (this is one of the requirement that Zhang imposes on his 'nice' functions);*
- *We are using test functions that are not necessarily compactly supported (since they are only rapidly decreasing, i.e. in the Schwartz space, at the archimedean places). This is necessary if we want to apply this theorem in conjunction with Xue's result (Theorem 3.4.3) as the dense subspace of 'transferable' test functions that he constructs has no reason to consist of compactly supported functions;*
- *We have drop the condition that π is supercuspidal at one split places but we add the assumption that π must be tempered at all non-split archimedean places.*

This last point is only minor: dropping the condition of π being supercuspidal at one split place can be done by slightly modifying Zhang's original argument when he separates contributions in the spectral side of the simple trace formulas and using the recent extension by Ramakrishnan of his 'mild Tchebotarev theorem for $GL(n)$ ' [Ra]. Moreover, here we assume that π is tempered at all non-split archimedean places in order to have a result independent of the local Gan-Gross-Prasad conjecture for generic L -packets at archimedean places (this conjecture is currently only known for tempered L -packets at these places, see §2.4), this conjecture was granted as a working hypothesis in [Zh2]. The first point is more serious and to get rid of this assumption on the support we have to use the recent works of Zydor [Zy] on regularization of the geometric side of the Jacquet-Rallis trace formulae and Chaudouard-Zydor [CZ] on extending the transfer to singular orbital integrals. Finally, the extension to rapidly decreasing functions (at the archimedean places) is an easy matter using basic estimates on these functions. A convenient way to do this is to introduce some norms on the automorphic quotient $[G] = G(F)\backslash G(\mathbb{A})$. We give definitions and basic properties of these norms in the appendix. Also, for convenience of the reader, we provide in appendix A a full proof of Theorem 3.5.1 using the aforementioned results of Zydor and Chaudouard-Zydor

Thanks to the theory of Rankin-Selberg convolution due to Jacquet, Piatetski-Shapiro and Shalika ([JPSS]), for every cuspidal automorphic representation π of $G(\mathbb{A})$ whose base change is cuspidal we know a factorization of the global spherical character $I_{BC(\pi)}$. More precisely, if $f' \in \mathcal{S}(G'(\mathbb{A}))$ is completely factorizable, we have (see [Zh2, Proposition 3.6])

$$(1) \quad I_{BC(\pi)}(f') = L(1, \eta_{E/F})^2 \frac{L(1/2, BC(\pi))}{L(1, \pi, Ad)} \prod_v I_{BC(\pi)_v}^\natural(f'_v)$$

An immediate consequence of this factorization of Theorem 3.5.1 and of the multiplicity one theorems of Aizenbud-Gourevitch-Rallis-Schiffmann [AGRS] or of Jiang-Sun-Zhu [JSZ] is the following:

Corollary 3.5.3 *Let π be a globally H -distinguished (i.e. such that $J_\pi \neq 0$) cuspidal automorphic representation of $G(\mathbb{A})$ satisfying the two following conditions:*

- *For all non-split archimedean place v of F , the representation π_v is tempered;*
- *There exists a split nonarchimedean place v_0 such that π_{v_0} is supercuspidal.*

Then for all place v_1 of F different from v_0 where π is tempered, there exists a constant $C(\pi_{v_1}) \in \mathbb{C}$ such that for all pair $(f_{v_1}, f'_{v_1}) \in \mathcal{S}(G(F_{v_1})) \times \mathcal{S}(G'(F_{v_1}))$ of matching functions with the property that f'_{v_1} has a matching test function $f_{v_1}^{W'} \in \mathcal{S}(G^{W'}(F_{v_1}))$ for all hermitian space W' of rank n , we have

$$J_{\pi_{v_1}}(f_{v_1}) = C(\pi_{v_1}) I_{BC(\pi_{v_1})}(f'_{v_1})$$

Remark 3.5.4 *Note that the condition of matching of the function f'_{v_1} is empty if v_1 is nonarchimedean (by Theorem 3.4.1) or splits in E .*

Proof: Let \mathbb{A}^{v_1} denote the adèles outside of v_1 and let $f^{v_1} = \prod_{v \neq v_1} f'_v \in \mathcal{S}(G(\mathbb{A}^{v_1}))$ be a factorizable test function. By the multiplicity one results of [AGRS] and [JSZ] and 3.2 (1), there exists a constant $C \in \mathbb{C}$ such that

$$J_\pi(f^{v_1} \otimes f_{v_1}) = C J_{\pi_{v_1}}(f_{v_1})$$

for all $f_{v_1} \in \mathcal{S}(G(F_{v_1}))$. Since $J_\pi \neq 0$ we may choose the function f^{v_1} so that $C \neq 0$. Moreover, up to replacing f_{v_0} by its projection to $\mathcal{S}(G(F_{v_0}))_{\Omega_0}$, where Ω_0 denotes the Bernstein component of π_{v_0} , we may assume that $f_{v_0} \in \mathcal{S}(G(F_{v_0}))_{\Omega_0}$. Then, for all $f_{v_1} \in \mathcal{S}(G(F_{v_1}))$ the function $f = f^{v_1} \otimes f_{v_1}$ is nice. By Theorem 3.4.3 of Xue we can also arrange f^{v_1} such that for all non-split archimedean place v the function f_v admits a transfer $f'_v \in \mathcal{S}(G'(F_v))$ which itself admits a transfer to $\mathcal{S}(G^{W'}(F_v))$ for all rank n hermitian space W' over E . By Theorem 3.4.1 of Zhang and our choice of f^{v_1} at non-split archimedean places, we can find a factorizable test function $f'^{v_1} = \prod_{v \neq v_1} f'_v \in \mathcal{S}(G'(\mathbb{A}^{v_1}))$ matching f^{v_1} and moreover admitting a matching to $\mathcal{S}(G^{W'}(\mathbb{A}^{v_1}))$ for all W' . Since v_0 splits in E , there exists an isomorphism $G'_{v_0} \simeq G_{v_0} \times G_{v_0}$ so that $BC(\pi_{v_0}) = \pi_{v_0} \boxtimes \pi_{v_0}^\vee$ and then every function $\varphi' = \varphi_1 \otimes \varphi_2 \in \mathcal{S}(G'(F_{v_0})) \simeq \mathcal{S}(G(F_{v_0})) \otimes \mathcal{S}(G(F_{v_0}))$ match the function $\varphi_1 * \varphi_2^\vee \in \mathcal{S}(G(F_{v_0}))$

where $\varphi_2^\vee(g) := \varphi_2(g^{-1})$ and $*$ denotes the convolution product. From this we easily infer that we can choose $f'_{v_0} \in \mathcal{S}(G'(F_{v_0}))_{\Omega_0 \times \Omega_0^\vee}$, where Ω_0^\vee is the Bernstein component dual to Ω_0 . Then, for all $f'_{v_1} \in \mathcal{S}(G'(F_{v_1}))$ the function $f' := f'^{v_1} \otimes f'_{v_1}$ is nice. Now the result follows from Theorem 3.5.1 applied to $f = f^{v_1} \otimes f_{v_1}$ and $f' = f'^{v_1} \otimes f'_{v_1}$ where $(f_{v_1}, f'_{v_1}) \in \mathcal{S}(G(F_{v_1})) \times \mathcal{S}(G'(F_{v_1}))$ is any pair of matching functions with f'_{v_1} satisfying the condition of the statement (Note that the assumption that π_{v_0} is supercuspidal implies that $BC(\pi)$ is cuspidal). ■

In [Zh2, conjecture 4.4], Zhang makes the following conjecture

Conjecture 3.5.5 *Let v be a place of F and let $\pi_v = \pi_{n,v} \boxtimes \pi_{n+1,v}$ be an irreducible tempered unitary H_v -distinguished representation of $G(F_v)$. Then, for all matching functions $f \in \mathcal{S}(G(F_v))$ and $f' \in \mathcal{S}(G'(F_v))$, we have*

$$I_{BC(\pi_v)}(f'_v) = \kappa_v(\pi_v) L(1, \eta_{E_v/F_v})^{-1} J_{\pi_v}(f_v)$$

the constant $\kappa_v(\pi_v)$ being given by

$$\kappa_v(\pi_v) = |\tau|_{E_v}^{(d_n+d_{n+1})/2} \left(\frac{\epsilon(1/2, \eta_{E_v/F_v}, \psi_v)}{\eta'_v(-2\tau)} \right)^{n(n+1)/2} \omega_{BC(\pi_{n,v})}(\tau)$$

where $\omega_{BC(\pi_{n,v})}$ denotes the central character of $BC(\pi_{n,v})$ and $d_n = \binom{n}{3}$, $d_{n+1} = \binom{n+1}{3}$.

Remark 3.5.6 *The above conjecture actually differs slightly from [Zh2, conjecture 4.4]. Indeed, there is a discrepancy in the definition of the constant $\kappa_v(\pi_v)$. In loc. cit. there is an extra factor $\eta_v(\text{disc}(W))$ and the factor $\eta'_v(-2\tau)$ is replaced by $\eta'_v(\tau)$. This discrepancy seems to come from lemma 9.1 of [Zh2] on the compatibility between the transfer factors on the group and on the Lie algebra (see §3.4(8)). Of course, this difference has no impact for global applications since in any case if π is an automorphic representation then $\prod_v \kappa_v(\pi_v) = 1$.*

Obviously, we may deduce from the conjunction of the above conjecture, of Theorems 3.5.1, 3.4.1 and 3.4.3 and of the factorization (1) some instances of the Ichino-Ikeda conjecture (as stated in the introduction). In [Zh2], Zhang was able to verify conjecture 3.5.5 in certain particular cases (see [Zh2, Theorem 4.6]). More precisely, he proves the conjecture when either

- The place v splits in E ;
- v is nonarchimedean, π_v is unramified and the residue characteristic of v is sufficiently large;
- v is nonarchimedean and π_v is supercuspidal.

The main goal of this paper is to prove conjecture 3.5.5 for all nonarchimedean places v . Namely, we prove

Theorem 3.5.7 *Conjecture 3.5.5 holds for all nonarchimedean place v of F .*

As in [Zh2], this theorem has consequences for the Ichino-Ikeda conjecture. Namely, we will deduce from it the following

Theorem 3.5.8 *Assume that all the archimedean places of F split in E and let π be a cuspidal automorphic representation of $G(\mathbb{A})$ which is everywhere tempered and such that there exists a nonarchimedean place v of F with $BC(\pi_v)$ is supercuspidal. Then conjecture 1.0.1 holds for π .*

The proofs of Theorems 3.5.7 and 3.5.8 will be given in §4.4 and §4.6.

3.6 A globalization result

Until the end of this paragraph, we make the following assumption

the hermitian space W is anisotropic

This means that $H = U(W)$ is an anisotropic group over F .

Let v_1 be a nonarchimedean place of F which is inert in E and let S be a finite set of nonarchimedean places of F which split in E . Let σ be a unitary supercuspidal representation of $G(F_S)$. Recall that $Temp_H(G_{v_1})$ denotes the set of (isomorphism classes of) tempered irreducible $H(F_{v_1})$ -distinguished representations π_1 of $G(F_{v_1})$ (see §2.4). Let $Irr_{v_1, \sigma, H}(G)$ be the set of irreducible representations $\pi_1 \in Irr(G_{v_1})$ for which there exists a cuspidal automorphic representation π of $G(\mathbb{A})$ which is globally H -distinguished (i.e. such that $J_\pi \neq 0$) such that $\pi_{v_1} \simeq \pi_1$ and $\pi_S \simeq \sigma \otimes \chi$ for some unramified character $\chi \in \Psi_{unit}(G_S)$. The goal of this section is to prove the following result:

Proposition 3.6.1 *The set $Irr_{v_1, \sigma, H}(G) \cap Temp(G_{v_1})$ is dense in $Temp_H(G_{v_1})$.*

The proof of this proposition follows closely that of Corollary A.8 in [ILM, appendix A]. We will need a lemma which is the analog of lemma A.2 of *loc. cit.* Before stating it we need to introduce some more notations.

Let $P = MN$ be a parabolic subgroup of G_{v_1} and let σ be a square-integrable representation of $M(F_{v_1})$. We will say that the tempered representation $i_P^{G_{v_1}} \sigma$ of $G(F_{v_1})$ is *regular* if for all $w \in W(G_{v_1}, M)$ we have $w\sigma \not\simeq \sigma$. Recall that this implies that the representation $i_M^{G_{v_1}} \sigma$ is irreducible. We will denote by $Temp_{reg}(G_{v_1})$ the set of all regular tempered representations of $G(F_{v_1})$. It is an open subset of $Temp(G_{v_1})$. Recall that in §2.3 for all $c > 0$ we have defined subsets $Irr_{\leq c}(U(W)_{v_1})$ and $Irr_{\leq c}(U(V)_{v_1})$ of $Irr(U(W)_{v_1})$ and $Irr(U(V)_{v_1})$ respectively. In what follows, we set $Irr_{\leq c}(G_{v_1}) = Irr_{\leq c}(U(W)_{v_1}) \boxtimes Irr_{\leq c}(U(V)_{v_1})$ and $Irr_{unit, \leq c}(G_{v_1}) = Irr_{unit}(G_{v_1}) \cap Irr_{\leq c}(G_{v_1})$.

Lemma 3.6.2 *Let $0 < c < \frac{1}{2}$. Then, $Temp_{reg}(G_{v_1})$ is open in $Irr_{unit, \leq c}(G_{v_1})$ (for the Fell topology).*

Proof: The proof is the same as in [ILM, lemma A.2] the key fact being that the real exponents of tempered representations of unitary groups are all half integers. In *loc. cit.* the authors use the work of Muic on generic square-integrable representations of classical groups ([Mui]) to deduce this fact for tempered generic representations. However, as already noted in [ILM, Remark A.3], the same result holds for all tempered representation thanks to the work of Mœglin and Mœglin-Tadic on classification of square-integrable representations of classical groups ([Moe1],[MT]). Note that the *basic assumption* made by Mœglin and Tadic (see [MT, §2] for a precise statement) to prove their classification is now known since it follows from the canonical normalization of intertwining operators for unitary groups due to Mok ([Mok, Proposition 3.3.1]) and Kaletha-Minguez-Shin-White ([KMSW, lemma 2.2.3]) together with the classical reducibility criterion of Silberger and Harish-Chandra ([Sil1, §5.4], [Sil2, lemma 1.2; lemma 1.3]). For quasi-split unitary groups, a different proof has been given by Mœglin ([Moe2]) using twisted endoscopy. For a proof of the basic assumption for quasisplit symplectic and orthogonal groups using the normalization of intertwining operators see [Xu, Proposition 3.2]. ■

Proof of Proposition 3.6.1: By [Beu1, Corollary 8.6.1], the closure of $Temp_H(G_{v_1})$ is an union of connected components of $Temp(G_{v_1})$. Hence $Temp_{H, reg}(G_{v_1}) := Temp_H(G_{v_1}) \cap Temp_{reg}(G_{v_1})$ is dense in $Temp_H(G_{v_1})$. Let $\pi_1 \in Temp_{H, reg}(G_{v_1})$. It is sufficient to show that π_1 belongs to the closure of $Irr_{v_1, \sigma, H}(G) \cap Temp(G_{v_1})$. Since σ is $H(F_S)$ -distinguished (see §3.2), by [SV] Theorem 6.2.1 and Theorem 6.4.1 we know that π_1 and σ belong to the support of the Plancherel measures for $L^2(H(F_{v_1}) \backslash G(F_{v_1}))$ and $L^2(H(F_S) \backslash G(F_S))$ respectively. From [SV, Theorem 16.3.2], it follows that there exists a sequence of globally H -distinguished automorphic representations π_n of $G(\mathbb{A})$ such that $\pi_{n, v_1} \rightarrow \pi_1$ and $\pi_{n, S} \rightarrow \sigma$ for the Fell topology. Since $\sigma \otimes \Psi_{unit}(G_S)$ is open in $Irr_{unit}(G_S)$ we have $\pi_{n, S} \in \sigma \otimes \Psi_{unit}(G_S)$ for n sufficiently large. This implies that π_{n, v_1} belongs to $Irr_{v_1, \sigma, H}(G)$ and $\pi_n, BC(\pi_n)$ are cuspidal for n sufficiently large.

Set $c = \frac{1}{2} - \frac{1}{(n+1)^2+1}$. By Lemma 2.3.2, π_{n, v_1} belongs to $Irr_{unit, \leq c}(G_{v_1})$ for n sufficiently large. Hence, by Lemma 3.6.2, $\pi_{n, v_1} \in Temp_{H, reg}(G_{v_1})$ for n sufficiently large and this ends the proof of the proposition. ■

4 Proof of Zhang's conjecture

In this section we will prove Theorem 3.5.7 and Theorem 3.5.8. As Theorem 3.5.7 has already been proved by Zhang at every split place v , we only need to prove it at every nonarchimedean place v of F which is inert in E . Fix such a place v . We will now drop all the index v : $E/F = E_v/F_v, G = G_v, H = H_v, G' = G'_v, H'_1 = H'_{1,v}, H'_2 = H'_{2,v}, \psi = \psi_v, \psi_E = \psi_{E,v}$ and so on. Also, to ease notation we will just write $\mathfrak{s} = \mathfrak{s}_{n+1}$. Finally, we will now use *unnormalized* Haar measures (see §2.5). In particular, Theorem 3.5.7 now takes the following form (see [Zh2, lemma 4.7]):

Theorem 4.0.1 *Let $\pi = \pi_n \boxtimes \pi_{n+1}$ be a H -distinguished irreducible tempered representation of $G(F)$. Then, for all matching functions $f \in \mathcal{S}(G(F))$ and $f' \in \mathcal{S}(G'(F))$, we have*

$$I_{BC(\pi)}(f') = \kappa(\pi) J_\pi(f)$$

where

$$\kappa(\pi) = |\tau|_E^{(d_n+d_{n+1})/2} \left(\frac{\epsilon(1/2, \eta_{E/F}, \psi)}{\eta'(-2\tau)} \right)^{n(n+1)/2} \omega_{BC(\pi_n)}(\tau)$$

4.1 A result of Zhang on truncated local expansion of the spherical character I_Π

In this section we recall a result of Zhang [Zh2] on the existence of truncated local expansion for the spherical characters I_Π . This result is the main ingredient in the proof by Zhang of some particular cases of conjecture 3.5.5. It will also play a crucial role in the proof of Theorem 3.5.7.

Let us set

$$\xi_- = \begin{pmatrix} 0 & \dots & \dots & \dots & 0 \\ \tau & \ddots & & & \vdots \\ 0 & \ddots & \ddots & & \vdots \\ \vdots & \ddots & \ddots & \ddots & \vdots \\ 0 & \dots & 0 & \tau & 0 \end{pmatrix} \in \mathfrak{s}(F)$$

It is a regular nilpotent element for the $GL_n(F)$ -action by conjugation (see [Zh2, lemma 6.1]). Zhang has defined a *regularized* orbital integral μ_{ξ_-} over the orbit of ξ_- (see [Zh2, definition 6.10]). It is a $GL_n(F)$ -invariant linear form $\varphi \in \mathcal{S}(\mathfrak{s}(F)) \mapsto \mu_{\xi_-}(\varphi)$ which coincide with the usual orbital integral when the support of φ intersect the orbit of ξ_- in a compact set.

For all $X = \begin{pmatrix} A & u \\ v & w \end{pmatrix} \in \mathfrak{s}_{n+1}(F)$, we define

$$\Delta_-(X) := \det(v, vA, \dots, vA^{n-1})$$

Note that (see §3.4 (7))

$$(1) \quad \omega(X) = \eta(-1)^n \eta'(\Delta_-(X))$$

for all $X \in \mathfrak{s}_{rs}(F)$ and

$$(2) \quad \Delta_-(\xi_-) = (-1)^{n(n-1)/2} \tau^{n(n+1)/2}$$

Let $r > m' > m > 0$ be positive integers. In [Zh2, definition 8.1], Zhang defines a notion of (m, m', r) -admissible test functions on $G'(F)$. They span a finite dimensional subspace of $\mathcal{S}(G'(F))$. In what follows when we say that (m, m', r) is *sufficiently large*, we shall mean that m is sufficiently large, that m' is sufficiently large depending on m and that r is sufficiently large depending on (m, m') . Recall that in §3.3 we have, using a Cayley map, associated to any function $f' \in \mathcal{S}(G'(F))$ a function $f'_\mathfrak{h}$ on $\mathfrak{s}(F)$. Also, in §2.5 we have defined a certain Fourier transform $\varphi \mapsto \widehat{\varphi}$ on $\mathcal{S}(\mathfrak{s}(F))$. We extract from [Zh2] the two following results (see Lemma 8.8, Theorem 8.5 of [Zh2] and their proofs):

Proposition 4.1.1 (Zhang) *Let U and \mathcal{Z} be relatively compact neighborhood of 1 and 0 in $G'(F)$ and $(\mathfrak{s}/GL_n)(F)$ respectively. Then, if (m, m', r) is sufficiently large, for every (m, m', r) -admissible function f' we have $\text{Supp}(f') \subseteq U$ and the function $X \in \mathcal{Z}_{rs} \mapsto \eta'(\Delta_-(X))O(X, \widehat{f'_\mathfrak{h}})$ is constant and equal to $\eta'(\Delta_-(\xi_-))\mu_{\xi_-}(\widehat{f'_\mathfrak{h}})$. Moreover, we can find a (m, m', r) -admissible function f' such that $\mu_{\xi_-}(\widehat{f'_\mathfrak{h}}) \neq 0$.*

Theorem 4.1.2 (Zhang) *Let $\Pi = \Pi_n \boxtimes \Pi_{n+1}$ be an irreducible tempered representation of $G'(F)$. Then, if (m, m', r) is sufficiently large (depending on Π) we have the equality*

$$I_\Pi(f') = |\tau|_E^{(d_n+d_{n+1})/2} \omega_{\Pi_n}(\tau) \mu_{\xi_-}(\widehat{f'_\mathfrak{h}})$$

for all (m, m', r) -admissible function f' , where $d_n = \binom{n}{3}$ and ω_{Π_n} denotes the central character of Π_n .

A direct consequence of Proposition 4.1.1 and Theorem 4.1.2 is the following:

Corollary 4.1.3 *Let $C \subseteq \text{Temp}(G')$ be a compact subset and let U and \mathcal{Z} be relatively compact neighborhood of 1 and 0 in $G'(F)$ and $(\mathfrak{s}/GL_n)(F)$ respectively.. Then, there exists a test function $f' \in \mathcal{S}(G'(F))$ satisfying the following conditions:*

(i) *Supp(f') $\subseteq U$ and the function $X \in \mathcal{Z}_{rs} \mapsto \eta'(\Delta_-(X))O(X, \widehat{f'_\mathfrak{h}})$ is constant and equal to $\eta'(\Delta_-(\xi_-))\mu_{\xi_-}(\widehat{f'_\mathfrak{h}})$;*

(ii) *$\mu_{\xi_-}(\widehat{f'_\mathfrak{h}}) \neq 0$;*

(iii) *For all $\Pi \in C$ we have the equality*

$$I_\Pi(f') = |\tau|_E^{(d_n+d_{n+1})/2} \omega_{\Pi_n}(\tau) \mu_{\xi_-}(\widehat{f'_\mathfrak{h}})$$

Proof: For all $r > m' > m > 0$, let us denote by $C[m, m', r]$ the set of $\Pi \in C$ such that the equality

$$I_\Pi(f') = |\tau|_E^{(d_n+d_{n+1})/2} \omega_{\Pi_n}(\tau) \mu_{\xi_-}(\widehat{f'_\mathfrak{h}})$$

holds for all (m, m', r) -admissible function f' . Note that $C[m, m', r]$ is a closed subset of C . Obviously, by Proposition 4.1.1, it suffices to show that if (m, m', r) is sufficiently large then $C[m, m', r] = C$ and for that we may assume C to be connected. By Theorem 4.1.2 we have

$$\bigcup_{m_0 > 0} \bigcap_{m \geq m_0} \bigcup_{m'_0 > m} \bigcap_{m' \geq m'_0} \bigcup_{r_0 > m'} \bigcap_{r \geq r_0} C[m, m', r] = C$$

Now, by Baire category theorem, this implies that for (m, m', r) sufficiently large the set $C[m, m', r]$ is not meager i.e. it has nonempty interior (since it is closed). By connectedness of C and analyticity of $\Pi \mapsto I_\Pi$ this implies $C[m, m', r] = C$ and this ends the proof. ■

4.2 Weak comparison of local spherical characters

Proposition 4.2.1 *For all $\pi \in \text{Temp}_H(G)$ there exists a nonzero constant $C(\pi) \in \mathbb{C}$ such that for all matching functions $f \in \mathcal{S}(G(F))$ and $f' \in \mathcal{S}(G'(F))$ we have*

$$J_\pi(f) = C(\pi) I_{BC(\pi)}(f')$$

Moreover, the function $\pi \in \text{Temp}_H(G) \mapsto C(\pi)$ is analytic.

Proof: Assume that we have proved the existence of a constant $C(\pi)$ as in the proposition for a dense set of π in $\text{Temp}_H(G)$. We claim that the proposition can be deduced from this. Indeed, for all $\pi \in \text{Temp}_H(G)$ we can define a constant $C(\pi)$ as follows: choose any set $(f, f') \in \mathcal{S}(G(F)) \times \mathcal{S}(G'(F))$ of matching functions such that $I_{BC(\pi)}(f') \neq 0$ (the existence of such a pair follows from Theorem 3.4.1 and Theorem 4.1.2) and set $C(\pi) = J_\pi(f) I_{BC(\pi)}(f')^{-1}$. Of course this constant may *a priori* depend on the choice of f and f' but it follows from the analyticity of $\pi \mapsto J_\pi$ and $\Pi \mapsto I_\Pi$ and our assumption that in fact it is independent of such a choice. Still by analyticity of the spherical characters the equality of the proposition is true for all $\pi \in \text{Temp}_H(G)$ and all pair of matching functions (f, f') and the function $\pi \in \text{Temp}_H(G) \mapsto C(\pi)$ is analytic. Moreover it is nowhere zero since for all $\pi \in \text{Temp}_H(G)$ there exists $f \in \mathcal{S}(G(F))$ such that $J_\pi(f) \neq 0$ and there exists a $f' \in \mathcal{S}(G'(F))$ matching f (by Theorem 3.4.1).

We now prove the existence of a dense subset of π satisfying the proposition. To this end we will use Proposition 3.6.1. We first need to globalize the situation at hand. Let

- \mathbb{E}/\mathbb{F} be a quadratic extension of number fields such that all archimedean places of \mathbb{F} are nonsplit in \mathbb{E} and v_1 be a place of \mathbb{F} such that $\mathbb{E}_{v_1}/\mathbb{F}_{v_1} \simeq E/F$;
- \mathbb{W} a n -dimensional hermitian space over \mathbb{E} such that
 - for all archimedean place v of \mathbb{F} the group $U(\mathbb{W})_v$ is anisotropic (in particular \mathbb{W} is anisotropic);
 - $\mathbb{W}_{v_0} \simeq W$.

We will set $\mathbb{V} = \mathbb{W} \oplus^\perp \mathbb{E}e$ where $(e, e) = 1$ (so that $\mathbb{V}_{v_1} \simeq V$), $\mathbb{H} = U(\mathbb{W})$ and $\mathbb{G} = U(\mathbb{V})$. Let v_0 be two nonarchimedean places of \mathbb{F} which split in \mathbb{E} and let σ be a supercuspidal representations of $\mathbb{G}(\mathbb{F}_{v_0})$. Applying Proposition 3.6.1 to $S = \{v_0\}$, we deduce the existence of a dense subset $D \subset \text{Temp}_H(G)$ such that for all $\pi \in D$ there exists a globally \mathbb{H} -distinguished cuspidal automorphic representation Π of $\mathbb{G}(\mathbb{A})$ such that $\Pi_{v_1} \simeq \pi$ and Π_{v_0} is supercuspidal. Applying Corollary 3.5.3 to such representations Π , which we remark are necessarily tempered at all archimedean places since $U(\mathbb{W})$ is anisotropic there, we deduce for all $\pi \in D$ there exists a constant $C(\pi)$ as in the proposition. ■

4.3 A local trace formula

Let $f_1, f_2 \in \mathcal{S}(G(F))$. Then

(1) The integral

$$J(f_1, f_2) = \int_{H(F)} \int_{H(F)} \int_{G(F)} f_1(h_1 g h_2) f_2(g) dg dh_1 dh_2$$

is absolutely convergent.

This follows from [Zh1, Lemma A.4].

By [Beu1, Proposition 8.2.1(v)], we have

$$(2) \quad J(f_1, f_2) = \int_{\text{Temp}_H(G)} J_\pi(f_1) J_{\pi^\vee}(f_2) d\mu_G(\pi)$$

where $d\mu_G(\pi)$ denotes the Harish-Chandra-Plancherel measure of $G(F)$. We also have (see §3.3 for the definition of \tilde{f}_i)

$$J(f_1, f_2) = \int_{U(W)(F)} \int_{U(V)(F)} \tilde{f}_1(h^{-1} x h) \tilde{f}_2(x) dx dh$$

Let us fix open and closed $U(W)(F_v)$ -invariant neighborhoods $\omega \subset \mathfrak{u}(V)(F_v)$ and $\Omega \subset U(V)(F_v)$ of 0 and 1 as in §3.3. Assume that \tilde{f}_2 is supported in Ω . Then, we have (see §3.3 for the definition of $f_{i, \mathfrak{h}}$)

$$J(f_1, f_2) = \int_{U(W)(F)} \int_{\mathfrak{u}(V)(F)} f_{1, \mathfrak{h}}(h^{-1} X h) f_{2, \mathfrak{h}}(X) dX dh$$

By Fourier transform, we also have

$$J(f_1, f_2) = \int_{U(W)(F)} \int_{\mathfrak{u}(V)(F)} \check{f}_{1, \mathfrak{h}}(h^{-1} X h) \hat{f}_{2, \mathfrak{h}}(X) dX dh$$

By [Zh1, Corollary 4.5] this expression is absolutely convergent so that we can switch the two integrals and we finally get

$$(3) \quad J(f_1, f_2) = \int_{\mathfrak{u}(V)(F)} \check{f}_{1, \mathfrak{h}}(X) O(X, \widehat{f}_{2, \mathfrak{h}}) dX$$

Summing up, from (2) and (3) we deduce that

$$(4) \quad \int_{Temp_H(G)} J_\pi(f_1) J_{\pi^\vee}(f_2) d\mu_G(\pi) = \int_{\mathfrak{u}(V)(F)} \check{f}_{1, \mathfrak{h}}(X) O(X, \widehat{f}_{2, \mathfrak{h}}) dX$$

for all functions $f_1, f_2 \in \mathcal{S}(G(F))$ with $Supp(\widehat{f}_2) \subseteq \Omega$.

We will also need the following formula (see [Beu1, Proposition 8.2.1(iv)]):

$$(5) \quad \check{f}(1) = \int_{Temp_H(G)} J_\pi(f) d\mu_G(\pi)$$

for all $f \in \mathcal{S}(G(F))$.

4.4 Proof of Theorem 4.0.1

We keep the notations of the previous paragraph. Let $f \in \mathcal{S}(G(F))$. Denote by $C \subseteq Temp_H(G)$ the support of the function $\pi \mapsto J_\pi(f)$. It is a compact set and so is its dual C^\vee . Let us denote by \mathcal{Y} the image of Ω in $(H \backslash G/H)(F) = (U(V)/U(W))(F)$ and by \mathcal{Z} the image of the support of $\check{f}_{\mathfrak{h}}$ in $(\mathfrak{u}(V)/U(W))(F)$. We will denote by the same letters the corresponding subsets in $(H'_1 \backslash G'/H'_2)(F)$ and $(\mathfrak{s}/GL_n)(F)$ respectively (see §3.4 (1) and (4)). By Corollary 4.1.3, there exists a function $f' \in \mathcal{S}(G'(F))$ such that

- f' is supported in the inverse image of \mathcal{Y} in $G'(F)$;
- The function $Y \in \mathcal{Z}_{r_s} \mapsto \eta'(\Delta_-(Y)) O(Y, \widehat{f}'_{\mathfrak{h}})$ is constant and equal $\eta'(\Delta_-(\xi_-)) \mu_{\xi_-}(\widehat{f}'_{\mathfrak{h}})$;
- $\mu_{\xi_-}(\widehat{f}'_{\mathfrak{h}}) \neq 0$;
- For all $\Pi \in BC(C^\vee)$ we have

$$I_\Pi(f') = |\tau|_E^{(d_n + d_{n+1})/2} \omega_{\Pi_n}(\tau) \mu_{\xi_-}(\widehat{f}'_{\mathfrak{h}})$$

Let $f_2 \in \mathcal{S}(G(F))$ be a function matching f' (whose existence is guaranteed by Theorem 3.4.1). Up to multiplying f_2 by the characteristic function of \mathcal{Y} we may assume that \check{f}_2 is supported in Ω . By §3.4(8) the functions $\eta(2)^{n(n+1)/2} f'_{\mathfrak{h}}$ and $f_{2, \mathfrak{h}}$ match. Hence, by Theorem 3.4.2 so do $\eta(2)^{n(n+1)/2} \widehat{f}'_{\mathfrak{h}}$ and $\epsilon(\frac{1}{2}, \eta_{E/F}, \psi)^{n(n+1)/2} \widehat{f}_{2, \mathfrak{h}}$. Thus, by 4.1(1) and 4.1(2), for all $X \in \Omega_{r_s}$, denoting by $Y \in \mathcal{Z}_{r_s}$ the corresponding element, we have

$$\begin{aligned}
\epsilon\left(\frac{1}{2}, \eta_{E/F}, \psi\right)^{n(n+1)/2} O(X, \widehat{f}_{2, \mathfrak{h}}) &= \eta(2)^{n(n+1)/2} \eta(-1)^n \eta'(\Delta_-(Y)) O(Y, \widehat{f}'_{\mathfrak{h}}) \\
&= \eta(2)^{n(n+1)/2} \eta(-1)^n \eta'(\Delta_-(\xi_-)) \mu_{\xi_-}(\widehat{f}'_{\mathfrak{h}}) \\
&= \eta'(-2\tau)^{n(n+1)/2} \mu_{\xi_-}(\widehat{f}'_{\mathfrak{h}})
\end{aligned}$$

Consequently, we have

$$\begin{aligned}
(1) \quad \left(\frac{\epsilon\left(\frac{1}{2}, \eta_{E/F}, \psi\right)}{\eta'(-2\tau)}\right)^{n(n+1)/2} \int_{\mathfrak{u}(V)(F)} \check{f}_{\mathfrak{h}}(X) O(X, \widehat{f}_{2, \mathfrak{h}}) dX &= \mu_{\xi_-}(\widehat{f}'_{\mathfrak{h}}) \int_{\mathfrak{u}(V)(F)} \check{f}_{\mathfrak{h}}(X) dX \\
&= \mu_{\xi_-}(\widehat{f}'_{\mathfrak{h}}) f_{\mathfrak{h}}(0) = \mu_{\xi_-}(\widehat{f}'_{\mathfrak{h}}) \check{f}(1) \\
&= \mu_{\xi_-}(\widehat{f}'_{\mathfrak{h}}) \int_{Temp_H(G)} J_{\pi}(f) d\mu_G(\pi)
\end{aligned}$$

where the last equality follows from 4.3(5).

On the other hand, by Proposition 4.2.1, for all $\pi \in C$ we have

$$J_{\pi^\vee}(f_2) = C(\pi^\vee) I_{BC(\pi^\vee)}(f') = C(\pi^\vee) |\tau|_E^{(d_n+d_{n+1})/2} \omega_{BC(\pi^\vee)_n}(\tau) \mu_{\xi_-}(\widehat{f}'_{\mathfrak{h}})$$

It follows that

$$\begin{aligned}
(2) \quad \int_{Temp_H(G)} J_{\pi}(f) J_{\pi^\vee}(f_2) d\mu_G(\pi) &= \\
&= \mu_{\xi_-}(\widehat{f}'_{\mathfrak{h}}) |\tau|_E^{(d_n+d_{n+1})/2} \int_{Temp_H(G)} J_{\pi}(f) C(\pi^\vee) \omega_{BC(\pi^\vee)_n}(\tau) d\mu_G(\pi)
\end{aligned}$$

Since $\mu_{\xi_-}(\widehat{f}'_{\mathfrak{h}}) \neq 0$, we deduce from (1), (2) and 4.3(4) that

$$(3) \quad \int_{Temp_H(G)} J_{\pi}(f) (\kappa(\pi^\vee) C(\pi^\vee) - 1) d\mu_G(\pi) = 0$$

for all $f \in \mathcal{S}(G(F))$ and where

$$\kappa(\pi) = |\tau|_E^{(d_n+d_{n+1})/2} \left(\frac{\epsilon\left(\frac{1}{2}, \eta_{E/F}, \psi\right)}{\eta'(-2\tau)}\right)^{n(n+1)/2} \omega_{BC(\pi)_n}(\tau)$$

Let $\mathcal{Z}(G)$ denotes the Bernstein center of G (see [BD]). We may see $\mathcal{Z}(G)$ as a unital subalgebra of the space of continuous functions on $Temp(G)$ which moreover acts on $\mathcal{S}(G(F))$ with the property that $J_{\pi}(z \star f) = z(\pi) J_{\pi}(f)$ for all $z \in \mathcal{Z}(G)$, all $f \in \mathcal{S}(G(F))$ and all $\pi \in Temp(G)$. Thus by (3), we get

$$(4) \quad \int_{Temp_H(G)} z(\pi) J_\pi(f) (\kappa(\pi^\vee) C(\pi^\vee) - 1) d\mu_G(\pi) = 0$$

for all $f \in \mathcal{S}(G(F))$ and all $z \in \mathcal{Z}(G)$. For all $\pi \in Temp(G)$ let us denote by χ_π the 'infinitesimal character' of π , that is the algebra homomorphism $\chi_\pi : \mathcal{Z}(G) \rightarrow \mathbb{C}$ given by $\chi_\pi(z) := z(\pi)$ for all $z \in \mathcal{Z}(G)$. Set $Y := Specmax \mathcal{Z}(G)$. Then, the map $Temp(G) \rightarrow Y$, $\pi \mapsto \chi_\pi$ is continuous and proper. Let $Y_{temp} \subseteq Y$ be the image of this map and μ_Y be the push-forward of the Plancherel measure μ_G to Y_{temp} . Then, by the disintegration of measures there exists a measurable mapping $\chi \mapsto \mu_\chi$ from Y_{temp} to the space of measures on $Temp(G)$ such that

$$(5) \quad \int_{Temp(G)} \varphi(\pi) d\mu_G(\pi) = \int_{Y_{temp}} \int_{Temp(G)} \varphi(\pi) d\mu_\chi(\pi) d\mu_Y(\chi)$$

for all continuous compactly-supported function $\varphi : Temp(G) \rightarrow \mathbb{C}$ and such that for all $\chi \in Y_{temp}$, μ_χ is supported on $Temp_\chi(G) := \{\pi \in Temp(G) \mid \chi_\pi = \chi\}$. By (4), we get

$$(6) \quad \int_{Y_{temp}} z(\chi) \int_{Temp_{H,\chi}(G)} J_\pi(f) (\kappa(\pi^\vee) C(\pi^\vee) - 1) d\mu_\chi(\pi) d\mu_Y(\chi) = 0$$

for all $f \in \mathcal{S}(G(F))$ and all $z \in \mathcal{Z}(G)$ where we have set $Temp_{H,\chi}(G) := Temp_H(G) \cap Temp_\chi(G)$. Since the restriction of $\mathcal{Z}(G)$ to Y_{temp} is self-adjoint (i.e. for all $z \in \mathcal{Z}(G)$ there exists $z^* \in \mathcal{Z}(G)$ such that $z^*(\chi) = \overline{z(\chi)}$ for all $\chi \in Y_{temp}$), separates points and for all $f \in \mathcal{S}(G(F))$ the function $\pi \in Temp(G) \mapsto J_\pi(f)$ is compactly supported, by (6) and the Stone-Weierstrass theorem for μ_Y -almost all $\chi \in Y_{temp}$, we get

$$\int_{Temp_{H,\chi}(G)} J_\pi(f) (\kappa(\pi^\vee) C(\pi^\vee) - 1) d\mu_\chi(\pi) = 0$$

for all $f \in \mathcal{S}(G(F))$. Since $Temp_\chi(G)$ is finite, we have

$$\int_{Temp_{H,\chi}(G)} J_\pi(f) (\kappa(\pi^\vee) C(\pi^\vee) - 1) d\mu_\chi(\pi) = \sum_{\pi \in Temp_{H,\chi}(G)} J_\pi(f) (\kappa(\pi^\vee) C(\pi^\vee) - 1) \mu_\chi(\pi)$$

for μ_Y -almost all $\chi \in Y_{temp}$ and all $f \in \mathcal{S}(G(F))$. Finally, as the spherical characters J_π for $\pi \in Temp_{H,\chi}(G)$ are linearly independent, we get that

$$(\kappa(\pi^\vee) C(\pi^\vee) - 1) \mu_\chi(\pi) = 0$$

for μ_Y -almost all $\chi \in Y_{temp}$ and all $\pi \in Temp_{H,\chi}(G)$ which by (5) means that $\kappa(\pi) C(\pi) = 1$ for μ_G -almost all $\pi \in Temp_H(G)$. Since $\pi \in Temp_H(G) \mapsto \kappa(\pi) C(\pi)$ is analytic and the support of μ_G is precisely $Temp(G)$, it follows that $\kappa(\pi) C(\pi) = 1$ for all $\pi \in Temp_H(G)$ which is what we wanted. ■

4.5 A first corollary

In this paragraph we prove the following corollary to Theorem 4.0.1. It will be needed for the proof of Theorem 3.5.8.

Corollary 4.5.1 *Let $f \in \mathcal{S}(G(F))$ and $f' \in \mathcal{S}(G'(F))$. Then f and f' match if and only if we have*

$$I_{BC(\pi)}(f') = \kappa(\pi)J_\pi(f)$$

for all $\pi \in \text{Temp}_H(G)$ and where as before we have set

$$\kappa(\pi) = |\tau|_E^{(d_n+d_{n+1})/2} \left(\frac{\epsilon(\frac{1}{2}, \eta_{E/F}, \psi)}{\eta'(-2\tau)} \right)^{n(n+1)/2} \omega_{BC(\pi)_n}(\tau)$$

Proof: The necessity follows from Theorem 4.0.1. Let us prove the sufficiency. Thus, we assume that

$$I_{BC(\pi)}(f') = \kappa(\pi)J_\pi(f)$$

for all $\pi \in \text{Temp}_H(G)$ and we want to prove that f and f' match. Let $f_2 \in \mathcal{S}(G(F))$ be a function which matches f' (such a function exists by Theorem 3.4.1). Then, by Theorem 4.0.1 and the assumption, for all $\pi \in \text{Temp}_H(G)$ we have $J_\pi(f) = J_\pi(f_2)$. Thus, by 4.3(2), for all $f_1 \in \mathcal{S}(G(F))$ we have

$$(1) \quad J(f_1, f) = J(f_1, f_2)$$

Let $x_0 \in U(V)_{rs}(F)$ and choose f_1 so that \tilde{f}_1 is supported in a small neighborhood of x_0 in $U(V)_{rs}(F)$. Then a formal manipulation, which is justified since everything is absolutely convergent here, yields

$$(2) \quad J(f_1, f) = \int_{U(V)(F)} f_1(x)O(x, f)dx$$

and

$$(3) \quad J(f_1, f_2) = \int_{U(V)(F)} f_1(x)O(x, f_2)dx$$

Since the functions $x \in U(V)_{rs}(F) \mapsto O(x, f)$ and $x \in U(V)_{rs}(F) \mapsto O(x, f_2)$ are locally constant (see [Zh1, Proposition 3.13]), we may choose f_1 such that $\int_{U(V)(F)} f_1(x)O(x, f)dx =$

$O(x_0, f)$ and $\int_{U(V)(F)} f_1(x)O(x, f_2)dx = O(x_0, f_2)$. For such a choice, it follows from (1), (2) and (3) that $O(x_0, f) = O(x_0, f_2)$. As x_0 was arbitrary we see that f and f_2 have the same regular semisimple orbital integrals and hence f and f' match. ■

4.6 Proof of Theorem 3.5.8

We may assume that π is abstractly $H(\mathbb{A})$ -distinguished (hence for all v , π_v is H_v -distinguished). By the multiplicity one theorems of Aizenbud-Gourevitch-Rallis-Schiffmann [AGRS] of Jiang-Sun-Zhu [JSZ], there exists a constant C such that

$$J_\pi(f) = C \prod_v J_{\pi_v}^\natural(f_v)$$

for all factorizable test function $f = \prod_v f_v \in \mathcal{S}(G(\mathbb{A}))$ and we only need to show that $C = 4^{-1}\mathcal{L}(\pi, 1/2)$. For this, it is sufficient to prove the existence of $f \in \mathcal{S}(G(\mathbb{A}))$ with

$$J_\pi(f) = 4^{-1}\mathcal{L}(\pi, \frac{1}{2}) \prod_v J_{\pi_v}^\natural(f_v)$$

and $J_{\pi_v}^\natural(f_v) \neq 0$ for all place v . Let v_1 be a (nonarchimedean) place of F such that $BC(\pi_{v_1})$ is supercuspidal (such a place exists by assumption). This implies in particular that $BC(\pi)$ is cuspidal and π_{v_1} supercuspidal (by Lemma 2.3.1). By Theorem 3.5.1, Theorem 3.5.7 and identity 3.5 (1), it suffices to show that there exist a nice function $f' \in \mathcal{S}(G'(\mathbb{A}))$ matching a tuple of nice functions $(f^{W'})_{W'}$, $f^{W'} \in \mathcal{S}(G^{W'}(\mathbb{A}))$, such that $I_{BC(\pi_v)}(f'_v) \neq 0$ for all v . Let Ω_1 be the Bernstein component of $BC(\pi_{v_1})$ in $G'(F_{v_1})$. Then, we can find a function $f'^{\circ}_{v_1} \in \mathcal{S}(G'(F_{v_1}))_{\Omega_1}$ such that $I_{BC(\pi_{v_1})}(f'^{\circ}_{v_1}) \neq 0$. Let $f' = \prod_v f'_v$ be a factorizable test function in $\mathcal{S}(G'(\mathbb{A}))$ such that $f'_{v_1} = f'^{\circ}_{v_1}$ and $I_{BC(\pi_v)}(f'_v) \neq 0$ for all other place v . By construction, the function f' is nice. Moreover, by the assumption on archimedean places, Theorem 3.4.1 and Theorem 3.4.4, we can find a tuple of functions $(f^{W'})_{W'}$, $f^{W'} \in \mathcal{S}(G^{W'}(\mathbb{A}))$, matching f' . Of course, the functions $f^{W'}$ have no reason of being nice. However, by Lemma 2.3.1 for all W' there exists a finite union $\Omega_1^{W'}$ of cuspidal Bernstein components of $G^{W'}(F_{v_1})$ such that $\Omega_1^{W'}$ contains all irreducible representation of $G^{W'}(F_{v_1})$ whose base change belongs to Ω_1 and by Corollary 4.5.1 up to replacing $f^{W'}$ by its projection $f^{W'}_{v_1, \Omega_1^{W'}}$ onto $\mathcal{S}(G^{W'}(F_{v_1}))_{\Omega_1^{W'}}$ we may assume that $f^{W'}_{v_1} = f^{W'}_{v_1, \Omega_1^{W'}}$. Then, for all W' the function $f^{W'}$ is nice and we are done. ■

A Comparison of relative trace formulae

The goal of this appendix is to provide a proof of Theorem 3.5.1. Inspired by [Kott, §18], we start by introducing a convenient notion of norms on the adelic points of a variety over F .

A.1 Norms on adelic varieties

We will use the following convenient although not very precise notations. If f_1, f_2 are positive valued functions on a set X we will write

$$f_1(x) \ll f_2(x), \quad \text{for all } x \in X$$

to mean that there exists a constant $C > 0$ such that $f_1(x) \leq Cf_2(x)$ for all $x \in X$. We will also write

$$f_1(x) < f_2(x), \quad \text{for all } x \in X$$

or just $f_1 < f_2$ if there exist constants $C, d > 0$ such that $f_1(x) \leq C f_2(x)^d$ for all $x \in X$. Finally, we will write

$$f_1(x) \sim f_2(x), \quad \text{for all } x \in X$$

or simply $f_1 \sim f_2$ if $f_1 < f_2$ and $f_2 < f_1$.

Let X be a set. By an abstract norm on X we will just mean a function $\|\cdot\| : X \rightarrow [1, +\infty[$. Let $\|\cdot\|_1$ and $\|\cdot\|_2$ be two abstract norms on X . We will say that $\|\cdot\|_1$ dominates $\|\cdot\|_2$ if $\|x\|_2 < \|x\|_1$ for all $x \in X$ and we will say that $\|\cdot\|_1$ and $\|\cdot\|_2$ are equivalent if $\|\cdot\|_1$ dominates $\|\cdot\|_2$ and $\|\cdot\|_2$ dominates $\|\cdot\|_1$ i.e. if $\|\cdot\|_1 \sim \|\cdot\|_2$. Let $f : X \rightarrow Y$ be a map between two sets and let $\|\cdot\|_Y$ be an abstract norm on Y . Then, we define an abstract norm $f^*\|\cdot\|_Y$ on X by

$$f^*\|x\|_Y := \|f(x)\|_Y$$

for all $x \in X$.

Let F be a number field, \mathbb{A} its ring of adeles and for every place v of F we will denote by F_v the corresponding completion. For every finite extension F' of F , we will write $\mathbb{A}_{F'} = \mathbb{A} \otimes_F F'$ for the adèle ring of F' . We fix algebraic closures \overline{F} of F and \overline{F}_v of F_v . For every place v of F , we will denote by $|\cdot|_v$ the normalized absolute value on F_v . This absolute value extends uniquely to an absolute value on \overline{F}_v that we will also denote by $|\cdot|_v$. We define

$$\mathbb{A}_{\overline{F}} = \overline{F} \otimes_F \mathbb{A} = \varinjlim_{F'} \mathbb{A}_{F'}$$

where the limits is taken over all finite subextension of \overline{F}/F . Let X be an algebraic variety over F (i.e. a reduced separated scheme of finite type over F). Since X is of finite type we have $X(\mathbb{A}_{\overline{F}}) = \varinjlim_{F'} X(\mathbb{A}_{F'})$. We are going to define certain (equivalence classes of) abstract

norms on $X(\mathbb{A}_{\overline{F}})$ and $X(\overline{F}_v)$, v a place of F . The definition of these abstract norms is mainly inspired by [Kott, §18]. First assume that X is affine and choose a set $\{P_1, \dots, P_k\}$ of generators for the F -algebra $F[X]$. For every place v of F we define an abstract norm $\|\cdot\|_{X_v}$ on $X(\overline{F}_v)$ by

$$\|x\|_{X_v} := \max(1, |P_1(x)|_v, \dots, |P_k(x)|_v)$$

for all $x \in X(\overline{F}_v)$. Choosing a different generating set $\{Q_1, \dots, Q_\ell\}$ would yield another family of abstract norms $(\|\cdot\|'_{X_v})_v$ with the following properties:

- For all place v , $\|\cdot\|'_{X_v} \sim \|\cdot\|_{X_v}$;
- There exists $d > 0$ such that for almost all place v , we have

$$\|\cdot\|_{X_v}^{1/d} \leq \|\cdot\|'_{X_v} \leq \|\cdot\|_{X_v}^d$$

In particular, for all v the equivalence class of the abstract norm $\|\cdot\|_{X_v}$ does not depend on the particular generating set chosen and by a *norm* on $X(\overline{F}_v)$ we will mean any abstract norm in this equivalence class. Note that the norms $(\|\cdot\|_{X_v})_v$ constructed above are Galois invariant in the sense that $\|\sigma x\|_{X_v} = \|x\|_{X_v}$ for all $x \in X(\overline{F}_v)$ and all $\sigma \in \text{Gal}(\overline{F}_v/F_v)$. This allows us to extend the norm $\|\cdot\|_{X_v}$ to $X(K)$ for any finite extension K of F_v : choosing any embedding $\iota : K \hookrightarrow \overline{F}_v$ we set

$$\|x\|_{X_v} := \|\iota(x)\|_{X_v}$$

for any $x \in X(K)$.

We now define an abstract norm $\|\cdot\|_X$ on $X(\mathbb{A}_{\overline{F}})$ as follows. Let $x \in X(\mathbb{A}_{\overline{F}})$ and choose a finite extension F'/F such that $x \in X(\mathbb{A}_{F'})$. Then, we may write x as a product $\prod_w x_w$, $x_w \in X(F'_w)$, indexed by the set of places of F' and we set

$$\|x\|_X := \prod_v \left(\prod_{w|v} \|x_w\|_{X_w}^{[F'_w:F_v]} \right)^{1/[F':F]}$$

where the first product is over the set of places v of F and the second product is over the set of places w of F' above v . Note that this definition does not depend on the choice of the finite extension F'/F such that $x \in X(\mathbb{A}_{F'})$. Moreover, choosing a different generating set would give an equivalent abstract norm. By a *norm* on $X(\mathbb{A}_{\overline{F}})$ we will mean any abstract norm in this equivalence class. We will assume from now on that for any affine variety X over F we have fixed norms $\|\cdot\|_X$ on $X(\mathbb{A}_{\overline{F}})$ and norms $\|\cdot\|_{X_v}$ on $X(\overline{F}_v)$, for all place v of F , as above (i.e. by choosing a finite generating set of $F[X]$). In the particular case $X = \mathbb{A}^1$ (the affine line) we will even take

$$\|x\|_{\mathbb{A}_v^1} = \max(1, |x|_v)$$

for all place v of F and for all $x \in X(\overline{F}_v) = \overline{F}_v$. Note that by the product formula we then have

$$(1) \quad \|x\|_{\mathbb{A}^1} = \|x^{-1}\|_{\mathbb{A}^1}$$

for every $x \in \overline{F}^\times$.

We continue to assume that X is affine. Let $\mathcal{U} = (U_i)_{i \in I}$ be a finite covering of X by affine open subsets. We can define another abstract norm $\|\cdot\|_{X_v, \mathcal{U}}$ on $X(\overline{F}_v)$ by

$$\|x\|_{X_v, \mathcal{U}} := \min\{\|x\|_{U_{i,v}}; i \in I \text{ such that } x \in U_i(\overline{F}_v)\}, \quad x \in X(\overline{F}_v)$$

Then we have (see [Kott, Proposition 18.1(6)])

- For all place v , $\|\cdot\|_{X_v, \mathcal{U}} \sim \|\cdot\|_{X_v}$;

- There exists $d > 0$ such that for almost all place v , we have

$$\|\cdot\|_{X_v}^{1/d} \leq \|\cdot\|_{X_v, \mathcal{U}} \leq \|\cdot\|_{X_v}^d$$

- For all place v , $\|\cdot\|_{X_v, \mathcal{U}}$ is Galois invariant.

We can also define an abstract norm $\|\cdot\|_{X, \mathcal{U}}$ on $X(\mathbb{A}_{\overline{F}})$ by sending $x \in X(\mathbb{A}_{F'})$, F'/F a finite extension, to

$$\|x\|_{X, \mathcal{U}} := \prod_v \left(\prod_{w|v} \|x_w\|_{X_w, \mathcal{U}}^{[F'_w:F_v]} \right)^{1/[F':F]}$$

Then $\|\cdot\|_{X, \mathcal{U}}$ is a norm on $X(\mathbb{A}_{\overline{F}})$ (i.e. $\|\cdot\|_{X, \mathcal{U}} \sim \|\cdot\|_X$). This allows us to extend the definition of the abstract norms $\|\cdot\|_X$ and $\|\cdot\|_{X_v}$ to any algebraic variety X over F as follows. Let X be such a variety and choose a finite covering $\mathcal{U} = (U_i)_{i \in I}$ of X by affine open subsets. Then the definitions of the abstract norms $\|\cdot\|_{X, \mathcal{U}}$ and $\|\cdot\|_{X_v, \mathcal{U}}$ as above still make sense and we will set $\|\cdot\|_X := \|\cdot\|_{X, \mathcal{U}}$, $\|\cdot\|_{X_v} := \|\cdot\|_{X_v, \mathcal{U}}$. Choosing a different covering \mathcal{V} of X would give abstract norms $(\|\cdot\|'_{X_v})_v$ and $\|\cdot\|'_X$ satisfying the following

- For all v , $\|\cdot\|'_{X_v} \sim \|\cdot\|_{X_v}$ and there exists $d > 0$ such that for almost all v we have

$$\|\cdot\|_{X_v}^{1/d} \leq \|\cdot\|'_{X_v} \leq \|\cdot\|_{X_v}^d$$

- $\|\cdot\|'_X \sim \|\cdot\|_X$.

In particular the equivalence class of $\|\cdot\|_X$ (resp. of $\|\cdot\|_{X_v}$ for v a place of F) doesn't depend on the particular choice of \mathcal{U} and by a *norm* on $X(\mathbb{A}_{\overline{F}})$ (resp. on $X(\overline{F}_v)$) we will mean any abstract norm in this equivalence class. From now on we assume that every algebraic variety over F has been equipped with a family of norms as above (i.e. by choosing a finite covering \mathcal{U} by affine open subsets). If X is affine we also assume that these norms have been defined using the trivial covering $\mathcal{U} = \{X\}$ so that they coincide with the ones we already fixed. If G is an affine algebraic group over F we also define a norm $\|\cdot\|_{[G]}$ on $[G] = G(F) \backslash G(\mathbb{A})$ by

$$\|x\|_{[G]} := \inf_{\gamma \in G(F)} \|\gamma x\|_G$$

for all $x \in [G]$.

Proposition A.1.1 *Let X and Y be algebraic varieties over F and let G be an affine algebraic group over F .*

- (i) *The function $x \mapsto \|x\|_X$ is locally bounded on $X(\mathbb{A})$.*
- (ii) *Let $f : X \rightarrow Y$ be a morphism of algebraic varieties. Then $f^* \|\cdot\|_Y < \|\cdot\|_X$. In particular we have $\|gg'\|_G < \|g\|_G \|g'\|_G$ and $\|g^{-1}\|_G \sim \|g\|_G$ for all $g, g' \in G(\mathbb{A}_{\overline{F}})$. If moreover f is a finite morphism (in particular if it is a closed embedding) then $f^* \|\cdot\|_Y \sim \|\cdot\|_X$.*

(iii) Let $f \in F[X]$ and let $X_f = D(f)$ be the principal open subset of X defined by the nonvanishing of f . Then, we have

$$\|x\|_{X_f} \sim \|x\|_X \|f(x)^{-1}\|_{\mathbb{A}^1}$$

for all $x \in X_f(\mathbb{A}_{\overline{F}})$.

(iv) Let $U \subset X$ be an open subset and assume that X is quasi-affine. Then, we have

$$\|x\|_U \sim \|x\|_X$$

for all $x \in U(\overline{F})$. More generally if $p : X \rightarrow Y$ is a regular map and Y is quasi-affine then for all open subset $V \subset Y$ we have

$$\|x\|_{p^{-1}(V)} \sim \|x\|_X$$

for all $x \in p^{-1}(V)(\mathbb{A}_{\overline{F}})$ such that $p(x) \in V(\overline{F})$.

(v) If X is quasi-affine then there exists $d > 0$ such that

$$\sum_{x \in X(F)} \|x\|_X^{-d}$$

converges.

(vi) Let $d_r g$ be a right Haar measure on $G(\mathbb{A})$. Then there exists $d > 0$ such that the two integrals

$$\int_{G(\mathbb{A})} \|g\|_G^{-d} d_r g, \quad \int_{[G]} \|x\|_{[G]}^{-d} dx$$

converge.

(vii) Assume that X carries a G -action and that we have a regular map $p : X \rightarrow Y$ making X into a G -torsor over Y . Fix a right Haar measure $d_r g$ on $G(\mathbb{A})$. Then for all $d > 0$ there exists $d' > 0$ such that

$$\int_{G(\mathbb{A})} \|gx\|_X^{-d'} d_r g \ll \|p(x)\|_Y^{-d}$$

for all $x \in X(\mathbb{A})$.

(viii) Assume that G is connected and reductive and let $\mathfrak{S} \subset G(\mathbb{A})$ be a Siegel domain (see [MW1, §I.2.1]). Then, we have

$$\|g\|_G \sim \|g\|_{[G]}$$

for all $g \in \mathfrak{S}$.

(ix) Let $H < G$ be a closed subgroup such that G/H is quasi-affine (this is the case if for example H is reductive or if there is no nontrivial morphism $H \rightarrow \mathbb{G}_m$). Then, we have

$$\|x\|_{[H]} \sim \|x\|_{[G]}$$

for all $x \in [H]$. In particular, by (vi) there exists $d > 0$ such that the integral

$$\int_{[H]} \|x\|_{[G]}^{-d} dx$$

converges.

Proof:

- (i) This follows from the fact that for all v the function $x \in X(F_v) \mapsto \|x\|_{X_v}$ is locally bounded and the fact that for almost all v we have $\|x_v\|_{X_v} = 1$ for all $x_v \in X(\mathcal{O}_v)$.
- (ii) It suffices to prove the following
 - For all place v , we have $f^*\|\cdot\|_{Y_v} < \|\cdot\|_{X_v}$ and if f is finite $\|\cdot\|_{X_v} < f^*\|\cdot\|_{Y_v}$;
 - There exists $d > 0$ such that for almost all place v , we have $f^*\|\cdot\|_{Y_v} \leq \|\cdot\|_{X_v}^d$ and if f is finite $\|\cdot\|_{X_v}^{1/d} \leq f^*\|\cdot\|_{Y_v}$.

Assume that the norms $(\|\cdot\|_{X_v})_v$ have been defined using the finite affine open covering $\mathcal{U} = (U_i)_{i \in I}$ of X and that the norms $(\|\cdot\|_{Y_v})_v$ have been defined using the finite affine open covering $\mathcal{V} = (V_j)_{j \in J}$ of Y . Up to refining \mathcal{U} , we may assume that for all $j \in J$ there exists a subset $I(j) \subset I$ such that $f^{-1}(V_j) = \bigcup_{i \in I(j)} U_i$. If moreover f is finite then for all $j \in J$, the open subset $f^{-1}(V_j)$ is affine so that we may assume that $\mathcal{U} = (f^{-1}(V_j))_{j \in J}$. This allows us to reduce to the case where both X and Y are affine in which case the statement can be proved much the same way as [Kott, Proposition 18.1(1)].

- (iii) Assume that the family of norms $(\|\cdot\|_{X_v})_v$ has been defined using the finite affine open covering $\mathcal{U} = (U_i)_{i \in I}$ of X . Set $U_{i,f} = U_i \cap X_f$ for all $i \in I$. Obviously, we may assume that the family of norms $(\|\cdot\|_{X_{f,v}})_v$ has been defined using the affine open covering $\mathcal{U}_f = (U_{i,f})_{i \in I}$ of X_f and that

$$\|x\|_{U_{i,f,v}} = \max(\|x\|_{U_{i,v}}, |f(x)|_v^{-1})$$

for all place v of F and all $x \in U_{i,f}(\overline{F}_v)$. Then we have

$$\sqrt{\|x\|_{U_{i,v}} \max(1, |f(x)|_v^{-1})} \leq \|x\|_{U_{i,f,v}} \leq \|x\|_{U_{i,v}} \max(1, |f(x)|_v^{-1})$$

for all place v of F and all $x \in U_{i,f}(\overline{F}_v)$. It follows that

$$\sqrt{\|x\|_{X_v} \max(1, |f(x)|_v^{-1})} \leq \|x\|_{X_{f,v}} \leq \|x\|_{X_v} \max(1, |f(x)|_v^{-1})$$

for all place v of F and all $x \in X_f(\overline{F}_v)$. Taking the product we get

$$\sqrt{\|x\|_X \|f(x)^{-1}\|_{\mathbb{A}^1}} \leq \|x\|_{X_f} \leq \|x\|_X \|f(x)^{-1}\|_{\mathbb{A}^1}$$

for all $x \in X_f(\mathbb{A}_{\overline{F}})$.

- (iv) We prove the second claim which is more general than the first. Let $p : X \rightarrow Y$ be a regular map, $V \subset Y$ an open subset and assume that Y is quasi-affine. It already follows from (ii) that we have

$$\|x\|_X < \|x\|_{p^{-1}(V)}$$

for all $x \in X(\mathbb{A}_{\overline{F}})$. Hence, it suffices to prove the reverse inequality for all $x \in p^{-1}(V)(\mathbb{A}_{\overline{F}})$ such that $p(x) \in V(\overline{F})$. As Y is quasi-affine, up to replacing V by a finite affine open cover we may assume that $V = Y_f$ for some $f \in F[Y]$. Still denoting by f its image in $F[X]$ we then have $p^{-1}(V) = X_f$. Then by (ii), (iii) and (1) we have

$$\|x\|_{X_f} \sim \|x\|_X \|f(x)^{-1}\|_{\mathbb{A}^1} = \|x\|_X \|f(x)\|_{\mathbb{A}^1} < \|x\|_X$$

for all $x \in X_f(\mathbb{A}_{\overline{F}})$ such that $f(x) \in \overline{F}^\times$. This implies the desired inequality.

- (v) As there exists an open embedding of X into an affine variety, by (iv) we immediately reduce to the case where X itself is affine. Then, we can find a closed embedding $\iota : X \hookrightarrow \mathbb{A}^n$ for some integer $n > 0$ and by (ii) we are reduced to prove the statement for $X = \mathbb{A}^n$ and then eventually for $X = \mathbb{A}^1$ in which case the statement is easily checked.
- (vi) Note that

$$\int_{[G]} \|x\|_{[G]}^{-d} dx \leq \int_{[G]} \sum_{\gamma \in G(F)} \|\gamma x\|_G^{-d} dx = \int_{G(\mathbb{A})} \|g\|_G^{-d} d_r g$$

for all $d > 0$. Hence it suffices to show that for d sufficiently large the last integral above is convergent. Assume that H is a closed distinguished subgroup of G isomorphic to \mathbb{G}_m or \mathbb{G}_a . We first show that if the statement is true for both H and G/H then it is true for G . For this we write

$$\int_{G(\mathbb{A})} \|g\|_G^{-d} d_r g = \int_{(G/H)(\mathbb{A})} \int_{H(\mathbb{A})} \|\dot{g}h\|_G^{-d} d_r h d_r \dot{g}$$

for all $d > 0$ and where $d_r h$, $d_r \dot{g}$ are suitable right Haar measures on $H(\mathbb{A})$ and $(G/H)(\mathbb{A})$ respectively (Note that $(G/H)(\mathbb{A}) = G(\mathbb{A})/H(\mathbb{A})$). Let $d_0, d_1 > 0$. Setting $d = d_0 + d_1$, we get

$$\int_{H(\mathbb{A})} \|gh\|_G^{-d} d_r h \leq \left(\inf_{h \in H(\mathbb{A})} \|gh\|_G \right)^{-d_0} \int_{H(\mathbb{A})} \|gh\|_G^{-d_1} d_r h$$

for all $g \in G(\mathbb{A})$. By (ii), there exists $c > 0$ such that $\|h\|_H \ll \|gh\|_G^c \|g\|_G^c$ for all $(h, g) \in H(\mathbb{A}) \times G(\mathbb{A})$. Hence,

$$\int_{H(\mathbb{A})} \|gh\|_G^{-d} d_r h \ll \left(\inf_{h \in H(\mathbb{A})} \|gh\|_G \right)^{-d_0} \|g\|_G^{d_1} \int_{H(\mathbb{A})} \|h\|_H^{-d_1/c} d_r h$$

for all $g \in G(\mathbb{A})$. As the left hand side above is, as a function of g , invariant by right translation by $H(\mathbb{A})$ we also get

$$\int_{H(\mathbb{A})} \|gh\|_G^{-d} d_r h \ll \left(\inf_{h \in H(\mathbb{A})} \|gh\|_G \right)^{d_1 - d_0} \int_{H(\mathbb{A})} \|h\|_H^{-d_1/c} d_r h$$

for all $g \in G(\mathbb{A})$. By assumption for d_1 sufficiently large the last integral above is convergent. Thus, it only remains to show that for $d' > 0$ sufficiently large the integral

$$\int_{(G/H)(\mathbb{A})} \left(\inf_{h \in H(\mathbb{A})} \|\dot{g}h\|_G \right)^{-d'} d_r \dot{g}$$

converges. By (ii), we have $\|\dot{g}\|_{G/H} < \inf_{h \in H(\mathbb{A})} \|\dot{g}h\|_G$ for all $\dot{g} \in G(\mathbb{A})/H(\mathbb{A})$. Consequently, the convergence of the last integral above for d' sufficiently large follows from the assumption on G/H .

Let P_0 be a minimal parabolic subgroup of G over F . Then, by the Iwasawa decomposition there exists a compact subgroup $K \subset G(\mathbb{A})$ such that $G(\mathbb{A}) = P_0(\mathbb{A})K$. As K is compact, by (i) the norm $\|\cdot\|_G$ is bounded on K . Moreover, we have

$$\int_{G(\mathbb{A})} \|g\|_G^{-d} d_r g = \int_{P_0(\mathbb{A})} \int_K \|kp_0\|_G^{-d} dk d_r p_0$$

for suitable (right) Haar measures $d_r p_0$ and dk on $P_0(\mathbb{A})$ and K respectively. By (i) and (ii), it follows that we may assume $G = P_0$. Let $P_0 = M_0 N_0$ be a Levi decomposition. Then the Haar measure $d_r p_0$ decomposes as $d_r p_0 = dn_0 dm_0$ according to the decomposition $P_0(\mathbb{A}) = N_0(\mathbb{A})M_0(\mathbb{A})$. Moreover we have $\|n_0 m_0\|_{P_0} \sim \|m_0\|_{M_0} \|n_0\|_{N_0}$ for all $(m_0, n_0) \in M_0(\mathbb{A}) \times N_0(\mathbb{A})$. This allows us to reduce to the case where $G = M_0$ or $G = N_0$. If $G = N_0$ then it admits a composition series whose successive quotients are isomorphic to \mathbb{G}_a and we are reduced to the case $G = \mathbb{G}_a$ where the statement can be checked directly. Assume now that $G = M_0$ and denote by A_0 the maximal split torus in the center of G . Then A_0 is isomorphic to a product of \mathbb{G}_m and M_0/A_0 is anisotropic. Thus, we only need to treat the cases $G = \mathbb{G}_m$ and G anisotropic. Once again if $G = \mathbb{G}_m$ the statement can be checked directly. Now if G is anisotropic we write

$$(2) \quad \int_{G(\mathbb{A})} \|g\|_G^{-d} dg = \int_{G(F) \backslash G(\mathbb{A})} \sum_{\gamma \in G(F)} \|\gamma g\|_G^{-d} dg$$

By (i), (ii) and (v) if d is sufficiently large the function

$$g \in G(\mathbb{A}) \mapsto \sum_{\gamma \in G(F)} \|\gamma g\|_G^{-d}$$

is locally bounded. Moreover by [BHC] the quotient $G(F) \backslash G(\mathbb{A})$ is compact. The result then follows from (2).

(vii) Let $d > 0$. As p is a G -torsor and Y is separated, the action of G on X is free i.e. the regular map

$$\begin{aligned} G \times X &\rightarrow X \times X \\ (g, x) &\mapsto (gx, x) \end{aligned}$$

is a closed embedding. By (ii), it follows that there exists $c > 0$ such that $\|g\|_G \ll \|gx\|_X^c \|x\|_X^c$ for all $(g, x) \in G(\mathbb{A}) \times X(\mathbb{A})$. Let $d_0, d_1 > 0$. Using the same trick as in the first part of the proof of (vi), we show that for $d' = d_0 + d_1$ we have

$$\int_{G(\mathbb{A})} \|gx\|_X^{-d'} d_r g \ll \left(\inf_{g \in G(\mathbb{A})} \|gx\|_X \right)^{d_1 - d_0} \int_{G(\mathbb{A})} \|g\|_G^{-d_1/c} d_r g$$

for all $x \in X(\mathbb{A})$. By (vi), the last integral above is convergent for d_1 sufficiently large. Moreover, by (ii) we have $\|p(x)\|_Y < \inf_{g \in G(\mathbb{A})} \|gx\|_X$ for all $x \in X(\mathbb{A})$. Thus, the statement follows by choosing d_0 sufficiently large (depending on d_1).

- (viii) Let T_0 be a maximal split torus in $R_{F/\mathbb{Q}}G$. Then, up to conjugating \mathfrak{S} by an element of $G(F)$, there exists a compact subset $\Omega \subset G(\mathbb{A})$ such that

$$\mathfrak{S} \subseteq T_0(\mathbb{R})\Omega$$

Hence, by (i) and (ii) it is sufficient to show that

$$\|a\|_G \sim \|a\|_{[G]}$$

for all $a \in T_0(\mathbb{R})$. The inequality $\|a\|_{[G]} \leq \|a\|_G$ is obvious so that we only need to show that $\|a\|_G < \|a\|_{[G]}$ for all $a \in T_0(\mathbb{R})$. Let χ_1, \dots, χ_n be a basis of $X^*(T_0)$. Then we have

$$\|a\|_G \sim \max(|\chi_1(a)|, |\chi_1(a)|^{-1}, \dots, |\chi_n(a)|, |\chi_n(a)|^{-1})$$

for all $a \in T_0(\mathbb{R})$. Thus, it suffices to show that for all character $\chi \in X^*(T_0)$ we have $|\chi(a)| < \|a\|_{[G]}$ for all $a \in T_0(\mathbb{R})$. Let χ be such a character and let V be a rational representation of $R_{F/\mathbb{Q}}G$ containing a nonzero vector v_0 such that $a.v_0 = \chi(a)v_0$ for all $a \in T_0$. Let v_1, \dots, v_r be a basis of V . Set $V_{\mathbb{A}} = V \otimes_{\mathbb{Q}} \mathbb{A}$ and define a nonnegative function $|\cdot|_V$ on $V_{\mathbb{A}}$ by

$$|\lambda_1 v_1 + \dots + \lambda_r v_r|_V = \prod_v \max(|\lambda_{1,v}|, \dots, |\lambda_{r,v}|)$$

for all $\lambda_1, \dots, \lambda_r \in \mathbb{A}$. Note that there exist nonzero vectors $v \in V_{\mathbb{A}}$ such that $|v|_V = 0$ but that, however, if $v \in V_F = V \otimes_{\mathbb{Q}} F$ is nonzero then $|v|_V \geq 1$. We have $|v|_V < \|v\|_{V_F}$ for all $v \in V_{\mathbb{A}}$, where V_F is considered as an algebraic variety over F . Note that G acts on V_F via the natural embedding $G \hookrightarrow (R_{F/\mathbb{Q}}G)_F$. Hence, by (ii) we have

$$|\chi(a)|^d \leq |\chi(a)|^d |\gamma v_0|_V = |\gamma a v_0|_V < \|\gamma a v_0\|_{V_F} < \|\gamma a\|_G$$

for all $a \in T_0(\mathbb{R})$, $\gamma \in G(F)$ and where we have set $d = [F : \mathbb{Q}]$. Taking the infimum over γ yields the desired inequality.

- (ix) By (ii), the inequality $\|x\|_{[G]} < \|x\|_{[H]}$ is obvious so that we only need to show that $\|x\|_{[H]} < \|x\|_{[G]}$ for all $x \in [H]$. We will need the following fact (which is where the assumption G/H quasi-affine is crucial):

(3) There exists a (set-theoretic) section $s : (H \backslash G)(\overline{F}) \rightarrow G(\overline{F})$ such that

$$\|s(x)\|_G < \|x\|_{H \backslash G} \text{ for all } x \in (H \backslash G)(\overline{F}).$$

Proof of (3): Let $p : G \rightarrow H \backslash G$ be the natural surjection. Since $H \backslash G$ is quasi-affine, by (iv), it suffices to find an open covering $(U_i)_{i \in I}$ of $H \backslash G$ and sections $s_i :$

$U_i(\overline{F}) \rightarrow p^{-1}(U_i)(\overline{F})$ such that $\|s_i(x)\|_{p^{-1}(U_i)} < \|x\|_{U_i}$ for all $i \in I$ and all $x \in U_i(\overline{F})$. It is even sufficient to construct one non-empty open subset $U \subseteq H \backslash G$ and a section $s_U : U(\overline{F}) \rightarrow p^{-1}(U)(\overline{F})$ such that $\|s_U(x)\|_{p^{-1}(U)} < \|x\|_U$ for all $x \in U(\overline{F})$. Indeed, if such a pair (U, s_U) exists, we can find a finite number of translates $U_i := U\gamma_i$, $\gamma_i \in G(\overline{F})$, $i \in I$, covering $H \backslash G$ and then the sections $s_i : U_i(\overline{F}) \rightarrow p^{-1}(U_i)(\overline{F})$ given by $s_i(x) := s(x\gamma_i^{-1})\gamma_i$, for all $i \in I$ and $x \in U_i(\overline{F})$, satisfy the desired condition. As $p : G \rightarrow H \backslash G$ is a torsor for the étale topology, we can find a non-empty open subset $U \subseteq H \backslash G$ and a finite étale map $U' \rightarrow U$ such that $U' \times_U G$ is the trivial G -torsor over U' . In particular there exists a regular section $s_{U'} : U' \rightarrow U' \times_U G$. Let $s_0 : U(\overline{F}) \rightarrow U'(\overline{F})$ be any set-theoretic section. Then, by (ii) and since $U' \rightarrow U$ is finite, the section $s_U := pr_2 \circ s_{U'} \circ s_0 : U(\overline{F}) \rightarrow p^{-1}(U)(\overline{F})$, where pr_2 denotes the projection $U' \times_U G \rightarrow G$, satisfies the desired condition.

Let $s : (G/H)(\overline{F}) \rightarrow G(\overline{F})$ be a section as in (3). We have $\|\gamma\|_{G/H} < \|\gamma h\|_G$, for all $(\gamma, h) \in G(F) \times H(\mathbb{A})$ (by (ii)) and thus

$$\inf_{\gamma' \in H(\overline{F})} \|\gamma' h\|_H \leq \|s(\gamma)^{-1} \gamma h\|_H < \|s(\gamma)\|_G \|\gamma h\|_G < \|\gamma\|_{G/H} \|\gamma h\|_G < \|\gamma h\|_G$$

for all $(\gamma, h) \in G(F) \times H(\mathbb{A})$. Taking the infimum over γ it follows that

$$\inf_{\gamma' \in H(\overline{F})} \|\gamma' h\|_H < \inf_{\gamma \in G(F)} \|\gamma h\|_G = \|h\|_{[G]}$$

for all $h \in H(\mathbb{A})$. Hence, it suffices to show

$$(4) \quad \|h\|_{[H]} < \inf_{\gamma \in H(\overline{F})} \|\gamma h\|_H$$

for all $h \in H(\mathbb{A})$. Denote by N_H the unipotent radical of H and let L_H be a Levi component of H (so that $H = L_H \times N_H$). As $[N_H]$ is compact we are easily infer from (i) and (ii) that

$$\|\ell n\|_{[H]} \sim \|\ell\|_{[L_H]} \quad \text{and} \quad \inf_{\gamma \in H(\overline{F})} \|\gamma \ell n\|_H \sim \inf_{\gamma_L \in L_H(\overline{F})} \|\gamma_L \ell\|_{L_H}$$

for all $\ell \in L_H(\mathbb{A})$ and all $n \in N_H(\mathbb{A})$. We are thus reduced to prove (4) in the case where H is reductive. Denote by H^0 the connected component of the identity in H . Since $H(F)/H^0(F)$, $H(\overline{F})/H^0(\overline{F})$ are finite and $H(\mathbb{A})/H^0(\mathbb{A})$ is compact we may assume that $H = H^0$. Let T_0 be a maximal split torus of $R_{F/\mathbb{Q}}H$ and let $\chi \in X^*(T_0)$. By (viii), it is sufficient to show that

$$(5) \quad |\chi(a)| < \|\gamma a\|_H$$

for all $a \in T_0(\mathbb{R})$ and all $\gamma \in H(\overline{F})$. Let V be a rational representation of $R_{F/\mathbb{Q}}H$ containing a nonzero vector v_0 such that $a.v_0 = \chi(a)v_0$ for all $a \in T_0$. Fix a basis v_1, \dots, v_n of V and let $|\cdot|_V$ be the nonnegative function on $V_{\mathbb{A}_{\overline{F}}} = V \otimes_{\mathbb{Q}} \mathbb{A}_{\overline{F}}$ defined by

$$|\lambda_1 v_1 + \dots + \lambda_r v_r|_V = \prod_v \left(\prod_{w|v} \max(|\lambda_{1,w}|_v, \dots, |\lambda_{r,w}|_v)^{[F'_w:F_v]} \right)^{1/[F':F]}$$

for all $\lambda_1, \dots, \lambda_r \in \mathbb{A}_{F'}$, F'/F a finite extension. Note that $|v|_V \geq 1$ for all nonzero vector $v \in V_{\overline{F}} = V \otimes_{\mathbb{Q}} \overline{F}$ and $|v|_V < \|v\|_{V_F}$ for all $v \in V_{\mathbb{A}_{\overline{F}}}$. It follows that

$$|\chi(a)|^d \leq |\chi(a)|^d |\gamma v_0|_V = |\gamma a v_0|_V < \|\gamma a v_0\|_{V_F} < \|\gamma a\|_H$$

for all $(a, \gamma) \in T_0(\mathbb{R}) \times H(\overline{F})$. Taking the infimum over γ we get (5) and this ends the proof of (ix). ■

Let G be a connected reductive group over F . Fix a maximal compact subgroup K_∞ of $G(\mathbb{A}_\infty)$ and a Haar measure dg on $G(\mathbb{A})$. We will denote by $\mathcal{U}(\mathfrak{g}_\infty)$ the universal enveloping algebra of (the complexification of) the Lie algebra of $G(\mathbb{A}_\infty)$. For simplicity we will assume that the split center of G is trivial. Denote by $\mathcal{A}([G])$ the space of automorphic functions on $[G]$ by which we mean functions $\phi : [G] \rightarrow \mathbb{C}$ satisfying the following conditions

- ϕ is smooth: there exists a compact-open subgroup K of $G(\mathbb{A}_f)$ such that ϕ is right K -invariant and for all $g_f \in G(\mathbb{A}_f)$ the function $g_\infty \in G(\mathbb{A}_\infty) \mapsto \phi(g_\infty g_f)$ is C^∞ ;
- ϕ is uniformly of moderate growth: there exists $d > 0$ such that for all $u \in \mathcal{U}(\mathfrak{g}_\infty)$ we have $|(R(u)\phi)(g)| \ll \|g\|_{[G]}^d$ for all $g \in G(\mathbb{A})$.

Note that we don't impose any condition of K_∞ -finiteness or \mathfrak{z}_∞ -finiteness (where \mathfrak{z}_∞ denotes the center of $\mathcal{U}(\mathfrak{g}_\infty)$). The space $\mathcal{A}([G])$ is naturally equipped with a LF topology (see [Beu1, appendix A] for basic facts about LF vector spaces). As usual, we define $\mathcal{A}_{cusp}([G])$ to be the subspace of cuspidal functions in the following sense: $\phi \in \mathcal{A}([G])$ is cuspidal if for all proper parabolic subgroup $P = MN$ of G we have

$$\int_{[N]} \phi(n g) dn = 0$$

for all $g \in G(\mathbb{A})$. The space $\mathcal{A}_{cusp}([G])$ is a closed subspace of $\mathcal{A}([G])$ from which it inherits a LF topology and moreover every cuspidal function $\phi \in \mathcal{A}_{cusp}([G])$ is of rapid decay in the following sense: for all $u \in \mathcal{U}(\mathfrak{g}_\infty)$ and for all $d > 0$ we have

$$|(R(u)\phi)(g)| \ll \|g\|_{[G]}^{-d}$$

for all $g \in [G]$ (see [MW1, Corollary I.2.12]). By the open mapping theorem, for all compact-open subgroup K of $G(\mathbb{A}_f)$ the topology on $\mathcal{A}_{cusp}([G])^K$ is also induced by the family of seminorms

$$\|\phi\|_{d,u} = \sup_{g \in [G]} |(R(u)\phi)(g)| \|g\|_{[G]}^d, \quad d > 0, u \in \mathcal{U}(\mathfrak{g}_\infty)$$

There is another natural family of seminorms inducing the given topology on $\mathcal{A}_{cusp}([G])^K$. Let $C_G \in \mathcal{U}(\mathfrak{g}_\infty)$ and $C_K \in \mathcal{U}(\mathfrak{k}_\infty)$ denote the Casimir elements of $G(\mathbb{A}_\infty)$ and K_∞ respectively and set $\Delta = C_G^2 + C_K^2$. Then the family of Sobolev seminorms

$$\|\phi\|_k = \|R(1 + \Delta)^k \phi\|_{L^2([G])}, \quad k \geq 0, \phi \in \mathcal{A}_{cusp}([G])$$

where $\|\cdot\|_{L^2([G])}$ denotes the L^2 -norm on $L^2([G])$, induce on $\mathcal{A}_{cusp}([G])^K$ its LF topology (this follows essentially from strong approximation together with the Sobolev lemma). We will denote $L_{cusp}^2([G])$ the completion of $\mathcal{A}_{cusp}([G])$ in $L^2([G])$. It is a unitary representation of $G(\mathbb{A})$ which decomposes discretely.

Let now $f \in \mathcal{S}(G(\mathbb{A}))$ be a Schwartz function on $G(\mathbb{A})$. We denote as usual by

$$K_f(x, y) = \sum_{\gamma \in G(F)} f(x^{-1}\gamma y), \quad x, y \in [G]$$

the automorphic kernel of f . Note that the sum is absolutely convergent by Proposition A.1.1. Let $\pi \subset \mathcal{A}_{cusp}([G])$ be a cuspidal automorphic representation and let \mathcal{B}_π be an orthonormal basis of (the completion of) π for the L^2 scalar product. We define

$$K_{f,\pi}(x, y) = \sum_{\phi \in \mathcal{B}_\pi} (R(f)\phi)(x) \overline{\phi(y)}, \quad x, y \in [G]$$

Then $K_{f,\pi}$ is the orthogonal projection of K_f , seen as a function in x , onto π or, what amounts to the same, the orthogonal projection of K_f , seen as a function of y , onto $\overline{\pi}$. Finally, letting $\mathcal{B} \subset \mathcal{A}_{cusp}([G])$ be an orthonormal basis of $L_{cusp}^2([G])$, we set

$$K_{f,cusp}(x, y) = \sum_{\phi \in \mathcal{B}} (R(f)\phi)(x) \overline{\phi(y)}, \quad x, y \in [G]$$

Note that

$$K_{f,cusp} = \sum_{\pi} K_{f,\pi}$$

where the sum is over a complete family of orthogonal cuspidal automorphic representations $\pi \subset \mathcal{A}_{cusp}([G])$ (all of them if there is multiplicity one).

Proposition A.1.2 *Let $H_1, H_2 \subset G$ be closed algebraic subgroups such that the quotients G/H_1 and G/H_2 are quasi-affine. Then the integral*

$$\int_{[H_1]} \int_{[H_2]} \sum_{\pi} |K_{f,\pi}(h_1, h_2)| dh_1 dh_2$$

the sum running over a complete family of orthogonal cuspidal automorphic representations, converges. We even have the stronger following result: let K_0 be a compact-open subgroup of $G(\mathbb{A}_f)$ such that f is right K_0 -invariant and let $\mathcal{B} \subset \mathcal{A}_{cusp}([G])^{K_0}$ be an orthonormal basis of $L^2_{cusp}([G])^{K_0}$ consisting of functions which are C_K and C_G eigenvectors, then the integral

$$\int_{[H_1]} \int_{[H_2]} \sum_{\phi \in \mathcal{B}} |(R(f)\phi)(h_1)| |\phi(h_2)| dh_1 dh_2$$

converges.

Proof: The second statement is obviously stronger than the first since for every cuspidal automorphic representation π we can find an orthonormal basis of π^{K_0} consisting of C_K - and C_G -eigenvectors. Let \mathcal{B} be an orthonormal basis of $L^2_{cusp}([G])^{K_0}$ as in the proposition. By Proposition A.1.1(ix) it suffices to prove that for all $d > 0$ we have

$$(7) \quad \sum_{\phi \in \mathcal{B}} |(R(f)\phi)(x)| |\phi(y)| \ll \|x\|_{[G]}^{-d} \|y\|_{[G]}^{-d}$$

for all $x, y \in [G]$. Let $d > 0$. Since the family of norms $(\|\cdot\|_k)_k$ generates the topology on $\mathcal{A}_{cusp}([G])^{K_0}$, there exists $k > 0$ such that

$$|\phi(x)| \ll \|\phi\|_k \|x\|_{[G]}^{-d}$$

for all $\phi \in \mathcal{A}_{cusp}([G])^{K_0}$ and all $x \in [G]$.

For all $\phi \in \mathcal{B}$, let us denote by $\lambda_K(\phi), \lambda_G(\phi) \in \mathbb{R}$ the eigenvalues of C_K and C_G acting on ϕ . Let N be a positive integer. For all $\phi \in \mathcal{B}$, we have

$$R(f)\phi = (1 + \lambda_G(\phi)^2 + \lambda_K(\phi)^2)^{-N} R(f^{(N)})\phi$$

where $f^{(N)} = (1 + \Delta)^N f$. Hence, we have

$$\begin{aligned} \sum_{\phi \in \mathcal{B}} |R(f)\phi(x)| |\phi(y)| &= \sum_{\phi \in \mathcal{B}} (1 + \lambda_G(\phi)^2 + \lambda_K(\phi)^2)^{-N} |R(f^{(N)})\phi(x)| |\phi(y)| \\ &\ll \|x\|_{[G]}^{-d} \|y\|_{[G]}^{-d} \sum_{\phi \in \mathcal{B}} (1 + \lambda_G(\phi)^2 + \lambda_K(\phi)^2)^{-N} \|R(f^{(N)})\phi\|_k \|\phi\|_k \\ &= \|x\|_{[G]}^{-d} \|y\|_{[G]}^{-d} \sum_{\phi \in \mathcal{B}} (1 + \lambda_G(\phi)^2 + \lambda_K(\phi)^2)^{k-N} \|R(f^{(N+k)})\phi\|_{L^2} \|\phi\|_{L^2} \\ &\leq \|x\|_{[G]}^{-d} \|y\|_{[G]}^{-d} \|f^{(N+k)}\|_{L^1} \sum_{\phi \in \mathcal{B}} (1 + \lambda_G(\phi)^2 + \lambda_K(\phi)^2)^{k-N} \end{aligned}$$

for all $x, y \in [G]$ and where $\|\cdot\|_{L^1}$ denotes the L^1 -norm on $L^1([G])$. By [Mu], for $N \gg 1$ the last sum above converges. This proves (7) and ends the proof of the proposition. ■

Remark A.1.3 • Fix $f^\infty \in \mathcal{S}(G(\mathbb{A}^\infty))$. Then, the proof of the proposition actually shows that for all $d > 0$ there exists a continuous seminorm ν_d on $\mathcal{S}(G(F_\infty))$ so that

$$(8) \quad |K_{f_\infty \otimes f^\infty, cusp}(x, y)| \leq \nu_d(f_\infty) \|x\|_{[G]}^{-d} \|y\|_{[G]}^{-d}$$

for all $f_\infty \in \mathcal{S}(G(F_\infty))$ and all $x, y \in [G]$.

- We can prove the first part of the proposition directly by using the Selberg trick. Indeed, it suffices to show that the series

$$\sum_{\pi} K_{f, \pi}$$

converges absolutely in $\mathcal{A}_{cusp}([G \times G])$ or, what amounts to the same, that it converges absolutely in $\mathcal{A}([G \times G])$. To prove this, we only need to show that the sum

$$\sum_{\pi} |K_{f, \pi}(x, y)|$$

converges absolutely for all $x, y \in [G]$ and is bounded uniformly in x, y . By a theorem of Dixmier-Malliavin ([DM]), we may write f as a finite sum of convolutions $f_{1,i} \star f_{2,i}$, $f_{1,i}, f_{2,i} \in \mathcal{S}(G(\mathbb{A}))$, $i = 1, \dots, k$. By the Cauchy-Schwarz inequality, we have

$$|K_{f, \pi}(x, y)| \leq \sum_{i=1}^k K_{h_{2,i}, \pi}(x, x)^{1/2} K_{h_{1,i}, \pi}(y, y)^{1/2}$$

for all π , all $x, y \in [G]$ and where we have set $h_{j,i} = f_{j,i}^* \star f_{j,i}$ where by definition $f_{j,i}^*(g) = \overline{f_{j,i}(g^{-1})}$. Thus, by another application of Cauchy-Schwarz, we get

$$\sum_{\pi} |K_{f, \pi}(x, y)| \leq \sum_{i=1}^k K_{h_{2,i}, cusp}(x, x)^{1/2} K_{h_{1,i}, cusp}(y, y)^{1/2}$$

for all $x, y \in [G]$ and the right hand side is uniformly bounded (even of rapid decay).

A.2 Relative trace formulae

We now return to the situation considered in section 3. We will use the same notations and normalization of measures as there (in particular our global Haar measures are Tamagawa measures). We define the following 'bases' (geometric quotients): $\mathcal{B} := H \backslash G / H$ and $\mathcal{B}' := H'_1 \backslash G' / H'_2$. These are affine varieties (as H , H'_1 and H'_2 are reductive). We set $p : G \rightarrow \mathcal{B}$ and $p' : G' \rightarrow \mathcal{B}'$ for the natural projections and p_F, p'_F for the corresponding maps at the level of F -points. For all test function $f \in \mathcal{S}(G(\mathbb{A}))$, $f' \in \mathcal{S}(G'(\mathbb{A}))$, $\delta \in \mathcal{B}(F)$, $\gamma \in \mathcal{B}'(F)$, we set

$$K_{f,\delta}(x, y) := \sum_{\underline{\delta} \in p_F^{-1}(\delta)} f(x^{-1} \underline{\delta} y), \quad x, y \in G(\mathbb{A})$$

$$K_{f',\gamma}(x, y) := \sum_{\underline{\gamma} \in (p'_F)^{-1}(\gamma)} f'(x^{-1} \underline{\gamma} y), \quad x, y \in G'(\mathbb{A})$$

Note that these sums are absolutely convergent by Proposition A.1.1 (v) and that

$$K_f(x, y) = \sum_{\delta \in \mathcal{B}(F)} K_{f,\delta}(x, y)$$

$$K_{f'}(x, y) = \sum_{\gamma \in \mathcal{B}'(F)} K_{f',\gamma}(x, y)$$

Whenever convergent, we define the following 'global orbital integrals'

$$O(\delta, f) := \int_{[H]} \int_{[H]} K_{f,\delta}(h_1, h_2) dh_1 dh_2$$

$$O(\gamma, f') := \int_{[H'_1]} \int_{[H'_2]} K_{f',\gamma}(h_1, h_2) \eta(h_2) dh_2 dh_1$$

and the following two expressions

$$J(f) = \int_{[H]} \int_{[H]} K_f(h_1, h_2) dh_1 dh_2$$

$$I(f') = \int_{[H'_1]} \int_{[H'_2]} K_{f'}(h_1, h_2) \eta(h_2) dh_2 dh_1$$

Proposition A.2.1 (i) *Assume that $f \in \mathcal{S}(G(\mathbb{A}))$ is a nice function (see §3.5). Then the expressions defining $J(f)$ and $O(\delta, f)$, $\delta \in \mathcal{B}(F)$, are absolutely convergent and we have the equalities*

$$\sum_{\delta \in \mathcal{B}(F)} O(\delta, f) = J(f) = \sum_{\pi} J_{\pi}(f)$$

where the left sum is absolutely convergent and the right sum is over the set of cuspidal automorphic representations π of $G(\mathbb{A})$.

(ii) Assume that $f' \in \mathcal{S}(G'(\mathbb{A}))$ is a nice function (see §3.5). Then the expressions defining $I(f')$ and $O(\gamma, f')$, $\gamma \in \mathcal{B}'(F)$, are absolutely convergent and we have the equalities

$$\sum_{\gamma \in \mathcal{B}'(F)} O(\gamma, f') = I(f') = \sum_{\Pi} 2^{-2} L(1, \eta_{E/F})^{-2} I_{\Pi}(f')$$

where the left sum is absolutely convergent and the right sum is over the set of cuspidal automorphic representations Π of $G'(\mathbb{A})$ whose central character is trivial on $Z_{H'_2}(\mathbb{A})$.

Proof: We only prove (ii) the proof of (i) being similar.

Set $\tilde{G}' = G'/Z_{H'_2}$ and define $\tilde{f}' \in \mathcal{S}(\tilde{G}'(\mathbb{A}))$ by

$$\tilde{f}'(\tilde{g}) = \int_{Z'_{H'_2}(\mathbb{A})} f'(z\tilde{g}) dz$$

Then we have, at least formally,

$$I(f') = \int_{[H'_1]} \int_{[H'_2/Z_{H'_2}]} K_{\tilde{f}'}(h_1, h_2) \eta(h_2) dh_2 dh_1$$

As f' is a nice function we have $K_{\tilde{f}'} = K_{\tilde{f}', \text{cusp}}$. Thus, by Proposition A.1.2, it follows that the expression defining $I(f')$ is absolutely convergent and that

$$I(f') = \sum_{\Pi} \int_{[H'_1]} \int_{[H'_2/Z_{H'_2}]} K_{\tilde{f}', \Pi}(h_1, h_2) \eta(h_2) dh_2 dh_1$$

where the sum is over the set of all cuspidal automorphic representations Π of $G'(\mathbb{A})$ with a central character trivial on $Z_{H'_2}(\mathbb{A})$. We would like to identify the term indexed by Π above with the global spherical character $I_{\Pi}(f')$. However, we don't have equality on the nose because the scalar products used to define $I_{\Pi}(f')$ and $K_{\tilde{f}', \Pi}$ are not the same. More precisely, $I_{\Pi}(f')$ is defined using the Petersson scalar product $(\cdot, \cdot)_{\text{Pet}}$ of section 3 whereas in the definition of $K_{\tilde{f}', \Pi}$ we have used the scalar product

$$\begin{aligned} (\phi, \phi')_{L^2([\tilde{G}'])} &= \int_{[\tilde{G}']} \phi(\tilde{g}) \overline{\phi'(\tilde{g})} d\tilde{g} \\ &= \text{vol}(Z_{G'}(F)Z_{H'_2}(\mathbb{A}) \backslash Z_{G'}(\mathbb{A})) (\phi, \phi')_{\text{Pet}} \end{aligned}$$

Thus, we get

$$\int_{[H'_1]} \int_{[H'_2/Z_{H'_2}]} K_{\tilde{f}', \Pi}(h_1, h_2) \eta(h_2) dh_2 dh_1 = \text{vol}(Z_{G'}(F)Z_{H'_2}(\mathbb{A}) \backslash Z_{G'}(\mathbb{A}))^{-1} I_{\Pi}(f')$$

By 2.5(1), we have

$$\text{vol} \left(Z_{G'}(F)Z_{H'_2}(\mathbb{A}) \backslash Z_{G'}(\mathbb{A}) \right) = \text{vol} \left(E^\times \mathbb{A}^\times \backslash \mathbb{A}_E^\times \right)^2 = 2^2 L(1, \eta_{E/F})^2$$

and the second equality of (ii) follows.

The first equality follows from standard formal manipulations. To justify these manipulations, we need to establish that the following expression is absolutely convergent (as a triple integral)

$$\int_{[H'_1]} \int_{[H'_2]} \sum_{\gamma \in \mathcal{B}'(F)} K_{f', \gamma}(h_1, h_2) \eta(h_2) dh_2 dh_1 = \int_{[H'_1]} \int_{[H'_2/Z_{H'_2}]} \sum_{\gamma \in \mathcal{B}'(F)} K_{\tilde{f}', \gamma}(h_1, h_2) \eta(h_2) dh_2 dh_1$$

where for all $\gamma \in \mathcal{B}'(F)$, $K_{\tilde{f}', \gamma}$ is defined the same way as $K_{f', \gamma}$. For this, by Proposition A.1.1 (v) and (ix), it suffices to show that for all $d > 0$ we have an inequality

$$(1) \quad \left| K_{\tilde{f}', \gamma}(h_1, h_2) \right| \ll \|\gamma\|_{\mathcal{B}'}^{-d} \|h_1\|_{[\tilde{G}']}^{-d} \|h_2\|_{[\tilde{G}']}^{-d}$$

for all $\gamma \in \mathcal{B}'(F)$ and all $h_1 \in [H'_1]$, $h_2 \in [H'_2/Z_{H'_2}]$. Since f' is factorizable, we may write $f' = f'_\infty \otimes f'^\infty$ with $f'_\infty \in \mathcal{S}(G'(F_\infty))$ and $f'^\infty \in \mathcal{S}(G'(\mathbb{A}_F^\infty))$. Let $\mathcal{B}' \hookrightarrow V = \mathbb{A}^r$ be a closed embedding of \mathcal{B}' into some affine space. For all $\varphi \in C_c^\infty(V(F_\infty))$, we define $f'_\varphi := (\varphi f'_\infty) \otimes f'^\infty$ where we identify φ with a function on $G'(F_\infty)$ by composition with the projection $G'(F_\infty) \rightarrow \mathcal{B}'(F_\infty)$ and the embedding $\mathcal{B}'(F_\infty) \hookrightarrow V(F_\infty)$. Then, f'_φ is again a nice function so that $K_{\tilde{f}'_\varphi} = K_{\tilde{f}'_\varphi, \text{cusp}}$ and, by A.1(8), for all $d > 0$ there exists a continuous semi-norm ν_d on $\mathcal{S}(G'(F_\infty))$ such that

$$(2) \quad \left| K_{\tilde{f}'_\varphi}(x, y) \right| \leq \nu_d(\varphi f'_\infty) \|x\|_{[\tilde{G}']}^{-d} \|y\|_{[\tilde{G}']}^{-d}$$

for all $\varphi \in C_c^\infty(V(F_\infty))$ and all $x, y \in [G]$. Let Γ denote the intersection of $\mathcal{B}'(F)$ with the projection of the support of f'^∞ (a compact subset of $\mathcal{B}'(\mathbb{A}_F^\infty)$). Note that for all $\varphi \in C_c^\infty(V(F_\infty))$, all $\gamma \in \mathcal{B}'(F)$ and all $(h_1, h_2) \in H'_1(\mathbb{A}_F) \times H'_2(\mathbb{A}_F)$, we have $K_{\tilde{f}'_\varphi, \gamma}(h_1, h_2) = \varphi(\gamma) K_{\tilde{f}', \gamma}(h_1, h_2)$ and $K_{\tilde{f}'_\varphi, \gamma}(h_1, h_2) = 0$ if $\gamma \notin \Gamma$. Hence, by (2), to show (1) it suffices to construct a family $(\varphi_\gamma)_{\gamma \in \Gamma}$ of functions in $C_c^\infty(V(F_\infty))$ satisfying the following two conditions

- For all $\gamma \in \Gamma$, we have $\text{Supp}(\varphi_\gamma) \cap \Gamma = \{\gamma\}$ and $\varphi_\gamma(\gamma) = 1$;
- For all $d > 0$, the function $\gamma \in \Gamma \mapsto \|\gamma\|_{\mathcal{B}'}^d \nu_d(\varphi_\gamma f'_\infty)$ is bounded.

There exists a lattice $L \subset V(F_\infty)$ containing Γ . Fix a function $\varphi_0 \in C_c^\infty(V(F_\infty))$ with the property that $\text{Supp}(\varphi_0) \cap L = \{0\}$ and $\varphi_0(0) = 1$. For all $v \in L$, define $\varphi_v \in C_c^\infty(V(F_\infty))$ by $\varphi_v(x) = \varphi_0(x - v)$. We claim that the family $(\varphi_\gamma)_{\gamma \in \Gamma}$ satisfies the two conditions above.

Indeed, the first condition is clear and for all $d > 0$, there exist $k > 0$ and two finite families $(u_i)_{i \in I}$ and $(v_i)_{i \in I}$ of elements of $\mathcal{U}(\mathfrak{g}'_\infty)$ such that

$$\nu_d(\varphi f'_\infty) \leq \sup_{i \in I} \sup_{g \in G'(F_\infty)} |(R(u_i)\varphi)(g)(R(v_i)f'_\infty)(g)| \|g\|_{G'}^k$$

for all $\varphi \in C_c^\infty(V(F_\infty))$. Then, for all $\ell > 0$ we have

$$\nu_d(\varphi f'_\infty) \leq \left(\sup_{i \in I} \sup_{g \in G'(F_\infty)} |(R(u_i)\varphi)(g)| \|g\|_{G'}^{-\ell} \right) \left(\sup_{i \in I} \sup_{g \in G'(F_\infty)} |(R(v_i)f'_\infty)(g)| \|g\|_{G'}^{k+\ell} \right)$$

for all $\varphi \in C_c^\infty(V(F_\infty))$. Thus, it suffices to show that for all $u \in \mathcal{U}(\mathfrak{g}'_\infty)$ there exists $\ell > 0$ so that the function

$$\gamma \in \Gamma \mapsto \|\gamma\|_{\mathcal{B}'}^d \sup_{g \in G'(F_\infty)} |(R(u)\varphi_\gamma)(g)| \|g\|_{G'}^{-\ell}$$

is bounded. Fix $u \in \mathcal{U}(\mathfrak{g}'_\infty)$. Then, we can find a finite family $(r_j)_{j \in J}$ of regular functions in $\mathbb{C}[R_{F/\mathbb{Q}}G']$ and a finite family $(X_j)_{j \in J}$ of elements of the symmetric algebra of $V(F_\infty)$ such that, denoting by $\partial(X_j)$ the corresponding constant coefficients differential operators on $V(F_\infty)$, we have

$$R(u)\varphi = \sum_{j \in J} r_j \partial(X_j)\varphi$$

for all $\varphi \in C_c^\infty(V(F_\infty))$. Since for each $j \in J$ the absolute value of r_j is bounded by a constant times a power of the norm $\|\cdot\|_{G'}$, we are reduced to prove that for all X in the symmetric algebra of $V(F_\infty)$ there exists $\ell > 0$ such that

$$\|\gamma\|_{\mathcal{B}'}^d \sup_{g \in G'(F_\infty)} |\partial(X)\varphi_\gamma(g)| \|g\|_{G'}^{-\ell} \ll 1$$

for all $\gamma \in \Gamma$. By our choice of the functions $(\varphi_\gamma)_{\gamma \in \Gamma}$ and Proposition A.1.1(ii) we have

$$\|\gamma\|_{\mathcal{B}'} < \|g\|_{G'}$$

for all $\gamma \in \Gamma$ and all $g \in G'(F_\infty)$ with $\varphi_\gamma(g) \neq 0$. Hence, we just need to show that for all X in the symmetric algebra of $V(F_\infty)$ the function

$$\gamma \in \Gamma \mapsto \sup_{g \in G'(F_\infty)} |\partial(X)\varphi_\gamma(g)| = \sup_{v \in \mathcal{B}'(F_\infty)} |\partial(X)\varphi_\gamma(v)|$$

is bounded. But this is obvious by the way we have defined the functions $(\varphi_\gamma)_{\gamma \in \Gamma}$. ■

A.3 Proof of Theorem 3.5.1

Let $f \in \mathcal{S}(G(\mathbb{A}))$ and $f' \in \mathcal{S}(G'(\mathbb{A}))$ be nice functions and assume that there exists a tuple $(f^{W'})_{W'}$, $f^{W'} \in \mathcal{S}(G^{W'}(\mathbb{A}))$, matching f' and such that $f^W = f$. By Theorem 3.4.4, we may assume that $f^{W'} = 0$ for almost all W' . There are natural isomorphisms $H^{W'} \backslash G^{W'} / H^{W'} \simeq \mathcal{B} \simeq \mathcal{B}'$ for all W' (see §3.4). In order to compare the trace formulas of Proposition A.2.1, we need to know that for all $\delta \in \mathcal{B}(F)$ and all $\gamma \in \mathcal{B}'(F)$ corresponding to each other via the previous bijection we have

$$(1) \quad \sum_{W'} O(\delta, f^{W'}) = O(\gamma, f')$$

Note that for γ regular semi-simple (i.e. such that the fiber over γ in $G'(F)$ consists of regular semi-simple elements), this is a direct consequence of the fact that $(f^{W'})_{W'}$ matches f' (in this case there is at most one nonzero contribution in the left sum). To treat the general case, we need to use recent results of Zydor and Chaudouard-Zydor. More precisely, in [Zy] Zydor has defined, for $\delta \in \mathcal{B}(F)$ and $\gamma \in \mathcal{B}'(F)$, certain distributions

$$g^{W'} \in \mathcal{S}(G^{W'}(\mathbb{A})) \mapsto O^Z(\delta, g^{W'})$$

$$g' \in \mathcal{S}(G'(\mathbb{A})) \mapsto O^Z(\gamma, g')$$

The definition of $O^Z(\gamma, \cdot)$ is (roughly) as follows. let A be the standard maximal split torus in GL_n and set $\mathfrak{a} := X_*(A) \otimes \mathbb{R}$. Then for $T \in \mathfrak{a}$, $g' \in \mathcal{S}(G'(\mathbb{A}))$ and $\gamma \in \mathcal{B}'(F)$, Zydor defines a certain 'truncated' kernel $K_{g', \gamma}^T$ on $[H'_1] \times [H'_2]$ (see [Zy] §5.5, note that the \mathfrak{o} of *loc. cit.* corresponds to our γ and that the function f of *loc. cit.* corresponds not to g' but rather to its descent \tilde{g}' to $S_{n+1}(\mathbb{A})$ as in §3.3) and he shows that for T in a certain cone the integral

$$O^{Z,T}(\gamma, g') := \int_{[H'_1] \times [H'_2]} K_{g', \gamma}^T(h_1, h_2) \eta(h_2) dh_2 dh_1$$

converges absolutely (see [Zy, Theorem 5.9]). The definition of $K_{g', \gamma}^T$ is as a sum the main term being $K_{g', \gamma}$ and the remaining terms depending only on g'_P for certain proper parabolic subgroup $P = MU$ of G' where $g'_P(x) := \int_{U(\mathbb{A})} g'(xu) du$. Since f' is a nice function, we have $f'_P = 0$ for all proper parabolic subgroup and thus $K_{f', \gamma}^T = K_{f', \gamma}$ and $O^{Z,T}(\gamma, f') = O(\gamma, f')$ for all T (remark that this also reproves the absolute convergence of $O(\gamma, f')$ of Proposition A.2.1). Finally, still for T in a certain cone, Zydor shows that the function $T \mapsto O^{Z,T}(\gamma, g')$ is an exponential-polynomial whose purely polynomial term is constant ([Zy, Theorem 5.9]) and he defines $O^Z(\gamma, g')$ to be this constant. Since $O^{Z,T}(\gamma, f') = O(\gamma, f')$ for all T , we also have $O^Z(\gamma, f') = O(\gamma, f')$. The definition of $O^Z(\delta, \cdot)$ is similar and since the functions $f^{W'}$ are nice the same argument shows that $O^Z(\delta, f^{W'}) = O(\delta, f^{W'})$ for all W' and all $\delta \in \mathcal{B}(F)$. Finally, the main result of [CZ] is that if $g' \in \mathcal{S}(G'(\mathbb{A}))$ match a tuple of functions $(g^{W'})_{W'}$, $g^{W'} \in \mathcal{S}(G^{W'}(\mathbb{A}))$, then we have

$$\sum_{W'} O^Z(\delta, g^{W'}) = O^Z(\gamma, g')$$

for all $\delta \in \mathcal{B}(F)$ and all $\gamma \in \mathcal{B}'(F)$ corresponding to each other (strictly speaking [CZ] only considers compactly supported functions but the proofs applies verbatim to Schwartz functions). Together with the previous equalities this shows (1).

Now (1) together with Proposition A.2.1 leads to the identity

$$(2) \quad \sum_{W'} \sum_{\pi_{W'}} J_{\pi_{W'}}(f^{W'}) = \sum_{\Pi} 2^{-2} L(1, \eta_{E/F})^{-2} I_{\Pi}(f')$$

where $\pi_{W'}$ runs over the set of all cuspidal automorphic representations of $G^{W'}(\mathbb{A})$ and Π runs over the set of all cuspidal automorphic representations of $G'(\mathbb{A})$ whose central character is trivial on $Z_{H'_2}(\mathbb{A})$. Fix a maximal compact subgroup $K^{W'} = \prod_v K_v^{W'}$ of $G^{W'}(\mathbb{A})$ for all W' and let Σ be the infinite set of places v of F which split in E and where π , f and f' are unramified. By Theorem 3.4.4, we may assume that for all W' and all $v \in \Sigma$ the function $f_v^{W'}$ is unramified (i.e. it equals $\text{vol}(K_v^{W'})^{-1} \mathbf{1}_{K_v^{W'}}$). Then, in the equality (2) only the $\pi_{W'}$ and the Π which are unramified at all places in Σ contribute. Define the Hecke algebra $\mathcal{H}_{G,\Sigma} = C_c(G(\mathbb{A}_{\Sigma})//K_{\Sigma})$ of compactly supported and K_{Σ} -biinvariant functions on $G(\mathbb{A}_{\Sigma})$. This is the restricted tensor product over $v \in \Sigma$ of the local Hecke algebras $\mathcal{H}_{G,v} = C_c(G(F_v)//K_v)$. We define similarly the Hecke algebra $\mathcal{H}_{G',\Sigma} = C_c(G'(\mathbb{A}_{\Sigma})//K'_{\Sigma})$ and the local Hecke algebras $\mathcal{H}_{G',v} = C_c(G'(F_v)//K'_v)$. Note that for all n -dimensional hermitian space W' over E we have an isomorphism $G(\mathbb{A}_{\Sigma}) \simeq G^{W'}(\mathbb{A}_{\Sigma})$ canonical up to conjugation which induces a canonical isomorphism $\mathcal{H}_{G,\Sigma} \simeq C_c(G^{W'}(\mathbb{A}_{\Sigma})//K_{\Sigma}^{W'})$. There is a base change homomorphism $\mathcal{H}_{G',\Sigma} \rightarrow \mathcal{H}_{G,\Sigma}$, $h \mapsto h^{bc}$ and for all $v \in \Sigma$, all W' and all $h_v \in \mathcal{H}_{G',v}$, $h_v \star f'_v$ and $h_v^{bc} \star f_v^{W'}$ match each other (see [Zh1, Proposition 2.5]). For all irreducible unitary representation Π of $G'(\mathbb{A})$ which is unramified at all places in Σ let us denote by $h \mapsto \widehat{h}(\Pi)$ the corresponding character of the Hecke algebra $\mathcal{H}_{G',\Sigma}$. Then, for all W' , all cuspidal automorphic representation $\pi_{W'}$ which is unramified at all places in Σ and all $h \in \mathcal{H}_{G',\Sigma}$ the element $h^{bc} \in \mathcal{H}_{G,\Sigma}$ acts on $\pi_{W'}^{K_{\Sigma}^{W'}}$ by $\widehat{h}(BC(\pi_{W'}))$. Let $h \in \mathcal{H}_{G',\Sigma}$. Since the functions $h \star f'$ and $(h^{bc} \star f^{W'})_{W'}$ are nice and match each other we can apply equality (2) to these functions to get

$$(3) \quad \sum_{W'} \sum_{\pi_{W'}} \widehat{h}(BC(\pi_{W'})) J_{\pi_{W'}}(f^{W'}) = \sum_{\Pi} 2^{-2} L(1, \eta_{E/F})^{-2} \widehat{h}(\Pi) I_{\Pi}(f')$$

Let $Irr_{unit,\Sigma}(G'(\mathbb{A}))$ be the set of all irreducible unitary representations of $G'(\mathbb{A})$ which are unramified at all places in Σ . Then the functions $\Pi \in Irr_{unit,\Sigma}(G'(\mathbb{A})) \mapsto \widehat{h}(\Pi)$, $h \in \mathcal{H}_{G',\Sigma}$, are bounded and we have $\widehat{h^*} = \overline{\widehat{h}}$ where $h^*(g) = \overline{h(g^{-1})}$. Hence by the Stone-Weierstrass theorem, from (3) we deduce

$$(4) \quad \sum_{W'} \sum_{\pi_{W'}} J_{\pi_{W'}}(f^{W'}) = \sum_{\Pi} 2^{-2} L(1, \eta_{E/F})^{-2} I_{\Pi}(f')$$

where this time $\pi_{W'}$ and Π run over the sets of cuspidal automorphic representations of $G^{W'}(\mathbb{A})$ and $G'(\mathbb{A})$ such that $BC(\pi_{W',v}) = \Pi_v = BC(\pi_v)$ for all $v \in \Sigma$. Recall the following *automorphic-Cebotarev-density* theorem due to Ramakrishnan ([Ra]):

Theorem A.3.1 (Ramakrishnan) *Let Π_1, Π_2 be two isobaric automorphic representations of $GL_d(\mathbb{A}_E)$ such that $\Pi_{1,v} \simeq \Pi_{2,v}$ for almost all places v of F that are split in E . Then, $\Pi_1 = \Pi_2$.*

As $BC(\pi_{W'})$ is always isobaric it follows from this theorem that the right hand side of (4) reduces to $2^{-2} L(1, \eta_{E/F})^{-2} I_{BC(\pi)}(f')$ and that if $\pi_{W'}$ contributes to the left hand side then $BC(\pi_{W'}) = BC(\pi)$. In particular, $\pi_{W'}$ and π belong to the same (global) Vogan L -packet. By the local Gan-Gross-Prasad conjecture (see §2.4), and since by assumption π is tempered at all archimedean places, we know that there is at most one abstractly $H^{W'}$ -distinguished representation in this L -packet. By assumption, π is such a representation. Hence, the left hand side of (4) reduces to $J_{\pi}(f)$ and this ends the proof of Theorem 3.5.1. ■

Bibliography

- [AMS] A.-M. Aubert, A. Moussaoui, M. Solleveld: *Generalizations of the Springer correspondence and cuspidal Langlands parameters*, prepublication 2015
- [AGRS] A. Aizenbud, D. Gourevitch, S. Rallis, G. Schiffmann: *Multiplicity one theorems*, Ann. of Math. (2) 172 (2010), no. 2, 1407-1434
- [BC] N. Bergeron, L. Clozel: *Spectre automorphe des variétés hyperboliques et applications topologiques*, Astérisque No. 303 (2005), xx+218 pp
- [BD] J. Bernstein, P. Deligne: *Le "centre" de Bernstein*, In "Représentations des groupes réductifs sur un corps local, Travaux en cours" (P.Deligne ed.), Hermann, Paris, 1-32 (1984)
- [BK] J. N. Bernstein and B. Krötz: *Smooth Fréchet globalizations of Harish-Chandra modules*, Israel J. Math. 199 (2014), no. 1, 45-111
- [Beu1] R. Beuzart-Plessis: *A local trace formula for the Gan-Gross-Prasad conjecture for unitary groups: the archimedean case*, prepublication 2015
- [Beu2] —————: *Endoscopie et conjecture locale raffinée de Gan-Gross-Prasad pour les groupes unitaires*, Compos. Math. 151 (2015), no. 7, 1309-1371
- [BHC] A. Borel, Harish-Chandra: *Arithmetic subgroups of algebraic groups*, Ann. of Math. 75 (1962), 485-535
- [Ca] W. Casselman: *Canonical extensions of Harish-Chandra modules to representations of G* , Canad. J. Math. 41 (1989), no. 3, 385-438
- [CZ] P.-H. Chaudouard, M. Zydor: *Le transfert singulier pour la formule des traces de Jacquet-Rallis*, preprint 2016, arXiv:1611.09656
- [Cl1] L. Clozel: *Orbital integrals on p -adic groups: a proof of the Howe conjecture*, Ann. of Math. (2) 129 (1989), no. 2, 237-251

- [Cl2] ———: *Characters of nonconnected, reductive p -adic groups*, *Canad. J. Math.* 39 (1987), no. 1, 149-167
- [De] P. Deligne: *Le support du caractère d'une représentation supercuspidale*, *C. R. Acad. Sci. Paris Sér. A-B* 283 (1976), no. 4, A155-A157
- [DM] J. Dixmier, P. Malliavin: *Factorisations de fonctions et de vecteurs indéfiniment différentiables*, *Bull. Sci. Math. (2)* 102 (1978), no. 4, 307-330
- [GGP] W. T. Gan, B. H. Gross, D. Prasad: *Symplectic local root numbers, central critical L -values and restriction problems in the representation theory of classical groups*, in "Sur les conjectures de Gross et Prasad. I" *Astérisque* No. 346 (2012), 1-109
- [GI] W. T. Gan, A. Ichino: *The Gross-Prasad conjecture and local theta correspondence*, prepublication 2014
- [Ha] R. Neal Harris: *A refined Gross-Prasad conjecture for unitary groups*, arXiv:1201.0518
- [HT] M. Harris, R. Taylor: *The geometry and cohomology of some simple Shimura varieties*, volume 151 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, 2001. With an appendix by Vladimir G. Berkovich
- [He] G. Henniart: *Une preuve simple des conjectures de Langlands pour $GL(n)$ sur un corps p -adique*, *Invent. Math.*, 139(2), 439-455, 2000
- [II] A. Ichino, T. Ikeda: *On the periods of automorphic forms on special orthogonal groups and the Gross-Prasad conjecture*, *Geom. Funct. Anal.* 19 (2010), no. 5, 1378-1425
- [ILM] A. Ichino, E. Lapid, Z. Mao: *On the formal degrees of square-integrable representations of odd special orthogonal and metaplectic groups*, prepublication 2014
- [Jac] H. Jacquet: *Archimedean Rankin-Selberg integrals*, *Automorphic forms and L-functions II. Local aspects*, 57-172, *Contemp. Math.*, 489, Amer. Math. Soc., Providence, RI, 2009
- [JPSS] H. Jacquet, I. I. Piatetski-Shapiro, J. A. Shalika: *Rankin-Selberg convolutions*, *Amer. J. Math.* 105 (1983), no. 2, 367-46
- [JR] H. Jacquet, S. Rallis: *On the Gross-Prasad conjecture for unitary groups*, in "On certain L-functions", 205-264, *Clay Math. Proc.*, 13, Amer. Math. Soc., Providence, RI, 2011
- [JS] H. Jacquet, J. A. Shalika: *On Euler products and the classification of automorphic representations I*, *Amer. J. Math.* 103 (1981), no. 3, 499-558
- [JSZ] D. Jiang, B. Sun, C.-B. Zhu: *Uniqueness of Bessel models: The archimedean case*, *Geom. Funct. Anal.* 20 (2010), no. 3, 690-709
- [KMSW] T. Kaletha, A. Minguez, S. W. Shin, P.-J. White: *Endoscopic Classification of Representations: Inner Forms of Unitary Groups*, prepublication 2014
- [Kott] R. E. Kottwitz: *Harmonic analysis on reductive p -adic groups and Lie algebras*, in "Harmonic analysis, the trace formula, and Shimura varieties", 393-522, *Clay Math. Proc.*, 4, Amer. Math. Soc., Providence, RI, 2005
- [Lan] R. P. Langlands: *On the classification of irreducible representations of real algebraic groups*, in "Representation theory and harmonic analysis on semisimple Lie groups", 101-170, *Math. Surveys Monogr.*, 31, Amer. Math. Soc., Providence, RI, 1989
- [LRS] W. Luo, Z. Rudnick, P. Sarnak: *On the generalized Ramanujan conjecture for*

$GL(n)$, in "Automorphic forms, automorphic representations, and arithmetic" (Fort Worth, TX, 1996), Proc. Sympos. Pure Math., vol. 66, Amer. Math. Soc., Providence, RI, 1999, pp. 301-310

[Moe1] C. Mœglin: *Sur la classification des séries discrètes des groupes classiques p -adiques: paramètres de Langlands et exhaustivité*, J. Eur. Math. Soc. (JEMS) 4 (2002), no. 2, 143-200

[Moe2]:—————: *Classification et changement de base pour les séries discrètes des groupes unitaires p -adiques*, Pacific J. Math. 233 (2007), no. 1, 159-204

[MT] C. Mœglin, M. Tadić: *Construction of discrete series for classical p -adic groups*, J. Amer. Math. Soc. 15 (2002), no. 3, 715-786

[MW1] C. Mœglin, J.-L. Waldspurger: *Spectral decomposition and Eisenstein series. Une paraphrase de l'Écriture*, Cambridge Tracts in Mathematics, 113. Cambridge University Press, Cambridge, 1995. xxviii+338 pp

[MW2]:—————: *Le spectre résiduel de $GL(n)$* , Ann. Sci. École Norm. Sup. (4) 22 (1989), no. 4, 605-674

[Mok] C. P. Mok: *Endoscopic Classification of representations of Quasi-Split Unitary Groups*, Mem. Amer. Math. Soc. 235 (2015), no. 1108

[Mou] A. Moussaoui: *Centre de Bernstein enrichi pour les groupes classiques*, prepublication 2015

[Mui] G. Muic: *Some results on square integrable representations; irreducibility of standard representations*, Internat. Math. Res. Notices (1998), no. 14, 705-726

[Mu] W. Müller: *The trace class conjecture without the K -finiteness assumption*, C. R. Acad. Sci. Paris Sér. I Math. 324 (1997), no. 12, 1333-1338

[MS] W. Müller, B. Speh: *Absolute convergence of the spectral side of the Arthur trace formula for GL_n* , Geom. Funct. Anal. 14 (2004), no. 1, 58-93, With an appendix by E. M. Lapid

[Ra] D. Ramakrishnan: *A theorem on $GL(n)$ a la Tchebotarev*, Preprint 2012

[SV] Y. Sakellaridis, A. Venkatesh: *Periods and harmonic analysis on spherical varieties*, prepublication 2012

[Sil1] A. Silberger: *Introduction to harmonic analysis on reductive p -adic groups*, Mathematical Notes, vol. 23, Princeton University Press, Princeton, N.J., 1979, Based on lectures by Harish-Chandra at the Institute for Advanced Study, 1971-1973

[Sil2]:—————: *Special representations of reductive p -adic groups are not integrable*, Ann. of Math., 111, 1980, pp. 571-587

[Tad] M. Tadić: *Geometry of dual spaces of reductive groups (non-Archimedean case)*, J. Analyse Math. 51 (1988), 139-181

[Wald1] J.-L. Waldspurger: *La formule de Plancherel pour les groupes p -adiques (d'après Harish-Chandra)*, J. Inst. Math. Jussieu 2 (2003), no. 2, 235-333

[Wald2]:—————: *La formule des traces locales tordue*, prepublication 2013

[Wald3]:—————: *Sur les valeurs de certaines fonctions L automorphes en leur centre de symétrie*, Compositio Math. 54 (1985), no. 2, 173-242

[Wall] N. Wallach: *Real reductive groups II*, Academic Press 1992

[Xu] B. Xu: *On the cuspidal support of discrete series for p -adic quasisplit $Sp(N)$ and $SO(N)$* , prepublication 2015

[Xue] H. Xue: *On the global Gan-Gross-Prasad conjecture for unitary groups: approximating smooth transfer of Jacquet-Rallis*, prepublication 2015

[Yu] Z. Yun: *The fundamental lemma of Jacquet and Rallis* With an appendix by Julia Gordon. *Duke Math. J.* 156 (2011), no. 2, 167-227

[Zh1] W. Zhang: *Fourier transform and the global Gan-Gross-Prasad conjecture for unitary groups*, *Ann. of Math. (2)* 180 (2014), no. 3, 971-1049

[Zh2]———: *Automorphic period and the central value of Rankin-Selberg L -function*, *J. Amer. Math. Soc.* 27 (2014), no. 2, 541-612

[Zh3]———: *On arithmetic fundamental lemmas*, *Invent. Math.*, Volume 188, Number 1 (2012), 197-252

[Zy] M. Zydor: *Les formules des traces relatives de Jacquet-Rallis grossières*, preprint 2015, arXiv:1510.04301