



Endosymbiont diversity and prevalence in herbivorous spider mite populations in South-Western Europe

Flore Zélé, Inês Santos, Isabelle Olivieri, Mylene Weill, Olivier Duron, Sara Magalhães

► To cite this version:

Flore Zélé, Inês Santos, Isabelle Olivieri, Mylene Weill, Olivier Duron, et al.. Endosymbiont diversity and prevalence in herbivorous spider mite populations in South-Western Europe. FEMS Microbiology Ecology, 2018, 94 (4), <10.1093/femsec/fiy015>. <hal-01943929>

HAL Id: hal-01943929

<https://hal.science/hal-01943929v1>

Submitted on 3 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

Endosymbiont diversity and prevalence in herbivorous spider mite populations in South-Western Europe

Flore Zélé^{1*}, Inês Santos¹, Isabelle Olivieri^{†2}, Mylène Weill², Olivier Duron³, Sara Magalhães¹

¹ Centre for Ecology, Evolution and Environmental Changes (cE3c), Faculdade de Ciências da Universidade de Lisboa, Edifício C2, Piso 3, Campo Grande, 1749016 Lisbon, Portugal

² Institut des Sciences de l'Evolution (CNRS-Université de Montpellier-IRD-EPHE), 34095 Montpellier, CEDEX 5, France

³ Maladies Infectieuses et Vecteurs: Ecologie, Génétique, Evolution et Contrôle (CNRS-Université de Montpellier-IRD), Centre de Recherche IRD, 911 Avenue Agropolis, 34394 Montpellier, France

[†] Isabelle Olivieri sadly passed away before the submission of this paper

* Corresponding author: e-mail: fezele@fc.ul.pt

Abstract

Bacterial endosymbionts are known as important players of the evolutionary ecology of their hosts. However, their distribution, prevalence and diversity are still largely unexplored. To this aim, we investigated infections by the most common bacterial reproductive manipulators in herbivorous spider mites of South-Western Europe. Across 16 populations belonging to three *Tetranychus* species, *Wolbachia* was the most prevalent (ca. 61%), followed by *Cardinium* (12-15%), while only few individuals were infected by *Rickettsia* (0.9-3%), and none carried *Arsenophonus* or *Spiroplasma*. These endosymbionts are here reported for the first time in *T. evansi* and *T. ludeni*, and showed variable infection frequencies between and within species, with several cases of coinfections. Moreover, *Cardinium* was more prevalent in *Wolbachia*-infected individuals, which suggests facilitation between these symbionts. Finally, sequence comparisons revealed no variation of the *Wolbachia* *wsp* and *Rickettsia* *gt1A* genes, but some diversity of the *Cardinium* 16S rRNA, both between and within populations of the three mite species. Some of the *Cardinium* sequences identified belonged to distantly-related clades, and the lack of association between these sequences and spider mite mitotypes suggests repeated host switching of *Cardinium*. Overall, our results reveal a complex community of symbionts in this system, opening the path for future studies.

Keywords

Endosymbiosis, maternally inherited bacteria, multiple infections, reproductive parasitism, two-spotted spider-mite, *Tetranychus urticae*.

1 Introduction

2 Terrestrial arthropods often harbour several bacterial microorganisms that occur in the
3 cytoplasm of host cells, and are maternally inherited from mother to offspring through the
4 egg cytoplasm (Moran *et al.*, 2008). In the last decade, intensive field surveys of distinct
5 arthropod species revealed that the most common of these endosymbionts are *Wolbachia*,
6 *Rickettsia*, *Cardinium*, *Arsenophonus* and *Spiroplasma* (Duron *et al.*, 2008, Weinert *et al.*,
7 2015). Whereas *Wolbachia* is present in most groups, *Rickettsia* and *Cardinium* have low
8 incidence in the hexapoda relative to other groups, with the exception of hemiptera (Weeks
9 *et al.*, 2003, Zchori-Fein & Perlman, 2004, Weinert *et al.*, 2015). These three bacterial
10 genera are also extremely prevalent in chelicerates, especially in spiders, mites and ticks
11 (Duron *et al.*, 2008, Weinert *et al.*, 2015, Zhang *et al.*, 2016a, Duron *et al.*, 2017).

12 The maternal inheritance of endosymbiotic bacteria has selected for a variety of
13 phenotypic changes in their hosts, such as cytoplasmic incompatibility (CI, *i.e.*, embryonic
14 mortality in crosses between infected males and uninfected females), feminization (*i.e.*,
15 genetic males that develop as females), thelytokous parthenogenesis (*i.e.*, asexual daughter
16 development), or male killing (*i.e.*, infected males are eliminated during embryogenesis or
17 late larval instars). These reproductive manipulations, by promoting the production and
18 fitness of infected daughters (*i.e.*, the transmitting sex) via negative effects on the fitness of
19 individuals not involved in the transmission, enhance symbiont transmission and allow their
20 spread within host populations (Werren *et al.*, 2008, Engelstadter & Hurst, 2009).
21 Moreover, such manipulations may also facilitate the transmission of other vertically-
22 transmitted symbionts co-occurring in the same host (Frank, 1998, Engelstadter *et al.*, 2004,
23 Vautrin *et al.*, 2008, Vautrin & Vavre, 2009, Hellard *et al.*, 2015).

24 Although traditionally viewed as vertically-transmitted symbionts, evidence of
25 horizontal transmission events stems from incongruences between host and symbiont
26 phylogenies (Schilthuizen & Stouthamer, 1997, Vavre *et al.*, 1999, Raychoudhury *et al.*,
27 2009, Gerth *et al.*, 2013). Indeed, a few recent studies have provided direct evidence for
28 horizontal transmission, and suggest parasitism and feeding as the two main routes of
29 transfer. For instance, *Wolbachia* can be transmitted between whiteflies via parasitoids
30 (Ahmed *et al.*, 2015), or to a parasitoid from its infected drosophila host (Heath *et al.*, 1999),
31 *Spiroplasma* may be transmitted between two *Drosophila* species via ectoparasitic mites
32 (Jaenike *et al.*, 2007, Brown & Lloyd, 2015), and *Rickettsia* infecting *Bemisia tabaci* may
33 be transferred to parasitoids, although it is not vertically transmitted in the latter (Chiel *et*

34 *al.*, 2009). Similarly, *Trichogramma* wasps of the same or different species have
35 exchanged symbionts when infecting the same butterfly egg (Huigens *et al.*, 2004).
36 Symbionts may also be transmitted via the resource that their hosts feed on (but see Faria
37 *et al.*, 2016). The most compelling evidence for this infection route comes from
38 herbivorous arthropods, in which horizontal transmission via plant feeding has been shown
39 for *Rickettsia* and *Wolbachia* in whiteflies (Caspi-Fluger *et al.*, 2012, Li *et al.*, 2017) and
40 for *Cardinium* in leaf hoppers (Gonella *et al.*, 2015). However, the incidence of horizontal
41 transmission for many vertically-transmitted arthropods endosymbionts is still under
42 debate and other studies do not corroborate these results (e.g. Gualtieri *et al.*, 2017).

43 Several studies have shown that herbivorous spider mites are often infected with
44 *Wolbachia* and *Cardinium* (Gotoh *et al.*, 2003, Liu *et al.*, 2006, Gotoh *et al.*, 2007b, Xie *et al.*,
45 *al.*, 2011, Ros *et al.*, 2012, Suh *et al.*, 2015, Zhang *et al.*, 2016b) and occasionally with
46 *Spiroplasma* (Zhang *et al.*, 2016b, Staudacher *et al.*, 2017) and *Rickettsia* (Hoy &
47 Jeyaprakash, 2005, Zhang *et al.*, 2016b). In spider mites, *Wolbachia* and *Cardinium* may
48 have diverse fitness effects. In particular, they may cause different degrees of CI (Gotoh *et al.*,
49 *al.*, 2003, Gotoh *et al.*, 2007a, Gotoh *et al.*, 2007b, Ros & Breeuwer, 2009, Xie *et al.*,
50 2011, Zhu *et al.*, 2012, Suh *et al.*, 2015, Xie *et al.*, 2016). Moreover, these bacteria have
51 been found in the same individual host (Ros *et al.*, 2012, Zhu *et al.*, 2012, Zhao *et al.*,
52 2013b, Zhao *et al.*, 2013a, Xie *et al.*, 2016), and multiple strains of these symbionts can be
53 found in spider-mite populations, although levels of nucleotide diversity depend on the
54 host species and/or populations (Gotoh *et al.*, 2003, Liu *et al.*, 2006, Gotoh *et al.*, 2007a,
55 Gotoh *et al.*, 2007b, Yu *et al.*, 2011, Ros *et al.*, 2012, Zhang *et al.*, 2016b).

56 This study aims to contribute to knowledge of the symbiont community of spider
57 mites, by characterizing the community of herbivorous spider-mite populations and their
58 endosymbionts in South-West Europe, a relatively unstudied area. By unravelling the
59 diversity and prevalence of endosymbionts in natural spider-mite populations this study
60 might provide some insight on potential facilitation in multiple infections and horizontal
61 transfers among hosts.

62 63 **Material and methods**

64 *Spider-mite collection and population rearing*

65 Populations of Tetranychidae mites were collected from September to December 2013 at
66 several locations in Central Portugal (Table 1). In addition, two populations were collected

from Southern Spain, and one population from Southern France for comparison with nearby countries. Each population in each location was collected from different leaves on different plants. Mite collection was done by placing infested leaves in closed paper bags. Subsequently, adult females ($n=25-600$) from all populations were transferred to insect-proof cages containing bean cv. Contender seedlings (obtained from Germisem, Oliveira do Hospital, Portugal) immediately after collection from the field. After species identification for each of these populations, Solanaceae specialist spider mites (*Tetranychus evansi*) were transferred to tomato cv. Money Maker seedlings (obtained from Mr. Fothergill's Seeds, Kentford, UK). The colonies were then maintained in the laboratory under standard conditions ($25 \pm 2^\circ\text{C}$, 60% RH, 16/8 h L/D; Clemente *et al.*, 2016).

Molecular species identification and screening for endosymbiont infection

From 0 to 100 days after collection, adult spider mite females ($n=11-16$) were randomly picked from each colony and individually analyzed for the presence of reproductive manipulators and for species identification. Total genomic DNA was extracted from each individual spider mite using the Sigma-Aldrich protocol and materials (GenElute™ Mammalian Genomic DNA Miniprep Kit, Sigma-Aldrich, St. Louis, MO, United States). Total DNA was eluted in the final step with 20µl RNase-free water (Qiagen NV, Venlo, The Netherlands) in order to increase the concentration of DNA obtained from these very tiny animals (c.a. 300µm long). The same DNA from each individual was used to screen for the presence of potential endosymbionts and to identify the spider-mite species. These identifications were performed by PCR amplification then sequencing of a fragment of the nuclear ribosomal DNA (rDNA) ITS2 (internal transcribed spacer 2) region, which is the most commonly used marker to distinguish among spider-mite species (Ben-David *et al.*, 2007, Hurtado *et al.*, 2008), as the mitochondrial DNA (mtDNA) Cytochrome Oxidase subunit I (*COI*) is more polymorph in spider mites (Navajas *et al.*, 1998). The presence of *Wolbachia*, *Cardinium*, *Rickettsia*, *Arsenophonus* and *Spiroplasma* in the each sample was determined using genus-specific primers, and the mitochondrial DNA (mtDNA) Cytochrome Oxidase subunit I (*COI*) was amplified and sequenced for 75 samples chosen depending of their infection status with *Cardinium* and *Wolbachia*. The list of primers, for both hosts and symbionts, is given in Table S1 (Additional file 1 in the Supplementary data). For each of the genes under study, the PCRs were done along with both positive and negative controls, in a 10µl reaction mixture containing 5µl of 2X QIAGEN Multiplex PCR Master Mix (Qiagen NV, Venlo, The Netherlands), 2µl RNase free water, 1µl of each primer at 2µM, and 1µl of

DNA solution. Amplification conditions were as follows: 15min at 94°C, then 35 cycles of 94°C for 30s, annealing (see primer-specific temperatures in Table S1) for 1min30s, 72°C for 1min and a final step at 72°C for 10min. PCR products were visualized on 2% agarose gel. To verify the identity of the amplified endosymbionts genes (for all *Cardinium*- and *Rickettsia*-infected individuals, and up to three *Wolbachia*-infected individuals per population haphazardly chosen within both single- and multiple-infected mites), the PCRs were repeated for positive samples in a 20µl final volume reaction, the PCR products were purified using Qiagen protocol and materials (QIAquick PCR Purification Kit) and sent for sequencing (Stabvida, Caparica, Portugal). All ITS2 and *COI* genes were also sequenced using the same protocol. The chromatograms were checked manually using MEGA version 5.1 beta (Tamura *et al.*, 2011) and the resulting sequences were run against the non-redundant nucleotide database using the BLAST algorithm of the National Center for Biotechnology Information (NCBI). New sequences have been deposited in GenBank (see Table S2 and S3 in the Supplementary data, Additional file 1).

Phylogenetic analyses of COI mitochondrial haplotypes

The phylogenetic relationships between *Tetranychus* spp. haplotypes were reconstructed using maximum-likelihood (ML) analyses based on *COI* mitochondrial nucleotide sequences (fragments of 392 unambiguously-aligned sites; best-fitting evolutionary model: HKY+I). The evolutionary model that best fitted the sequence data was determined using the Akaike information criterion with the program MEGA (Tamura *et al.*, 2011). ML heuristic searches using a starting tree obtained by neighbor joining were conducted in MEGA (Tamura *et al.*, 2011). Clade robustness was assessed by bootstrap analysis using 1000 replicates.

Statistical analyses

Analyses were carried out using the R statistical package (v. 3.3.2). The different statistical models built to analyse the effects of species, or of population within species, on the prevalence of *Wolbachia* and of *Cardinium* are described in the electronic supplementary material, Table S4 (Additional file 1, Supplementary data). The general procedure for building the statistical models was as follows: time (number of days after collection from the field at which DNA extraction was performed), species and the status of infection of the females by *Wolbachia* or *Cardinium* were fit as fixed explanatory variables, whereas population was either fit as a random explanatory variable when analysing the complete dataset, or as a fixed effect when analysing the differences between populations within each

spider-mite species. The effect of spider-mite species (models 1 and 5) on the proportion of *Wolbachia*- or *Cardinium*-infected females, a binary response variable (1: infected, or 0: uninfected), was analysed using a generalized linear mixed model with a binomial distribution (glmer, lme4 package), and the effect of population within species (models 2 to 4, and 6 to 8) was analysed using a generalized linear model with a binomial distribution (glm). When the variables species or population were found to be significant, *a posteriori* contrasts (Crawley, 2007) were carried out by aggregating factor levels and testing the fit of the simplified model using ANOVA. For all analyses, maximal models were simplified by sequentially eliminating non-significant terms to establish a minimal model (Crawley, 2007), and the significance of the explanatory variables was established using chi-squared tests (Bolker, 2008). The significant X^2 values given in the text are for the minimal model (Crawley, 2007). Finally, the effect of *Wolbachia* and *Cardinium* infection on sequence divergence among *COI* haplotypes, and, for *Cardinium*-infected individuals, the effect of *Cardinium* strains on *COI* sequence divergence, were assessed by a hierarchical analysis of molecular variance (AMOVA; Excoffier *et al.*, 1992). The statistical significance of the variance components of the AMOVA was determined by nonparametric procedures using 10000 random permutations. Full dataset are given in Additional files 2, 3 and 4 (Supplementary data).

Results

Species identification: ITS ribosomal types and COI mitochondrial haplotypes

Amplification and sequencing of a portion of the ITS2 gene allowed assigning 3 out of the 16 populations collected to *T. evansi*, 3 populations to *T. ludeni* and the 10 remaining populations to *T. urticae* (red morph; Table 1). All sequences obtained from populations of *T. urticae* were strictly identical and matched (100% identity at the nucleotide level) a sequence obtained from *T. urticae* collected in southern Spain (e.g. GenBank: GU565314). Similarly all sequences obtained for *T. ludeni* (GenBank: MF428439) were strictly identical and matched (99% identity) several sequences obtained for this species worldwide (e.g. New Zealand, Japan, India, or Israel; GenBank: KP744529, AB076371, KU759957 and DQ656451, respectively). For *T. evansi*, however, both ITS2 ribosomal type 1 (GenBank: FJ440674) and type 2 (GenBank: FJ440673; Boubou *et al.*, 2011) were found in the population “QL”, but only the type 1 in “GRA” and “GH”.

Analysis of mitochondrial *COI* DNA sequences from 75 individuals revealed 64

polymorphic sites within 392 bases sequenced (Table S2), and allowed distinguishing eight different mtDNA haplotypes (each individual within the same haplotype sharing exactly the same *COI* sequence; Table 2; Fig. 1). In *T. urticae*, almost all mites collected in Spain and Portugal belonged to the same haplotype (GenBank: HM565901, and MF428440, respectively), and shared a high degree of similarity with each other (99% identity; the two sequences differed by one single nucleotide mutation; Table S2). One female from the population “DC” collected in Portugal, however, belonged to a *COI* haplotype (GenBank: MF428441; Table 2) more similar to a haplotype found in the population “FR” from France (GenBank: HM486513) than to the one from the Spanish populations (96% and 94% identity, respectively; Table S2). In the French population, two other haplotypes were found: the main haplotype found in Portugal (GenBank: MF428440), and another, highly similar to this one (99% identity; GenBank: HM565899). In *T. evansi*, two different haplotypes were found (GenBank: MF428442 and MF428443), which were strictly identical (100% identity, query cover: 91%; Table S2) to the *COI* haplotypes I and II, respectively, originally recorded in France and Spain (GenBank: FJ440676 and FJ440678; Gotoh *et al.*, 2009), but also previously found in Portugal (Boubou *et al.*, 2011). Conversely, only one haplotype of *T. ludeni* was found and was identical to a haplotype found in China (GenBank: GQ141911; Table 2). Importantly, one *T. ludeni* female from the population “Assaf”, and 4 females from all *T. evansi* populations (from both ITS types), harbored two different *T. urticae COI* haplotypes, but the reverse (*i.e.*, *T. ludeni* or *T. evansi COI* haplotypes associated to *T. urticae* ITS type) was not observed (Table 2).

Inter- and intraspecific variation of endosymbionts prevalence and diversity

Overall, our sampling revealed infections by *Wolbachia*, *Cardinium* and *Rickettsia* within the three spider mites species, but none of the tested populations were found infected by *Arsenophonus* or *Spiroplasma* (Fig. 2).

Wolbachia was the most prevalent endosymbiont with an average of $60.7 \pm 3.3\%$ of infected females (Fig. 2). This prevalence differed significantly between host species (model 1, $X^2_2 = 5.92$, $p=0.05$), with no significant difference between infection prevalence in *T. ludeni* and *T. evansi* ($31.6 \pm 7.6\%$ and $41.5 \pm 7.8\%$, respectively; $X^2_1 = 0.24$, $p=0.63$), but the infection frequency in *T. urticae* being significantly higher than in these two species ($74.3 \pm 3.7\%$; $X^2_1 = 5.69$, $p=0.02$). Within species, no significant difference was found in *Wolbachia* prevalence among populations of *T. evansi* (model 2, $X^2_2 = 3.62$, $p=0.16$), but populations differed significantly for *Wolbachia* infection frequencies in *T. ludeni* and *T. urticae* (model

3, $X^2_2 = 11.51$, $p=0.003$, and model 4, $X^2_9 = 93.58$, $p<0.0001$, respectively). In *T. ludeni*, no *Wolbachia*-infected mites were found in “Assaf”, while both “Alval” and “CVM” harbored 46.2 ± 10.0 % of infected mites. In *T. urticae*, the lowest prevalences were found in the populations collected in Spain (on average 14.8 ± 7.0 %) and France (46.7 ± 13.3 %; contrast between Spanish-French populations: $X^2_1 = 5.28$, $p=0.02$), whereas *Wolbachia* infection was fixed in 5 out of 7 of the populations collected in Portugal, and the remaining two Portuguese populations (CH and DC) had an average prevalence of 83.3 ± 6.9 % (contrasts between CH-DC and the other Portuguese populations: $X^2_1 = 13.05$, $p<0.001$; contrast between CH-DC and the French populations, $X^2_1 = 6.99$, $p=0.008$).

At the molecular level, no diversity of *Wolbachia* was found (Table 3). Of the 42 *Wolbachia*-infected mites for which the *wsp* gene was sequenced, 41 were strictly identical and matched (100% identity at the nucleotide level) several sequences available in GenBank for *Wolbachia* infecting several *Tetranychus* species worldwide (e.g. GenBank: DQ910771 and JX844806 in *T. urticae*, AB096218 and AF358417 in *T. kanzawai*, AB096226 in *T. pueraricola*, AY585711 in *T. cinnabarinus*) and previously attributed to members of the *Wolbachia* B supergroup and Ori subgroup (Gotoh *et al.*, 2003). Only one of the sequences obtained (GenBank: MF428430), from a *Wolbachia*-infected mites of the *T. evansi* population “GRA”, differed from the others (82% identity at the nucleotide level), but was highly similar (99% identity at the nucleotide level) to many sequences found in diverse arthropods species such as fruit flies (GenBank: FJ403332, KU870673, DQ834379, or EU116319), beetles (GenBank: KX436089), moths (GenBank: GU166594), or head lice (GenBank: AY596782). Note, however, that strains with similar *wsp* sequences can have very different MLST allelic profiles (Baldo *et al.*, 2006), but the amount of DNA obtained from each individual spider mite strongly restrained the possible number of PCRs and an in-depth study of *Wolbachia* variation within *Tetranychus* mites was beyond the scope of this study.

The prevalence of *Cardinium* was 14.6 ± 2.4 % on average (total of 32 infected mites over 219). However, we could obtain the *16S rRNA* sequences only for 27 of 32 infected individuals: for 5 individuals we failed to amplify *Cardinium* through three successive PCR following the first diagnostic of infection. Two main possibilities could explain this failure: First, very low densities of *Cardinium* within these individuals could lead to PCR-based imperfect detection, as shown for other parasites (Zélé *et al.*, 2014). Second, we cannot totally exclude the possibility of contaminations or of unspecific amplifications of other bacteria when using *16S rRNA* primers for *Cardinium*. For this latter reason, we did not

consider these five individuals as infected in the following analyses. After correction, the prevalence of *Cardinium* was overall of $12.3 \pm 2.2\%$ ($9.8 \pm 4.7\%$ in *T. evansi*, $10.5 \pm 5.0\%$ in *T. ludeni*, and $13.6 \pm 2.9\%$ in *T. urticae*) with no significant difference in prevalence among the three species (model 5, $X^2_2 = 0.32$, $p=0.85$; Fig. 2). Within species, no differences among populations of *T. evansi* and *T. ludeni* were found (models 6 and 7, $X^2_2 = 0.55$, $p=0.76$ and $X^2_2 = 3.25$, $p=0.19$, respectively), but *Cardinium* prevalences in populations of *T. urticae* differed significantly (model 8, $X^2_9 = 31.19$, $p<0.001$). Indeed, in 3 populations (“COL”, “DF” and “LOU”), none of the mites tested were infected, but *a posteriori* contrast analyses among the remaining (infected) populations revealed no significant difference in *Cardinium* prevalence ($X^2_6 = 6.62$, $p=0.36$).

At the molecular level, we found some degrees of inter-individual variations in the *Cardinium* 16S rRNA sequences (Table 3). Of the 27 *Cardinium* 16S rRNA sequences obtained, we could discriminate 6 different alleles, which, despite sharing a high degree of similarity (97-99% identity at the nucleotide level), differed by single nucleotide mutations over 13 different nucleotide sites (Table S3). The most prevalent allele (GenBank: MF428434) was found in 17 *Cardinium*-infected mites over the three species (c.a. 63%; Table 3). The corresponding sequence is very close to several sequences found for *Cardinium* in diverse arthropod species, including spider mites (e.g. 99% identity at the nucleotide level with a sequence obtained in *T. urticae* in Japan; GenBank: AB241134; Gotoh *et al.*, 2007a). Three additional mites harbored highly similar *Cardinium* alleles (99% identity at the nucleotide level; GenBank: MF428435, MF428436, and MF428437, respectively; Tables 3 and S3). Interestingly, we also found alleles identical or highly similar (99% identity at the nucleotide level), to an allele previously found for *Cardinium* endosymbiont of *Scaphoideus titanus*, the American grapevine leafhopper (GenBank: AM042540; Marzorati *et al.*, 2006), for two *T. urticae* (GenBank: MF428431), and one *T. ludeni* (GenBank: MF428432) individuals, respectively. Finally, four individuals (one *T. evansi* and three *T. urticae*) harbored *Cardinium* sequences (GenBank: MF428433) with double nucleotide peaks at the nucleotide positions that differ between AB241134 and MF428434 (Table S3). This result suggests co-infection with two different *Cardinium* strains within these mites.

Infection by *Rickettsia* was rare. Indeed, this symbiont was found in only 6 of 219 individuals belonging to the three spider mite species ($2.7 \pm 1.1\%$; Fig. 2): $7.3 \pm 4.1\%$ of *T. evansi* individuals were infected, belonging to a single population, $2.6 \pm 2.6\%$ of *T. ludeni* individuals, also from a single population, and $1.4 \pm 1.0\%$ of *T. urticae* individuals, stemming from two populations. However, for 4 of the 6 infected individuals we failed to

obtain the *gltA* sequences of *Rickettsia*. After correction, the prevalence of *Rickettsia* was overall of 0.9 ± 0.6 % among the three mite species (2.4 ± 2.4 % in *T. evansi*, 0% in *T. ludeni*, and 0.7 ± 0.7 % in *T. urticae*). The two sequences obtained, one in *T. evansi*, and one in *T. urticae* (GenBank: MF428438; Table 3), were strictly identical to a sequence previously found in *T. urticae* in China (GenBank: KP828066; Zhang *et al.*, 2016b). The most closely related sequences found in other arthropod hosts were from *Rickettsia* in the parasitoid wasp *Pnigalio sp.* (97% identity at the nucleotide level; GenBank: GU559856), and in *Bemisia tabaci* (96% identity; GenBank: DQ077708).

Multiple infections and covariation of Wolbachia and Cardinium prevalence

Several individuals within the three mites species, and across different populations, were infected with more than one endosymbiont. We detected 25 mites (11.4 ± 2.2 % in average) coinfecting by *Wolbachia* and *Cardinium*, 4 mites (ca. 1.8 ± 0.9 %) coinfecting by *Wolbachia* and *Rickettsia* and one individual (ca. 0.5 ± 0.5 %) infected by the three endosymbionts (population “CVM”; Fig. 2). Moreover, our analyses revealed a co-variation between the prevalence of *Wolbachia* and that of *Cardinium*, both symbionts affecting significantly the probability of infection by each other (model 1, $X^2_1 = 4.99$, $p=0.03$, and model 5, $X^2_1 = 4.56$, $p=0.03$, respectively; Fig. 3). However, no particular pattern of association between *Cardinium 16S rRNA* gene sequences, *Wolbachia* infection, mites species or population emerged (Table 3).

Effects of Wolbachia and Cardinium infection on mtDNA variation

No particular pattern of association was found between the mitochondrial *COI* DNA sequences obtained from individual mites and their *Wolbachia* and/or *Cardinium* infection status ($F_{1,74} = 0.05$; $p=0.07$, and $F_{1,74} = 0.009$; $p=0.26$, respectively; Fig. 1), or between their mitochondrial *COI* DNA sequences and the *Cardinium 16S rRNA* type ($F_{1,75} = 0.22$; $p=0.18$, Table 4). For instance, the most prevalent *Cardinium* sequence obtained (MF428434; c.a. 63% of *Cardinium* infected-mites; Table 3) was found for all mitotypes, and, reciprocally, several mitotypes harbored different *Cardinium* sequences (Table 4).

Discussion

In this study, we analysed the prevalence and diversity of South-West European populations of spider mites and their endosymbionts. The populations collected mainly belonged to three

species of spider mites: *T. urticae*, *T. evansi* and *T. ludeni*, the most prevalent being *T. urticae*. Our sampling effort was such that we did not pre-select the plants sampled. Consequently, this result may reflect the host range of *T. urticae*, which is larger than that of *T. evansi* and *T. ludeni* (Migeon & Dorkeld, 2006-2017). However, our results contrast with those of another study conducted in the Eastern part of the Iberian Peninsula, near Valencia, where *T. evansi* is the most abundant species (Ferragut *et al.*, 2013). A study on invasion scenarios of *T. evansi* predicts that populations around Valencia stem from an invasion route that differs from that leading to the establishment of *T. evansi* in Portugal (Boubou *et al.*, 2012). Possibly, the invasive success of this species differs depending on the invasion event. The *T. evansi* clades that we recovered match those found earlier for the same region in Portugal (Boubou *et al.*, 2011).

Analysis of mitochondrial and nuclear DNA revealed a high *COI* diversity in comparison to that of ITS for *T. urticae*, as found earlier (Navajas *et al.*, 1998). Overall, these two markers yielded concordant species identifications in most individuals tested. However, we found mitochondrial DNA of *T. urticae* in the other two species, albeit at low frequencies. Screening contaminations seems unlikely, as mitochondrial and nuclear DNA were extracted from the same individuals, PCR controls were all correct and these unexpected ITS-*COI* associations occurred for two different mitotypes, and in four different populations (one of *T. ludeni* and the three *T. evansi*). Therefore, it seems that introgression may occur among these species, and this suggest that they co-occur on the same host plants and locations, although we did not detect species co-occurrence in this study. In laboratory studies using these species, viable hybrids have been found at very low frequencies as well (Sato *et al.*, 2014, Clemente *et al.*, 2016, but see Clemente *et al.*, in press), and assortative mating was either low (Clemente *et al.*, 2016) or absent (Sato *et al.*, 2014). These findings are compatible with the possible introgression found here. Nevertheless, the low frequencies observed here might also be due to the detection of F1 hybrids only, as in most cases such hybrids are sterile (Clemente *et al.*, 2016). Curiously, however, this introgression seemed to be unidirectional, with the mtDNA of *T. ludeni* or *T. evansi* never being found in individuals with nuclear *T. urticae* DNA. It would thus be interesting to corroborate these results in future laboratory experiments, along with broader analyses of natural populations of these species.

Of the five endosymbionts screened, only *Wolbachia*, *Cardinium* and *Rickettsia* were found to infect spider mites, whereas none of the tested individuals harbored *Spiroplasma* or *Arsenophonus*. However, this is the first report of *Wolbachia*, *Cardinium* and *Rickettsia* in *T. evansi* and *T. ludeni*. Overall, *Wolbachia* prevalence was higher than that of the two other

symbionts, even though it was lower in *T. evansi* and *T. ludeni*, and in non-portuguese *T. urticae* populations. Such variable levels of *Wolbachia* prevalence across species and populations have already been reported in other regions worldwide (Gotoh *et al.*, 2003, Gotoh *et al.*, 2007b, Zhang *et al.*, 2016b). Interestingly, however, we found *Wolbachia* in the invasive species *T. evansi*. This runs counter other studies that have shown the loss of *Wolbachia* in invasive populations of fire ants (Shoemaker *et al.*, 2000), Argentine ants (Reuter *et al.*, 2005), and Australian thrips (Nguyen *et al.*, 2016). In invasive species, the loss of *Wolbachia* is expected because introduced populations are likely to harbor only a subset of the symbionts of their parental population(s), possibly because a bottleneck effect during invasion. Spider mites may thus be used to test hypotheses concerning the loss or maintenance of symbionts in invasive populations (see below). *Cardinium* was found in all three spider mites species, which corroborates previous findings of a relatively high occurrence of this symbiont in mites (Nakamura *et al.*, 2009, Zhang *et al.*, 2016a) compared with other arthropods groups (Weeks *et al.*, 2003, Zchori-Fein & Perlman, 2004, Duron *et al.*, 2008, Nakamura *et al.*, 2009). However, its prevalence was overall much lower than that of *Wolbachia*, a result not always found in spider mites (Zhang *et al.*, 2016b). The prevalence of *Rickettsia* was very low, as found by Zhang *et al.* (2016b), but this study is the first to report *Rickettsia* infection in *T. ludeni* and *T. evansi*. Finally, although *Spiroplasma* has been previously found in *T. urticae* (Enigl & Schausberger, 2007), we did not detect it in our study, as reported by Zhang *et al.* (2016b). Similarly, we did not find *Arsenophonus*. Although the latter has been found in the Acari (Clay *et al.*, 2008, Dergousoff & Chilton, 2010, Clayton *et al.*, 2015, Duron *et al.*, 2017), it has, to our knowledge, never been reported in spider mites.

The maintenance and spread of reproductive manipulators in nature, hence their prevalence, are mainly affected by the type and strength of reproductive manipulation, the fitness costs/benefits of the symbionts to their hosts, and the efficiency of vertical transmission (Werren *et al.*, 2008, Engelstadter & Hurst, 2009). Variability in any of these factors might explain the observed lower prevalence of *Cardinium* than that of *Wolbachia*. Both endosymbionts induce CI in spider mites (Gotoh *et al.*, 2007a, Gotoh *et al.*, 2007b, Ros & Breeuwer, 2009, Zhu *et al.*, 2012, Zhao *et al.*, 2013b). Although some studies found that *Cardinium* induces a higher level of CI than *Wolbachia* in several mites species (e.g. *T. pierci* and *Bryobia sarothamni*; Ros & Breeuwer, 2009, Zhu *et al.*, 2012), another showed that *Cardinium* may not induce CI in *T. urticae* (Gotoh *et al.*, 2007a). It is thus possible that its lower prevalence is due to low reproductive manipulation. In addition, *Cardinium* may

exhibit imperfect maternal transmission, as found in thrips (Nguyen *et al.*, 2017), or in the parasitic wasp *Encarsia pergandiella* (although near-perfect in this species; Perlman *et al.*, 2008). Conversely, the high prevalence of a single *Wolbachia* strain found here may be due to this strain inducing low fitness costs, strong reproductive manipulation and having nearly perfect maternal transmission. An in-depth study of the fitness effects of these endosymbionts, along with their ability to manipulate the reproduction of their host in these populations would thus allow testing these hypotheses.

These general predictions, coming from epidemiological modelling, may be tempered by environmental factors. Indeed, a large number of other factors have been identified, such as symbiont-host genotype interactions, host diet, or temperature effects (e.g. Van Opijnen & Breeuwer, 1999, Gotoh *et al.*, 2007b, Mouton *et al.*, 2007, Wilkinson *et al.*, 2007, Pan *et al.*, 2013, Zhang *et al.*, 2013b, Boivin *et al.*, 2014, Suh *et al.*, 2015). For instance, Gotoh *et al.* (2007a) showed that *Wolbachia* is much more sensitive to elevated temperature than *Cardinium* in *T. pueraricola*. In our study, the higher summer temperatures in Almeria (Spain; c.a. 26.7°C on average in August; <http://www.aemet.es/es/serviciosclimaticos/datosclimatologicos/valoresclimatologicos?l=63250&k=and>) compared to that of the region of Lisbon (Portugal; c.a. 23.5°C on average in August; <http://www.ipma.pt/pt/oclima/normais.clima/1981-2010/012/>) might explain the lower prevalence of *Wolbachia*, but not that of *Cardinium* in the *T. urticae* populations collected in this region. The fact that a single *Wolbachia* *wsp* sequence was found in this region, which contrasts with other studies of spider mites in other regions (Gotoh *et al.*, 2003, Gotoh *et al.*, 2007b, Ros *et al.*, 2012, Zhang *et al.*, 2013a, Zhang *et al.*, 2013b), may indicate specific adaptation of this strain to the environmental conditions of South-Western Europe. Host plants of herbivorous arthropods may also affect symbiont diversity and identity (e.g. Ferrari *et al.*, 2004, Ahmed *et al.*, 2010, Brady & White, 2013, Guidolin & Consoli, 2017; but see Ji *et al.*, 2015). Moreover, bacterial endosymbionts may also be conditional mutualists in some host species, conferring advantages under certain environmental conditions. For instance, they may compensate for the nutritionally deficient diets of herbivore hosts and promote dietary specialization by affecting their performance on plants, they may also increase their ability to tolerate high temperatures, or may protect them against natural enemies (reviewed in Haine, 2008, Moran *et al.*, 2008, Frago *et al.*, 2012). Unfortunately, our sampling design, largely unbalanced with respect to these factors, does not allow testing their relevance for the system under study.

Interaction between symbionts within a single host may also be an important factor

affecting their prevalence, either negatively (competition), or positively (facilitation) (Vautrin *et al.*, 2008, Vautrin & Vavre, 2009, Hellard *et al.*, 2015). As previously found in several spider mites species in China (Liu *et al.*, 2006, Zhao *et al.*, 2013b, Zhang *et al.*, 2016b), and in Northern Europe (Ros & Breeuwer, 2009, Ros *et al.*, 2012), in this study *Cardinium* prevalence was higher in populations also infected by *Wolbachia*, which may suggest facilitation between these endosymbionts. The maintenance of *Cardinium* in the spider mites populations collected in this study might thus be due to a hitchhiking effect, whereby *Wolbachia*-induced phenotypic changes favor their own transmission, but also that of other co-infecting symbionts (Frank, 1998, Engelstadter *et al.*, 2004, Vautrin & Vavre, 2009). Moreover, the effects of each symbiont on their host may be exacerbated in presence of the other. Indeed, several studies showed that *Wolbachia* can strengthen the induction of cytoplasmic incompatibility by *Cardinium* in several spider-mite (Zhu *et al.*, 2012, Zhao *et al.*, 2013b, Zhao *et al.*, 2013a), or arthropods (White *et al.*, 2009) species (but see Nguyen *et al.*, 2017). Moreover, Ros & Breeuwer (2009) found a higher fecundity advantage conferred by *Wolbachia-Cardinium* co-infections in *Bryobia sarothamni* mites, than by *Wolbachia* or *Cardinium* single infections (although this result may be confounded by the host genetic background, as acknowledged by the authors). Studies of the fitness effects of both *Wolbachia* and *Cardinium*, in single and multiple infections in these populations would allow corroborating or rejecting this hypothesis.

Symbionts usually spread within species, but they may also cross species boundaries via two main routes. First, cytoplasmic introgression, through interspecific hybridization events, may favor the spread of vertically-transmitted symbionts among species (Raychoudhury *et al.*, 2009, Jackel *et al.*, 2013). The incongruences between the mites *COI* and ITS sequences obtained for some individuals in this study (see above), go in that direction. Second, intimate ecological associations, such as trophic interactions and plant feeding in particular, may provide horizontal transmission opportunities. It has been shown, for instance, for *Rickettsia* and *Wolbachia* in whiteflies (Caspi-Fluger *et al.*, 2012, Li *et al.*, 2017) and for *Cardinium* in leaf hoppers (Gonella *et al.*, 2015), and could explain the genetic diversity patterns observed here. Indeed, multiple events of horizontal transfer of *Cardinium* in the field would explain both its maintenance at low frequency within populations, the high sequence variation (*i.e.*, potential strains) found for this symbiont, as well as the lack of association with the host mtDNA. In addition, phylogenetic similarity of donor and recipient host species should facilitate horizontal transmission (Boivin *et al.*, 2014), and incongruences between phylogenies of host and parasites, suggesting occasional

horizontal transmission of *Cardinium* and/or *Wolbachia* between hosts, have been found at many occasions in spider mites (Yu *et al.*, 2011, Ros *et al.*, 2012, Zhang *et al.*, 2016b; but see Zhang *et al.*, 2013a). However, although these symbionts might have unexpected tolerance towards stressful conditions (e.g. *Wolbachia* can be maintained alive for at least a week in synthetic liquid medium out of the host cells; Rasgon *et al.*, 2006), the acquisition of a symbiont by horizontal transfer might be transient and not subsequently vertically transmitted in the new host species (Chiel *et al.*, 2009). Further studies should thus be conducted to determine the occurrence of both intra- and interspecific horizontal transfers of these endosymbionts under controlled laboratory conditions in spider mites.

In conclusion, this study highlights the richness and complexity of the symbiont community in spider mites. Although the factors affecting the spread and persistence of these endosymbionts in host populations remain largely unexplored, this system has the potential to become a model system in the study of host-symbiont interactions.

Authors' contributions

Experimental conception and design: FZ, OD, MW, SM, IO; field collections: FZ, IS, SM; molecular analyses: FZ, IS; statistical analyses: FZ; phylogenetic analyses: OD; paper writing: FZ, SM, with input from all authors. Funding: SM, IO. All authors have read and approved the final version of the manuscript.

Conflicts of interest

We declare that we do not have any conflicts of interest.

Funding

This work was funded by an FCT-ANR project (FCT-ANR//BIA-EVF/0013/2012) to SM and IO, and by an FCT-Tubitak project (FCT-TUBITAK/0001/2014) to SM and Ibrahim Cakmak. FZ was funded through an FCT Post-Doc fellowships (SFRH/BPD/125020/2016). Funding agencies did not participate in the design or analysis of experiments.

Acknowledgments

We are grateful to J. Moya-Laraño and F. Vavre for useful discussions and comments. We also thank D. Godinho, R. Ponce, and L. Rodrigues, for their help in spider-mite collection.

References

- Ahmed MZ, Ren SX, Mandour NS, Greeff JM & Qiu BL (2010) Prevalence of *Wolbachia* supergroups A and B in *Bemisia tabaci* (Hemiptera: Aleyrodidae) and some of its natural enemies. *J Econ Entomol* **103**: 1848-1859.
- Ahmed MZ, Li SJ, Xue X, Yin XJ, Ren SX, Jiggins FM, Greeff JM & Qiu BL (2015) The intracellular bacterium *Wolbachia* uses parasitoid wasps as phoretic vectors for efficient horizontal transmission. *Plos Pathogens* **11**: e1004672.
- Baldo L, Hotopp JCD, Jolley KA, Bordenstein SR, Biber SA, Choudhury RR, Hayashi C, Maiden MCJ, Tettelin H & Werren JH (2006) Multilocus sequence typing system for the endosymbiont *Wolbachia pipientis*. *Appl Environ Microbiol* **72**: 7098-7110.
- Ben-David T, Melamed S, Gerson U & Morin S (2007) ITS2 sequences as barcodes for identifying and analyzing spider mites (Acari : Tetranychidae). *Exp Appl Acarol* **41**: 169-181.
- Boivin T, Henri H, Vavre F, Gidoin C, Veber P, Candau JN, Magnoux E, Roques A & Auger-Rozenberg MA (2014) Epidemiology of asexuality induced by the endosymbiotic *Wolbachia* across phytophagous wasp species: host plant specialization matters. *Molecular Ecology* **23**: 2362-2375.
- Bolker BM (2008) *Ecological models and data in R* Princeton University Press, New Jersey.
- Boubou A, Migeon A, Roderick GK & Navajas M (2011) Recent emergence and worldwide spread of the red tomato spider mite, *Tetranychus evansi*: genetic variation and multiple cryptic invasions. *Biological Invasions* **13**: 81-92.
- Boubou A, Migeon A, Roderick GK, Auger P, Cornuet JM, Magalhães S & Navajas M (2012) Test of colonisation scenarios reveals complex invasion history of the red tomato spider mite *Tetranychus evansi*. *Plos One* **7**: e35601.
- Brady CM & White JA (2013) Cowpea aphid (*Aphis craccivora*) associated with different host plants has different facultative endosymbionts. *Ecological Entomology* **38**: 433-437.
- Brown AN & Lloyd VK (2015) Evidence for horizontal transfer of *Wolbachia* by a *Drosophila* mite. *Exp Appl Acarol* **66**: 301-311.
- Caspi-Fluger A, Inbar M, Mozes-Daube N, Katzir N, Portnoy V, Belausov E, Hunter MS & Zchori-Fein E (2012) Horizontal transmission of the insect symbiont *Rickettsia* is plant-mediated. *Proc Roy Soc B-Biol Sci* **279**: 1791-1796.

- Chiel E, Zchori-Fein E, Inbar M, Gottlieb Y, Adachi-Hagimori T, Kelly SE, Asplen MK & Hunter MS (2009) Almost there: transmission routes of bacterial symbionts between trophic levels. *Plos One* **4**: e4767.
- Clay K, Klyachko O, Grindle N, Civitello D, Oleske D & Fuqua C (2008) Microbial communities and interactions in the lone star tick, *Amblyomma americanum*. *Molecular Ecology* **17**: 4371-4381.
- Clayton KA, Gall CA, Mason KL, Scoles GA & Brayton KA (2015) The characterization and manipulation of the bacterial microbiome of the Rocky Mountain wood tick, *Dermacentor andersoni*. *Parasite Vector* **8**: 632.
- Clemente SH, Rodrigues LR, Ponce R, Varela SAM & Magalhães S (2016) Incomplete species recognition entails few costs in spider mites, despite first-male precedence. *Behav Ecol Sociobiol* **70**: 1161-1170.
- Clemente SH, Santos I, Ponce R, Rodrigues LR, Varela SAM & Magalhães S (2018) Despite reproductive interference, the net outcome of reproductive interactions among spider mite species is not necessarily costly. *Behavioral Ecology* **29**: 321-327.
- Crawley MJ (2007) *The R Book* John Wiley & Sons, Ltd, Chichester, England.
- Dergousoff SJ & Chilton NB (2010) Detection of a new *Arsenophonus*-type bacterium in Canadian populations of the Rocky Mountain wood tick, *Dermacentor andersoni*. *Exp Appl Acarol* **52**: 85-91.
- Duron O, Bouchon D, Boutin S, Bellamy L, Zhou LQ, Engelstadter J & Hurst GD (2008) The diversity of reproductive parasites among arthropods: *Wolbachia* do not walk alone. *BMC Biol* **6**: 27.
- Duron O, Binetruy F, Noël V, *et al.* (2017) Evolutionary changes in symbiont community structure in ticks. *Molecular Ecology* **26**: 2905-2921.
- Engelstadter J & Hurst GDD (2009) The ecology and evolution of microbes that manipulate host reproduction. *Annu Rev Ecol Evol Syst* **40**: 127-149.
- Engelstadter J, Telschow A & Hammerstein P (2004) Infection dynamics of different *Wolbachia*-types within one host population. *Journal of Theoretical Biology* **231**: 345-355.
- Enigl M & Schausberger P (2007) Incidence of the endosymbionts *Wolbachia*, *Cardinium* and *Spiroplasma* in phytoseiid mites and associated prey. *Exp Appl Acarol* **42**: 75-85.
- Excoffier L, Smouse PE & Quattro JM (1992) Analysis of molecular variance inferred from metric distances among DNA haplotypes: Application to human mitochondrial DNA restriction data. *Genetics* **131**: 479-491.

- Faria VG, Paulo TF & Sucena E (2016) Testing cannibalism as a mechanism for horizontal transmission of *Wolbachia* in *Drosophila*. *Symbiosis* **68**: 79-85.
- Ferragut F, Garzon-Luque E & Pekas A (2013) The invasive spider mite *Tetranychus evansi* (Acari: Tetranychidae) alters community composition and host-plant use of native relatives. *Exp Appl Acarol* **60**: 321-341.
- Ferrari J, Darby AC, Daniell TJ, Godfray HCJ & Douglas AE (2004) Linking the bacterial community in pea aphids with host-plant use and natural enemy resistance. *Ecological Entomology* **29**: 60-65.
- Frago E, Dicke M & Godfray HCJ (2012) Insect symbionts as hidden players in insect-plant interactions. *Trends in Ecology & Evolution* **27**: 705-711.
- Frank SA (1998) Dynamics of cytoplasmic incompatibility with multiple *Wolbachia* infections. *Journal of Theoretical Biology* **192**: 213-218.
- Gerth M, Rothe J & Bleidorn C (2013) Tracing horizontal *Wolbachia* movements among bees (Anthophila): a combined approach using multilocus sequence typing data and host phylogeny. *Molecular Ecology* **22**: 6149-6162.
- Gonella E, Pajoro M, Marzorati M, *et al.* (2015) Plant-mediated interspecific horizontal transmission of an intracellular symbiont in insects. *Sci Rep* **5**: 15811.
- Gotoh T, Noda H & Hong XY (2003) *Wolbachia* distribution and cytoplasmic incompatibility based on a survey of 42 spider mite species (Acari : Tetranychidae) in Japan. *Heredity* **91**: 208-216.
- Gotoh T, Noda H & Ito S (2007a) *Cardinium* symbionts cause cytoplasmic incompatibility in spider mites. *Heredity* **98**: 13-20.
- Gotoh T, Sugawara J, Noda H & Kitashima Y (2007b) *Wolbachia*-induced cytoplasmic incompatibility in Japanese populations of *Tetranychus urticae* (Acari : Tetranychidae). *Exp Appl Acarol* **42**: 1-16.
- Gotoh T, Araki R, Boubou A, Migeon A, Ferragut F & Navajas M (2009) Evidence of co-specificity between *Tetranychus evansi* and *Tetranychus takafujii* (Acari: Prostigmata, Tetranychidae): Comments on taxonomic and agricultural aspects. *Int J Acarol* **35**: 485-501.
- Gualtieri L, Nugnes F, Nappo AG, Gebiola M & Bernardo U (2017) Life inside a gall: closeness does not favour horizontal transmission of *Rickettsia* between a gall wasp and its parasitoid. *FEMS Microbiol Ecol* **93**: fix087.
- Guidolin AS & Consoli FL (2017) Symbiont diversity of *Aphis (Toxoptera) citricidus* (Hemiptera: Aphididae) as Influenced by host plants. *Microbial Ecology* **73**: 201-210.

570 Haine ER (2008) Symbiont-mediated protection. *Proc Roy Soc B-Biol Sci* **275**: 353-361.

571 Heath BD, Butcher RDJ, Whitfield WGF & Hubbard SF (1999) Horizontal transfer of
 572 *Wolbachia* between phylogenetically distant insect species by a naturally occurring
 573 mechanism. *Current Biology* **9**: 313-316.

574 Hellard E, Fouchet D, Vavre F & Pontier D (2015) Parasite-parasite interactions in the wild:
 575 How to detect them? *Trends Parasitol* **31**: 640-652.

576 Hoy MA & Jeyaprakash A (2005) Microbial diversity in the predatory mite *Metaseiulus*
 577 *occidentalis* (Acari : Phytoseiidae) and its prey, *Tetranychus urticae* (Acari :
 578 Tetranychidae). *Biol Control* **32**: 427-441.

579 Huigens ME, de Almeida RP, Boons PAH, Luck RF & Stouthamer R (2004) Natural
 580 interspecific and intraspecific horizontal transfer of parthenogenesis-inducing *Wolbachia*
 581 in *Trichogramma* wasps. *Proc Roy Soc Lond B Bio* **271**: 509-515.

582 Hurtado MA, Ansaloni T, Cros-Arteil S, Jacas JA & Navajas M (2008) Sequence analysis of
 583 the ribosomal internal transcribed spacers region in spider mites (Prostigmata:
 584 Tetranychidae) occurring in citrus orchards in Eastern Spain: use for species
 585 discrimination. *Ann Appl Biol* **153**: 167-174.

586 Jackel R, Mora D & Dobler S (2013) Evidence for selective sweeps by *Wolbachia* infections:
 587 phylogeny of Altica leaf beetles and their reproductive parasites. *Molecular Ecology* **22**:
 588 4241-4255.

589 Jaenike J, Polak M, Fiskin A, Helou M & Minhas M (2007) Interspecific transmission of
 590 endosymbiotic *Spiroplasma* by mites. *Biology Letters* **3**: 23-25.

591 Ji HL, Qi LD, Hong XY, Xie HF & Li YX (2015) Effects of host sex, plant species, and
 592 putative host species on the prevalence of *Wolbachia* in natural populations of *Bemisia*
 593 *tabaci* (Hemiptera: Aleyrodidae): A modified nested PCR study. *J Econ Entomol* **108**:
 594 210-218.

595 Li SJ, Ahmed MZ, Lv N, Shi PQ, Wang XM, Huang JL & Qiu BL (2017) Plant-mediated
 596 horizontal transmission of *Wolbachia* between whiteflies. *Isme Journal* **11**: 1019-1028.

597 Liu Y, Miao H & Hong XY (2006) Distribution of the endosymbiotic bacterium *Cardinium*
 598 in Chinese populations of the carmine spider mite *Tetranychus cinnabarinus* (Acari :
 599 Tetranychidae). *J Appl Entomol* **130**: 523-529.

600 Marzorati M, Alma A, Sacchi L, *et al.* (2006) A novel bacteroidetes symbiont is localized in
 601 *Scaphoideus titanus*, the insect vector of flavescence doree in *Vitis vinifera*. *Applied and*
 602 *Environmental Microbiology* **72**: 1467-1475.

- Migeon A & Dorkeld F (2006-2017) Spider Mites Web: a comprehensive database for the Tetranychidae. <http://www.montpellier.inra.fr/CBGP/spmweb/>.
- Moran NA, McCutcheon JP & Nakabachi A (2008) Genomics and evolution of heritable bacterial symbionts. *Annual Review of Genetic* **42**: 165-190.
- Mouton L, Henri H, Charif D, Bouletrea M & Vavre F (2007) Interaction between host genotype and environmental conditions affects bacterial density in *Wolbachia* symbiosis. *Biol Lett* **3**: 210-213.
- Nakamura Y, Kawai S, Yukuhiro F, Ito S, Gotoh T, Kisimoto R, Yanase T, Matsumoto Y, Kageyama D & Noda H (2009) Prevalence of *Cardinium* bacteria in planthoppers and spider mites and taxonomic revision of "candidatus *Cardinium hertigii*" based on detection of a new *Cardinium* group from biting midges. *Applied and Environmental Microbiology* **75**: 6757-6763.
- Navajas M, Lagnel J, Gutierrez J & Boursot P (1998) Species-wide homogeneity of nuclear ribosomal ITS2 sequences in the spider mite *Tetranychus urticae* contrasts with extensive mitochondrial COI polymorphism. *Heredity* **80**: 742-752.
- Nguyen DT, Spooner-Hart RN & Riegler M (2016) Loss of *Wolbachia* but not *Cardinium* in the invasive range of the Australian thrips species, *Pezothrips kellyanus*. *Biological Invasions* **18**: 197-214.
- Nguyen DT, Morrow JL, Spooner-Hart RN & Riegler M (2017) Independent cytoplasmic incompatibility induced by *Cardinium* and *Wolbachia* maintains endosymbiont coinfections in haplodiploid thrips populations. *Evolution* **71**: 995-1008.
- Pan HP, Chu D, Liu BM, Xie W, Wang SL, Wu QJ, Xu BY & Zhang YJ (2013) Relative Amount of Symbionts in Insect Hosts Changes with Host-Plant Adaptation and Insecticide Resistance. *Environ Entomol* **42**: 74-78.
- Perlman SJ, Kelly SE & Hunter MS (2008) Population biology of cytoplasmic incompatibility: maintenance and spread of *Cardinium* symbionts in a parasitic wasp. *Genetics* **178**: 1003-1011.
- Rasgon JL, Gamston CE & Ren XX (2006) Survival of *Wolbachia pipientis* in cell-free medium. *Appl Environ Microbiol* **72**: 6934-6937.
- Raychoudhury R, Baldo L, Oliveira D & Werren JH (2009) Modes of acquisition of *Wolbachia*: horizontal transfer, hybrid introgression, and codivergence in the *Nasonia* species complex. *Evolution* **63**: 165-183.
- Reuter M, Pedersen JS & Keller L (2005) Loss of *Wolbachia* infection during colonisation in the invasive Argentine ant *Linepithema humile*. *Heredity* **94**: 364-369.

637 Ros VID & Breeuwer JAJ (2009) The effects of, and interactions between, *Cardinium* and
638 *Wolbachia* in the doubly infected spider mite *Bryobia sarothamni*. *Heredity* **102**: 413-
639 422.

640 Ros VID, Fleming VM, Feil EJ & Breeuwer JAJ (2012) Diversity and recombination in
641 *Wolbachia* and *Cardinium* from *Bryobia* spider mites. *BMC Microbiol* **12**: S13.

642 Sato Y, Alba JM & Sabelis MW (2014) Testing for reproductive interference in the
643 population dynamics of two congeneric species of herbivorous mites. *Heredity* **113**: 495-
644 502.

645 Schilthuizen M & Stouthamer R (1997) Horizontal transmission of parthenogenesis-inducing
646 microbes in *Trichogramma* wasps. *Proc Roy Soc B-Biol Sci* **264**: 361-366.

647 Shoemaker DD, Ross KG, Keller L, Vargo EL & Werren JH (2000) *Wolbachia* infections in
648 native and introduced populations of fire ants (*Solenopsis* spp.). *Insect Mol Biol* **9**: 661-
649 673.

650 Staudacher H, Schimmel BCJ, Lamers MM, Wybouw N, Groot AT & Kant MR (2017)
651 Independent effects of a herbivore's bacterial symbionts on its performance and induced
652 plant defences. *Int J Mol Sci* **18**: 182.

653 Suh E, Sim C, Park J-J & Cho K (2015) Inter-population variation for *Wolbachia* induced
654 reproductive incompatibility in the haplodiploid mite *Tetranychus urticae*. *Exp Appl*
655 *Acarol* **65**: 55-71.

656 Tamura K, Peterson D, Peterson N, Stecher G, Nei M & Kumar S (2011) MEGA5: Molecular
657 Evolutionary Genetics Analysis using maximum likelihood, evolutionary distance, and
658 maximum parsimony methods. *Molecular Biology and Evolution* **28**: 2731-2739.

659 Van Opijnen T & Breeuwer JAJ (1999) High temperatures eliminate *Wolbachia*, a
660 cytoplasmic incompatibility inducing endosymbiont, from the two-spotted spider mite.
661 *Exp Appl Acarol* **23**: 871-881.

662 Vautrin E & Vavre F (2009) Interactions between vertically transmitted symbionts:
663 cooperation or conflict? *Trends in Microbiology* **17**.

664 Vautrin E, Genieys S, Charles S & Vavre F (2008) Do vertically transmitted symbionts co-
665 existing in a single host compete or cooperate? A modelling approach. *J Evol Biol* **21**:
666 145-161.

667 Vavre F, Fleury F, Lepetit D, Fouillet P & Bouletreau M (1999) Phylogenetic evidence for
668 horizontal transmission of *Wolbachia* in host-parasitoid associations. *Molecular Biology*
669 *and Evolution* **16**: 1711-1723.

- Weeks AR, Velten R & Stouthamer R (2003) Incidence of a new sex-ratio-distorting endosymbiotic bacterium among arthropods. *Proc Roy Soc B-Biol Sci* **270**: 1857-1865.
- Weinert LA, Araujo-Jnr EV, Ahmed MZ & Welch JJ (2015) The incidence of bacterial endosymbionts in terrestrial arthropods. *Proc Roy Soc London* **282**: 20150249.
- Werren JH, Baldo L & Clark ME (2008) *Wolbachia*: master manipulators of invertebrate biology. *Nat Rev Microbiol* **6**: 741-751.
- White JA, Kelly SE, Perlman SJ & Hunter MS (2009) Cytoplasmic incompatibility in the parasitic wasp *Encarsia inaron*: disentangling the roles of *Cardinium* and *Wolbachia* symbionts. *Heredity* **102**: 483-489.
- Wilkinson TL, Koga R & Fukatsu T (2007) Role of host nutrition in symbiont regulation: Impact of dietary nitrogen on proliferation of obligate and facultative bacterial endosymbionts of the pea aphid *Acyrtosiphon pisum*. *Applied and Environmental Microbiology* **73**: 1362-1366.
- Xie RR, Chen XL & Hong XY (2011) Variable fitness and reproductive effects of *Wolbachia* infection in populations of the two-spotted spider mite *Tetranychus urticae* Koch in China. *Appl Entomol Zool* **46**: 95-102.
- Xie RR, Sun JT, Xue XF & Hong XY (2016) Cytoplasmic incompatibility and fitness benefits in the two-spotted spider mite *Tetranychus urticae* (red form) doubly infected with *Wolbachia* and *Cardinium*. *Syst Appl Acarol* **21**: 1161-1173.
- Yu MZ, Zhang KJ, Xue XF & Hong XY (2011) Effects of *Wolbachia* on mtDNA variation and evolution in natural populations of *Tetranychus urticae* Koch. *Insect Mol Biol* **20**: 311-321.
- Zchori-Fein E & Perlman SJ (2004) Distribution of the bacterial symbiont *Cardinium* in arthropods. *Molecular Ecology* **13**: 2009-2016.
- Zélé F, Vézilier J, L'Ambert G, Nicot A, Gandon S, Rivero A & Duron O (2014) Dynamics of prevalence and diversity of avian malaria infections in wild *Culex pipiens* mosquitoes: the effects of *Wolbachia*, filarial nematodes and insecticide resistance. *Parasite Vector* **7**: 437.
- Zhang YK, Ding XL, Zhang KJ & Hong XY (2013a) *Wolbachia* play an important role in affecting mtDNA variation of *Tetranychus truncatus* (Trombidiformes: Tetranychidae). *Environ Entomol* **42**: 1240-1245.
- Zhang YK, Chen YT, Yang K & Hong XY (2016a) A review of prevalence and phylogeny of the bacterial symbiont *Cardinium* in mites (subclass: Acari). *Syst Appl Acarol* **21**: 978-990.

704 Zhang YK, Chen YT, Yang K, Qiao GX & Hong XY (2016b) Screening of spider mites
 705 (Acari: Tetranychidae) for reproductive endosymbionts reveals links between co-
 706 infection and evolutionary history. *Sci Rep* **6**: 27900.

707 Zhang YK, Zhang KJ, Sun JT, Yang XM, Ge C & Hong XY (2013b) Diversity of *Wolbachia*
 708 in natural populations of spider mites (genus Tetranychus): Evidence for complex
 709 infection history and disequilibrium distribution. *Microb Ecol* **65**: 731-739.

710 Zhao DX, Zhang XF & Hong XY (2013a) Host-symbiont interactions in spider mite
 711 *Tetranychus truncates* doubly infected with *Wolbachia* and *Cardinium*. *Environ Entomol*
 712 **42**: 445-452.

713 Zhao DX, Chen DS, Ge C, Gotoh T & Hong XY (2013b) Multiple Infections with *Cardinium*
 714 and two strains of *Wolbachia* in the spider mite *Tetranychus phaselus* Ehara: Revealing
 715 new forces driving the spread of *Wolbachia*. *Plos One* **8**: e54964.

716 Zhu LY, Zhang KJ, Zhang YK, Ge C, Gotoh T & Hong XY (2012) *Wolbachia* strengthens
 717 *Cardinium*-induced cytoplasmic incompatibility in the spider mite *Tetranychus piercei*
 718 McGregor. *Curr Microbiol* **65**: 516-523.

Table 1. Tetranychidae mite populations collected in Portugal (P), France (F) and Spain (S) from September to December 2013.

| Name | Species | Collection date | Location | Coordinates | Host plant |
|-------|---------------------|-----------------|----------------------------|----------------------|-----------------------------|
| FR | <i>T. urticae</i> * | 11/10/2013 | Montpellier (F) | 43.614951, 3.859846 | <i>Solanum lycopersicum</i> |
| AlBe | | 09/11/2013 | Almería (S) | 36.855725, -2.320374 | <i>Solanum melongena</i> |
| AlRo | | 09/11/2013 | Almería (S) | 36.855725, -2.320374 | <i>Rosa spp.</i> |
| DC | | 10/09/2013 | S. Domingos (P) | 39.058742, -9.135427 | <i>Cucurbita pepo</i> |
| DF | | 10/09/2013 | S. Domingos (P) | 39.058742, -9.135427 | <i>Phaseolus vulgaris</i> |
| COL | | 08/09/2013 | Colares (P) | 38.799517, -9.448335 | <i>Phaseolus vulgaris</i> |
| LOU | | 03/10/2013 | Lourinhã (P) | 39.248145, -9.276321 | <i>Solanum melongena</i> |
| CH | | 31/10/2013 | Casal Hortelão (P) | 38.851962, -9.393918 | <i>Solanum lycopersicum</i> |
| RF | | 04/11/2013 | Ribeira de Fráguas (P) | 39.366415, -8.851037 | <i>Solanum lycopersicum</i> |
| AMP | | 18/11/2013 | Aldeia da Mata Pequena (P) | 38.534363, -9.191163 | <i>Datura stramonium</i> |
| GH | <i>T. evansi</i> | 03/10/2013 | University of Lisbon (P) | 38.757852, -9.158221 | <i>Solanum lycopersicum</i> |
| QL | | 26/11/2013 | Quinta das Lameiras (P) | 39.085837, -8.991478 | <i>Datura stramonium</i> |
| GRA | | 03/12/2013 | Graça, Lisbon (P) | 38.71392, -9.129537 | <i>Solanum lycopersicum</i> |
| Assaf | <i>T. ludeni</i> | 20/09/2013 | Assafora (P) | 38.904743, -9.408592 | <i>Datura stramonium</i> |
| CVM | | 03/10/2013 | Casal Vale do Medo (P) | 39.248450, -9.294550 | <i>Ipomoea purpurea</i> |
| Alval | | 05/11/2013 | Alvalade, Lisbon (P) | 38.75515, -9.14685 | <i>Ipomoea purpurea</i> |

*red form (or *T. cinnabarinus*)

Table 2. Mitochondrial *COI* alleles in spider mite populations. A total of 75 sequences were obtained from individual female spider mites. For each allele, we give the genbank accession number and the corresponding number of mites per population. We also give the total number of mites carrying each allele and the corresponding proportion relative to the total number of mites.

| Species | Genbank accession number [Ref] | <i>T. ludeni</i> | | | <i>T. evansi</i> | | | <i>T. urticae</i> | | | | | | | | | | Total number (proportion) |
|-------------------|--------------------------------|------------------|-----|-------|------------------|----|----|-------------------|------|----|-----|-----|-----|----|----|----|----|---------------------------|
| | | Assaf | CVM | Alval | GRA | GH | QL | AlRo | AlBe | DF | LOU | COL | AMP | RF | DC | CH | FR | |
| <i>T. ludeni</i> | GQ141911 [*] | 3 | 5 | 5 | - | - | - | - | - | - | - | - | - | - | - | - | - | 13 (0.17) |
| <i>T. evansi</i> | MF428442 [**] | - | - | - | 3 | 4 | 1 | - | - | - | - | - | - | - | - | - | - | 8 (0.11) |
| | MF428443 [**] | - | - | - | - | - | 3 | - | - | - | - | - | - | - | - | - | - | 3 (0.04) |
| <i>T. urticae</i> | HM486513 [49] | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | 1 | 1 (0.01) |
| | HM565899 [96] | - | - | - | 1 | 1 | - | - | - | - | - | - | - | - | - | - | 1 | 3 (0.04) |
| | HM565901 [96] | - | - | - | - | - | - | 5 | 5 | - | - | - | - | - | - | - | - | 10 (0.13) |
| | MF428440 [**] | 1 | - | - | 1 | - | 1 | - | - | 4 | 4 | 4 | 4 | 5 | 4 | 5 | 3 | 36 (0.48) |
| | MF428441 [**] | - | - | - | - | - | - | - | - | - | - | - | - | - | 1 | - | - | 1 (0.01) |

* unpublished

** this study

Table 3. Occurrence and co-occurrence of *Wolbachia wsp*, *Cardinium 16S rRNA* and *Rickettsia gtlA* alleles. A total of 72 sequences were obtained from 80 positive samples through PCR. For each allele, we give the genbank accession number, the corresponding number of infected mites per population, and indicate the symbiont species involved in a multiple infection with superscripts (a: *Cardinium*, b: *Wolbachia*, and c: *Rickettsia*). For each allele, we also give the total number of infected mites and the corresponding proportion relative to the total number of mites infected by the same endosymbiont.

| Gene (fragment length) | Genbank accession numbers [Ref] | <i>T. ludeni</i> | | <i>T. evansi</i> | | | <i>T. urticae</i> | | | | | | | | | | Total number (proportion) |
|--|---------------------------------|-------------------|-------|------------------|-------------------|---------------------|-------------------|-------------------|----|-----|-------------------|-------------------|-------------------|-------------------|---------------------|---------------------|---------------------------|
| | | CVM | Alval | GRA | GH | QL | AlRo | AlBe | DF | LOU | COL | AMP | RF | DC | CH | FR | |
| <i>Wolbachia wsp</i> (524bp) | DQ910771 [72] | 3 ^(1a) | 3 | 1 | 3 ^(1a) | 3 ^(1a1c) | 2 ^(1a) | 2 ^(1a) | 3 | 3 | 3 ^(1a) | 3 ^(2a) | 3 ^(2a) | 3 ^(2a) | 3 ^(2a1c) | 3 ^(1a1c) | 41 (0.98) |
| | MF428430 [*] | - | - | 1 | - | - | - | - | - | - | - | - | - | - | - | - | 1 (0.02) |
| <i>Cardinium 16S rRNA</i> (325bp) | MF428431 [*] | - | - | - | - | - | 1 | - | - | - | - | - | - | - | 1 ^b | 1 ^b | 3 (0.11) |
| | MF428432 [*] | 1 ^{b,c} | - | - | - | - | - | - | - | - | - | - | - | - | - | - | 1 (0.04) |
| | MF428433 [*] | - | - | 1 | - | - | 1 | - | - | - | - | 1 ^b | - | - | - | - | 3 (0.11) |
| | MF428434 [*] | - | 2 | - | 2 ^(2b) | - | 1 ^b | 2 ^(1b) | - | - | - | - | 4 ^(4b) | 3 ^(2b) | 1 ^b | 2 ^(2b) | 17 (0.63) |
| | MF428435 [*] | 1 ^b | - | - | - | - | - | - | - | - | - | - | - | - | - | - | 1 (0.04) |
| | MF428436 [*] | - | - | - | - | - | - | - | - | - | - | - | 1 ^b | - | - | - | 1 (0.04) |
| | MF428437 [*] | - | - | - | - | 1 ^b | - | - | - | - | - | - | - | - | - | - | 1 (0.04) |
| <i>Rickettsia gtlA</i> (431bp) | MF428438 [*] | - | - | - | - | 1 ^b | - | - | - | - | - | - | - | - | - | 1 ^b | 2 (1.00) |

* this study

Table 4. Co-occurrence of *Cardinium 16S rRNA* and mitochondrial *COI* alleles of spider mites. *Cardinium 16S rRNA* and mite *COI* sequences were obtained from a total of 29 *Cardinium*-infected individual female spider mites. For each *Cardinium 16S rRNA* allele, we give the corresponding number of infected mites per *COI* allele (Genbank accession numbers [Ref]), and the species to which they belong between brackets. Te: *Tetranychus evansi*, Tl: *T. ludeni*, Tu: *T. urticae*.

| <i>Cardinium</i> 16S rRNA | Mites' COI | | | | | | |
|---------------------------|---------------|---------------|---------------|--------------|--------------|--------------|---------------|
| | HM486513 [49] | HM565899 [96] | HM565901 [96] | MF428440 [*] | MF428441 [*] | MF428442 [*] | GQ141911 [**] |
| MF428431 [*] | - | 1 (Tu) | 1 (Tu) | - | - | - | - |
| MF428432 [*] | - | - | - | - | - | - | 1 (Tl) |
| MF428433 [*] | - | - | 1 (Tu) | 2 (Tu) | - | 1 (Te) | - |
| MF428434 [*] | 1 (Tu) | 2 (Te-Tu) | 3 (Tu) | 9 (Tu) | 1 (Tu) | 1 (Te) | 2 (Tl) |
| MF428435 [*] | - | - | - | - | - | - | 1 (Tl) |
| MF428436 [*] | - | - | - | 1 (Tu) | - | - | - |
| MF428437 [*] | - | - | - | 1 (Te) | - | - | - |

* this study

** unpublished




Figure 1. Phylogeny and distribution of *Wolbachia* and *Cardinium* (co)infections among *T. evansi*, *T. ludeni* and *T. urticae* mitochondrial *COI* haplotypes. The left panel shows the phylogeny of the 8 mtDNA haplotypes via maximum likelihood. Numbers on branches indicate percentage bootstrap support for major branches (1000 replicates). GenBank accession numbers are given between brackets. The scale bar is in units of substitutions/site. The right panel gives the distribution of symbiont infection observed with each mtDNA haplotype (black: *Wolbachia*-infected, n = 20; grey: *Cardinium*-infected, n = 7; dashed: *Wolbachia*-*Cardinium* coinfecting individuals, n = 20; white: uninfected specimens, n = 28).




Figure 2. Prevalence of endosymbionts for each spider-mite population of (A) *T. ludeni*, (B) *T. evansi*, and (C) *T. urticae*. Each graph represents a population, in which the rows correspond to individual mites and the columns to their infection status by W: *Wolbachia*; C: *Cardinium*; and R: *Rickettsia*. White cells: uninfected; Grey cells: infected; Hatched cells: *Cardinium* and *Rickettsia* failed sequences but positive through PCR. None of the tested colonies were found infected by *Spiroplasma* or *Arsenophonus*.




Figure 3. Mixed logistic regression of individual probability of *Wolbachia* infection on the probability of *Cardinium* infection. The solid line represents the mean predicted values; the dark shaded area represents the range of the 95% confidence intervals based on the fixed-effects uncertainty only, while the light shaded area also includes the variance of a random factor “population”. The observed data at the individual level are not represented as they all take the values ‘0’ or ‘1’ for both *Wolbachia* and *Cardinium* infection. Open circles are the observed mean prevalence of *Cardinium* per population plotted against the observed mean prevalence of *Wolbachia*, while filled circles are their corresponding linear unbiased predictors.

Additional file 1: Electronic supplementary materials (Tables S1 to S4).

Additional file 2: Complete dataset for the inter- and intraspecific variations of endosymbiont prevalences.

Additional file 3: Complete *COI* sequences dataset. Sequences name: species_population_*COI* GenBank accession number_*Wolbachia* infection status (W+: infected; W-: uninfected)_ *Cardinium* infection status (C+: infected; C-: uninfected); *Cardinium 16S rRNA* GenBank accession number for infected individuals.

Additional file 4: Complete *Cardinium 16S rRNA* sequences dataset. Sequences name: species_population_*COI* GenBank accession number_*Wolbachia* infection status (W+: infected; W-: uninfected)_ *Cardinium* infection status (C+: infected; C-: uninfected); *Cardinium 16S rRNA* GenBank accession number for infected individuals.