



HAL
open science

Strong disorder RG approach – a short review of recent developments

Ferenc Iglói, Cécile Monthus

► **To cite this version:**

Ferenc Iglói, Cécile Monthus. Strong disorder RG approach – a short review of recent developments. Eur.Phys.J.B, 2018, 91 (11), pp.290. <10.1140/epjb/e2018-90434-8>. <hal-01952677>

HAL Id: hal-01952677

<https://hal.science/hal-01952677v1>

Submitted on 11 Jan 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

Strong Disorder RG approach - a short review of recent developments

Ferenc Iglói*

Wigner Research Centre for Physics, Institute for Solid State Physics and Optics, H-1525 Budapest, P.O. Box 49, Hungary and
Institute of Theoretical Physics, Szeged University, H-6720 Szeged, Hungary

Cécile Monthus†

Institut de Physique Théorique, Université Paris Saclay, CNRS, CEA, 91191 Gif-sur-Yvette, France

(Dated: November 26, 2018)

CONTENTS

I. Introduction	2	D. Relations between SDRG and Entanglement-Algorithms	10
II. Random Quantum Ising Model in various dimensions d	2	VI. Localized and Many-Body-Localized Phases of quantum spin chains	11
A. Strong Disorder RG rules	2	A. RSRG-X for excited eigenstates	11
B. Solution in one dimension - a reminder[3]	3	B. RSRG-t for the unitary dynamics	11
1. Critical point - infinite disorder fixed point	3	C. Non-equilibrium dynamical scaling of observables	12
2. Griffiths phases - strong disorder fixed points	3	D. Comparison with other RG procedures existing in the field of Many-Body-Localization	13
3. Consequences for the ground-state wavefunction	3	VII. Floquet dynamics of periodically driven chains in their localized phases	13
C. Higher dimensions $d > 1$	4	VIII. Open dissipative quantum spin chains	13
D. Approximate RG methods	5	A. Quantum spin chains coupled to a bath of quantum oscillators	13
III. Random quantum systems with long-range interactions	5	B. Lindblad dynamics for random quantum spin chains	14
A. 1D - solution of the primary model [36]	5	IX. Anderson localization models	14
B. 3D - numerical Strong Disorder RG study	6	X. Random contact process	14
C. Other quantum models with LR interactions	6	A. Strong disorder RG rules	15
1. Ising chains with other type of LR interactions	6	B. Long range spreading	15
2. Random Heisenberg chains with LR interactions	7	C. Temporal disorder	15
IV. Other random quantum models	7	XI. Classical master equations	16
A. Models with discrete symmetry	7	A. Real-space renormalization for random walks in random media	16
1. Antiferromagnetic random quantum Ising chain	7	B. RG in configuration-space for the dynamics of classical many-body models	16
2. Ashkin-Teller chains	7	XII. Random classical oscillators	17
B. Models with continuous symmetry	7	A. Random elastic networks	17
C. Disordered non-abelian anyonic chains	8	B. Synchronisation of interacting non-linear dissipative classical oscillators	17
D. Superfluid-insulator transition	8	XIII. Other classical models	17
E. Superconductor-Metal transition	9	A. Equilibrium properties of random systems	17
F. The Rainbow Spin Chain	9	B. Extremes of stochastic processes	18
V. Entanglement properties	9	XIV. Conclusion	18
A. Random quantum chains	9	Acknowledgments	18
B. RTIM in higher dimensions	10	References	18
C. Bond diluted quantum Ising model	10		

* igloi.ferenc@wigner.mta.hu

† cecile.monthus@cea.fr

I. INTRODUCTION

The Strong Disorder Renormalization Group (SDRG) approach has been introduced by Ma, Dasgupta and Hu [1] and later further developed by D. Fisher [2] to study the low-energy excitations and spatial and temporal correlations of random quantum spin chains. In these systems, the critical properties are controlled by so called Infinite-Disorder-Fixed-Points (IDFPs), at which disorder fluctuations dominate over quantum fluctuations, so that the calculated properties (critical exponents and scaling functions) are asymptotically exact. Soon after Fisher's results, the RG approach has been applied to a large number of disordered systems, either quantum (in one and higher dimensions) or classical (Sinai random walk, classical random spin chains and polymers, stochastic systems with quenched disorder, etc.). For the models where exact results are available via other approaches, the RG method has been not only able to reproduce them correctly, but has allowed in addition to predict many new critical exponents and scaling functions. All the developments that have occurred before 2005 have been already summarised in our Review [3]. The goal of the present colloquium paper is thus to give an overview of the various developments since 2005. We will stress the physical ideas and the SDRG-way-of-thinking for each type of problem, but we will usually omit the detailed derivation of results that can be found in the original papers.

This colloquium paper is organized as follows. We begin with the ground state properties of random quantum systems, such as the random transverse-field Ising model in different dimensions d in Section II, the effects of Long-Ranged interactions in Section III, as well as various other quantum models (antiferromagnets, Ashkin-Teller models, anyonic models, the superfluid-insulator transition) in Section IV. The scaling properties of the entanglement in critical ground states are described in Section V. We then focus on the dynamical properties of Localized and Many-Body-Localized models within three different settings, namely the unitary dynamics of isolated models in Section VI, the Floquet dynamics of periodically driven models in Section VII, and the dissipative dynamics of open quantum models in Section VIII. Section IX is then devoted to Anderson Localization tight-binding models for electrons. We then turn to classical disordered models, with the random contact process for epidemic spreading in section X, the renormalization of general master equations with randomness in section XI, the dynamics of random classical oscillators with or without dissipation in section XII, as well as some other miscellaneous topics in section XIII.

II. RANDOM QUANTUM ISING MODEL IN VARIOUS DIMENSIONS d

The quantum Ising model with random couplings and/or with random transverse fields (RTIM) is the prototype of disordered quantum magnets having discrete symmetry. The model is defined by the Hamiltonian:

$$\mathcal{H} = -\frac{1}{2} \sum_{ij} J_{ij} \sigma_i^x \sigma_j^x - \frac{1}{2} \sum_i h_i \sigma_i^z, \quad (1)$$

where the $\sigma_i^{x,z}$ are Pauli-matrices and i, j denote sites of a lattice. In Eq.(1) the couplings J_{ij} and the transverse fields h_i are independent random variables, which are taken from the distributions, $p(J)$ and $q(h)$, respectively. In the following, we will discuss the case of ferromagnetic models $J_{ij} \geq 0$, where the order parameter is the magnetization, but it should be stressed that this is not restrictive : the SDRG approach has been applied as well to the spin-glass case where the sign of the couplings is also random. The numerical computations are usually done either with two box-distributions $p(J) = \Theta(J)\Theta(1-J)$ and $q(h) = \frac{1}{h_b} \Theta(h)\Theta(h_b-h)$ ($\Theta(x)$ being the Heaviside step-function), or with the fixed-h distribution $q(h) = \delta(h-h_f)$, while $p(J)$ follows the box-distribution as before. The quantum control parameter is defined as $\theta = \overline{\ln h} - \overline{\ln J}$, where \overline{x} stands for the average value of the variable x over quench disorder. In any dimension the RTIM has a zero-temperature quantum phase transition, located at some θ_c , separating a paramagnetic disordered ($\theta > \theta_c$) and a ferromagnetic ordered phase ($\theta < \theta_c$).

The simplest geometry that one can consider is the one-dimensional chain with nearest-neighbour couplings, which has been solved analytically through the Strong Disorder RG method by Fisher [2], as described in details in Review [3]. Later, more general geometries have been considered, such as ladders; star-like objects; hypercubic lattices in dimensions $d = 2, 3, 4$; Bethe lattices, Erdős-Rényi random graphs and other type of complex networks, as well as systems with long-range interactions. Besides the bulk quantities, the critical behaviour of several observables at surfaces, edges, wedges and cones have also been calculated.

A. Strong Disorder RG rules

The basic steps of the renormalization procedure are identical in every geometry: at each step the largest term of the Hamiltonian in Eq.(1), denoted by Ω , is eliminated and new terms are generated through a second-order perturbation calculation between the remaining degrees of freedom.

i) If the largest term is a coupling, say $J_{ij} = \Omega$, then the two connected sites, i and j are coupled to form a cluster. This spin-cluster then perceives an effective transverse

field of strength

$$\tilde{h}_{ij} \approx \frac{h_i h_j}{J_{ij}}, \quad (2)$$

and the magnetic moments transform additively: $\mu_{ij} = \mu_i + \mu_j$, while originally $\mu_i = \mu_j = 1$.

ii) If the largest term is a transverse field, say $h_i = \Omega$, then this site has negligible contribution to the (longitudinal) susceptibility, and is therefore decimated out. At the same time, new effective couplings are generated between all sites, say j and k , which were nearest neighbours of i . The new contributions to the couplings are given by: $\tilde{J}_{jk} \approx \frac{J_{ij} J_{ik}}{h_i}$.

In higher dimensions $d > 1$, the topology of the lattice is modified during the renormalization and often the new contribution $\tilde{J}_{jk} \approx \frac{J_{ij} J_{ik}}{h_i}$ concerns a pair of sites that were already coupled via some coupling J_{jk} . In this case, one can use their sum (*sum rule*), or their maximum value (*maximum rule*) as the renormalized coupling

$$\tilde{J}_{jk} \approx \max \left[J_{jk}, \frac{J_{ij} J_{ik}}{h_i} \right]. \quad (3)$$

The maximum rule is justified at an IDFP, furthermore the numerical algorithms are more efficient in this case, as explained in Refs.4-7. If, however, the critical behaviour is controlled by a conventional random fixed point, such as in the superfluid-insulator transition (see section IVD), the sum rule can lead to more accurate results [8].

B. Solution in one dimension - a reminder[3]

In one dimension with nearest neighbour couplings, the topology does not change during renormalization. The cut-off (Ω) dependence of the distribution functions (couplings, transverse fields, lengths, moments, etc) is written in the form of integro-differential equations, which are solved analytically. At the fixed point $\Omega \rightarrow 0$, the distribution of transverse fields and that of the couplings are given in the form:

$$P_0(h, \Omega) = \frac{p_0(\Omega)}{\Omega} \left(\frac{\Omega}{h} \right)^{1-p_0(\Omega)} \quad (4)$$

$$R_0(J, \Omega) = \frac{r_0(\Omega)}{\Omega} \left(\frac{\Omega}{J} \right)^{1-r_0(\Omega)}, \quad (5)$$

1. Critical point - infinite disorder fixed point

At the critical point where the decimation of couplings and transverse fields are symmetric, we have

$$p_0 = r_0 = \frac{1}{\ln(\Omega_0/\Omega)}, \quad (6)$$

where Ω_0 is a reference energy scale. This is an infinite disorder fixed point, since the ratio of typical couplings and transverse fields is going to zero or to infinity. Therefore the decimation steps are asymptotically correct and consequently the calculated critical properties are asymptotically exact. At this fixed point the relation between length-scale, L , and energy-scale, Ω is given by the activated scaling

$$\ln \left(\frac{\Omega_0}{\Omega} \right) \sim L^\psi, \quad \psi = 1/2, \quad (7)$$

and the average moment of clusters scales as:

$$\bar{\mu} = \bar{\mu}_0 \left[\ln \left(\frac{\Omega_0}{\Omega} \right) \right]^\phi, \quad \phi = \frac{1 + \sqrt{5}}{2}. \quad (8)$$

leading to the fractal dimension d_f

$$\bar{\mu} \sim L^{d_f}, \quad d_f = \psi\phi = \frac{1 + \sqrt{5}}{4}. \quad (9)$$

2. Griffiths phases - strong disorder fixed points

Outside the critical point, the decimation of the couplings and the transverse fields is asymmetric, which is characterized by the parameter $\Delta = (p_0 - r_0)/2$. Δ is expressed in terms of the original distributions as:

$$\left[\left(\frac{J^2}{h^2} \right)^\Delta \right]_{\text{av}} = 1, \quad (10)$$

and close to the critical point: $\Delta \sim \theta$, since $\theta_c = 0$. In the paramagnetic or disordered Griffiths phase, almost exclusively transverse fields are decimated out and the solution close to the line of fixed points, i.e. in the limit $\Omega/\Omega_0 \ll 1$ is given by:

$$p_0 = 2\Delta, \quad r_0 \sim (\Omega/\Omega_0)^{2\Delta}. \quad (11)$$

In the ferromagnetic or ordered Griffiths phase the expressions for p_0 and r_0 are reversed.

Relation between the length scale (distance between non-decimated sites) and the energy-scale reads as:

$$\Omega \sim L^{-\frac{1}{2|\Delta|}}, \quad (12)$$

thus Δ is simply related to the dynamical exponent as $z = 1/2|\Delta|$.

The properties of the Griffiths phases can be interpreted in terms of rare regions, as reviewed in [9, 10].

3. Consequences for the ground-state wavefunction

The SDRG procedure allows to evaluate any observable in each given disordered sample. Besides all the observables of interest reviewed in [3], a particular attention has been paid in recent years towards the characterization of the ground-state wavefunction, via its fidelity [11] and its multifractality [12], while the entanglement properties will be discussed in detail in Section V.

C. Higher dimensions $d > 1$




Figure 1. Structure of connected clusters at the critical point of the 2D RTIM with $L = 64$. Left panel: fixed- h distribution, right panel: box-distribution. The colour-code is used to indicate the moment of the clusters.

In more complicated geometries, in particular in higher dimensions, the SDRG can in principle be performed numerically and the first results are reviewed in [3]. In practical applications, a large finite sample with N sites is renormalized up to the last effective spin, and the original sites of the sample are parts of effective clusters of different sizes. In the *paramagnetic phase*, the clusters have a finite typical linear extent ξ , which characterizes the correlation length of the system. As the critical point is approached, ξ diverges as $\xi \sim |\theta - \theta_c|^{-\nu}$. In the *ferromagnetic phase* $\theta < \theta_c$, there is a huge connected cluster, which is compact and contains a finite fraction m of the sites, that directly represents the average longitudinal magnetization. At the *critical point*, the giant cluster is a fractal: its total moment μ scales with the linear size L of the system as $\mu \sim L^{d_f}$, d_f being the fractal dimension as in Eq.(9). This is related to the scaling dimension of the longitudinal magnetization as $x = d - d_f$. Renormalizing sites at a special position (surfaces, edges, wedges and cones), one obtains the scaling behaviour of the system at the special local environment, which is characterised with the actual local scaling exponent[13]. One can also consider more special geometries, such as multiple-junctions, at which several semi-infinite chains or plans meet [14, 15].

Transverse-spin correlations are defined as: $\langle \sigma_i^z \sigma_j^z \rangle$, which are of $\mathcal{O}(1)$, if both sites i and j are one-site clusters, (i.e. both are immediately decimated out, and these are the white sites in Fig.1) otherwise the correlation is negligible. At the critical point one should consider the connected part of the average transverse-spin correlation function, which decays with the distance as a power with the exponent η_t . Numerical SDRG results indicate[16], that this exponent is approximately $\eta_t \approx 2 + 2d$, for $1 \leq d \leq 3$, see in Table I.

One important point is the relation between the energy-scale Ω (or inverse time-scale) and the length-scale: in the numerical procedure, Ω is the transverse field associated with the last effective spin. If the scaling is controlled by an IDFP, then the scaling relation is activated $\ln \Omega \sim L^{-\psi}$ as in Eq.(7). For strong disorder Fixed Point, the scaling relation is in the conventional power-law form $\Omega \sim L^{-z}$, which generally holds in the Griffiths phase, and the dynamical exponent z depends on the value of the control parameter, see also in Eq.(12). Distribution of the smallest gap in different random samples, $P(\Omega, L)$ depends on the variable $u = \Omega L^z$: it is universal and given by the limit distribution of extremes of independent and identically distributed random numbers[17], as explained in the framework of the SDRG approach in Ref. [18].

Table I. Critical exponents of the RTIM in different dimensions. In 1D the analytical results are from[2], in 2D the numerical results are taken from[5], in 3D and 4D these are from[6, 7]. The surface magnetization exponent x_s and the decay exponent of the transverse magnetization, η_t are from [13] and [16], respectively.

	1D	2D	3D	4D
ν	2.	1.24(2)	0.98(5)	0.79(5)
x	$\frac{3-\sqrt{5}}{4}$	0.982(15)	1.840(15)	2.72(12)
x_s	0.5	1.60(2)	2.65(15)	3.7(1)
η_t	4.1(2)	6.0(2)	8.1(2)	
ψ	1/2	0.48(2)	0.46(2)	0.46(2)

The numerical implementation of the SDRG procedure needs some care. Naive application of the decimation rules leads to a computational time which scales as $t \sim \mathcal{O}(N^3)$. Such type of procedure has been used for ladders [4] and for square samples of linear size up to $L \approx 160$ [19, 20]. In Ref. [21], the so called planar approximation is introduced and in this way they could go up to $L = 500$. Finally, using the maximum rule, an efficient numerical procedure has been introduced in Refs. [6, 7] which works as $t \sim \mathcal{O}(N \ln N + E)$, where E stands for the number of edges of the lattice. In models with nearest-neighbour interaction $E \sim N$ and one can go up to $N = 4 \times 10^6$, c.f. in 2D up to $L = 2048$. Numerical studies of the RTIM in 2D, 3D and 4D indicate, that in each case the critical behaviour is governed by an IDFP, like in the one-dimensional case. The probably most accurate values of the critical exponents are collected in Table I. The critical exponents of the RTIM in any studied dimension have found universal, i.e. they do not depend on the actual form of the initial disorder.

To decide about the upper critical dimension of the IDFP, the critical properties of the RTIM have been studied on Erdős-Rényi random graphs [6, 7], which are formally infinite-dimensional objects. The numerical results have indicated that the critical behaviour is controlled by a logarithmically infinite disorder fixed point, pointing towards an infinite upper critical dimension for this type

of problem.

D. Approximate RG methods

Simpler approximation methods have been developed and applied to the RTIM [22–28]. One of those [22] is based on the quantum cavity approach [29], which is found to reproduce some of the exact results in 1D. The quantum cavity method is equivalent to a linearized transfer matrix approach [24]. If no linearization is performed, i.e. in the so called non-linear transfer matrix approach, one obtains IDFP behavior for $d \geq 2$ [24]. Also approximate renormalization group schemes have been suggested [25–28], during which the order of the RG steps is changed in such a way that the proliferation of new couplings is avoided. These methods have reproduced some exact 1D results and also provide IDFP behavior for $d \geq 2$, in agreement with the standard SDRG method.

III. RANDOM QUANTUM SYSTEMS WITH LONG-RANGE INTERACTIONS

In nature there are magnetic materials in which ordering is due to long-range (LR) interactions which decay as a power $\alpha = d + \sigma$ with the distance. The best known examples are dipolar systems, such as the LiHoF₄. Putting this compound into an appropriate external magnetic field we obtain an experimental realisation of a dipolar quantum ferromagnet [30]. Similar systems have been experimentally realised recently by ultracold atomic gases in optical lattices [31–35]. Here we consider quantum magnets with LR interactions in the presence of quenched disorder. Such type of a system is realised by the compound LiHo_xY_{1-x}F₄, in which a fraction of $(1 - x)$ of the magnetic Ho atoms is replaced by non-magnetic Y atoms [30]. A related, but somewhat simplified quantum model which describes the low-energy properties of this system is the random transverse-field Ising model with LR interactions given by the Hamiltonian:

$$\mathcal{H} = - \sum_{i \neq j} \frac{b_{ij}}{r_{ij}^\alpha} \sigma_i^x \sigma_j^x - \sum_i h_i \sigma_i^z. \quad (13)$$

where the $b_{ij} > 0$ parameters and the $h_i > 0$ transverse fields are i.i.d. random variables with given initial distributions.

In the LR model couplings and transverse fields play a different role and this asymmetry is manifested in the SDRG trajectories. At the critical trajectory couplings are very rarely decimated and the renormalised transverse fields follow Eq.(2). On the contrary almost always transverse fields are decimated out, and according to the maximum rule in Eq.(3) we have typically: $\tilde{J}_{jk} \approx J_{jk}$. Using these observations a primary model has been formulated, which has an exact solution in one dimension.

A. 1D - solution of the primary model [36]

In the primary model, the transverse fields are random, but the couplings are non-random, i.e. $b_{ij} = b = 1$. In the paramagnetic phase and at the critical point, almost always transverse fields are decimated. After decimating h_i , the effective coupling between nearest clusters $i - 1$ and $i + 1$ will always be smaller than the deleted ones, $J_{i-1,i}$ and $J_{i,i+1}$ and we assume that according to the maximum rule $\tilde{J}_{i-1,i+1} = J_{i-1,i+1}$. Then the renormalization rule of couplings between nearest clusters can be expressed in terms of the length variables as $\tilde{J}_{i-1,i+1}^{-1/\alpha} = J_{i-1,i}^{-1/\alpha} + J_{i,i+1}^{-1/\alpha} + w_i$, where w_i is the extension of the cluster, which will be neglected in the following. Using reduced variables $\zeta = \left(\frac{\Omega}{J}\right)^{1/\alpha} - 1$ and $\beta = \frac{1}{\alpha} \ln \frac{\Omega}{h}$, the approximate renormalization rules are

$$\tilde{\zeta} = \zeta_{i-1,i} + \zeta_{i,i+1} + 1 \quad (14)$$

and

$$\tilde{\beta} = \beta_i + \beta_{i+1} \quad (15)$$

for field and bond decimation, respectively. Since, in the ferromagnetic phase, the effective couplings between remote clusters may be stronger than those between adjacent ones due to the large mass of clusters, this approach is justified only in the paramagnetic phase and at the critical point.

The decimation equations in Eqs.(14) and (15) are identical to those of the 1d disordered $O(2)$ quantum rotor model of granular superconductors [37, 38] with the grain charging energy U_i and Josephson coupling $\mathcal{J}_{i,i+1}$ corresponding to $U_i \leftrightarrow J_{i,i+1}^{1/\alpha}$ and $\mathcal{J}_{i,i+1} \leftrightarrow h_i^{1/\alpha}$, see in Sec.IV D.

The fixed-point solution for the distributions ζ and β can be found in [36], here we recapitulate the basic results. The solutions are parameterised with a variable a , which is positive $a > 0$ in the paramagnetic phase and vanishes $a = 0$ at the critical point. The relation between the average distance of clusters, L , and the energy-scale Ω is given by:

$$L \sim \left(\frac{\Omega_0}{\Omega}\right)^{\frac{1+a}{\alpha}}, \quad (16)$$

with an additional factor $\ln^2(\Omega/\Omega_0)$ for $a = 0$. Thus the dynamical exponent $z = \alpha/(1 + a)$ is a continuous function of a , and it is maximal but finite at the critical point: $z_c = \alpha$. The limit distribution of the transverse fields for $\Gamma \rightarrow \infty$ follows the power law $g(h) \sim h^{1/z-1}$, thus the transition is controlled by a strong disorder fixed point. The average correlation length scales at the vicinity of the critical point as:

$$\xi \sim \exp(C'/a), \quad a = (\theta - \theta_c), \quad (17)$$

which is similar to that at a Kosterlitz-Thouless transition point. The ratio of decimated couplings and decimated fields scales at the critical point with the size of

the system, L , as:

$$r(L) \simeq 2 \ln^{-\omega}(L/L_0), \quad \omega = 2. \quad (18)$$

The typical magnetic moment of critical clusters scales also logarithmically:

$$\mu(L) \sim \ln^\chi L, \quad \chi = 2, \quad (19)$$

thus the fractal dimension is formally $d_f = 0$. Finally, the entanglement entropy S_L of a finite block of size L in an infinite system is found to approach a finite limiting value as $L \rightarrow \infty$, even at the critical point.

Results of the numerical SDRG analysis (with the maximum rule) of the LR random transverse-field Ising chain are in agreement with the findings of the primary model, even if the couplings, i.e. the parameters b_{ij} are random.

B. 3D - numerical Strong Disorder RG study

The 3D LR model has been studied by the SDRG approach with the maximum rule [39] and similar behaviour of the RG trajectories are observed as in 1D. At a given energy scale, Ω , the renormalization is characterised by the ratio of decimated couplings and decimated fields, r .

In the paramagnetic phase and at the critical point, where the maximum rule is expected to hold, at the line of fixed points, $\Omega \rightarrow 0$ we have $r \rightarrow 0$, and these fixed points are stable. In the vicinity of this fixed line almost exclusively transverse fields are decimated, the distribution of which is given by a power-law:

$$g(h) = \frac{d}{z} h^{-1+d/z}, \quad (20)$$

with an effective (Ω , i.e. r dependent) dynamical exponent z . At the fixed line z is maximal at the critical point, having a value $z_c \approx \alpha$, as in 1D. The other critical parameters (correlation length, decimation ratio and cluster moment) have similar behaviour as in 1D, thus the relations in Eqs.(17,18,19) are valid, only the exponents of the logarithm in Eqs.(18) and (19) are somewhat different.

The RG phase-diagram can be extended to include the ferromagnetic phase, too, where the RG-flow scales to $r \rightarrow \infty$. This is shown in Fig.2. The line of fixed points at $r = 0$ at other side of the critical point with $\alpha/z < 1$ are unstable and the RG-flow scales to $r \rightarrow \infty$. In this regime the maximum rule in the SDRG procedure is certainly not valid. The two regimes of fixed points are separated by the critical fixed point at $\alpha/z = 1$.

In the vicinity of the line of fixed points at $r = 0$, almost exclusively transverse fields are decimated out. In a finite cluster of linear size ℓ however, there are the few smallest ones which remain intact. From the distribution of the fields in Eq.(20) one can estimate the value of these non-decimated fields through extreme value statistics. This type of analysis leads to basically identical results, as obtained through the SDRG approach. Using




Figure 2. (Color online) Schematic SDRG phase diagram obtained through the maximum rule as a function of the ratio r and the effective dynamical exponent defined in Eq.(20). The arrows indicate the direction the parameters evolve as the energy-scale is reduced. Fixed points (blue circles) are at $r = 0$: the attractive fixed points of the paramagnetic phase ($\alpha/z > 1$) and the repulsive ones ($\alpha/z < 1$) are separated by the critical fixed point (red circle).

this type of analysis it has been argued that the extrapolated value of the magnetization from the ferromagnetic side has a finite limiting value and thus the transition is of mixed-order.

C. Other quantum models with LR interactions

1. Ising chains with other type of LR interactions

The case where the Long-ranged interactions are dilute are analyzed in [40]. The Dyson hierarchical version of the quantum Ising chain with LR power-law ferromagnetic couplings and pure or random transverse fields is studied via real-space renormalization in [41], while the Dyson Hierarchical LR Quantum Spin-Glasses are investigated in [42].

The critical properties of random quantum systems, in particular the RTIM in one dimension in the presence of LR interactions which decay in a stretched exponential fashion, as $J(r) \sim \exp(-Cr^a)$ has been studied in Ref. [43]. Using a variant of the SDRG approach similar to the primary model presented in Sec.III A, the critical behaviour is found to depend on the parameter a . For $0 < a < 1/2$ the critical behaviour is controlled by an IDFP, in which the critical exponents are a -dependent and these are different from those in the SR model. For example one obtains $\psi = a$, which is understandable, since the relation between energy-scale and length-scale is dominated here the form of LR interactions. On the contrary for $a > 1/2$ the LR interactions are irrelevant and the critical properties of the model are the same as its SR variant.

2. Random Heisenberg chains with LR interactions

In Refs. [44, 45] the Hamiltonian of N LR interacting $S = 1/2$ spins is considered:

$$\mathcal{H}_{Heis} = \sum_{i \neq j, \beta} J_{ij}^{\beta} S_i^{\beta} S_j^{\beta}, \quad (21)$$

which are placed randomly on a periodic lattice of length L and lattice spacing s . The interaction between all pairs of sites i, j are antiferromagnetic:

$$J_{ij}^{\beta} = J |r_i - r_j|^{-\alpha} \exp(-|(r_i - r_j)|/\xi), \quad (22)$$

having a power-law decay with exponent α and an exponential cut off with a length scale ξ . This model is aimed to better understand the magnetic properties of doped semiconductors. The singular properties of the model are studied by the SDRG method with the use of the sum rule. In the LR limit, $\xi \rightarrow \infty$, the distribution function of the low energy excitations are studied by various values of α and at a critical value $\alpha_c = 1.6$ it coincides with a critical function pointing towards the existence of a many body localization transition.

In Ref. [45] the magnetic susceptibility $\chi(T)$ is studied as a function of the temperature T . In the LR limit, $\xi \rightarrow \infty$ a crossover is observed at $\alpha^* = 1.066$ between a phase with a divergent low-temperature susceptibility $\chi(T \rightarrow 0)$ for $\alpha > \alpha^*$ to a phase with a vanishing $\chi(T \rightarrow 0)$ for $\alpha < \alpha^*$.

IV. OTHER RANDOM QUANTUM MODELS

The RTIM is one of the most studied random quantum models, for which some new quantities, such as the transverse magnetization [16] and the magnetic Grüneisen ratio [46] have been studied by the SDRG method. Besides models that can be directly mapped onto the RTIM, such as for instance the Majorana model considered in [47], many other short-ranged random quantum models have been analyzed via SDRG since the Review [3].

A. Models with discrete symmetry

1. Antiferromagnetic random quantum Ising chain

The antiferromagnetic random quantum Ising chain is defined by the Hamiltonian:

$$\mathcal{H}_{AF} = \sum_i J_i \sigma_i^z \sigma_{i+1}^z - \sum_i h_i \sigma_i^x - \sum_i H_i \sigma_i^z, \quad (23)$$

with $J_i, h_i, H_i > 0$. The clean model has paramagnetic and antiferromagnetic ordered phases, which are separated by a critical line, which for $h > 0$ is controlled by the Ising quantum fixed point at $h/J = 1$ and $H/J = 0$.

At $h = 0$, when the model is classical there is a multicritical point at $H/J = 2$ separating the antiferromagnetic phase from the ferromagnetic one.

In the random chain, where the couplings are distributed uniformly in $0 < J < 1$ and the random transverse-fields are distributed uniformly in $0 < h < h_0$ (but the longitudinal fields were non-random, $H_i = H$), infinite disorder scaling is observed only at $H = 0$, which is equivalent to the RTIM in 1D. For any finite value of $H > 0$ the numerical results indicate strong disorder scaling, thus a paramagnetic phase with Griffiths singularities [48]. More recent studies indicate, that the region of infinite disorder criticality is extended to $H > 0$, provided the distributions of the couplings and that of the transverse fields have a finite bounding value from below and above [49].

2. Ashkin-Teller chains

The N -colour Ashkin-Teller chain is defined in terms of $\alpha = 1, 2, \dots, N$ -sets Pauli-matrices by the Hamiltonian:

$$\begin{aligned} \mathcal{H}_{AT} = & - \sum_{\alpha=1}^N \sum_i (J_i \sigma_{\alpha,i}^z \sigma_{\alpha,i+1}^z + h_i \sigma_{\alpha,i}^x) \\ & - \sum_{\alpha < \beta}^N \sum_i (K_i \sigma_{\alpha,i}^z \sigma_{\alpha,i+1}^z \sigma_{\beta,i}^z \sigma_{\beta,i+1}^z + g_i \sigma_{\alpha,i}^x \sigma_{\beta,i}^x) \end{aligned} \quad (24)$$

which is the generalization of the standard Ashkin-Teller chain with $N = 2$, which has been studied before and reviewed in [3]. In terms of positive random parameters, however keeping the ratios $\epsilon_{h,i} = g_i/h_i$ and $\epsilon_{J,i} = K_i/J_i$ spatially homogeneous: $\epsilon_{h,i} = \epsilon_{J,i} = \epsilon_i$ the model has been investigated by the SDRG method in a series of papers [50, 52, 53, 83] Different phases and various critical and multicritical points have been identified, which all are of the infinite disorder type. Various numerical studies have been performed to confirm the SDRG predictions [54–57].

B. Models with continuous symmetry

The random Heisenberg Antiferromagnetic spin chain is the first model where SDRG has been introduced [1]. After the various works already reviewed in [3], more recent studies include the effects of next-nearest-neighbor interaction in $d = 1$ [58], the case of weakly coupled chains [59], models in dimension $d = 2$ [60–62], as well as the generalizations to various type continuous symmetry ($SU(3)$, $SU(N)$, $SO(N)$) considered in the series of papers [63–67], where different types of random singlet phases are identified via SDRG and the low-energy behaviour is controlled by infinite disorder fixed points.

The $S = 1$ random spin chain is studied by the numerical application of the SDRG method [68], finding that the cross-over in the critical behaviour with the strength of disorder is in agreement with the analytical theory. Other studies of random spin chains with string-order

parameter include the random Majumdar-Ghosh Chain [69] and the random Cluster-Ising-Model [70].

The one-dimensional Hubbard model with random-hopping matrix-elements, and with random onsite Coulomb repulsion terms is studied by the SDRG method [71]. Two critical phases are identified, which correspond to an infinite disorder spin random singlet for strong interactions and to an orbital infinite disorder fixed point for vanishing interactions. To each critical infinite disorder fixed point is connected a Griffiths phase.

Layers and bilayers of $S=1/2$ Heisenberg antiferromagnets with different types of disorder: bond randomness, site dilution, and dimer dilution are studied by the numerical application of the SDRG method [72]. Generally the systems exhibit an ordered and a disordered phase separated by a phase boundary on which the static critical exponents appear to be independent of bond randomness in the strong-disorder regime, while the dynamical exponent is a continuous function of the bond disorder strength. The low-energy fixed points of the off-critical phases are affected by the actual form of the disorder, and the disorder-induced dynamical exponent depends on the disorder strength. As the strength of the bond disorder is increased, there is a set of crossovers in the properties of the low-energy singularities.

C. Disordered non-abelian anyonic chains

Non-abelian order is a particular feature in two-dimensional quantum systems and non-abelian excitations are present in fractional quantum Hall states. Chains of interacting anyonic quasiparticles are introduced recently and their properties in the presence of quenched disorder has been studied through the SDRG method [73–76].

Let us consider a simple example of anyonic models, the Fibonacci or golden chain, which is introduced on the analogy of the $S = 1/2$ Heisenberg model with $SU(2)$ symmetry: $\mathcal{H} = \sum_i J_i \mathbf{S}_i \mathbf{S}_{i+1}$, for which the energy of a nearest neighbour pair is different in the singlet and in the triplet channels, and according to tensor product or the fusion rule we have: $\frac{1}{2} \otimes \frac{1}{2} = 0 \oplus 1$. The Fibonacci chain consists of non-abelian anyons carrying a topological charge, τ , and their interaction is described by the fusion rule: $\tau \otimes \tau = 1 \oplus \tau$, which means that the Hilbert space of two neighbouring anyons is the direct sum of unity and a copy of τ . This type of construction can only be described by a truncated tensor product, where the $SU(2)$ representations are truncated at a level k . The Hamiltonian of the Fibonacci chain is given in the form:

$$\mathcal{H} = \sum_{i=1}^N J_i P_i^A \quad (25)$$

where J_i are the random interactions and P_i^A is the singlet projection operator between site i and $i + 1$. Generally the Hilbert space of $N + 1$ anyons is given by the

N^{th} Fibonacci number. Other type of non-abelian anyonic chains can be constructed in similar way, important class being the $SU(2)_k$ anyonic chains, the Fibonacci chain corresponds to $k = 3$.

In the SDRG procedure we choose the strongest coupling in the chain, $\Omega = |J_i|$, and decimate it. For a strong antiferromagnetic coupling the sites i and $i + 1$ form a singlet, and an effective coupling $\tilde{J} \approx \kappa \frac{J_{i-1} J_{i+1}}{J_i}$ is formed between sites $i - 1$ and $i + 2$, with $\kappa = \phi^2$ and $\phi = \frac{1+\sqrt{5}}{2}$, the golden mean ratio. For a strong ferromagnetic bond sites i and $i + 1$ form a cluster, having effective couplings to the nearby sites: $\tilde{J} \approx -\frac{J_{i\pm 1}}{\phi}$.

The SDRG transformation has an infinite disorder fixed point, so that the ground state is a random singlet phase. The relation between energy-scale and length-scale is activated, with the critical exponent $\psi_{AF} = 1/2$ if the original model is antiferromagnetic, and $\psi_{AF/F} = 1/3$ if in the original model there is a finite fraction of ferromagnetic bonds. The SDRG analysis has been extended for other non-abelian anyonic chains, a detailed analysis of the results can be found in Ref. [76]. Relation with $SU(N)$ symmetric random chains has been noticed in [63–67].

D. Superfluid-insulator transition

One-dimensional Josephson junction array with random couplings J_i and random charging energies U_i are described by a quantum rotor Hamiltonian:

$$\mathcal{H} = \sum_i U_i (\hat{n}_i - \bar{n}_i)^2 - \sum_i J_i \cos(\hat{\theta}_i - \hat{\theta}_{i+1}), \quad (26)$$

with charges \hat{n}_i and phases $\hat{\theta}_i$ at site i , which satisfy the commutation relations: $[\hat{n}_k, \hat{\theta}_l] = -i\delta_{kl}$. At each site there is a random offset charge (or chemical potential), which is taken as $-1/2 < \bar{n}_i < 1/2$ and the integer part is absorbed into the definition of \hat{n}_i . This model is considered as an effective theory of interacting bosons propagating in a random potential and the type of the insulating phase is found to depend on the symmetry properties of the offset charge distribution.

The SDRG treatment of this model is described in Refs. [8, 77–81] and has been recently reviewed in [82]. Here we just recapitulate the main ideas. The energy gap due to charging energy is $\Delta_i = U_i(1 - 2|\bar{n}_i|)$, that should be compared with the interaction coupling J_i , and their maximum term defines the energy-scale Ω .

If $\Omega = \Delta_i$ corresponds to a large gap, this site is fixed to its lowest energy state and an effective coupling is generated between sites $i - 1$ and $i + 1$:

$$\tilde{J} = \frac{J_{i-1} J_i}{\Omega(1 + 2|\bar{n}_i|)}, \quad (27)$$

and in the denominator the term $1 + 2|\bar{n}_i|$ is set to unity. On the contrary, if $\Omega = J_i$ corresponds to a large Josephson coupling, a cluster with a coherent phase is formed

with an effective capacitance: $\tilde{C} = C_i + C_{i+1}$, thus the effective charging energy follows the rule:

$$\frac{1}{\tilde{U}} = \frac{1}{U_i} + \frac{1}{U_{i+1}}. \quad (28)$$

Introducing scaling variables: $\beta_i = \ln(\Omega/J_i)$ and $\xi_i = \Omega/U_i - 1$ in the vicinity of the fixed point, where the expressions for the offset charges are simplified, one arrives at the same form of the flow equation of the distribution functions as for the primary model of the one-dimensional RTIM with LR interactions of Sec.III A. The paramagnetic phase of the LR RTIM corresponds to the superfluid phase, while the ferromagnetic one to the insulator phase. The superfluid phase is just a Griffiths phase with a dynamical exponent $z = 1/(1+a)$, the superfluid-insulator transition is of strong disorder type with $z = 1$ and with an exponentially diverging correlation length. Details of the solution and a review of the numerical results are given in Ref. [82]. As a final remark, let us stress that the nature of Superfluid-Bose Glass Transition has remained controversial over the years, as summarized in recent numerical investigations[83, 84].

E. Superconductor-Metal transition

An Infinite-Randomness Fixed Point has also been found for the Superconductor-Metal Quantum Phase Transition [85–88].

F. The Rainbow Spin Chain

Although the so-called 'Rainbow spin chain' is not random, the spatial structure of its inhomogeneity allows to apply iteratively the Ma-Dasgupta SDRG rule to construct the ground state and analyze its entanglement properties [89–93]. The addition of disorder in this rainbow spin chain has been also studied recently via SDRG [94].

V. ENTANGLEMENT PROPERTIES

The entanglement of quantum many body systems is a promising concept to understand their topological and universal properties, in particular in the vicinity of a quantum phase-transition point (see the reviews [95–97]). The entanglement of the ground state $|\Psi\rangle$ between the subsystem \mathcal{A} and the rest \mathcal{B} of the system is quantified by the von Neumann entropy $\mathcal{S} = -\text{Tr}_{\mathcal{A}}(\rho_{\mathcal{A}} \log_2 \rho_{\mathcal{A}})$ of the reduced density matrix $\rho_{\mathcal{A}} = \text{Tr}_{\mathcal{B}}|\Psi\rangle\langle\Psi|$. Generally \mathcal{S} scales with the area of the interface separating \mathcal{A} and \mathcal{B} . In some cases however, there are singular corrections to the area law. In one-dimensional pure systems, \mathcal{S} is logarithmically divergent at a quantum critical point[98–100]: $\mathcal{S} = \frac{c}{3} \log_2 \ell + cst$. Here ℓ is the size of the subsystem \mathcal{A} and the prefactor is universal, c being the central

charge of the conformal field theory. Besides this entanglement entropy, it is interesting to consider also the full entanglement spectrum [101].

In random quantum systems the entanglement properties are conveniently studied by the SDRG approach and many of the obtained results, mainly in one-dimension are thoroughly reviewed in Refs. [76, 97]. Therefore here we just shortly mention the known results in 1D and we concentrate on the higher dimensional results, as well as on the more recent developments.

A. Random quantum chains

The ground state of the $S = 1/2$ random-bond Heisenberg chain in the SDRG approach is a random singlet phase and each singlet bond which connects the two subsystems \mathcal{A} and \mathcal{B} has a contribution 1 to the entanglement entropy. The entanglement entropy scales logarithmically: $\mathcal{S} = \frac{c_{eff}}{3} \log_2 \ell + cst$, with an effective central charge $c_{eff} = \ln 2$. The entanglement across a weakened link is studied in Ref. [102], while the multifractal Orthogonality Catastrophe produced by a local cut is analyzed in [103].

The ground state of random $S > 1/2$ chains depends on the strength of disorder. For strong enough disorder it is a spin- S random singlet phase having an effective central charge $c_{eff} = \ln(2S + 1)$ [104–107]. For weaker disorder there are a set of multi-critical points, at which the central charge has not yet been calculated exactly. The case $S = 1$ is studied in Ref. [105].

For the RTIM the ground state in the SDRG approach consists of a set of clusters of different sizes (see in Fig.1) and each cluster which has points in both subsystems \mathcal{A} and \mathcal{B} has a contribution 1 to the entanglement entropy. In one dimension, according to SDRG calculations in the off-critical region, \mathcal{S} is finite and thus satisfies the area law. At the critical point, it is logarithmically divergent, and the effective central charge is $c_{eff} = \ln 2/2$, i.e. just the half of that in the random XX- and $S=1/2$ Heisenberg chains, which follows also from an exact mapping [108]. In a given sample of finite length, the position of the maximum of the average entanglement entropy (the average is made over all possible positions of the subsystem) can be used to define a sample dependent pseudo-critical point [109].

The effective central charge of random one-dimensional systems calculated from the average entanglement entropy is generally smaller than its analogous value in the pure model. One can however construct models, in which the effective central charge of the random model is the larger [110]. One can also construct models with (locally) correlated disorder, so that the local control parameter stays constant [111]. In such models the critical behaviour, as well as scaling of the critical entanglement entropy follows the same form as in the pure systems [111, 112].

The entanglement entropy between a random and a

clean part of a system, such in the XX-chain has also been considered recently. In this case, at the critical point, a very weak, double-logarithmic dependence is observed [113]: $\mathcal{S} \sim \ln \ln \ell$. For a related study of entanglement across extended random defects see [114]. Entanglement entropy at multiple junctions of random quantum chains has been studied in [115]. The SDRG approach has also been used to study the critical properties of aperiodic quantum spin chains [116–119] and their entanglement entropy is calculated in the strong aperiodicity limit [120, 121].

Other entanglement measures have been also studied in random quantum chains, like the entanglement [122, 123] or the concurrence [124] between distant pairs of q-bits, the full entanglement spectrum of random singlet critical points [125], the Rényi entropies [126], the fluctuations of the entanglement entropy [127], the full probability distribution of the entanglement entropy [128], the Schmidt-gap (i.e. the difference between the two largest eigenvalues of the entanglement spectrum) for the RTIM and for the $S = 1$ random spin chain [129]. Using the SDRG method the entanglement negativity in random singlet phases are shown to scale logarithmically with the size of the system [130].

Motivated by the entanglement entropy in the random singlet phase of the random $S = 1/2$ spin chain, entanglement measure through valence bond entanglement has been proposed for $SU(2)$ quantum systems [131], which can be efficiently measured through quantum MC calculations both in one- and two-dimensions [132, 133].

B. RTIM in higher dimensions

The entanglement entropy of the RTIM is given by the number of such clusters in the ground state which have points both in \mathcal{A} and \mathcal{B} . This quantity has been considered first in Ref. [19] and at the critical point, a singular behavior in the form of $\mathcal{S} \sim \ell \ln \ln \ell$ has been found in finite systems with linear size $L = 64$. Soon after the calculation has been extended up to $L = 160$ [20] and the numerical results have been interpreted as a logarithmic correction to the area law: $\mathcal{S} \approx a\ell + b \ln \ell$. To decide between the two suggested singular forms, a calculation has been performed with the very efficient numerical algorithm up to $L = 2028$ and by two different forms of disorder. These calculations have been performed also for 3D and 4D [134].

At the critical point of a D-dimensional system, the entanglement entropy when the subsystem \mathcal{A} is a cube is found to be in the form:

$$\mathcal{S}_{\text{cube}}^{(D)}(\ell) = a_{D-1} f_{D-1} + \sum_{E=1}^{D-2} a_E f_E + \mathcal{S}_{\text{cr}}^{(D)}(\ell), \quad (29)$$

where the first term represents the area-law, the second terms are analytical corrections due to E -dimensional edges and the last term is the corner contribution, which

is logarithmically divergent: $\mathcal{S}_{\text{cr}}^{(D)}(\ell) = b^{(D)} \ln \ell + \text{const.}$ According to numerical estimates the prefactors are universal, i.e. disorder independent and given by $b^{(2)} = -0.029(1)$, $b^{(3)} = 0.012(2)$ and $b^{(4)} = -0.006(2)$.

The corner-entropy has also been studied in the vicinity of the critical point and found to be extremal at the critical point. Outside the critical point, $\mathcal{S}_{\text{cr}}^{(D)}(\ell, \delta)$ is finite and can be obtained by replacing ℓ with the finite correlation length ξ , where the divergence of ξ at the critical point is characterised by the given exponent in Table I.

C. Bond diluted quantum Ising model

The bond diluted quantum Ising model is defined by the Hamiltonian in Eq.(1) with $h_i = h$ and with nearest-neighbour couplings which are $J > 0$ with probability p and $J = 0$ with probability $1 - p$. At the percolation transition point p_c , for small transverse field h , there is a line of phase transition, the critical properties of which are controlled by the percolation fixed point [135], for a review see [136]. The ground state of \mathcal{H} is given by a set of ordered clusters, which are in the same form as for percolation. Now consider a subsystem \mathcal{A} with boundary Γ and calculate the entanglement entropy between the subsystem and the environment, which is given by the number of clusters in \mathcal{A} which intersect Γ and contain also at least one point of the environment.

In two dimensions it is given in the same form as for the RTIM [137]:

$$\mathcal{S}_{\Gamma} = aL_{\Gamma} + b \log L_{\Gamma}, \quad (30)$$

where L_{Γ} is the length of Γ . The prefactor of the logarithm in Eq.(30) is given by the Cardy-Peschel formula [138]:

$$b = -\frac{5\sqrt{3}}{96\pi} \sum_k \left[\left(\frac{\pi}{\gamma_k} \right) - \left(\frac{\gamma_k}{\pi} \right) + \left(\frac{\pi}{2\pi - \gamma_k} \right) - \left(\frac{2\pi - \gamma_k}{\pi} \right) \right], \quad (31)$$

where γ_k is the interior angle at each corner. In the special case of the square subsystem one has $b = -5\sqrt{3}/(36\pi) = -0.07657$. The conformal prediction in Eq.(31) has been confirmed by numerical calculations for different shapes of Γ [137].

In $d = 3$, numerical calculations have confirmed that the singular contributions to the entanglement entropy of the bond diluted quantum Ising model are due to corners, and the prefactor of the logarithm is given by $b = 1.72(3)$, which is different from that in the RTIM [139].

D. Relations between SDRG and Entanglement-Algorithms

Since Tensor-Networks have become very popular in recent years, it is interesting to point out

that the SDRG actually corresponds to a special type of Multi-scale-Entanglement-Renormalization-Ansatz (MERA) (see section IV of the review [140]) and has been integrated into various tensor-network algorithms [48, 141–144]. Recently, in analogy with SDRG, a Strong-Disorder-Disentangling procedure [145] has been introduced : at each step, one chooses the most strongly entangled pair of sites, in order to construct iteratively the appropriate unitary circuit that transforms a given quantum state into an unentangled product state. The goal is to reveal the emergent entanglement geometry.

VI. LOCALIZED AND MANY-BODY-LOCALIZED PHASES OF QUANTUM SPIN CHAINS

For random quantum spin chains that can be mapped onto free-fermions via the Jordan-Wigner transformation, the presence of disorder in this one-dimensional geometry leads to the Anderson-real-space-localization of all fermionic modes. In the presence of interactions, the issue of Many-Body-Localization in isolated random quantum spin chains has attracted a lot of attention recently, as reviewed in [146–150]. Among the various methods that have been proposed to construct the Local Integrals of Motion (LIOMs) that characterize the Many-Body-Localized-Phase (see the reviews [151, 152] on LIOMs), Strong Disorder RG procedures have been introduced under the names of RSRG-X and RSRG-t, in order to construct the excited eigenstates or the effective dynamics respectively.

A. RSRG-X for excited eigenstates

In order to construct the whole set of eigenstates, the main idea of the RSRG-X procedure [153] is to keep the two possible local-energy-branches at each step, instead of projecting systematically onto the lowest local energy branch when the goal is to construct only the ground-state. The RSRG-X can be formulated for the most general Hamiltonian involving Pauli matrices [154]

$$H = \sum_{[\mu]} h_{[\mu]} \sigma^{[\mu]} = \sum_{\mu_1, \dots, \mu_N} h_{[\mu_1, \dots, \mu_N]} \sigma_1^{\mu_1} \sigma_2^{\mu_2} \dots \sigma_N^{\mu_N} \quad (32)$$

where $\mu_i = 0, 1, 2, 3$ is the index of the Pauli matrix acting on spin i . One chooses the maximum $\Omega = \max(h_{[\mu]}) = h_{[\mu_0]}$ among the real couplings $h_{[\mu]}$ of the Hamiltonian. The corresponding term

$$H_0 = h_{[\mu_0]} \sigma^{[\mu_0]} \quad (33)$$

has two levels ($\pm h_{[\mu_0]}$) corresponding to the high/low energy sectors. The rest of the Hamiltonian can be classified according to the commutativity or anticommutativity with H_0

$$H - H_0 = H_1^{comm} + H_1^{anti} \quad (34)$$

The part H_1^{comm} that commutes with H_0 is kept to describe its effect withing each energy-level of H_0 . The part H_1^{anti} that anticommutes with H_0 and that couples the two sectors is taken into account by second-order perturbation theory to obtain the renormalized Hamiltonian within each energy sector of H_0

$$H^R = H_0 + H_1^{comm} - H_1^{anti} \frac{1}{2H_0} H_1^{anti} \quad (35)$$

These rules are thus formally very similar to the Fisher SDRG rules for the ground state. An alternative formulation of the RSRG-X rules in terms of Majorana fermions is described in [155] with its advantages.

While the writing of RSRG-X rules is a direct generalization of the SDRG-rules for the ground-state, it should be stressed that their numerical implementation is much more involved. Indeed, the exact construction of the 2^N eigenstates for a chain of N spins is limited to small sizes as a consequence of the exponential cost. To overcome this limitation, the authors of Ref [153] have thus proposed to replace the exact application of the renormalization rules on all branches by a Monte Carlo sampling of the typical branches of the tree. The RSRG-X procedure has been applied to many random models, including the XX chain [156, 157], the XXX chain [158], the XXZ chain [159], the XYZ chain [160], the three-state quantum clock model [161] and anyonic spin chains [162, 163]. Variants of the RSRG-X procedure have been also introduced to analyze the phase transition between different MBL-phases in the Long-Ranged quantum spin-glass model [164] and for the random Transverse Field Spin-Glass Model on the Cayley tree [165].

B. RSRG-t for the unitary dynamics

The RSRG-t for the effective unitary dynamics of isolated Quantum Spin chains [166, 167] is based on the iterative elimination of the highest local frequency Ω . The idea is that the local degree of freedom with the two energy-levels e_1 and e_2 and corresponding projectors $P_{1,2}$

$$H_0 = e_1 P_1 + e_2 P_2 \quad (36)$$

that is associated to the highest frequency $\Omega = e_2 - e_1$ oscillates freely. In the interaction picture, the rest of the Hamiltonian

$$V \equiv H - H_0 \quad (37)$$

becomes the time-periodic Hamiltonian of high frequency Ω

$$V^{int}(t) = e^{iH_0 t} V e^{-iH_0 t} = \mathcal{V}_0 + \mathcal{V}_1 e^{i\Omega t} + \mathcal{V}_{-1} e^{-i\Omega t} \quad (38)$$

with the three Fourier coefficients

$$\begin{aligned} \mathcal{V}_0 &= P_1 V P_1 + P_2 V P_2 \\ \mathcal{V}_1 &= P_2 V P_1 \\ \mathcal{V}_{-1} &= P_1 V P_2 = \mathcal{V}_1^\dagger \end{aligned} \quad (39)$$

The high-frequency-expansion for Floquet dynamics [168] then yields that the effective Hamiltonian for the remaining degrees of freedom reads [169]

$$V_{eff} = (P_1 V P_1 + P_2 V P_2) + \frac{1}{\Omega} (P_2 V P_1 V P_2 - P_1 V P_2 V P_1) + O\left(\frac{1}{\Omega^2}\right) \quad (40)$$

The first ligne corresponds simply to the projection of V onto the energy levels of H_0 , while the second ligne of order $1/\Omega$ contains virtual processes between the two energy levels. This formula for the RSRG-t rules is thus equivalent to the RSRG-X rules of Eq 35 based on the two first order perturbation theory for energy-levels, but this dynamical point of view shed a different light on the interpretation of the RG procedure. At some given time t , the degrees of freedom are separated into two groups with respect to $\Omega_t = \frac{1}{t}$:

(i) the local degrees of freedom that would have had higher eigenfrequencies $|\Omega| > \Omega_t$ have been converted into Local Integrals of Motions (LIOMs) via the projectors $P_{1,2}$ that commute with H_{eff} , i.e. they have converged towards their asymptotic state described by the diagonal ensemble of their local Hamiltonian H_0 , while the off-diagonal contributions have been time-averaged-out.

(ii) the remaining degrees of freedom that are characterized by renormalized eigenfrequencies $|\Omega| < \Omega_t$ have not yet converged towards their asymptotic state, since they have not had enough time to oscillate with their eigenfrequency.

The application of the RSRG-t procedure to various models is described in [166, 167, 170]. As a final remark, let us mention that another type of Nonequilibrium dynamical renormalization group has been studied in [171, 172].

C. Non-equilibrium dynamical scaling of observables

Following the experimental progress in non-equilibrium dynamics of ultracold-atomic gases in optical lattices, there are tremendous theoretical efforts aimed at understanding the time-evolution of certain observables in closed quantum systems after a sudden or smooth change of Hamiltonian parameters. In a quench process, both the functional form of the relaxation and the properties of the stationary state are of interest. Here we report numerical results obtained about the non-equilibrium relaxation process in random quantum systems, almost exclusively in one dimension.

Concerning the functional form of the time-dependence of the entanglement entropy [173–177], the results depends on whether the random quantum system can be described in terms of free fermions or not. If the system consists of non-interacting fermions - such as the XX-spin chain with bond disorder or the critical random

transverse-field Ising chain - the dynamical entanglement entropy grows ultraslowly in time as

$$\mathcal{S}(t) \sim a \ln \ln t, \quad (41)$$

and saturates in a finite system at a value

$$\mathcal{S}(\ell) \sim b \ln \ell, \quad (42)$$

where ℓ denotes the size of a block in a bipartite system and can be chosen to be proportional to the size of the system L [174, 177]. Similar scaling forms have been observed for the non-equilibrium relaxation of the full counting statistics in a disordered free-fermion system [175]. By the strong disorder RSRG-t method [166, 167] of Sec.VIB, the ratio of the prefactors in (41) and (42) is predicted as $b/a = \psi_{ne}$, where $\psi_{ne} = 1/2$ is a critical exponent in the non-equilibrium process and describes the relation between time-scale and length-scale as

$$\ln t \sim L^{\psi_{ne}}. \quad (43)$$

Numerical estimates of b/a are somewhat larger, being in the range 0.69–0.59. This discrepancy may be due to the fact that for disordered systems, because of the necessity of calculation of some extremely small eigenvalues, standard eigenvalue solvers would fail to converge for some large-size samples, leading to significant numerical errors.

For interacting fermion models in the Many-Body-Localized phase, the time-dependence of the dynamical entropy is $\mathcal{S}(t) \sim \ln^\omega t$ with $\omega \geq 1$, while the saturation value follows the volume law, $\mathcal{S}(\ell) \sim \ell$ [176]. In this case SDRG theory and numerical results are mainly consistent.

The time evolution of the average magnetization, $\overline{m}(t)$, of the one-dimensional RTIM after global quenches is studied numerically by using multiple precision arithmetic [178]. In this way, the numerical inaccuracies observed in the computation of the entanglement entropy are circumvented. Starting from a fully ordered initial state, the relaxation to the critical point is logarithmically slow described by $\overline{m}(t) \sim \ln^{a'} t$, and in a finite sample of length L , the average magnetization saturates at a size-dependent plateau $\overline{m}_p(L) \sim L^{-b'}$; here the two exponents satisfy the relation $b'/a' = \psi_{ne} = 1/2$. This result is consistent with the SDRG prediction. Starting from a fully disordered initial state, the magnetization stays at zero for a period of time until $t = t_d$ with $\ln t_d \sim L^{\psi_{ne}}$ and then starts to increase until it saturates to an asymptotic value $\overline{m}_p(L) \sim L^{-b''}$, with $b'' \approx 1.5$. The distribution of long-time limiting values of the magnetization shows that the typical and the average values scale differently and the average is governed by rare events.

For the random Heisenberg chain, the dynamical properties at finite temperature have been studied numerically and compared with SDRG predictions in [179, 180]. The non-equilibrium quench dynamics in quantum spin chains with aperiodic interactions have been studied numerically in [181–183].

Besides the quantum quenches discussed up to now, the opposite limit of adiabatic changes of the parameters of the Hamiltonian has also been studied recently via SDRG to analyze the Kibble-Zurek dynamics through the critical point [184].

D. Comparison with other RG procedures existing in the field of Many-Body-Localization

Since the purpose of the RSRG-X and RSRG-t procedures is to produce an extensive number of Local Integrals of Motion (LIOMS), it is clear that their validity is limited to Many-Body-Localized Phases : they allow to analyse the long-ranged order of the excited eigenstates and to study the phase transitions between different Many-Body-Localized phases. To analyze the MBL-transition towards the ergodic delocalized phase, various other RG procedures have been introduced, such as the Aoki exact RG procedure in configuration space [185], and real-space RG procedures based on entanglement [186] or resonances [187, 188], while the phenomenological RG procedure based on the decomposition into insulating and thermal blocks [189] is related to some coarsening models that can be exactly solved by Strong Disorder RG (see Appendix E of the review [3]). An exactly-solvable generalization that takes into account the asymmetry between insulating and thermal blocks corresponds to some Kosterlitz-Thouless scenario [190]. Finally, the Wegner-RG flow [191] or variants thereof have been applied recently to various MBL models [192–196].

VII. FLOQUET DYNAMICS OF PERIODICALLY DRIVEN CHAINS IN THEIR LOCALIZED PHASES

The Floquet dynamics of periodically driven quantum systems has attracted a lot of attention recently (see the reviews [168, 197]). The stroboscopic dynamics can be analyzed via the diagonalization of the time-evolution-operator over one period T

$$U(T, 0) \equiv \mathcal{T} \left(e^{-i \int_0^T dt H(t)} \right) = \sum_{n=1}^{\mathcal{N}} e^{-i\theta_n} |u_n\rangle \langle u_n| \quad (44)$$

The phases $\theta_n \in]-\pi, +\pi]$ characterize the eigenvalues $e^{-i\theta_n}$ of this unitary operator, while the $|u_n\rangle$ are the corresponding eigenvectors.

To have an explicit evolution operator (instead of the implicit time ordering of Eq 44), it is convenient to consider periodic switching between two Hamiltonians H_0 and H_1 during T_0 and T_1 respectively

$$U(T = T_1 + T_0, 0) = e^{-iT_1 H_1} e^{-iT_0 H_0} \quad (45)$$

The simplest example is

$$H(0 \leq t \leq T_0) = H_0 \equiv - \sum_{n=1}^{N-1} J_n \sigma_n^z \sigma_{n+1}^z$$

$$H(T_0 \leq t \leq T = T_0 + T_1) = H_1 \equiv - \sum_{n=1}^N h_n \sigma_n^x \quad (46)$$

since the time-averaged Hamiltonian is the random transverse field Ising chain. The RSRG-X rules for the corresponding Floquet evolution operator of Eq. 45 are analyzed in [198] and can be considered as a direct generalization of the Fisher RG rules.

More generally, the phase-transitions between different Floquet-Localized-phases are expected to be controlled by Infinite-Disorder-Fixed-Points that can be studied via SDRG [199, 200].

VIII. OPEN DISSIPATIVE QUANTUM SPIN CHAINS

In the field of open quantum systems, the interplay between quantum coherence and dissipation can be analyzed within various frameworks [201, 202].

A. Quantum spin chains coupled to a bath of quantum oscillators

The dynamics of a single two-level system coupled to a bath of quantum oscillators is the famous 'spin-boson model' [203]. The generalization for the random quantum Ising chain is described by the Hamiltonian

$$H^{spins} = - \sum_i h_i \sigma_i^x - \sum_{\langle i,j \rangle} J_{ij} \sigma_i^z \sigma_j^z$$

$$H^{bosons} = \sum_i \sum_k \omega_{i,k} \left(a_{i,k}^\dagger a_{i,k} + \frac{1}{2} \right)$$

$$H^{coupling} = \sum_i \sigma_i^z \sum_k \lambda_{i,k} \left(a_{i,k}^\dagger + a_{i,k} \right) \quad (47)$$

where each spin is coupled to its own local bath of oscillators described by its spectral density

$$\mathcal{E}_i(\omega) = \pi \sum_k \lambda_{i,k}^2 \delta(\omega - \omega_{i,k}) \underset{\omega \rightarrow 0}{\propto} \omega^s \quad (48)$$

The case $s = 1$ is called Ohmic dissipation, while $s > 1$ is called super-Ohmic and $s < 1$ is called sub-Ohmic.

To obtain the appropriate SDRG rules [204–209], the idea is to supplement the Fisher SDRG rules of the chain by the adiabation renormalization of the bath developed for the spin-boson model [203]. The main conclusion is the smearing of the quantum phase transition by the dissipation for Ohmic and sub-Ohmic baths, with freezing of large magnetic clusters, while super-Ohmic is irrelevant. These predictions have been tested via Monte-Carlo simulations [210]. Related studies based on Landau-Ginzburg functionals can be found in [211–213].

B. Lindblad dynamics for random quantum spin chains

Another popular description of open quantum systems [201, 202] is the Lindblad dynamics for the density matrix ρ

$$\frac{\partial \rho}{\partial t} = -i[H, \rho] + \mathcal{D}[\rho] \quad (49)$$

where the unitary dynamics governed by the Hamiltonian H is supplemented by the dissipative contribution defined in terms of some set of operators L_α that describe the interaction with the reservoirs

$$\mathcal{D}[\rho] = \sum_{\alpha} \gamma_{\alpha} \left(L_{\alpha} \rho L_{\alpha}^{\dagger} - \frac{1}{2} L_{\alpha}^{\dagger} L_{\alpha} \rho - \frac{1}{2} \rho L_{\alpha}^{\dagger} L_{\alpha} \right) \quad (50)$$

In the field of quantum spin chains, it is interesting to consider two reservoirs acting only on the two boundary spins in order to impose a current-carrying Non-Equilibrium Steady-State (NESS).

For the XX chain with random fields, a strong disorder boundary renormalization has been introduced [214], in order to describe the strong hierarchy of relaxation times as a function of the distance to the boundaries, and to compute explicitly the sample-dependent step-profile of the magnetization. This step profile is expected in other localized chains [215], in stark contrast with the usual linear profile for diffusive dynamics. Note however that the addition of dephasing acting on all spins of the bulk destroys the phase coherence responsible for the localization properties, and produces an effective dynamics described by a classical exclusion process with randomness [216]. Another study concerning the Lindblad dynamics with a contact between random and pure quantum XX spin chains can be found in [217].

IX. ANDERSON LOCALIZATION MODELS

In the field of Anderson localization (see the review [218]), one is interested into the localization/delocalization properties of the eigenstates of tight-binding Hamiltonian of the form

$$H = \sum_i H_{ii} |i\rangle\langle i| + \sum_{i \neq j} H_{ij} (|i\rangle\langle j| + |j\rangle\langle i|) \quad (51)$$

that can be defined for various geometries.

The goal of the SDRG [219, 220] is to analyze the properties at zero-energy $E = 0$ corresponding to the middle of the spectrum. The iterative elimination of the strongest on-site energy H_{ii} or the strongest off-diagonal hopping H_{ij} (in absolute value) leads to the following decimation rules. The decimation of the on-site energy $\Omega = |H_{i_0 i_0}|$ yields the RG rule (even for $k = p$)

$$H_{kp}^R = H_{kp} - \frac{H_{ki_0} H_{i_0 p}}{H_{i_0 i_0}} \quad (52)$$

while the decimation of the off-diagonal coupling $\Omega = |H_{i_0 j_0}|$ produces the RG rule (even for $k = p$)

$$H_{kp}^R = H_{kp} + \frac{H_{i_0 j_0} (H_{ki_0} H_{j_0 p} + H_{kj_0} H_{i_0 p})}{H_{i_0 j_0}^2 - H_{i_0 j_0} H_{j_0 i_0}} - \frac{H_{i_0 i_0} H_{kj_0} H_{j_0 p} + H_{j_0 j_0} H_{ki_0} H_{i_0 p}}{H_{i_0 j_0}^2 - H_{i_0 j_0} H_{j_0 i_0}} \quad (53)$$

These SDRG rules actually coincide with the exact Aoki RG rules at zero-energy $E = 0$ [221–223]. As a consequence, if one focuses on zero-energy $E = 0$, the SDRG rules are exact in both phases (localized and delocalized) as well as at the Anderson phase transition between them.

The SDRG rules have been applied to analyze the localization properties in $d = 1$ [219] and $d = 2$ [223] and to characterize the critical properties of the Anderson delocalization transition in $d = 3$ [220, 223, 224], while the application in higher dimensions $d > 3$ [220, 224] points towards an infinite upper critical dimension $d_{upper} = +\infty$ for the Anderson transition. Note that another SDRG rules based on the Inverse Participation Ratios of eigenstates have been proposed in [225], while the effects of rare resonances on various observables is discussed in [226].

As a final remark, let us mention that several other real-space renormalization approaches have been introduced to analyze the multifractality of eigenstates at the localization-delocalization transition in various models, in particular the Levitov RG reviewed in [218], the block-RG [227, 228] and the Wegner flow approach [229].

X. RANDOM CONTACT PROCESS

The contact process [230, 231] is a basic model in the fields of epidemic spreading and population dynamics. It is defined on a lattice, the sites of which are either active (infected) or inactive (healthy). The time evolution is a continuous-time Markov process with the following independent transition rates. Site i , if it is active, becomes spontaneously inactive with a rate μ_i or it activates site j , provided the latter is inactive, with a rate λ_{ij} . This model in the simplest case with homogeneous parameters and nearest neighbour spreading falls into the universal class of directed percolation.

In experimental realizations of directed percolation, quenched disorder is observed to play an inevitable role, therefore different variants of randomness in the contact process have been studied theoretically. According to the Harris criterion, quenched disorder (both spatial and temporal) is a relevant perturbation, therefore new type of critical behaviour is expected to be present in these systems.

A. Strong disorder RG rules

The contact process with random short range interactions (both the λ_{ij} and the μ_i are i.i.d. random variables) has been studied by the SDRG method, the basic results can be found in Review [3]. The elementary decimation steps are the following. Having a very strong activation rate $\lambda_{ij} = \Omega$, the two connected sites form a cluster in the presence of an effective recovery rate $\tilde{\mu} = 2\mu_i\mu_j/\lambda_{ij}$. On the contrary, for a strong recovery rate $\mu_i = \Omega$, this site is almost always inactive, and there are effective branching rates between all sites j, k , which are nearest neighbours to i , as $\tilde{\lambda}_{jk} = \lambda_{ji}\lambda_{ik}/\mu_i$. Supplementing this relation with the maximum rule we arrive to the elementary decimation rules which are very similar to that of the RTIM. Indeed, for nearest neighbour interactions and for strong enough initial disorder, the extra prefactor is unimportant, and the critical behaviour is controlled by the IDFP of the RTIM. The infinite-randomness scenario has been checked by large scale Monte-Carlo simulations in $d = 2$ [232, 233] and in $d = 3$ [234]. For weaker disorder, the cross-over between the weak- and the strong disorder scaling regions is analyzed in [235]. Detailed results on the distribution of dynamical observables can be found in [236]. The contact process with asymmetric spreading has also been studied by SDRG [237]. In dimension $d = 5$, the Griffiths singularities are analyzed in [238], as an example where they can co-exist with a clean critical behavior predicted by the Harris criterion [239]. The effects of long-ranged correlated disorder is studied in [240]. The contact process on aperiodic chains (instead of random chains) has been found to display double-logarithmic periodic oscillations via real-space renormalization [241]. The numerical study of the contact process on complex networks has revealed the importance of Griffiths phases and other rare region effects as a consequence of topological heterogeneity of the network [242]. Finally, the effect on random-field disorder on the Generalized contact process has been studied in [243].

B. Long range spreading

Spreading of epidemics with long range infections - which has a power-law distribution - can happen in different situations. This type of process can be modelled by the contact process in which the activation rates are parametrised as:

$$\lambda_{ij} = \Lambda_{ij} r_{ij}^{-(d+\sigma)}, \quad (54)$$

where r_{ij} is the Euclidean distance between site i and j , and Λ_{ij} are $O(1)$ i.i.d. quenched random variables, while the recovery rates μ_i are also i.i.d. quenched random variables as before.

The SDRG trajectories have been analysed [244] in the same way as that of the RTIM with LR interaction

of Sec.III. Analytical solution of the primary model in 1D, as well as numerical implementation of the renormalization with the maximum rule in 1D and 2D lead to identical critical scaling behavior as illustrated in Fig.2. In the language of the contact process, the following consequences have been obtained. Starting from a single infected site, the average survival probability is found to decay as $P(t) \sim t^{-d/z}$ up to multiplicative logarithmic corrections. Below the epidemic threshold, a Griffiths phase emerges, where the dynamical exponent z varies continuously with the control parameter and tends to $z_c = d + \sigma$ as the threshold is approached. At the threshold, the spatial extension of the infected cluster (in surviving trials) is found to grow as $R(t) \sim t^{1/z_c}$ with a multiplicative logarithmic correction, and the average number of infected sites in surviving trials is found to increase as $N_s(t) \sim (\ln t)^\chi$ with $\chi = 2$ in one dimension. These results have been confirmed by numerical Monte Carlo simulations [244]. We note that on a long-range connected network, the contact process has infinite disorder criticality [245].

C. Temporal disorder

The contact process in time-varying environmental noise, i.e. temporal disorder, has been considered in Refs. [246–248]. The system is spatially homogeneous, but the (nearest neighbour) activation and recovery rates are time dependent:

$$\lambda(t) = \lambda_n, \quad \mu(t) = \mu_n \quad (t_n < t < t_{n+1}). \quad (55)$$

In the mean-field approximation the time evolution of the density ρ of the active sites follows the differential equation:

$$\dot{\rho}(t) = [\lambda(t) - \mu(t)]\rho(t) - \lambda(t)\rho^2(t), \quad (56)$$

the solution of which in the interval $t_n < t < t_{n+1}$ for a given disorder realization is given by:

$$\rho_{n+1}^{-1} = a_n \rho_n^{-1} + c_n. \quad (57)$$

Here $\rho_n = \rho(t_n)$, $a_n = \exp[(\mu_n - \lambda_n)\Delta t]$ and the constant $c_n = (a_n - 1)\lambda_n/(\mu_n - \lambda_n)$.

The strong disorder (or strong noise) RG consists of iteratively decimating the weakest spreading and recovery segments, characterised by $a_i^\uparrow > 1$ and $a_i^\downarrow < 1$, respectively and $\Omega = \min(a_i^\uparrow, 1/a_i^\downarrow)$. The decimation equations are given by: $\tilde{a}^\uparrow = a_{i+1}^\uparrow a_i^\uparrow / \Omega$ (for $\Omega = 1/a_i^\downarrow$) and $1/\tilde{a}^\downarrow = (1/a_i^\downarrow)(1/a_{i-1}^\downarrow)/\Omega$ (for $\Omega = a_i^\uparrow$), which are equivalent to those of the RTIM in one dimension. Thus the critical behavior in the mean-field approximation is controlled by the IDFP of this model.

In finite dimensions the decimation equations are $\tilde{a}^\uparrow = a_{i+1}^\uparrow a_i^\uparrow / \Omega$ (for $\Omega = 1/a_i^\downarrow$) and $(1/\tilde{a}^\downarrow)^{1/D} = (1/a_i^\downarrow)^{1/D} + (1/a_{i-1}^\downarrow)^{1/D} - \Omega^{1/D}$ (for $\Omega = a_i^\uparrow$). Here in

the second equation, one takes into account that in finite dimensions by decimating $\Omega = a_i^\dagger$ the radii of the combined active clusters grows linearly in time. The elementary RG steps in this case are equivalent to that of the RTIM in one dimension with long-range interactions, see in Sec.III A and the RG trajectories are illustrated in Fig.2. The singular properties of the observables in the contact process can be found in [246, 248] and numerical calculations are performed in Ref. [248].

Temporal disorder at first-order non-equilibrium phase transitions has been studied in [249]. More generally, the effect of spatio-temporal disorder on various equilibrium and nonequilibrium critical points is discussed in [250].

XI. CLASSICAL MASTER EQUATIONS

The stochastic dynamics in random classical models usually displays a broad continuum of relevant time scales. The scaling between the characteristic time t and the linear length L can be either activated with some exponent ψ characterizing some Infinite Disorder Fixed Point

$$\ln t = L^\psi \quad (58)$$

or power-law with some dynamical exponent z that may vary continuously as a function of the model parameters

$$t = L^z \quad (59)$$

The limit $z \rightarrow \infty$ corresponds to the activated scaling (Eq 58) of the Infinite Disorder Fixed Point, but the whole region

$$1 < z < +\infty \quad (60)$$

can usually be well described by the Strong Disorder approximation.

A. Real-space renormalization for random walks in random media

As reviewed in [3], Strong Disorder RG procedures have been applied to various models of random walks in random media. The main principle is the iterative elimination of the fastest degree of freedom to obtain the effective dynamics for the slowest ones. In this field, new developments since 2005 include one-dimensional random walks with dilute absorbers [251] or with long-range connections [252], random walks on strips [253, 254] and on arbitrary networks [255], and in two-dimensional self-affine random potentials [256].

B. RG in configuration-space for the dynamics of classical many-body models

The real-space SDRG for random walks in random media has been generalized into the configuration-space

SDRG [257] for any classical master equation governing the dynamics of the probability $P_t(\mathcal{C})$ to be in configuration \mathcal{C} at time t

$$\frac{dP_t(\mathcal{C})}{dt} = \sum_{\mathcal{C}'} P_t(\mathcal{C}') W(\mathcal{C}' \rightarrow \mathcal{C}) - P_t(\mathcal{C}) W_{out}(\mathcal{C}) \quad (61)$$

where $W(\mathcal{C}' \rightarrow \mathcal{C})$ represents the transition rate per unit time from configuration \mathcal{C}' to \mathcal{C} while

$$W_{out}(\mathcal{C}) \equiv \sum_{\mathcal{C}'} W(\mathcal{C} \rightarrow \mathcal{C}') \quad (62)$$

denotes the total exit rate out of configuration \mathcal{C} . The SDRG rule consists in the elimination of the configuration \mathcal{C}^* with the highest total exit rate $W_{out}(\mathcal{C}^*)$ to obtain the new renormalized transition rates between surviving configuration

$$W^{new}(\mathcal{C}_i \rightarrow \mathcal{C}_j) = W^{old}(\mathcal{C}_i \rightarrow \mathcal{C}_j) + W^{old}(\mathcal{C}_i \rightarrow \mathcal{C}^*) \frac{W^{old}(\mathcal{C}^* \rightarrow \mathcal{C}_j)}{W_{out}(\mathcal{C}^*)} \quad (63)$$

and the new exit rates

$$W_{out}^{new}(\mathcal{C}_i) = W_{out}^{old}(\mathcal{C}_i) - W^{old}(\mathcal{C}_i \rightarrow \mathcal{C}^*) \frac{W^{old}(\mathcal{C}^* \rightarrow \mathcal{C}_i)}{W_{out}(\mathcal{C}^*)} \quad (64)$$

The physical interpretation of this procedure is as follows : the time spent in the decimated configuration \mathcal{C}^* is neglected with respects to the other time scales remaining in the system; the remaining configurations represents some 'valleys' in configuration space that takes into account all the previously decimated configurations. As a consequence of the multiplicative structure of the renormalization rule of Eq 63, the renormalized rates $W(\mathcal{C} \rightarrow \mathcal{C}')$ can rapidly become very small and the appropriate variables are the logarithms of the transition rates, called 'barriers'

$$B(\mathcal{C} \rightarrow \mathcal{C}') \equiv -\ln W(\mathcal{C} \rightarrow \mathcal{C}') \quad (65)$$

This SDRG procedure has been applied numerically to interfaces in two-dimensional random media [257–259] with possibly driving [260]. Note that the idea to eliminate fast degrees of freedom in classical master equations is very natural and has been thus developed independently in many other contexts (see the recent review [261] and references therein). The SDRG rules above are actually similar to the exact RG rules concerning first-passage times [262] where the application to spin-glasses is discussed.

As a final remark, it is important to stress that for many-body classical models, the fact that the above RG rules are defined in configuration space clearly limits the numerical implementation to small sizes. As a consequence, various other types of real-space RG procedures have been developed for the dynamics of classical spin models, in particular boundary-RG for the dynamics in $d = 1$ [263] and on the Cayley tree [264] or block-RG for the dynamics of Long-Ranged ferromagnetic [265] or Spin-Glass models [266, 267].

XII. RANDOM CLASSICAL OSCILLATORS

A. Random elastic networks

The model of random masses m_i connected by random springs K_{ij} is one of the oldest problem in the field of localization of classical disordered models [268]. One is interested into the Newton equations of motions for the displacements $u_i(t)$

$$m_i \frac{d^2 u_i}{dt^2} = \sum_j K_{ij} (u_j - u_i) \quad (66)$$

for various geometries. The issue of the localization properties of the eigenmodes (phonon localization) is related to the Anderson localization properties of tight-binding models discussed in Section IX even if they are some differences (see for instance [269] and references therein).

Various slightly different SDRG rules have been proposed : the idea is to eliminate iteratively either only the masses [270], or only the couplings [271], or both [272] as we now describe. One first needs to identify the local degree of freedom oscillating with the highest frequency.

The frequency $\Omega_{i,j}$ associated to the spring $K_{i,j}$ between two masses m_i and m_j is defined by [272]

$$\Omega_{i,j}^2 \equiv K_{i,j} \left(\frac{1}{m_i} + \frac{1}{m_j} \right) \quad (67)$$

while the frequency Ω_i associated to the mass m_i connected to the springs K_{ij} is given by [272]

$$\Omega_i^2 \equiv \frac{1}{m_i} \sum_j K_{i,j} \quad (68)$$

The renormalization scale Ω is defined as the highest local frequency remaining in the system among all the frequencies associated with masses or spring constants.

$$\Omega \equiv \max\{\Omega_i, \Omega_{i,j}\} \quad (69)$$

If the highest frequency $\Omega = \Omega_{i_0, j_0}$ is associated with the spring constant K_{i_0, j_0} , the two masses m_{i_0} and m_{j_0} are replaced by their center of mass $G(i_0, j_0)$ of mass

$$m_{G(i_0, j_0)} = m_{i_0} + m_{j_0} \quad (70)$$

and the spring constants are replaced by spring constants linked to their center of mass

$$K_{j, G(i_0, j_0)} = K_{j, i_0} + K_{j, j_0} \quad (71)$$

This renormalization step thus constructs a cluster of strongly-coupled masses oscillating together.

If the highest frequency $\Omega = \Omega_{i_0}$ is associated with the mass i_0 , the mass m_{i_0} is eliminated, and the spring constants are renormalized according to

$$K_{i,j}^{new} = K_{i,j} + \frac{K_{i, i_0} K_{i_0, j}}{\sum_n K_{i_0, n}} \quad (72)$$

This renormalization step thus constructs an isolated localized oscillating mode.

These SDRG rules coincide with the exact Aoki RG rules at zero frequency $\omega = 0$ [272], so they are expected to remain a good approximation at low frequency.

The SDRG approach has been applied to complex networks to analyze their localization properties [270] as well as to some matrix model in relation with slow relaxation in glasses [271].

B. Synchronisation of interacting non-linear dissipative classical oscillators

In the field of emergent collective structures in nonequilibrium systems, the spontaneous synchronization of interacting nonlinear oscillators is one of the most studied phenomenon [273–275]. Each oscillator is characterized by its mass m_i and its own frequency ω_i , while the interactions between oscillators are described by couplings K_{ij} that define the geometry of the network of oscillators. The dynamical equations for the phases $\theta_i(t)$ of the oscillators are written in the dissipative limit (first order in time)

$$m_i \frac{d\theta_i}{dt} = m_i \omega_i + \sum_j K_{ij} \sin(\theta_j - \theta_i) \quad (73)$$

The aim of the SDRG procedure [276, 277] is to construct clusters of frequency-synchronized-oscillators. The two decimation possibilities are as follows. The decimation of a coupling K_{ij} corresponds to the synchronization of the two corresponding oscillators and its replacement by a single renormalized oscillator. The decimation of a frequency ω_i means that the corresponding oscillator rotates freely and does not contribute to the global synchronization. The details of the SDRG rules and the numerical results are described in [276, 277].

XIII. OTHER CLASSICAL MODELS

A. Equilibrium properties of random systems

SDRG has been also used to analyze the equilibrium phase transitions of various classical systems, as reviewed in [3], while more recent applications include the randomly layered Heisenberg magnet [278], the wetting transition on the Cayley tree [279], the DNA denaturation transition [280]. In the field of classical spin-glasses, some SDRG procedure have been also introduced to study the spin-glass phase of the Long-Ranged Spin-Glass chain [281] or the fractal dimension of interfaces in Short-Ranged Spin-Glasses as a function of the dimension d [282–284].

B. Extremes of stochastic processes

As explained in detail in the Review [3], SDRG procedures are closely related to the statistics of extrema of some random processes associated to the disorder variables : for instance, the Fisher solution is directly related to the statistics of extrema of the Brownian motion[18]. Reciprocally, the Extreme Value Statistics of various stochastic processes can be analyzed via SDRG [285] (while the Extreme Value Statistics of independent variables is analyzed via RG in the series of works [286]). Some coagulation model with extremal dynamics has been also studied [287], in relation with previous works reviewed in Appendix E of [3]).

XIV. CONCLUSION

In summary, we have reviewed the new developments of Strong Disorder RG methods since 2005. For the quantum phase transitions of ground states, the critical properties have been described for short-ranged models in higher dimensions $d > 1$ and for long-ranged models. The scaling of the entanglement entropy has been discussed both for critical ground-states and after quantum quenches. In Many-Body-Localized phases, we have explained how the SDRG procedure has been extended into RSRG-X procedure to construct the whole set excited states and into the RSRG-t procedure for the unitary dynamics. Other generalizations of the SDRG approach concern non-isolated quantum models, namely periodically driven models (Floquet dynamics) or dissipa-

tive models (coupling to external baths). We have then focused on the recent progress for classical disordered models, with the contact process for epidemic spreading, the strong disorder renormalization procedure for general master equations, the localization properties of random elastic networks and the synchronization of interacting non-linear dissipative oscillators.

In conclusion, SDRG methods have flourished over the years well beyond their initial scopes, and we thus expect that they will continue to be developed even further in the future.

ACKNOWLEDGMENTS

It is a pleasure to thank collaborations and discussions with several colleagues: F. Alet, E. Altman, G. Biroli, P. Calabrese, J. Cardy, C. Chatelain, L. Cugliandolo, U. Divakaran, D. Fisher, T. Garel, J.A. Hoyos, D. Huse, R. Juhász, I.A. Kovács, N. Laflorencie, P. Lajkó, P. Le Doussal, Y.-C. Lin, J. M. Luck, R. Mélin, J. E. Moore, M.A. Moore, G. Ódor, F. Pollmann, G. Refael, H. Rieger, G. Roósz, A. Sandvik, L. Santen, G. Schehr, M. Schiro, Zs. Szatmári, L. Turban, R. Vasseur, T. Vojta and Z. Zimborás.

The work of F.I. has been supported by the Hungarian Scientific Research Fund under Grants No. K109577, No. K115959 and No. KKP-126749. He thanks the IPhT Saclay for hospitality.

Author contribution statement:

F.I. and C.M. have contributed equally by planning the structure of the review and by writing the text.

-
- [1] S.-K. Ma, C. Dasgupta, and C.-k. Hu, Random Antiferromagnetic Chain, Phys. Rev. Lett. 43, 1434 (1979) ; C. Dasgupta and S.-K. Ma Low-temperature properties of the random Heisenberg antiferromagnetic chain Phys. Rev. B 22, 1305 (1980).
 - [2] D. S. Fisher, Random transverse field Ising spin chains, Phys. Rev. Lett. 69, 534 (1992) ; D. S. Fisher, Critical behavior of random transverse-field Ising spin chains Phys. Rev. B 51, 6411 (1995).
 - [3] F. Iglói and C. Monthus, Strong disorder RG approach of random systems, Phys. Rep. 412, 277 (2005).
 - [4] I. A. Kovács and F. Iglói, Critical behavior and entanglement of the random transverse-field Ising model between one and two dimensions, Phys. Rev. B 80, 214416 (2009).
 - [5] I. A. Kovács and F. Iglói, Renormalization group study of the two-dimensional random transverse-field Ising model, Phys. Rev. B 82, 054437 (2010).
 - [6] I. A. Kovács and F. Iglói, Infinite disorder scaling of random quantum magnets in three and higher dimensions, Phys. Rev. B 83, 174207 (2011).
 - [7] I. A. Kovács and F. Iglói, Renormalization group study of random quantum magnets, J. Phys.: Condens. Matter 23, 404204 (2011).
 - [8] S. Iyer, D. Pekker and G. Refael, A Mott Glass to Superfluid Transition for Random Bosons in Two Dimensions Phys. Rev. B 85, 094202 (2012).
 - [9] T. Vojta Rare region effects at classical, quantum and nonequilibrium phase transitions J. Phys. A: Math. Gen. 39 R143 (2006).
 - [10] T. Vojta, Phases and phase transitions in disordered quantum systems, arxiv: 1301.7746.
 - [11] S. Garnerone, N. Jacobson, S. Haas and P. Zanardi, Fidelity Approach to the Disordered Quantum XY Model Phys. Rev. Lett. 102 057205 (2009); N. T. Jacobson, S. Garnerone, S. Haas and P. Zanardi, Scaling of the fidelity susceptibility in a disordered quantum spin chain Phys. Rev. B 79, 184427 (2009).
 - [12] C. Monthus, Pure and Random Quantum Ising Chain : Shannon and Rényi entropies of the ground state via real space renormalization J. Stat. Mech. P04007 (2015).
 - [13] I. A. Kovács and F. Iglói, Boundary critical phenomena of the random transverse Ising model in $D \geq 2$ dimensions, Phys. Rev. B 87, 024204 (2013).

- [14] R. Juhász, Critical behavior of models with infinite disorder at a star junction of chains, *J. Stat. Mech.* P08005 (2014)
- [15] C. Monthus, Star junctions and watermelons of pure or random quantum Ising chains : finite-size properties of the energy gap at criticality, *J. Stat. Mech.* P06036 (2015)
- [16] F. Iglói and I. A. Kovács, Transverse-spin correlations of the random transverse-field Ising model, *Phys. Rev. B* 97, 094205 (2018)
- [17] J. Galambos, *The Asymptotic Theory of Extreme Order Statistics* (Wiley, New York, 1978).
- [18] R. Juhász, Y-C. Lin, F. Iglói, Strong Griffiths singularities in random systems and their relation to extreme value statistics, *Phys. Rev. B* 73, 224206 (2006)
- [19] Y-C. Lin, F. Iglói and H. Rieger, Entanglement entropy at infinite randomness fixed points in higher dimensions, *Phys. Rev. Lett.* 99, 147202 (2007)
- [20] R. Yu, H. Saleur and S. Haas, Entanglement Entropy in the Two-Dimensional Random Transverse Field Ising Model, *Phys. Rev. B* 77, 140402(R) (2008)
- [21] C.R. Laumann, D.A. Huse, A.W.W. Ludwig, G. Refael, S. Trebst and M. Troyer, Strong-disorder renormalization for interacting non-Abelian anyon systems in two dimensions, *Phys. Rev. B* 85, 224201 (2012)
- [22] O. Dimitrova and M. Mézard, The cavity method for quantum disordered systems: from transverse random field ferromagnets to directed polymers in random media, *J. Stat. Mech.* P01020 (2011).
- [23] C. Monthus and T. Garel, Random Transverse Field Ising Model in dimension $d > 1$: scaling analysis in the disordered phase from the Directed Polymer model, *J. Phys. A: Math. Theor.* 45, 095002 (2012).
- [24] C. Monthus and T. Garel, Random Transverse Field Ising Model in dimension $d = 2, 3$: Infinite Disorder scaling via a non-linear transfer approach *J. Stat. Mech.* P01008 (2012).
- [25] C. Monthus and T. Garel, Strong Disorder RG principles within a fixed cell-size real space renormalization : application to the Random Transverse Field Ising model on various fractal lattices, *J. Stat. Mech.* P05002 (2012).
- [26] C. Monthus and T. Garel, Random Transverse Field Ising model on the Cayley Tree : analysis via Boundary Strong Disorder Renormalization, *J. Stat. Mech.* P10010 (2012).
- [27] C. Monthus and T. Garel, Random Transverse Field Ising model in $d = 2$: analysis via Boundary Strong Disorder Renormalization, *J. Stat. Mech.* P09016 (2012).
- [28] R. Miyazaki and H. Nishimori, Real-space renormalization-group approach to the random transverse-field Ising model in finite dimensions, *Phys. Rev. E* 87, 032154 (2013).
- [29] L. B. Ioffe and M. Mézard, Disorder-Driven Quantum Phase Transitions in Superconductors and Magnets, *Phys. Rev. Lett.* 105, 037001 (2010); M.V. Feigel'man, L. B. Ioffe and M. Mézard, Superconductor-Insulator transition and energy localization, *Phys. Rev. B* 82, 184534 (2010).
- [30] For a review see, A. Dutta, G. Aeppli, B. K. Chakrabarti, U. Divakaran, T. F. Rosenbaum and D. Sen, *Quantum Phase Transitions in Transverse Field Spin Models: From Statistical Physics to Quantum Information*, Cambridge University Press, Cambridge, (2015)
- [31] A. Friedenauer, H. Schmitz, J. T. Glueckert, D. Porras, and T. Schaetz, Simulating a quantum magnet with trapped ions, *Nat. Phys.* 4, 757 (2008).
- [32] K. Kim, M.-S. Chang, S. Korenblit, R. Islam, E. E. Edwards, J. K. Freericks, G.-D. Lin, L.-M. Duan, and C. Monroe, Quantum simulation of frustrated Ising spins with trapped ions, *Nature (London)* 465, 590 (2010).
- [33] R. Islam, E. E. Edwards, K. Kim, S. Korenblit, C. Noh, H. Carmichael, G.-D. Lin, L.-M. Duan, C.-C. J. Wang, J. Freericks, and C. Monroe, Onset of a Quantum Phase Transition with a Trapped Ion Quantum Simulator, *Nat. Commun.* 2, 377 (2011).
- [34] J.W. Britton, B. C. Sawyer, A. C. Keith, C. J. Wang, J. K. Freericks, H. Uys, M. J. Biercuk, and J. J. Bollinger, Engineered two-dimensional Ising interactions in a trapped-ion quantum simulator with hundreds of spins, *Nature (London)* 484, 489 (2012).
- [35] R. Islam, C. Senko, W. C. Campbell, S. Korenblit, J. Smith, A. Lee, E. E. Edwards, C.-C. J. Wang, J. K. Freericks, and C. Monroe, Emergence and Frustration of Magnetism with Variable-Range Interactions in a Quantum Simulator, *Science* 340, 583 (2013).
- [36] R. Juhász, I. A. Kovács, and F. Iglói, Random transverse-field Ising chain with long-range interactions *Europhys. Lett.* 107, 47008 (2014).
- [37] E. Altman, Y. Kafri, A. Polkovnikov, and G. Refael, Phase transition of one dimensional bosons with strong disorder *Phys. Rev. Lett.* 93, 150402 (2004)
- [38] E. Altman, Y. Kafri, A. Polkovnikov, and G. Refael, Superfluid-insulator transition of disordered bosons in one-dimension *Phys. Rev. B* 81, 174528 (2010).
- [39] I. A. Kovács, R. Juhász and F. Iglói, Long-range random transverse-field Ising model in three dimensions, *Phys. Rev. B* 93, 184203 (2016).
- [40] A. Dutta and R. Loganayagam, Effect of long-range connections on an infinite randomness fixed point associated with the quantum phase transitions in a transverse Ising model *Phys. Rev. B* 75 052405 (2007); U. Divakaran and A. Dutta, Long-range connections, quantum magnets and dilute contact processes *Physica A* 384 39 (2007)
- [41] C. Monthus, Dyson hierarchical quantum ferromagnetic Ising chain with pure or random transverse fields *J. Stat. Mech.* P05026 (2015).
- [42] C. Monthus, Dyson Hierarchical Long-Ranged Quantum Spin-Glass via real-space renormalization *J. Stat. Mech.* P10024 (2015).
- [43] R. Juhász, Infinite-disorder critical points of models with stretched exponential interactions, *J. Stat. Mech.* P09027 (2014).
- [44] N. Moure, S. Haas, S. Kettemann, Many-Body Localization Transition in Random Quantum Spin Chains with Long-Range Interactions, *Europhysics Letters* 111, 27003 (2015).
- [45] N. Moure, H.Y. Lee, S. Haas, R. N. Bhatt and S. Kettemann, Disordered Quantum Spin Chains with Long-Range Antiferromagnetic Interactions, *Phys. Rev. B* 97, 014206 (2018)
- [46] T. Vojta and J. A. Hoyos, Magnetic Grüneisen ratio of the random transverse-field Ising chain, *phys. stat. sol (b)* 247, 525 (2010)
- [47] V. Shivamoggi, G. Refael and J. E. Moore, Majorana fermion chain at the Quantum Spin Hall edge, *Phys. Rev. B* 82, 041405(R) (2010).

- [48] Y.P. Lin, Y.J. Kao, P. Chen, Y.C. Lin, Griffiths Singularities in the Random Quantum Ising Antiferromagnet: A Tree Tensor Network Renormalization Group Study, *Phys. Rev. B* 96, 064427 (2017).
- [49] P. Lajkó and F. Iglói, Numerical study of the random quantum Ising antiferromagnetic chain (to be published)
- [50] P. Goswami, D. Schwab and S. Chakravarty, Rounding by Disorder of First-Order Quantum Phase Transitions: Emergence of Quantum Critical Points, *Phys. Rev. Lett.* 100, 015703 (2008).
- [51] F. Hrahsheh, J. A. Hoyos and T. Vojta, Rounding of a first-order quantum phase transition to a strong-coupling critical point, *Phys. Rev. B* 86, 214204 (2012).
- [52] F. Hrahsheh, R. Narayanan, J. A. Hoyos and T. Vojta, Strong-randomness infinite-coupling phase in a random quantum spin chain, *Phys. Rev. B* 89, 014401 (2014).
- [53] H. Barghathi, F. Hrahsheh, J. A. Hoyos, R. Narayanan and T. Vojta, Strong-randomness phenomena in quantum Ashkin-Teller models, *Physica Scripta* T165, 014040 (2015).
- [54] Q. Zhu, X. Wan, R. Narayanan, J. A. Hoyos and T. Vojta, Emerging criticality in the disordered three-color Ashkin-Teller model, *Phys. Rev. B* 91, 224201 (2015)
- [55] A. Bellafard and S. Chakravarty Activated scaling in disorder-rounded first-order quantum phase transitions *Phys. Rev. B* 94, 094408 , (2016)
- [56] C. Chatelain and D. Voliotis, Numerical evidence of the double-Griffiths phase of the random quantum Ashkin-Teller chain, *The European Physical Journal B* 89, 18 (2016).
- [57] A. K. Ibrahim and T. Vojta, Monte Carlo simulations of the disordered three-color quantum Ashkin-Teller chain, *Phys. Rev. B* 95, 054403 (2017).
- [58] C. A. Lamas, D. C. Cabra, M. D. Grynberg, and G. L. Rossini, Comparison between disordered quantum spin-1/2 chains, *Phys. Rev. B* 74, 224435 (2006).
- [59] J. Kokalj, J. Herbrych, A. Zheludev and P. Prelovsek, Antiferromagnetic order in weakly coupled random spin chains *Phys. Rev. B* 91, 155147 (2015)
- [60] R. Yu, T. Roscilde and S. Haas, Quantum disorder and Griffiths singularities in bond-diluted two-dimensional Heisenberg antiferromagnets, *Phys. Rev. B* 73, 064406 (2006).
- [61] N. Ma, A. W. Sandvik and D.X. Yao Criticality and Mott glass phase in a disordered two-dimensional quantum spin system *Phys. Rev. B* 90 104425 (2014)
- [62] L. Liu, H. Shao, Y.C. Lin, W. Guo and A. W. Sandvik, Random-Singlet Phase in Disordered Two-Dimensional Quantum Magnets, arxiv 1804.06108.
- [63] S. Zhou, J. A. Hoyos, V. Dobrosavljevic and E. Miranda, Valence-bond theory of highly disordered quantum antiferromagnets, *Europhys. Lett.* 87, 27003 (2009).
- [64] V. L. Quito, J. A. Hoyos and E. Miranda, Emergent SU(3) symmetry in random spin-1 chains, *Phys. Rev. Lett.* 115, 167201 (2015).
- [65] V. L. Quito, J. A. Hoyos and E. Miranda, Random SU(2)-symmetric spin-S chains, *Phys. Rev. B* 94, 064405 (2016).
- [66] V. L. Quito, Pedro L. S. Lopes, J. A. Hoyos and E. Miranda Highly-symmetric random one-dimensional spin models, arXiv:1711.04783.
- [67] V. L. Quito, Pedro L. S. Lopes, J. A. Hoyos and E. Miranda Emergent SU(N) symmetry in disordered SO(N) spin chains, arXiv:1711.04781.
- [68] P. Lajkó, Renormalization-group investigation of the S=1 random antiferromagnetic Heisenberg Chain, *Int. J. Mod. Phys. C* 17, 1739 (2006).
- [69] A. Lavarello and G. Roux, Localization of Spinons in Random Majumdar-Ghosh Chains, *Phys. Rev. Lett.* 110, 087204 (2013).
- [70] M. C. Strinati, D. Rossini, R. Fazio and A. Russomanno, Resilience of hidden order to symmetry-preserving disorder, *Phys. Rev. B* 96, 214206 (2017).
- [71] R. Mélin and F. Iglói, Strongly disordered Hubbard model in one dimension: spin and orbital infinite randomness and Griffiths phases, *Phys. Rev. B* 74, 155104 (2006).
- [72] Y.C. Lin, H. Rieger, N. Laflorencie and F. Iglói, Strong disorder renormalization group study of S=1/2 Heisenberg antiferromagnet layers/bilayers with bond randomness, site dilution and dimer dilution, *Phys. Rev. B* 74, 024427 (2006).
- [73] N.E. Bonesteel and K. Yang, Infinite-Randomness Fixed Points for Chains of Non-Abelian Quasiparticles, *Phys. Rev. Lett.* 99, 140405 (2007).
- [74] L. Fidkowski, G. Refael, N. Bonesteel and J. Moore, c-theorem violation for effective central charge of infinite-randomness fixed points, *Phys. Rev. B* 78, 224204 (2008).
- [75] L. Fidkowski, G. Refael, H.H. Lin and P. Titum, Permutation Symmetric Critical Phases in Disordered Non-Abelian Anyonic Chains, *Phys. Rev. B* 79, 155120 (2009).
- [76] G. Refael, J. E. Moore, Criticality and entanglement in random quantum systems, *J. Phys. A: Math. Theor.* 42 504010 (2009).
- [77] E. Altman, Y. Kafri, A. Polkovnikov and G. Refael The insulating phases and superfluid-insulator transition of disordered boson chains, *Phys. Rev. Lett.* 100, 170402 (2008)
- [78] V. Gurarie, G. Refael and J. T. Chalker, Excitations of the One Dimensional Bose-Einstein Condensates in a Random Potential, *Phys. Rev. Lett.* 101, 170407 (2008).
- [79] E. Altman, Y. Kafri, A. Polkovnikov and G. Refael, Superfluid-insulator transition of disordered bosons in one-dimension, *Phys. Rev. B* 81, 174528 (2010).
- [80] R. Vosk and E. Altman, Superfluid-insulator transition of ultracold bosons in disordered one-dimensional traps, *Phys. Rev. B* 85 024531 (2012).
- [81] S. Iyer, D. Pekker and G. Refael, Susceptibility at the Superfluid-Insulator Transition for One-Dimensional Disordered Bosons, *Phys. Rev. B* 88, 220501 (2013).
- [82] G. Refael and E. Altman, Strong disorder renormalization group primer and the superfluid-insulator transition, *Comptes Rendus Physique* 14 , 725 (2013).
- [83] F. Hrahsheh and T. Vojta, Disordered bosons in one dimension: from weak to strong randomness criticality, *Phys. Rev. Lett.* 109, 265303 (2012)
- [84] E. V. H. Doggen, G. Lemari, S. Capponi and N. Laflorencie, Weak Versus Strong Disorder Superfluid-Bose Glass Transition in One Dimension, *Phys. Rev. B* 96, 180202 (2017).
- [85] A. Del Maestro, B. Rosenow, M. Muller and S. Sachdev, Infinite Randomness Fixed Point of the Superconductor-Metal Quantum Phase Transition, *Phys. Rev. Lett.* 101, 035701 (2008).
- [86] A. Del Maestro, B. Rosenow, J. A. Hoyos and T. Vojta,

- Dynamical Conductivity at the Dirty Superconductor-Metal Quantum Phase Transition, *Phys. Rev. Lett.* 105, 145702 (2010).
- [87] D. Nozadze and T. Vojta, Numerical method for disordered quantum phase transitions in the large- N limit, *Physica Status Solidi* 251 675 (2014).
- [88] A. K. Ibrahim and T. Vojta, Monte Carlo simulations of a disordered superconductor-metal quantum phase transition, arXiv:1808.00094
- [89] G. Ramirez, J. Rodriguez-Laguna and G. Sierra, From conformal to volume-law for the entanglement entropy in exponentially deformed critical spin $1/2$ chains, *J. Stat. Mech.* P10004 (2014).
- [90] G. Ramirez, J. Rodriguez-Laguna and G. Sierra, Entanglement over the rainbow, *J. Stat. Mech.* P06002 (2015).
- [91] J. Rodriguez-Laguna, S. N. Santalla, G. Ramirez and G. Sierra, Entanglement in correlated random spin chains, RNA folding and kinetic roughening, *New J. Phys.* 18 073025 (2016).
- [92] J. Rodriguez-Laguna, J. Dubail, G. Ramirez, P. Calabrese, G. Sierra, More on the rainbow chain: entanglement, space-time geometry and thermal states, *J. Phys. A: Math. Theor.* 50 164001 (2017).
- [93] G. Ramirez, Quantum Entanglement In Inhomogeneous 1D Systems, AIP Conference Proceedings 1950, 030007 (2018).
- [94] V. Alba, S. N. Santalla, P. Ruggiero, J. Rodriguez-Laguna, P. Calabrese and G. Sierra, Unusual area-law violation in random inhomogeneous systems, arxiv : 1807.04179
- [95] P. Calabrese, J. Cardy and B. Doyon (Eds. of the special issue) Entanglement entropy in extended quantum systems, *J. Phys. A* 42 500301 (2009).
- [96] L. Amico, R. Fazio, A. Osterloh, and V. Vedral, Entanglement in many-body systems *Rev. Mod. Phys.* 80, 517 (2008).
- [97] N. Laflorencie, Quantum entanglement in condensed matter systems, *Phys. Rep.* 643, 1 (2016).
- [98] C. Holzhey, F. Larsen, and F. Wilczek, Geometric and Renormalized Entropy in Conformal Field Theory *Nucl. Phys. B* 424, 443 (1994).
- [99] G. Vidal, J. I. Latorre, E. Rico, and A. Kitaev, Entanglement in Quantum Critical Phenomena, *Phys. Rev. Lett.* 90, 227902 (2003).
- [100] P. Calabrese and J. Cardy, Entanglement Entropy and Quantum Field Theory, *J. Stat. Mech.* (2004) P06002.
- [101] P. Calabrese and A. Lefevre, Entanglement spectrum in one-dimensional systems, *Phys. Rev. A* 78, 32329 (2008).
- [102] R. Vasseur, A. Roshani, S. Haas and H. Saleur, Healing of Defects in Random Antiferromagnetic Spin Chains, *EPL* 119 50004 (2017).
- [103] R. Vasseur and J. E. Moore, Multifractal Orthogonality Catastrophe in 1D Random Quantum Critical Points, *Phys. Rev. B* 92, 054203 (2015)
- [104] J. A. Hoyos, A. P. Vieira, N. Laflorencie, E. Miranda, Correlation amplitude and entanglement entropy in random spin chains, *Phys. Rev. B* 76, 174425 (2007).
- [105] G. Refael and J. E. Moore, Entanglement entropy of the random spin-1 Heisenberg chain, *Phys. Rev. B* 76, 024419 (2007).
- [106] A. Saguia, M. S. Sarandy, B. Boechat and M. A. Continentino, Entanglement Entropy in Random Quantum Spin- S Chains, *Phys. Rev. A* 75, 052329 (2007).
- [107] A. Saguia and M. S. Sarandy, Nonadditive entropy for random quantum spin- S chains, *Phys. Lett. A* 374, 3384 (2010).
- [108] F. Iglói and R. Juhász, Exact relationship between the entanglement entropies of XY and quantum Ising chains, *Europhys. Lett.* 81, 57003 (2008).
- [109] F. Iglói, Y.C. Lin, H. Rieger and C. Monthus, Finite-size scaling of pseudo-critical point distributions in the random transverse-field Ising chain, *Phys. Rev. B* 76, 064421 (2007).
- [110] R. Santachiara, Increasing of entanglement entropy from pure to random quantum critical chains, *J. Stat. Mech.* L06002 (2006).
- [111] J. A. Hoyos, N. Laflorencie, A. P. Vieira and T. Vojta, Protecting clean critical points by local disorder correlations, *Europhys. Lett.* 93, 30004 (2011).
- [112] J. C. Getelina, F. C. Alcaraz, J. A. Hoyos, Entanglement properties of correlated random spin chains and similarities with conformal invariant systems, *Phys. Rev. B* 93, 045136 (2016).
- [113] R. Juhász, I. A. Kovács, G. Roósz and F. Iglói, Entanglement between random and clean quantum spin chains, *J. Phys. A: Math. Theor.* 50 , 324003 (2017).
- [114] R. Juhász, Entanglement across extended random defects in the XX spin chain, *J. Stat. Mech.* 083107 (2017).
- [115] R. Juhász, J. M. Oberreuter and Z. Zimborás, Entanglement Entropy of Disordered Quantum Wire Junctions, arXiv:1808.02576
- [116] A. P. Vieira, Aperiodic quantum XXZ chains: Renormalization-group results, *Phys. Rev. B* 71, 134408 (2005).
- [117] F. J. O. Filho, M. S. Faria and A. P. Vieira, Strong disorder renormalization group study of aperiodic quantum Ising chains, *J. Stat. Mech.* P03007 (2012).
- [118] H. L. Casa Grande, N. Laflorencie, F. Alet and A. P. Vieira, Analytical and numerical studies of disordered spin-1 Heisenberg chains with aperiodic couplings, *Phys. Rev. B* 89, 134408 (2014).
- [119] A. P. Vieira and J. A. Hoyos, Localization and emergent dimerization in aperiodic quantum spin chains, arXiv:1804.05108
- [120] F. Iglói, R. Juhász and Z. Zimborás, Entanglement entropy of aperiodic quantum spin chains, *Europhys. Lett.* 79, 37001 (2007).
- [121] R. Juhász and Z. Zimborás, Entanglement entropy in aperiodic singlet phases, *J. Stat. Mech.* P04004 (2007).
- [122] J. A. Hoyos and G. Rigolin, Quantum channels in random spin chains *Phys. Rev. A* 74, 062324 (2006).
- [123] J. C. Getelina, T. R. de Oliveira, J. A. Hoyos Violation of the Bell inequality in quantum critical random spin- $1/2$ chains arXiv:1711.10005
- [124] J. Hide, Concurrence in disordered systems *J. Phys. A* 45 115302 (2012).
- [125] M. Fagotti, P. Calabrese and J. E. Moore, Entanglement spectrum of random-singlet quantum critical points, *Phys. Rev. B* 83, 045110 (2011).
- [126] G. Ramirez, J. Rodriguez-Laguna and G. Sierra, Entanglement in low-energy states of the random-hopping model, *J. Stat. Mech.* P07003 (2014).
- [127] H. Tran and N. E. Bonesteel Valence bond entanglement and fluctuations in random singlet phases *Phys. Rev. B* 84 144420 (2011)
- [128] T. Devakul, S. N. Majumdar and D. A. Huse, Probability distribution of the entanglement across a cut at

- an infinite-randomness fixed point, *Phys. Rev. B* 95, 104204 (2017).
- [129] G. Torlai, K. D. McAlpine, G. De Chiara Schmidt gap in random spin chains, arXiv:1805.07404.
- [130] P. Ruggiero, V. Alba and P. Calabrese, The entanglement negativity in random spin chains, *Phys. Rev. B* 94, 035152 (2016).
- [131] F. Alet, S. Capponi, N. Laflorencie and M. Mambrini, Valence Bond Entanglement Entropy, *Phys. Rev. Lett.* 99, 117204 (2007).
- [132] Y.C. Lin and A. W. Sandvik, Definitions of entanglement entropy of spin systems in the valence-bond basis, *Phys. Rev. B* 82, 224414 (2010).
- [133] Y.-R. Shu, D.-X. Yao, C.-W. Ke, Y.-Ch. Lin and A. W. Sandvik, Properties of the random-singlet phase: from the disordered Heisenberg chain to an amorphous valence-bond solid, *Phys. Rev. B* 94, 174442 (2016).
- [134] I. A. Kovács and F. Iglói, Universal logarithmic terms in the entanglement entropy of 2d, 3d and 4d random transverse-field Ising models, *EPL* 97, 67009 (2012).
- [135] T. Senthil and S. Sachdev, Higher Dimensional Realizations of Activated Dynamic Scaling at Random Quantum Transitions, *Phys. Rev. Lett.* 77, 5292 (1996).
- [136] T. Vojta and J. A. Hoyos, Quantum Phase Transitions on Percolating Lattices, arXiv:0707.0658
- [137] I. A. Kovács, F. Iglói and J. Cardy, Corner contribution to percolation cluster numbers, *Phys. Rev. B* 86, 214203 (2012).
- [138] J. Cardy and I. Peschel, Finite-size dependence of the free energy in two-dimensional critical systems *Nucl. Phys. B*, 300 , 377 (1988).
- [139] I. A. Kovács and F. Iglói, Corner contribution to percolation cluster numbers in three dimensions, *Phys. Rev. B*. 89 174202 (2014).
- [140] J. I. Cirac and F. Verstraete, Renormalization and tensor product states in spin chains and lattices, *J. Phys. A: Math. Theor.* 42, 504004 (2009).
- [141] O. Gittsovich, R. Hubener, E. Rico and H.J. Briegel, Local renormalization method for random systems *New J. Phys.* 12, 025020 (2010).
- [142] A. M. Goldsborough and R. A. Romer, Self-assembling tensor networks and holography in disordered spin chains, *Phys. Rev. B* 89, 214203 (2014).
- [143] A. M. Goldsborough and G. Evenbly, Entanglement renormalization for disordered systems, *Phys. Rev. B* 96, 155136 (2017).
- [144] C. Chatelain, Quantifying and improving the accuracy of the Matrix Product Operator Renormalization Group of random spin chains, arxiv: 1807.08984
- [145] K. Hyatt, J. R. Garrison and B. Bauer, Extracting entanglement geometry from quantum states, *Phys. Rev. Lett.* 119, 140502 (2017).
- [146] R. Nandkishore and D. A. Huse, Many body localization and thermalization in quantum statistical mechanics, *Ann. Review of Cond. Mat. Phys.* 6, 15 (2015).
- [147] E. Altman and R. Vosk, Universal dynamics and renormalization in many body localized systems, *Ann. Review of Cond. Mat. Phys.* 6, 383 (2015).
- [148] S. A. Parameswaran, A. C. Potter and R. Vasseur, Eigenstate phase transitions and the emergence of universal dynamics in highly excited states, *Annalen der Physik* 529, 1600302 (2017).
- [149] F. Alet and N. Laflorencie, Many-body localization: an introduction and selected topics, *Comptes Rendus Physique* (2018).
- [150] D. A. Abanin, E. Altman, I. Bloch and M. Serbyn, Ergodicity, Entanglement and Many-Body Localization, arxiv 1804.11065
- [151] J. Z. Imbrie, V. Ros, A. Scardicchio, Review: Local Integrals of Motion in Many-Body Localized systems, *Annalen der Physik* 529, 1600278 (2017).
- [152] L. Rademaker, M. Ortuno and A.M. Somoza, Many-body localization from the perspective of Integrals of Motion, *Annalen der Physik* 529, 1600322 (2017).
- [153] D. Pekker, G. Refael, E. Altman, E. Demler and V. Oganesyan, Hilbert-Glass Transition: New Universality of Temperature-Tuned Many-Body Dynamical Quantum Criticality, *Phys. Rev. X* 4, 011052 (2014).
- [154] Y.Z. You, X.L. Qi and C. Xu, Entanglement Holographic Mapping of Many-Body Localized System by Spectrum Bifurcation Renormalization Group, *Phys. Rev. B* 93, 104205 (2016).
- [155] C. Monthus, Strong Disorder Real-Space Renormalization for the Many-Body-Localized phase of random Majorana models, *J. Phys. A: Math. Theor.* 51 115304 (2018).
- [156] Y. Huang, Joel E. Moore, Excited-state entanglement and thermal mutual information in random spin chains, *Phys. Rev B*90, 220202 (2014).
- [157] M. Pouranvari and K. Yang, Entanglement spectrum and entangled modes of highly excited states in random XX spin chains, *Phys. Rev. B* 92, 245134 (2015).
- [158] K. Agarwal, E. Demler and I. Martin, $1/f^\alpha$ noise and generalized diffusion in random Heisenberg spin systems, *Phys. Rev. B* 92, 184203 (2015).
- [159] R. Vasseur, A. J. Friedman, S. A. Parameswaran and A. C. Potter, Particle-hole symmetry, many-body localization, and topological edge modes, *Phys. Rev. B* 93, 134207 (2016).
- [160] K. Slagle, Y. Z. You, and C. Xu, Disordered XYZ spin chain simulations using the spectrum bifurcation renormalization group, *Phys. Rev. B* 94, 014205 (2016).
- [161] A. J. Friedman, R. Vasseur, A. C. Potter, S. A. Parameswaran, Localization-protected order in spin chains with non-Abelian discrete symmetries, arxiv 1706.00022
- [162] R. Vasseur, A. C. Potter and S.A. Parameswaran, Quantum Criticality of Hot Random Spin Chains, *Phys. Rev. Lett.* 114, 217201 (2015).
- [163] B. Kang, A. C. Potter and R. Vasseur, Universal crossover from ground state to excited-state quantum criticality, *Phys. Rev. B* 95, 024205 (2017).
- [164] C. Monthus, Many-Body Localization : construction of the emergent local conserved operators via block real-space renormalization, *J. Stat. Mech.* 033101 (2016)
- [165] C. Monthus, Random Transverse Field Spin-Glass Model on the Cayley tree : phase transition between the two Many-Body-Localized Phases, *J. Stat. Mech.* 123304 (2017).
- [166] R. Vosk and E. Altman, Many-body localization in one dimension as a dynamical renormalization group fixed point, *Phys. Rev. Lett.* 110, 067204 (2013).
- [167] R. Vosk and E. Altman, Dynamical Quantum Phase Transitions in Random Spin Chains, *Phys. Rev. Lett.* 112, 217204 (2014).
- [168] M. Bukov, L. D'Alessio and A. Polkovnikov, Universal High-Frequency Behavior of Periodically Driven Systems: from Dynamical Stabilization to Floquet Engi-

- neering, *Advances in Physics*, Vol. 64, No. 2, 139 (2015).
- [169] C. Monthus, Strong Disorder Renormalization for the dynamics of Many-Body-Localized systems : iterative elimination of the fastest degree of freedom via the Floquet expansion, *J. Phys. A: Math. Theor.* 51 275302 (2018).
- [170] Y. Huang, Entanglement dynamics in critical random quantum Ising chain with perturbations, *Annals of Physics* 380, 224 (2017).
- [171] M. Heyl and M. Vojta, Nonequilibrium dynamical renormalization group: Dynamical crossover from weak to infinite randomness in the transverse-field Ising chain, *Phys. Rev. B* 92, 104401 (2015).
- [172] P. Hauke and M. Heyl, Many-body localization and quantum ergodicity in disordered long-range Ising models, *Phys. Rev. B* 92, 134204 (2015).
- [173] G. De Chiara, S. Montangero, P. Calabrese, R. Fazio, Entanglement Entropy dynamics in Heisenberg chains, *J. Stat. Mech.*, L03001 (2006).
- [174] F. Iglói, Z. Szatmári, and Y.-C. Lin, Entanglement entropy dynamics of disordered quantum spin chains *Phys. Rev. B* 85, 094417 (2012).
- [175] G. C. Levine, M. J. Bantegui, J. A. Burg, Full counting statistics in a disordered free fermion system, *Phys. Rev. B* 86, 174202 (2012).
- [176] J. H. Bardarson, F. Pollmann, and J. E. Moore, Unbounded Growth of Entanglement in Models of Many-Body Localization, *Phys. Rev. Lett.* 109, 017202 (2012).
- [177] Y. Zhao, F. Andraschko, J. Sirker, Entanglement entropy of disordered quantum chains following a global quench, *Phys. Rev. B* 93, 205146 (2016).
- [178] G. Roósz, Y.C. Lin and F. Iglói, Critical quench dynamics of random quantum spin chains: Ultra-slow relaxation from initial order and delayed ordering from initial disorder, *New J. Phys.* 19, 023055 (2017).
- [179] J. Herbrych, J. Kokalj, and P. Prelovsek, Local Spin Relaxation within the Random Heisenberg Chain, *Phys. Rev. Lett.* 111 147203 (2013).
- [180] Y.R. Shu, M. Dupont, D.X. Yao, S. Capponi and A. W. Sandvik Dynamical properties of the $S=1/2$ random Heisenberg chain, *Phys. Rev. B* 97, 104424 (2018).
- [181] F. Iglói, G. Roósz and Y.C. Lin, Nonequilibrium quench dynamics in quantum quasicrystals, *New J. Phys.* 15, 023036 (2013).
- [182] G. Roósz, U. Divakaran, H. Rieger and F. Iglói, Nonequilibrium quantum relaxation across a localization-delocalization transition *Phys. Rev. B* 90, 184202 (2014).
- [183] U. Divakaran, Sudden quenches in quasiperiodic Ising model, arXiv:1805.07977
- [184] P. Mason, A. M. Zagoskin and J. J. Betouras, Time-dependent Real-space Renormalization-Group Approach: application to an adiabatic random quantum Ising model, arxiv 1708.05948
- [185] C. Monthus and T. Garel, Many-body localization transition in a lattice model of interacting fermions: statistics of renormalized hoppings in configuration space, *Phys. Rev. B* 81, 134202 (2010).
- [186] R. Vosk, D.A. Huse, and E. Altman, Theory of the many-body localization transition in one dimensional systems, *Phys. Rev. X* 5, 031032 (2015).
- [187] A. C. Potter, R. Vasseur and S.A. Parameswaran, Universal properties of many-body delocalization transitions, *Phys. Rev. X* 5, 031033 (2015).
- [188] P. T. Dumitrescu, R. Vasseur, A. C. Potter, Scaling Theory of Entanglement at the Many-Body Localization Transition, *Phys. Rev. Lett.* 119, 110604 (2017).
- [189] L. Zhang, B. Zhao, T. Devakul and D. A. Huse, Many-body localization phase transition: A simplified strong-randomness approximate renormalization group, *Phys. Rev. B* 93, 224201 (2016).
- [190] A. Goremykina, R. Vasseur and M. Serbyn, Analytically solvable renormalization group for the many-body localization transition, arxiv: 1807.04285
- [191] S. Kehrein, *The flow equation approach to many-particle systems*, Springer-Verlag Berlin (2006).
- [192] L. Rademaker and M. Ortuno, Explicit Local Integrals of Motion for the Many-Body Localized State, *Phys. Rev. Lett.* 116, 010404 (2016).
- [193] C. Monthus, Flow towards diagonalization for Many-Body-Localization models : adaptation of the Toda matrix differential flow to random quantum spin chains, *J. Phys. A: Math. Theor.* 49 305002 (2016).
- [194] D. Pekker, B. K. Clark, V. Oganesyan and G. Refael, Fixed points of Wegner-Wilson flows and many-body localization, *Phys. Rev. Lett.* 119, 075701 (2017).
- [195] S. Savitz and G. Refael, Stable Unitary Integrators for the Numerical Implementation of Continuous Unitary Transformations, *Phys. Rev. B* 96, 115129 (2017).
- [196] S. J. Thomson and M. Schiro, Time Evolution of Many-Body Localized Systems with the Flow Equation Approach, *Phys. Rev. B* 97, 060201 (2018).
- [197] R. Moessner and S. L. Sondhi, Equilibration and Order in Quantum Floquet Matter, *Nature Physics* 13, 424 (2017)
- [198] C. Monthus, Periodically driven random quantum spin chains : Real-Space Renormalization for Floquet localized phases, *J. Stat. Mech.* (2017) 073301
- [199] W. Berdanier, M. Kolodrubetz, S. A. Parameswaran and R. Vasseur, Floquet Quantum Criticality, arXiv:1803.00019
- [200] W. Berdanier, M. Kolodrubetz, S. A. Parameswaran and R. Vasseur, Strong-Disorder Renormalization Group for Periodically Driven Systems, arxiv : 1807.09767
- [201] U. Weiss, *Quantum dissipative systems*, Singapore: World Scientific, 2nd edition (1999).
- [202] H. P. Breuer and F. Petruccione, *The theory of open quantum systems* Oxford (2002).
- [203] Leggett, A., S. Chakravarty, A. Dorsey, M. Fisher, A. Garg, and W. Zwerger, Dynamics of the dissipative two-state system, *Rev. Mod. Phys.* 59, 1 (1987).
- [204] G. Schehr and H. Rieger, Strong Randomness Fixed Point in the Dissipative Random Transverse Field Ising Model, *Phys. Rev. Lett.* 96, 227201 (2006).
- [205] G. Schehr and H. Rieger, Finite temperature behavior of strongly disordered quantum magnets coupled to a dissipative bath *J. Stat. Mech.* P04012 (2008).
- [206] J. A. Hoyos and T. Vojta, Dissipation effects in percolating quantum Ising magnets, *Physica B* 403, 1245 (2008).
- [207] J. A. Hoyos and T. Vojta, Theory of Smeared Quantum Phase Transitions, *Phys. Rev. Lett.* 100, 240601 (2008).
- [208] T. Vojta and J. A. Hoyos, Smeared quantum phase transition in the dissipative random quantum Ising model, *Physica E* 42, 383 (2010).
- [209] J. A. Hoyos and T. Vojta, Dissipation effects in random transverse-field Ising chains, *Phys. Rev. B* 85, 174403

- (2012).
- [210] M. Al-Ali and T. Vojta, Monte-Carlo simulations of the dissipative random transverse-field Ising chain arxiv 1307.7166
- [211] J. A. Hoyos, C. Kotabage and T. Vojta, Effects of dissipation on a quantum critical point with disorder, *Phys. Rev. Lett.* 99, 230601 (2007).
- [212] T. Vojta, C. Kotabage and J. A. Hoyos, Infinite-randomness quantum critical points induced by dissipation, *Phys. Rev. B* 79, 024401 (2009).
- [213] T. Vojta, J. A. Hoyos, P. Mohan and R. Narayanan, Influence of superohmic dissipation on a disordered quantum critical point, *J. Phys.: Condens. Matter* 23, 094206 (2011).
- [214] C. Monthus, Boundary-driven Lindblad dynamics of random quantum spin chains : strong disorder approach for the relaxation, the steady state and the current, *J. Stat. Mech.* 043303 (2017).
- [215] W. De Roeck, A. Dhar, F. Huveneers and M. Schutz, Step Density Profiles in Localized Chains *J. Stat. Phys.* 167 , 1143 (2017).
- [216] C. Monthus, Dissipative random quantum spin chain with boundary-driving and bulk-dephasing: magnetization and current statistics in the Non-Equilibrium-Steady-State, *J. Stat. Mech.* 043302 (2017).
- [217] C. Chatelain, Diverging conductance at the contact between random and pure quantum XX spin chains, *J. Stat. Mech.* 113301 (2017).
- [218] F. Evers and A. D. Mirlin, Anderson transitions *Rev. Mod. Phys.* 80 1355 (2008).
- [219] H. J. Mard, J. A. Hoyos, E. Miranda and V. Dobrosavljevic, Strong-disorder renormalization-group study of the one-dimensional tight-binding model, *Phys. Rev. B* 90,125141 (2014).
- [220] H. J. Mard, J. A. Hoyos, E. Miranda and V. Dobrosavljevic, Strong-disorder approach for the Anderson localization transition, *Phys. Rev. B* 96, 045143 (2017).
- [221] H. Aoki, Real-space renormalisation-group theory for Anderson localisation: decimation method for electron systems, *J. Phys. C: Solid State Phys.* 13 3369 (1980).
- [222] H. Aoki, Decimation method of real-space renormalization for electron systems with application to random systems, *Physica A* 114, 538 (1982).
- [223] C. Monthus and T. Garel, Statistics of renormalized on-site energies and renormalized hoppings for Anderson localization models in dimensions $d=2$ and $d=3$, *Phys. Rev. B* 80, 024203 (2009).
- [224] E. Tarquini, G. Biroli and M. Tarzia, Critical properties of the Anderson localization transition and the high dimensional limit, *Phys. Rev. B* 95, 094204 (2017).
- [225] S. Johri and R. N. Bhatt, Large Disorder Renormalization Group Study of the Anderson Model of Localization *Phys. Rev. B* 90, 060205(R) (2014).
- [226] R.N. Bhatt and S. Johri, Rare Fluctuation Effects in the Anderson Model of Localization, *Int. J. Mod. Phys. Conf. Ser.* 11 79 (2012).
- [227] C. Monthus and T. Garel, Random cascade models of multifractality : real-space renormalization and travelling-waves, *J. Stat. Mech.* P06014 (2010).
- [228] C. Monthus and T. Garel, A critical Dyson hierarchical model for the Anderson localization transition *J. Stat. Mech.* P05005 (2011).
- [229] V. L. Quito, P. Titum, D. Pekker and G. Refael, Localization transition in one dimension using Wegner flow equations *Phys. Rev. B* 94, 104202 (2016).
- [230] T.E. Harris, Contact Interactions on a Lattice, *Ann. Prob.*, 2, 969 (1974).
- [231] T.M. Liggett, Stochastic interacting systems: contact, voter, and exclusion processes, Berlin Springer (2005).
- [232] T. Vojta, A. Farquhar and J. Mast, Infinite-randomness critical point in the two-dimensional disordered contact process, *Phys. Rev. E* 79 011111 (2009)
- [233] T. Vojta, A. Farquhar and J. Mast, Infinite-randomness critical point in the two-dimensional disordered contact process, *Phys. Rev. E* 79, 011111 (2009)
- [234] T. Vojta, Monte-Carlo simulations of the clean and disordered contact process in three dimensions, *Phys. Rev. E* 86, 051137 (2012)
- [235] J. A. Hoyos, Weakly disordered absorbing-state phase transitions, *Phys. Rev. E* 78, 032101 (2008).
- [236] R. Juhász, Distribution of dynamical quantities in the contact process, random walks, and quantum spin chains in random environments, *Phys. Rev. E* 89, 032108 (2014)
- [237] R. Juhász, Disordered contact process with asymmetric spreading, *Phys. Rev. E* 87, 022133 (2013).
- [238] T. Vojta, J. Igo and J. A. Hoyos, Rare regions and Griffiths singularities at a clean critical point: The five-dimensional disordered contact process *Phys. Rev. E* 90, 012139 (2014).
- [239] T. Vojta and J. A. Hoyos, Criticality and quenched disorder: rare regions vs. Harris criterion *Phys. Rev. Lett.* 112, 075702 (2014)
- [240] A. K. Ibrahim, H. Barghathi and T. Vojta, Enhanced rare region effects in the contact process with long-range correlated disorder *Phys. Rev. E* 90, 042132 (2014).
- [241] H. Barghathi, D. Nozadze and T. Vojta, Contact process on generalized Fibonacci chains: infinite-modulation criticality and double-log periodic oscillations, *Phys. Rev. E* 89, 012112 (2014)
- [242] M. A. Munoz, R. Juhász, C. Castellano and G. Ódor, Griffiths Phases on Complex Networks *Phys. Rev. Lett.* 105 128701 (2010).
- [243] H. Barghathi and T. Vojta, Random fields at a nonequilibrium phase transition *Phys. Rev. Lett.* 109, 170603 (2012) ; H. Barghathi and T. Vojta, Random field disorder at an absorbing state transition in one and two dimensions, *Phys. Rev. E* 93, 022120 (2016).
- [244] R. Juhász, I. A. Kovács and F. Iglói, Long-range epidemic spreading in a random environment, *Phys. Rev. E* 91, 032815 (2015).
- [245] R. Juhász, I. A. Kovács, Infinite randomness critical behavior of the contact process on networks with long-range connections, *J. Stat. Mech.* P06003 (2013).
- [246] T. Vojta and J. A. Hoyos, Infinite-noise criticality: Nonequilibrium phase transitions in fluctuating environments, *Europhys. Lett.* 112, 30002 (2015).
- [247] F. Vazquez, J. A. Bonachela, C. Lopez and M. A. Munoz, Temporal Griffiths Phases, *Phys. Rev. Lett.* 106 235702 (2011)
- [248] H. Barghathi, J. A. Hoyos and T. Vojta, Contact process with temporal disorder, *Phys. Rev. E* 94, 022111 (2016).
- [249] C. E. Fiore, M. M. de Oliveira, J. A. Hoyos, Temporal disorder in discontinuous non-equilibrium phase transitions: general results, arXiv:1806.10421
- [250] T. Vojta and R. Dickman Spatio-temporal generalization of the Harris criterion and its application to diffu-

- sive disorder Phys. Rev. E 93, 032143 (2016).
- [251] P. Le Doussal, The Sinai model in the presence of dilute absorbers, J. Stat. Mech. P07032 (2009).
- [252] R. Juhász, Competition between quenched disorder and long-range connections: A numerical study of diffusion, Phys. Rev. E 85, 011118 (2012).
- [253] R. Juhász, Random walks in a random environment on a strip: a renormalization group approach, J. Phys. A: Math. Theor. 41 315001 (2008).
- [254] R. Juhász and F. Iglói, Anomalous diffusion in disordered multi-channel systems, J. Stat. Mech. P03012 (2010).
- [255] R. Juhász, The effect of asymmetric disorder on the diffusion in arbitrary networks, Europhys. Lett. 98 30001 (2012).
- [256] C. Monthus and T. Garel, Random walk in two-dimensional self-affine random potentials: Strong-disorder renormalization approach, Phys. Rev. E 81, 011138 (2010)
- [257] C. Monthus and T. Garel, Non equilibrium dynamics of disordered systems : understanding the broad continuum of relevant time scales via a strong-disorder RG in configuration space, J. Phys. A: Math. Theor. 41 , 255002 (2008).
- [258] C. Monthus and T. Garel, Non-equilibrium dynamics of finite-dimensional disordered systems : RG flow towards an "infinite disorder" fixed point at large times, J. Stat. Mech. P07002 (2008).
- [259] C. Monthus and T. Garel, Equilibrium of disordered systems : constructing the appropriate valleys in each sample via strong disorder renormalization in configuration space, J. Phys. A: Math. Theor. 41, 375005 (2008).
- [260] C. Monthus and T. Garel, Driven interfaces in random media at finite temperature : is there an anomalous zero-velocity phase at small external force ? Phys. Rev. E 78, 041133 (2008).
- [261] S. Bo and A. Celani, Multiple-scale stochastic processes: Decimation, averaging and beyond, Physics Reports 670, 1 (2017).
- [262] C. Monthus and T. Garel, Statistics of first-passage times in disordered systems using backward master equations and their exact renormalization rules, J. Phys. A: Math. Theor. 43 095001 (2010).
- [263] C. Monthus and T. Garel, Dynamics of Ising models near zero temperature : Real Space Renormalization Approach, J. Stat. Mech. P02037 (2013).
- [264] C. Monthus and T. Garel, Dynamical barriers for the random ferromagnetic Ising model on the Cayley tree : traveling-wave solution of the real space renormalization flow, J. Stat. Mech. P05012 (2013).
- [265] C. Monthus and T. Garel, Dynamical Barriers in the Dyson Hierarchical model via Real Space Renormalization, J. Stat. Mech. P02023 (2013).
- [266] C. Monthus, Low-temperature dynamics of Long-Ranged Spin-Glasses : full hierarchy of relaxation times via real-space renormalization, J. Stat. Mech. P08009 (2014).
- [267] C. Monthus, Real-space renormalization for the finite temperature statics and dynamics of the Dyson Long-Ranged Ferromagnetic and Spin-Glass models, J. Stat. Mech. 043302 (2016).
- [268] F. J. Dyson, The Dynamics of a Disordered Linear Chain, Phys. Rev. 92, 1331 (1953).
- [269] C. Monthus and T. Garel, Anderson localization of phonons in dimension $d=1,2,3$: finite-size properties of the Inverse Participation Ratios of eigenstates, Phys. Rev. B 81, 224208 (2010).
- [270] M. B. Hastings, Random Vibrational Networks and the Renormalization Group, Phys. Rev. Lett. 90, 148702 (2003).
- [271] A. Amir, Y. Oreg and Y. Imry, Localization, Anomalous Diffusion, and Slow Relaxations: A Random Distance Matrix Approach, Phys. Rev. Lett. 105, 070601 (2010).
- [272] C. Monthus and T. Garel, Random elastic networks : strong disorder renormalization approach, J. Phys. A: Math. Theor. 44 085001 (2011).
- [273] S. H. Strogatz, From Kuramoto to Crawford: exploring the onset of synchronization in populations of coupled oscillators, Physica D 143, 1 (2000).
- [274] A. Pikovsky, M. Rosenblum, and J. Kurths, Synchronization: A Universal Concept in Nonlinear Science, Cambridge University Press, New York, 2001.
- [275] A. Pikovsky and M. Rosenblum, Dynamics of globally coupled oscillators: progress and perspectives Chaos 25, 097616 (2015).
- [276] O. Kogan, J. L. Rogers, M. C. Cross, and G. Refael, Renormalization group approach to oscillator synchronization, Phys. Rev. E 80, 036206 (2009).
- [277] T. E. Lee, G. Refael, M. C. Cross, O. Kogan and J. L. Rogers, Universality in the one-dimensional chain of phase-coupled oscillators, Phys. Rev. E 80, 046210 (2009).
- [278] P. Mohan, R. Narayanan and T. Vojta, Infinite randomness and quantum Griffiths effects in a classical system: the randomly layered Heisenberg magnet, Phys. Rev. B 81, 144407 (2010)
- [279] C. Monthus and T. Garel, Random wetting transition on the Cayley tree: a disordered first-order transition with two correlation length exponents, J. Phys. A: Math. Theor. 42 165003 (2009).
- [280] C. Monthus, Strong Disorder Renewal Approach to DNA denaturation and wetting : typical and large deviation properties of the free energy, J. Stat. Mech. 013301 (2017).
- [281] C. Monthus, One-dimensional Ising spin-glass with power-law interaction : real-space renormalization at zero temperature, J. Stat. Mech. P06015 (2014).
- [282] C. Monthus, Fractal dimension of spin glasses interfaces in dimensions $d=2$ and $d=3$ via strong disorder renormalization at zero temperature, Fractals, 23, 1550042 (2015).
- [283] W. Wang, M. A. Moore and H. G. Katzgraber, The Fractal Dimension of Interfaces in Edwards-Anderson and Long-range Ising Spin Glasses: Determining the Applicability of Different Theoretical Descriptions, Phys. Rev. Lett. 119, 100602 (2017).
- [284] W. Wang, M. A. Moore and H. G. Katzgraber, Fractal dimension of interfaces in Edwards-Anderson spin glasses for up to six space dimensions, Phys. Rev. E 97, 032104 (2018).
- [285] G. Schehr and P. Le Doussal, Extreme value statistics from the real space renormalization group: Brownian motion, Bessel processes and continuous time random walks, J. Stat. Mech. (2010) P01009 (2010).
- [286] G. Györgyi, N. R. Moloney, K. Ozogány, Z. Rácz, and M. Droz, Renormalization-group theory for finite-size scaling in extreme statistics Phys. Rev. E 81, 041135 (2010); E. Bertin and G. Györgyi, Renormalization

flow in extreme value statistics, *J. Stat. Mech.* P08022 (2010); I. Calvo, J. C. Cuch, J. G. Esteve, and F. Falceto, Extreme-value distributions and renormalization group, *Phys. Rev. E* 86 041109 (2012); F. Angeletti, E.

Bertin, and P. Abry, Renormalization flow for extreme value statistics of random variables raised to a varying power, *J. Phys. A: Math. Theor.* 45 115004 (2012).
[287] R. Juhász, A non-conserving coagulation model with extremal dynamics, *J. Stat. Mech.* P03033 (2009).