



**HAL**  
open science

## Micro(plastics) in the Environment of Some French rivers

Vincent Verney, Florence Jestin, Dominik Zdybal, Camila Mota-Moteiro, Lola Desvignes, Haroutioun Askanian, Alexandre Schaal, Lise Durantou, Max Liboiron

### ► To cite this version:

Vincent Verney, Florence Jestin, Dominik Zdybal, Camila Mota-Moteiro, Lola Desvignes, et al.. Micro(plastics) in the Environment of Some French rivers. Micro 2018. FATE AND IMPACT OF MICROPLASTICS: KNOWLEDGE, ACTIONS AND SOLUTIONS, Nov 2018, Lanzarote, Spain. hal-01963924

**HAL Id: hal-01963924**

**<https://hal.science/hal-01963924>**

Submitted on 21 Dec 2018

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# MICRO(PLASTICS) IN THE ENVIRONMENT OF SOME FRENCH RIVERS

Vincent Verney<sup>1</sup>, Florence Delor Jestin<sup>1</sup>, Dominik Zdybal<sup>1</sup>, Camila Mota-Moteiro<sup>1</sup>, Lola Desvignes<sup>1</sup>, Haroutioun Askanian<sup>1</sup>, **Alexandre Schaal**<sup>2</sup>, Lise Durantou<sup>2</sup>, Max Liboiron<sup>3</sup>

(1) CNRS, Université Clermont Auvergne, UMR 6296, Institut de Chimie de Clermont-Ferrand (ICCF), Clermont Ferrand, FRANCE


(2) Microplastics Observatory, La Pagaie Sauvage, Anglet, FRANCE

(3) Civic Laboratory for Environmental Action Research (CLEAR), Memorial University of Newfoundland, St John's, CANADA

It is now well known that the vast majority of microplastics (MPs) found in the seas and oceans come from lands and as such freshwater environments (rivers and riverbanks) play a major role in this process. We must therefore imagine the scenario that a used plastic, become a waste after use, is found in the environment if it left, for one reason or another from a collection and recycling stream. Its stay in the environment can last a wild and this waste will then be exposed to a set of environmental constraints (UV, rain, wind, mechanical erosion, ..) which will continue and amplify its degradation, leading to its fragmentation since the macroplastics into MPs, even into nanoplastics (NPs). Riverpast Project consists in collecting and analyzing the composition of MPs at the surface waters of different French rivers. To this, we rely on citizen science operations, in particular thanks to the contribution of the BabyLegs sampling net [1], which makes it possible to multiply samples and analyzes [2]. A specific focus has been done on the Garonne River during spring 2018 with a sampling campaign of more than 120 samples analyzed. Collected data and results are reported on open web access map [3].

## EXPERIMENTAL

Once received, the contents of the babyleg sampling are sieved in different steps using decreasing mesh sizes. First, it is sieved at 4.57 mm to separate the bigger particles (not considered as MPs). The content is rinsed with distilled water and used water is kept apart. A first manual separation is done to eliminate the vegetal big particles (pieces of branches, leaves, lichens,...). After what, the rest is sieved again at 1mm and 250 µm mesh sizes. The filtered content of 1 mm and 250 µm is dispersed in water. An optical observation is done under a binocular light device. Suspected plastic particles are taken away from the liquid medium using tweezers and disposed in a Petri dish. After half day in an oven regulated at 60°C, the samples are observed and captured with a photo using a 5 Megapixel resolution PCE-MM200 digital microscope along with a measuring tape. Finally, the particles are observed with a FTIR spectrometer (Thermo-Nicolet 380 FTIR) in ATR (Attenuated Total Reflection, 4000 - 643 cm<sup>-1</sup>, 4 cm<sup>-1</sup> resolution, 32 scans accumulated) mode using a diamond crystal. Spectra are recorded and analyzed using OMNIC software. The chemical identification is ensured by using a polymer database spectra.



## RESULTS

Since the beginning of our analysis (September 2017), a total number of **196 samplings** has been carried out. Over these samplings, **123** have been analyzed giving rise to the identification of **246 MP** particles (5 mm < size < 250 µm) c.a. an average of **2 particles/sampling**.




Results are saved on a datadrive and an updated map is accessible online [5]

- ③ 120 particles have been identified as Polyethylene c.a. 49%
- ③ 45 particles have been identified as Polystyrene c.a. 18%
- ③ 30 particles have been identified as Polypropylene c.a. 12%
- ③ 13 as EPDM (Etylene-Propylene Diene monomer), 9 as cellophane (4%), 4 as PET (Polyethylene Terephthalate), 2 as PVC (Polyvinyl Chloride) and 23 as Others (Polyamide, Ethylene Vinyl acetate, Polymethyle methacrylate, Polydimethyl Siloxane)

## FOCUS ON GARONNE RIVER

A specific focus has been done onto the Garonne river during **spring 2018** (3 weeks in may 2018) in the field of the **NERRI** campaign [5]. During this sampling period **116 samplings** have been carried out. Over these 116 samplings **46** have been analyzed and **137 MPs** have been identified representing thus an average of **3 MPs/sampling** (about 50% more than the overall ratio see upper)

For a better reliability, some samplings have been performed in duplicate or triplicate



137 MPs have been identified and the average size of most of them is comprised between 1 and 5 mm.

## CONCLUSIONS AND PERSPECTIVES

- ③ During the first year of analysis (2017-2018) we built up:
  - a methodology based upon citizen scientists participation
  - a physico chemical methodology called the **WYSIWYG methodology** (What You See Is What You Get) allowing us to separate, to count and to identify MP with sizes greater than 250 µm (both mesh size of Manta nets and babylegs is around 300 µm + the minimum size for ATR-FTIR is around 100 µm)
- ③ The future development of this research is in the improvements of the WYSIWYG protocol sensitivity. As MP characterization implies three inseparable analytical steps: Collecting/Separating/Analysis, efforts must be done one each of them to increase the WYSIWYG resolution. we shall develop new:
  - methodology for collection (hydrocyclonic)
  - method of density separation by using specific chemical liquids such as Ionic liquids
  - FTIR analysis by using a Focus Plane Array detector (FPA-FTIR) allowing us to go down to 10 µm resolution.
  - microscopic-thermal coupled methods for the quantification of the MPs fraction

## REFERENCES

- 1- Schmidt, C.; Krauth, T.; Wagner, S., Export of Plastic Debris by Rivers into the Sea. Environ Sci Technol 2017, 51 (2), 12246-12253.
- 2- Siegfried, M.; Koelmans, A. A.; Besseling, E.; Kroeze, C., Export of microplastics from land to sea. A modelling approach. Water Res 2017, 127, 249-257
- 3- Rech, S.; Macaya-Caquigan, V.; Pantoja, J. F.; Rivadeneira, M. M.; Campodonico, C. K.; Thiel, M., Sampling of riverine litter with citizen scientists--findings and recommendations. Environ Monit Assess 2015, 187 (6), 335
- 4- Max Liboiron, Compromise agency, the case of babylegs. Engaging Science, Technology and Society 3(2017), 499-527
- 5- <http://lapagaiesauvage.org/laboratoirecitoyen>
- 6- <https://www.researchgate.net/project/RiverPlast>

