



Corrigendum to "Improvement of the PR-CPA equation of state for modelling of acid gases solubilities in aqueous alkanolamine solutions" [Fluid Phase Equilibria, Vol. 471, 2018, 74-87]

Tianyuan Wang, Philippe Guittard, Christophe Coquelet, Elise El Ahmar, Olivier O. Baudouin, Georgios Kontogeorgis

► **To cite this version:**

Tianyuan Wang, Philippe Guittard, Christophe Coquelet, Elise El Ahmar, Olivier O. Baudouin, et al.. Corrigendum to "Improvement of the PR-CPA equation of state for modelling of acid gases solubilities in aqueous alkanolamine solutions" [Fluid Phase Equilibria, Vol. 471, 2018, 74-87]. Fluid Phase Equilibria, 2019, 485, pp.126-127. hal-01968780

HAL Id: hal-01968780

<https://hal.science/hal-01968780>

Submitted on 3 Jan 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Corrigendum to “Improvement of the PR-CPA equation of state for modelling of acid gases solubilities in aqueous alkanolamine solutions” [Fluid Phase Equilibria, Vol. 471, 2018, 74-87]

Tianyuan Wang^{1,2}, , Philippe Guittard³, Christophe Coquelet¹ Elise El Ahmar¹, Olivier Baudouin³ and Georgios M. Kontogeorgis²

1: Mines ParisTech-PSL University Centre of Thermodynamics of Processes CTP, 35 rue Saint-Honore, 77305, Fontainebleau, France

2: Center for Energy Resources Engineering (CERE), Department of Chemical and Biochemical Engineering, Technical University of Denmark, DK-2800 Kgs. Lyngby, Denmark

3: ProSim SA Immeuble Stratège A, 51 rue Ampère 31670, Labège, France

After careful comparison with the results of implanted PR-CPA EoS in Prosim software (Simulis™ Thermodynamics) with the presented data in the paper, the authors found that some parameters in the table “PR-CPA EoS parameters for compounds for association compounds considered in this work “and the table “BIP values and ARD of total pressure between PR-CPA EoS results and experimental data for acid gas -alkanolamine binary system” were misprinted. The average absolute deviations and results presented in the paper are not influenced by this misprint.

Moreover, these parameters were also adjusted by Prosim using their algorithms. The correct and new parameter values are presented in Tables 1 and 2 respectively. Figure 1 shows a comparison between the calculations using the two set of parameters. We can easily observe that with new parameters, the prediction is improved.

Table 1. PR-CPA EoS parameters for compounds for association compounds considered in this work.

Compound	scheme	a_0	b	c_1	ϵ^{AIBj}	β^{AIBj}	T_c	Range of Tr		ARD% ^a	
		/bar L ² mol ⁻² ($\times 10^{-1}$)	/L mol ⁻¹ ($\times 10^2$)		/bar.L.mol ⁻¹		/K				
								For P_{sat}	For ρ_L	P_{sat}	ρ_L
MEA	4C	1.333	5.467	0.763	168.23	0.0142	671	0.42-0.92	0.43-0.61	1.8	0.6
MDEA	4C	3.339	11.346	0.695	201.76	0.0083	742	0.39-0.9	0.38-0.63	0.9	2
water	4C	0.123	1.445	0.674	170.48	0.0698	647	0.43-0.95	0.43-0.95	1	1.6
CO ₂		0.397	2.661	0.701			304				
H ₂ S		0.492	2.696	0.517			373				

^a ARD% = $1/n \times \sum |1 - \chi_i^{calc}/\chi_i^{exp}| \times 100\%$, χ : P_{sat} or ρ_L

Table 2 BIP values and ARD of total pressure between PR-CPA EoS results and experimental data for acid gas -alkanolamine binary system.

Binary	T/K	$ARD_{Pt}\%$	k_{ij}^a			β^{AiBj}	ε^{AiBj} /bar.L.mol ⁻¹	β^{AiBj^*}	ε^{AiBj^*} /bar.L.mol ⁻¹	Source of BIP
			a	b×10 ³ /K ⁻¹	c×10 ⁶ /K ⁻²					
CO ₂ -MEA	298-393	12	-6.99	29.9	-23.73	0.00411	408.62	0.00102	361.17	[1]
CO ₂ -MEA	298-393	12	-6.68816	30.7772	-23.2	0.00353698	441.451	0.00114328	351.619	PROSIM
CO ₂ -MDEA	313-413	11	6.53	-40.0	69.3	0.00487	345.69	NA	NA	[1]
H ₂ S-MEA	298-413	30 [*]	-6.426	38.15	-49.94	0.00380	411.51	0.0170	178.26	[1]
H ₂ S-MDEA	311-388	13	0.8489	-2.834	8.461	0.01846	316.42	NA	NA	[1]

a: $k_{ij} = a + b \cdot T + c \cdot T^2$, ARD_{Pt} ARD on total pressure of acid gas-water-alkanolamine ternary systems, BIP: Binary Interaction Parameter

*: ARD% on partial pressure of H




Figure 1: CO₂ solubility into H₂O-MEA mixture (30wt %): comparison of results using the two sets of parameters. Symbols: experimental data from Jou et al. [2]. (▲)=298 K, (●)=333 K, (×)=353 K, (+)=393 K

References

- [1] T. Wang, E. El Ahmar, C. Coquelet, G. M. Kontogeorgis, Improvement of the PR-CPA equation of state for modelling of acid gases solubilities in aqueous alkanolamine solutions, 2018, *Fluid Phase Equilibria*, 471, 74-87.
- [2] F.-Y. Jou, A.E. Mather, F.D. Otto, The solubility of CO₂ in a 30 mass percent monoethanolamine solution, *Can. J. Chem. Eng.* 73 (1995) 140–147.