



**HAL**  
open science

## What is the effect of cloud inhomogeneities on actinic fluxes and chemical species concentrations?

Christel Bouet, Frédéric Szczap, Maud Leriche, Albert Benassi

### ► To cite this version:

Christel Bouet, Frédéric Szczap, Maud Leriche, Albert Benassi. What is the effect of cloud inhomogeneities on actinic fluxes and chemical species concentrations?. *Geophysical Research Letters*, 2006, 33 (1), pp.n/a-n/a. hal-01971911

**HAL Id: hal-01971911**

**<https://hal.science/hal-01971911v1>**

Submitted on 15 Aug 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Copyright

# What is the effect of cloud inhomogeneities on actinic fluxes and chemical species concentrations?

Christel Bouet, Frédéric Szczap, Maud Leriche, and Albert Benassi

Laboratoire de Météorologie Physique, Centre National de la Recherche Scientifique, Université Blaise Pascal, Aubière, France

Received 22 September 2005; revised 25 November 2005; accepted 1 December 2005; published 14 January 2006.

[1] The purpose of the present paper is to evaluate the bias introduced by the commonly used homogeneous plane-parallel cloud hypothesis in the computation of actinic fluxes, photolysis coefficients, and main tropospheric species concentrations. Accordingly, these quantities obtained for an inhomogeneous cloud field generated with a stochastic model are compared to their homogeneous plane-parallel equivalent. Results show that neglecting cloud inhomogeneities has a significant impact on actinic flux. For instance, the bias introduced by the homogeneous plane-parallel cloud hypothesis can reach more than 100% below the cloud leading to a comparable bias in photolysis coefficients and, in turn, creating an enhancement of the oxidizing capacity of the system. **Citation:** Bouet, C., F. Szczap, M. Leriche, and A. Benassi (2006), What is the effect of cloud inhomogeneities on actinic fluxes and chemical species concentrations?, *Geophys. Res. Lett.*, 33, L01818, doi:10.1029/2005GL024727.

## 1. Introduction

[2] Small changes in the cloud-radiative forcing (which is the effects of clouds on the radiation balance of the Earth fields) can play a significant role as a climate feedback mechanism [Ramanathan *et al.*, 1989]. Observations show that clouds are far from being homogeneous revealing autosimilar and fractal statistical properties [Lovejoy, 1982; Cahalan and Snider, 1989; Davis *et al.*, 1999]. They also exhibit numerous structures in their shapes, aspect ratio and cloud coverage as well as a high horizontal and vertical variability in their microphysical and optical properties [Stephens and Platt, 1987; Korolev *et al.*, 2001]. Numerous studies were conducted to estimate the radiative impact of cloud inhomogeneities in order to improve cloud properties retrieval from various radiometers and to better estimate the role of clouds on Earth's climate [Harshvardhan, 1982; Welch and Wielicki, 1984, 1985, 1989; Barker *et al.*, 1998, 1999]. Barker *et al.* [1999] showed that it was crucial to consider horizontal fluctuations of cloud extinction in order not to underestimate surface absorption and not to overestimate reflected fluxes.


[3] Daytime atmospheric chemistry is driven by photolysis processes. Due to their impact on radiation, clouds can disrupt these processes. Photolysis rate coefficients are derived from actinic flux [Madronich, 1987] and numerous studies have already demonstrated the impact of clouds on actinic flux [Madronich, 1987; Junkermann *et al.*, 2002; Kanaya *et al.*, 2003; Monks *et al.*, 2004]. For example,

Madronich [1987] showed that actinic fluxes inside the clouds can frequently exceed the clear-sky values (up to 5 times greater). However, all of these studies as well as those concerning the impact of clouds on photolysis coefficients [Van Weele and Duynkerke, 1993; Crawford *et al.*, 1999, 2003; Fröh *et al.*, 2000; Tie *et al.*, 2003], considered clouds as a single homogeneous, plane-parallel layer. A recent study of Stockwell and Goliff [2004] showed that NO<sub>2</sub> photolysis coefficients derived from actinic fluxes simulated with the TUV model (Tropospheric Ultraviolet Visible model [Madronich and Flocke, 1998]) using the delta-Eddington method [Joseph *et al.*, 1976] could be up to 56% greater than photolysis coefficients derived from measured actinic fluxes. However, a few studies have already investigated the three-dimensional (3D) effects of inhomogeneous clouds on actinic fluxes and photolysis coefficients [Los *et al.*, 1997; Trautmann *et al.*, 1999; Várnai and Davis, 1999; Brasseur *et al.*, 2002; Trentmann *et al.*, 2003]. Brasseur *et al.* [2002] demonstrated that, in presence of a deep convective cloud, actinic fluxes can be increased by a factor of 2 to 5 compared to clear-sky values, leading to changes in ozone production rates (+15%) and enhancement in OH concentrations (120–200%) in the upper troposphere. They also showed that photolysis coefficients are inhomogeneously distributed throughout the cloud.

[4] Since numerical chemistry transport models usually use the homogeneous plane-parallel cloud assumption, it is critical to investigate the bias introduced in tropospheric chemistry compared to inhomogeneous clouds, which are closer to real clouds. The aims of this numerical modeling study are firstly to quantify the effects of cloud optical and geometrical inhomogeneities on actinic flux in order to estimate, in a second step, the impact of these cloud inhomogeneities on tropospheric photochemistry.

## 2. Methodology

[5] A two-dimensional (2D) inhomogeneous cumulus cloud field is generated with the stochastic tdMAP model (tree driven Mass Accumulation Process [Benassi *et al.*, 2004]), which is able to provide stratocumulus and cumulus cloud fields with properties close to those observed in real clouds. In our case, the simulated 2D cloud field represents a horizontal extent of  $12.8 \times 12.8 \text{ km}^2$  ( $256 \times 256$  pixels) with a 58 % cloud cover. Each  $50 \times 50 \text{ m}^2$  cloud pixel has the same vertical extent and is assumed to be vertically homogeneous leading to a 3D cloud field. The cloud layer is located between 1 and 1.5 km high. The SHDOM model (Spherical Harmonics Discrete Ordinate Method [Evans,



**Figure 1.**  $\Delta F$  as a function of solar angle and mean optical depth below the cloud at  $z = 200$  m (a), in the cloud at  $z = 1050$  m (b) and  $1450$  m (c), and above the cloud at  $z = 2$  km (d).

1998]) is used to solve the radiative transfer equation (RTE) in the inhomogeneous 3D cloud field. Thereby, actinic fluxes at a wavelength of 400 nm corresponding to the wavelength of maximum dissociation of  $\text{NO}_2$  are computed for several cloud mean optical depths (1, 2, 5, 10, and 30) and for different solar zenith angles (0, 20, 40, and  $60^\circ$ ). The RTE is also solved using SHDOM for the homogeneous plane-parallel equivalent clouds in order to estimate the bias introduced by the plane-parallel homogeneous cloud approximation. For all cases, droplets are assumed to follow a lognormal distribution with an effective radius of  $10 \mu\text{m}$  and a standard deviation of  $0.35 \mu\text{m}$ . Optical parameters are calculated using Mie theory. Then, the actinic fluxes computed by SHDOM for the inhomogeneous case are averaged over the whole cloud field. These averaged fluxes and the ones computed for the plane-parallel case are parameterized as a function of the solar zenith angle and the wavelength in order to estimate photolysis coefficients with the TUV model [Madronich and Flocke, 1998] for the main tropospheric chemical species. Photolysis coefficients are computed only for the cloud of optical depth 10 at location ( $51^\circ\text{N}$ ;  $0^\circ\text{E}$ ). Finally, the M2C2 box model (Model of Multiphase Cloud Chemistry [Leriche et al., 2003]) is used to simulate the multiphase cloud chemistry for two different scenarios (remote and urban) from Ervens et al. [2003]. These scenarios include emissions and dry deposition with variable photolysis for three days in order to assess the bias introduced in tropospheric photochemistry by the homogeneous plane-parallel assumption.

### 3. Results and Discussion

#### 3.1. Effect of 2D Inhomogeneities on Actinic Fluxes

[6] The bias between actinic fluxes calculated with SHDOM in the inhomogeneous ( $F_{\text{SHDOM}3\text{D}}$ ) and homogeneous plane-parallel ( $F_{\text{SHDOM}PP}$ ) cases is computed as:

$$\Delta F = \frac{F_{\text{SHDOM}3\text{D}} - F_{\text{SHDOM}PP}}{F_{\text{SHDOM}PP}} \times 100 \quad (1)$$

This bias is shown on Figure 1 as a function of mean cloud optical depth and solar zenith angle below the cloud at  $z =$

200 m, in the cloud at  $z = 1050$  m and  $1450$  m, and above the cloud at  $z = 2$  km. Cloud inhomogeneities significantly enhance the actinic flux below the cloud and in its lower part whereas they slightly decrease it above. For instance, for an optical depth of 10, which is a typical value for stratocumulus clouds, and a solar zenith angle of  $60^\circ$ , below the cloud,  $\Delta F$  is about 70% whereas above, it is around  $-10\%$ . Figure 1 also points out that  $|\Delta F|$  increases with both solar zenith angle and cloud optical depth. Indeed, the greater cloud optical depth or solar zenith angle is, the more important multiple scattering from the cloud sides is, which leads to an enhancement in actinic fluxes. Finally, this means that the closer the sun is to the zenith, the more the inhomogeneous cloud becomes closer to the homogeneous one.

[7] Considering the cloud effect toward clear sky onto actinic fluxes as shown for instance by Madronich [1987], our results imply that the assumption of the homogeneous plane-parallel cloud overestimates this negative effect.

#### 3.2. Effect of 2D Inhomogeneities on Photolysis Coefficients

[8] Prior to the computing of photolysis coefficients with the TUV model, actinic fluxes at 400 nm need to be extended to the visible spectrum. The parameterization used is based upon the statement that, as the mean free path, defined by the mean distance between two collisions undergone by the photon, is constant in our simulations, the effects of cloud inhomogeneities are assumed to be constant with wavelength.

[9] The bias between photolysis coefficients for inhomogeneous ( $J_{3\text{D}}$ ) and homogeneous ( $J_{PP}$ ) cases is defined in the same way as for the actinic fluxes:

$$\Delta J = \frac{J_{3\text{D}} - J_{PP}}{J_{PP}} \times 100 \quad (2)$$

$\Delta J$  is the same for a given hour for all the photolysis reactions because the bias in photolysis coefficients is directly related to the ones on actinic fluxes by definition. Figure 2 presents this bias below the cloud, at  $z = 200$  m, where the bias computed on actinic fluxes is the highest, as a function of time and solar zenith angle. As  $\Delta F$ ,  $\Delta J$  is minimum at noon for a solar zenith angle of  $27^\circ$  with a value around 5%. Thus, the impact of the bias introduced by the homogeneous plane-parallel assumption on the photochemistry varies over time. At noon, when photochemistry is the most active, the impact will be the lowest. However, the most important emissions of pollutants occur in the morning around 8 a.m. where the bias is around 50% and where some important photolysis reactions take place, such as the photolysis of nitrous acid which is the largest source of OH radicals in the morning [Lammel and Cape, 1996].

#### 3.3. Impact on Tropospheric Trace Gases Concentrations

[10] As the first day of simulation is the time for the chemical system to reach the equilibrium, results are presented averaged over days 2 and 3.

[11] In the same way as for the actinic flux, the bias between concentrations for inhomogeneous ( $C_{3\text{D}}$ ) and ho-


homogeneous ( $C_{PP}$ ) cases below the cloud, at  $z = 200$  m, is defined as:

$$\Delta C = \frac{C_{3D} - C_{PP}}{C_{PP}} \times 100 \quad (3)$$

Table 1 shows this bias averaged over days 2 and 3 for main gas phase tropospheric species in the morning, at noon and in the afternoon for the two chemical scenarios, remote and urban. The bias introduced by the use of the homogeneous plane-parallel approximation is different from one species to another: either positive or negative, it is high for radicals (between 5.4 and 54%) whereas it is negligible for methane (less than 0.5%). Neglecting cloud inhomogeneities will have a significant impact on tropospheric chemistry because radicals drive tropospheric chemistry; for instance, ozone concentrations are underestimated using the homogeneous plane-parallel assumption. The stronger photolysis for the 3D case (cf. Figure 2) and the positive bias on OH radicals lead to a larger oxidizing capacity of the atmosphere. Moreover, the minimum values of the positive bias on radicals concentrations is at noon when the bias on actinic fluxes is the lowest. For methane, the bias is close to zero because of its low reactivity driven by its slow oxidation by OH radicals. For  $\text{NO}_2$ , the bias is negative due to a more efficient photolysis in the 3D case while for NO, the bias is negative in the morning and at noon and positive in the afternoon due to the combined effects of the continuous NO emission, the  $\text{NO}_2$  photolysis and the NO to  $\text{NO}_2$  conversion by  $\text{RO}_2$  radicals. For organic compounds, the behavior is more complicated with either positive or negative impact depending on the time of the day and on chemical species considered underlying the non-linearity of the tropospheric chemistry system.

#### 4. Conclusions

[12] The aim of this study was to quantify the bias introduced by the commonly used homogeneous plane-parallel cloud hypothesis on actinic flux, photolysis coef-



**Figure 2.**  $\Delta J$  as a function of solar zenith angle and time below the cloud, at  $z = 200$  m.

**Table 1.**  $\Delta C$  (in %), Averaged Over Days 2 and 3 of the Simulation, for Main Tropospheric Species in Gaseous Phase at 8 a.m., Noon and 4 p.m. for the Two Cases, Remote and Urban

Species	Remote			Urban		
	8 a.m.	Noon	4 p.m.	8 a.m.	Noon	4 p.m.
$\text{O}_3$	11	7.5	5.8	13	9.2	7.6
OH	42	8.3	50	46	9.7	54
$\text{HO}_2$	28	5.4	23	32	7.5	26
$\text{H}_2\text{O}_2$	13	11	7.3	19	17	13
NO	-13	-8.2	12	-7.4	-9.3	14
$\text{NO}_2$	-31	-6.0	-15	-27	-6.1	-15
$\text{HNO}_2$	-16	-5.7	9.8	-6.1	-5.8	14
$\text{HNO}_3$	6.4	-3.3	-1.4	6.9	-2.2	1.1
$\text{CH}_4$	-0.2	-0.3	-0.3	-0.2	-0.2	-0.3
HCHO	0.4	1.3	0.3	-10	-1.4	-2.2
$\text{CH}_3\text{OOH}$	16	16	8.9	8.8	21	13
$\text{CH}_3\text{O}_2$	51	16	32	48	19	35

ficients, and tropospheric chemistry. It was shown that cloud inhomogeneities have a significant impact on actinic flux: for example, for a solar zenith angle of  $60^\circ$  and an optical cloud depth of 10, differences greater than 100% are found. It was also shown that the bias introduced neglecting cloud inhomogeneities depends on both solar zenith angle and cloud optical depth. The same behavior was found for photolysis coefficients. However, due to the non-linearity of tropospheric chemistry system, the bias on chemical species concentrations showed a more complex behavior with either positive or negative bias depending on the chemical species considered and on the time of the day. Finally, for the remote and urban chemical scenarios used, the effect of the cloud inhomogeneities on tropospheric chemistry in comparison with homogeneous clouds is to enhance the oxidizing capacity of the system by higher concentrations of radicals and stronger photolysis.

[13] The next step of this study will be to parameterize the effects due to cloud inhomogeneities in radiative transfer models in order to compare the photolysis coefficients obtained with measurements.

[14] **Acknowledgments.** This work was supported by the "Programme Atmosphère et Océans Multi-échelles" (PATOM) of the INSU (Institut des Sciences de l'Univers). Computer resources were provided by I.D.R.I.S. (Institut du Développement et des Ressources en Informatique Scientifique, project 000187). We gratefully acknowledge Olivier Jourdan for his patience in revising the paper.

#### References

- Barker, H. W., J.-J. Morcrette, and G. D. Alexander (1998), Broadband solar fluxes and heating rates for atmospheres with 3D broken clouds, *Q. J. R. Meteorol. Soc.*, *124*, 1245–1271.
- Barker, H. W., G. L. Stephens, and Q. Fu (1999), The sensitivity of domain-averaged solar fluxes to assumptions about cloud geometry, *Q. J. R. Meteorol. Soc.*, *125*, 2127–2152.
- Benassi, A., F. Szczap, A. Davis, M. Masbou, C. Cornet, and P. Bleuyard (2004), Thermal radiative fluxes through inhomogeneous cloud fields: A sensitivity study using a new stochastic cloud generator, *Atmos. Res.*, *72*, 291–315.
- Brasseur, A.-L., R. Ramarason, A. Delanoy, W. Skamarock, and M. Barth (2002), Three-dimensional calculation of photolysis frequencies in the presence of clouds and impact on photochemistry, *J. Atmos. Chem.*, *41*, 211–237.
- Cahalan, R. F., and J. B. Snider (1989), Marine stratocumulus structure, *Remote Sens. Environ.*, *28*, 95–107.
- Crawford, J., D. Davis, G. Chen, R. Shetter, M. Müller, J. Barrick, and J. Olson (1999), An assessment of cloud effects on photolysis rate coefficients: Comparison of experimental and theoretical values, *J. Geophys. Res.*, *104*(D5), 5725–5734.

- Crawford, J., R. E. Shetter, B. Lefer, C. Cantrell, W. Junkermann, S. Madronich, and J. Calvert (2003), Cloud impacts on UV spectral actinic flux observed during the International Photolysis Frequency Measurement and Model Intercomparison (IPMMI), *J. Geophys. Res.*, *108*(D16), 8545, doi:10.1029/2002JD002731.
- Davis, A., A. Marshak, H. Gerber, and W. Wiscombe (1999), Horizontal structure of marine boundary layer clouds from centimeter to kilometer scales, *J. Geophys. Res.*, *104*(D6), 6123–6144.
- Ervens, B., C. George, J. E. Williams, G. V. Buxton, G. A. Salomon, M. Bydder, F. Wilkinson, F. Dentener, P. Mirabel, R. Wolke, and H. Herrmann (2003), CAPRAM2.4 (MODAC mechanism): An extended and condensed tropospheric aqueous phase mechanism and its application, *J. Geophys. Res.*, *108*(D14), 4426, doi:10.1029/2002JD002202.
- Evans, K. F. (1998), The spherical harmonics discrete ordinate method for three-dimensional atmospheric radiative transfer, *J. Atmos. Sci.*, *55*, 429–446.
- Früh, B., T. Trautmann, M. Wendisch, and A. Keil (2000), Comparison of observed and simulated NO<sub>2</sub> photodissociation frequencies in a cloudless atmosphere and in continental boundary layer clouds, *J. Geophys. Res.*, *105*(D8), 9843–9857.
- Harshvardhan (1982), The effect of brokenness on cloud-climate sensitivity, *J. Atmos. Sci.*, *39*, 1853–1861.
- Joseph, J. H., W. J. Wiscombe, and J. A. Weinman (1976), The delta-Eddington approximation for radiative flux transfer, *J. Atmos. Sci.*, *33*, 2452–2459.
- Junkermann, W., et al. (2002), Actinic radiation and photolysis in the lower troposphere: Effect of clouds and aerosols, *J. Atmos. Chem.*, *42*, 413–441.
- Kanaya, Y., Y. Kajii, and H. Akimoto (2003), Solar actinic flux and photolysis frequency determinations by radiometers and a radiative transfer model at Rishiri Island: Comparisons, cloud effects, and detection of an aerosol plume from Russian forest fires, *Atmos. Environ.*, *37*, 2463–2475.
- Korolev, A. V., G. A. Isaac, I. P. Mazin, and H. W. Barker (2001), Microphysical properties of continental clouds from in-situ measurements, *Q. J. R. Meteorol. Soc.*, *127*, 2117–2151.
- Lammel, G., and J. N. Cape (1996), Nitrous acid and nitrite in the atmosphere, *Chem. Soc. Rev.*, *25*, 361–369.
- Leriche, M., L. Deguillaume, and N. Chaumerliac (2003), Modelling study of strong acids formation and partitioning in a polluted cloud during wintertime, *J. Geophys. Res.*, *108*(D14), 4433, doi:10.1029/2002JD002950.
- Los, A., M. van Weele, and P. G. Duynkerke (1997), Actinic fluxes in broken cloud fields, *J. Geophys. Res.*, *102*(D4), 4257–4266.
- Lovejoy, S. (1982), Area-perimeter relation for rain and cloud areas, *Science*, *216*, 185–187.
- Madronich, S. (1987), Photodissociation in the atmosphere I. Actinic flux and the effects of ground reflections and clouds, *J. Geophys. Res.*, *92*(D8), 9740–9752.
- Madronich, S., and S. Flocke (1998), The role of solar radiation in atmospheric chemistry, in *Handbook of Environmental Chemistry*, edited by P. Boule, pp. 1–26, Springer, New York.
- Monks, P. S., A. R. Rickard, S. L. Hall, and N. A. D. Richards (2004), Attenuation of spectral actinic flux and photolysis frequencies at the surface through homogeneous cloud fields, *J. Geophys. Res.*, *109*, D17206, doi:10.1029/2003JD004076.
- Ramanathan, V., R. D. Cess, E. F. Harrison, P. Minnis, B. R. Barkstrom, E. Ahmad, and D. Hartmann (1989), Cloud-radiative forcing and climate: Results from the Earth radiation budget experiment, *Science*, *243*, 57–63.
- Stephens, G. L., and C. M. R. Platt (1987), Aircraft observations of the radiative and microphysical properties of stratocumulus and cumulus cloud fields, *J. Climatol. Appl. Meteorol.*, *26*, 1243–1269.
- Stockwell, W. R., and W. S. Goliff (2004), Measurement of actinic flux and calculation of photolysis rate parameters for the Central California Ozone Study, *Atmos. Environ.*, *38*, 5169–5177.
- Tie, X., S. Madronich, S. Walters, R. Zhang, P. Raes, and W. Collins (2003), Effect of clouds on photolysis and oxidants in the troposphere, *J. Geophys. Res.*, *108*(D20), 4642, doi:10.1029/2003JD003659.
- Trautmann, T., I. Podgorny, J. Landgraf, and P. J. Crutzen (1999), Actinic fluxes and photodissociation coefficients in cloud fields embedded in realistic atmospheres, *J. Geophys. Res.*, *104*(D23), 30,173–30,192.
- Trentmann, J., B. Früh, O. Boucher, T. Trautmann, and M. O. Andreae (2003), Three-dimensional solar radiation effects on the actinic flux field in a biomass-burning plume, *J. Geophys. Res.*, *108*(D17), 4558, doi:10.1029/2003JD003422.
- Van Weele, M., and P. G. Duynkerke (1993), Effect of clouds on the photodissociation of NO<sub>2</sub>: Observations and modeling, *J. Atmos. Chem.*, *16*, 231–255.
- Várnai, T., and R. Davis (1999), Effects of cloud heterogeneities on short-wave radiation: Comparison of cloud-top variability and internal heterogeneity, *J. Atmos. Sci.*, *56*, 4206–4224.
- Welch, R. M., and B. A. Wielicki (1984), Stratocumulus cloud field reflected fluxes: The effect of cloud shape, *J. Atmos. Sci.*, *41*, 3085–3103.
- Welch, R. M., and B. A. Wielicki (1985), A radiative parameterization of stratocumulus cloud fields, *J. Atmos. Sci.*, *42*, 2888–2897.
- Welch, R. M., and B. A. Wielicki (1989), Reflected fluxes for broken clouds over a lambertian surface, *J. Atmos. Sci.*, *46*, 1385–1395.

---

A. Benassi, C. Bouet, M. Leriche, and F. Szczap, Laboratoire de Météorologie Physique, Centre National de la Recherche Scientifique, Université Blaise Pascal, 24 avenue des Landais, F-63177 Aubière Cedex, France. (c.bouet@opgc.univ-bpclermont.fr)