



HAL
open science

Modelling of trace metals leaching behaviour in sludge deposits with high organic contents.

Camille Banc, Denise Blanc, Mathieu Gautier, Maria Lupsea-Toader, Rémy Gourdon

► **To cite this version:**

Camille Banc, Denise Blanc, Mathieu Gautier, Maria Lupsea-Toader, Rémy Gourdon. Modelling of trace metals leaching behaviour in sludge deposits with high organic contents.. European Meeting on Environmental Chemistry, Dec 2018, Clermont, France. hal-01973320

HAL Id: hal-01973320

<https://hal.science/hal-01973320>

Submitted on 8 Jan 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.


L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The matrix considered in this study comes from a small wastewater treatment plant called Vertical Flow Constructed Wetland. These are increasingly used in France with some 5000 plants already installed in 2018. Sludge deposits accumulated at the top of the filter are composed by organic matter but also by inorganic pollutants such as trace metals. Improving the understanding of the interactions between minerals and organic matter is necessary to predict the pollutant retention / release dynamics.

Context


Purpose

- Definition of a solid organo-mineral assemblage representative of the matrix;
- Evaluate the impact of different scenarios on the release and speciation of pollutants into the matrix;
- Better predict the behavior of trace metals.



Experimental


Results of elemental analysis



Others analysis :
 . Acid and Base neutralisation (CENTS/TS 14429)
 . Sequential extraction (method from Tessier et al., 1979)
 . X-Ray diffraction



Trace elements	Concentrations (mg/kgDM)	Thresholds limits (mg/kgDM)
Zn	1219	3000
Cu	623	1000
Pb	64.2	800
Cr	57.4	1000
Ni	30.5	200
Cd	1.7	20
...		

Conceptual model



Pure mineral phases

By their dissolution / precipitation properties, these phases will influence the concentration available in major elements but also in pollutant. The following example shows the CaCO₃ dissolution reaction :



Mineral assemblage



PhreeQc software (Parkhurst and Appelo, 1999) allows to simulate the dissolution of the different mineral phases implemented. For this, the software uses the thermodynamic data available in the database that has been specified. In our case, pure and mineral-bearing phases were implemented.

Electrostatic part

Depends on coulombic interactions that results from the **negative charge density** present on the molecule. Model VII considers that the charges of humic substances are located **only on the surface**.

Mineral-bearing phases

These matrices are also composed of mineral surfaces that can adsorb / release some pollutants through their binding sites. Example of Iron oxyhydroxides :




Organic assemblage


Model VII, developed by Tipping and its teams in 2011, is a database for simulating the adsorption capacity of humic substances (HS) via carboxylic and phenolic sites. The HS affinity for cations has two components: the **intrinsic part** and the **electrostatic part**.

Intrinsic part


It takes into account the specific binding properties of humic substances. 50 different sites are specified including **mono, bi and tridentate** site. Example of binding of Lead with a **Fulvic acid** site :




Evolution of Zn-leaching as a function of pH



Evolution dissolved organic carbon leaching concentration as a function of pH



Evolution of Cu-leaching as a function of pH



- . Importance of the sorption capacity of HS in basic solutions;
- . Copper is mainly related to HS;
- . Zinc is only partly bound to HS.

Results

Conclusions

- Combining PhreeQc and Model VII allows to reconstruct a complex organo mineral assemblage;
- The implementation of humic substances to the model provides good predictions of trace metals behavior ;
- Test on other type of matrices such as sediment from stormwater management pond.

References

Avena, M. J., Koopal, L. K., & van Riemsdijk, W. H. (1999). Proton binding to humic acids: electrostatic and intrinsic interactions. *Journal of colloid and interface science*, 217(1), 37-48.

Kim, B., Gautier, M., Michel, P., & Gourdon, R. (2013). Physical-chemical characterization of sludge and granular materials from a vertical flow constructed wetland for municipal wastewater treatment. *Water Science and Technology*, 68(10), 2257-2263.

Manon, K., Mathieu, G., Denise, B., Maria, L. T., Laurent, M., Maria-Chiara, Q., & Rémy, G. (2019). Leaching behavior of major and trace elements from sludge deposits of a French vertical flow constructed wetland. *Science of The Total Environment*, 649, 544-553.

Marsac, R., Davranche, M., Gruau, G., Bounnik-Le Coz, M., & Dia, A. (2011). An improved description of the interactions between rare earth elements and humic acids by modeling: PHREEQC-Model VI coupling. *Geochimica et Cosmochimica Acta*, 75(19), 5625-5637.