HAL
open science

The use of biogeochemical models to evaluate mitigation of greenhouse gas emissions from managed grasslands

Renata Sandor, Fiona Ehrhardt, Lorenzo Brilli, Marco Carozzi, Sylvie Recous, Pete Smith, Val Snow, Jean-François Soussana, Christopher D. Dorich, Kathrin Fuchs, et al.

To cite this version:

Renata Sandor, Fiona Ehrhardt, Lorenzo Brilli, Marco Carozzi, Sylvie Recous, et al.. The use of biogeochemical models to evaluate mitigation of greenhouse gas emissions from managed grasslands. Science of the Total Environment, 2018, 642, pp.292-306. 10.1016/j.scitotenv.2018.06.020 . hal-01976249

HAL Id: hal-01976249

https://hal.science/hal-01976249v1

Submitted on 9 Jan 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Science of the Total Environment 642 (2018) 292–306

Contents lists available at ScienceDirect

Science of the Total Environment

journal homepage: www.elsevier.com/locate/scitotenv

The use of biogeochemical models to evaluate mitigation of greenhouse gas emissions from managed grasslands

Renáta Sándor [a,b], Fiona Ehrhardt [c], Lorenzo Brilli [d,e], Marco Carozzi [f], Sylvie Recous [g], Pete Smith [h], Val Snow [i], Jean-François Soussana [c], Christopher D. Dorich [j], Kathrin Fuchs [k], Nuala Fitton [h], Kate Gongadze [l], Katja Klumpp [a], Mark Liebig [m], Raphaël Martin [a], Lutz Merbold [k,n], Paul C.D. Newton [o], Robert M. Rees [p], Susanne Rolinski [q], Gianni Bellocchi [a,*]

[a] INRA, VetAgro Sup, UCA, Unité Mixte de Recherche sur l'Écosystème Prairial (UREP), 63000 Clermont-Ferrand, France
[b] Agricultural Institute, CAR HAS, 2462 Martonvásár, Hungary
[c] INRA, 75007 Paris, France
[d] University of Florence, DISPAA, 50144 Florence, Italy
[e] IBIMET-CNR, 50145 Florence, Italy
[f] Agroscope Research Station, Climate Agriculture Group, Zurich, Switzerland
[g] FARE Laboratory, INRA, Université de Reims Champagne-Ardenne, 51100 Reims, France
[h] Institute of Biological & Environmental Sciences, School of Biological Sciences, University of Aberdeen, Aberdeen AB24 3UU, United Kingdom
[i] AgResearch - Lincoln Research Centre, Private Bag 4749, Christchurch 8140, New Zealand
[j] NREL, Colorado State University, Fort Collins, CO, USA
[k] Department of Environmental Systems Science, Institute of Agricultural Sciences, ETH Zürich, 8092 Zurich, Switzerland
[l] Rothamsted Research, Sustainable Soil and Grassland Systems Department, United Kingdom
[m] USDA-ARS, NGPRL, Mandan, ND, USA
[n] Mazingira Centre, International Livestock Research Institute, 00100 Nairobi, Kenya
[o] AgResearch Grasslands Research Centre, Private Bag 11008, Palmerston North 4442, New Zealand
[p] Scotland's Rural College, EH9 3JG Edinburgh, United Kingdom
[q] Potsdam Institute for Climate Impact Research, 14473 Potsdam, Germany

HIGHLIGHTS

- We perform multi-model simulations of C and N fluxes at five grassland sites.
- We assess modelled greenhouse gas emissions with alternative management practices.
- We use multi-model medians to reduce the uncertainty of the responses.
- We identify some shift towards a C sink with decreasing inputs.
- We show the considerable effect of N fertilizer reduction on C and N emissions.

GRAPHICAL ABSTRACT



ARTICLE INFO

Article history:
Received 24 February 2018
Received in revised form 15 May 2018
Accepted 2 June 2018

ABSTRACT

Simulation models quantify the impacts on carbon (C) and nitrogen (N) cycling in grassland systems caused by changes in management practices. To support agricultural policies, it is however important to contrast the responses of alternative models, which can differ greatly in their treatment of key processes and in their response

* Corresponding author.
E-mail address: gianni.bellocchi@inra.fr (G. Bellocchi).

https://doi.org/10.1016/j.scitotenv.2018.06.020
0048-9697/© 2018 Elsevier B.V. All rights reserved.

Available online 12 June 2018

Editor: P. Kassomenos

Keywords:
GHG emission intensity
Livestock density
Nitrogen fertilization
Process-based model
Sensitivity analysis

to management. We applied eight biogeochemical models at five grassland sites (in France, New Zealand, Switzerland, United Kingdom and United States) to compare the sensitivity of modelled C and N fluxes to changes in the density of grazing animals (from 100% to 50% of the original livestock densities), also in combination with decreasing N fertilization levels (reduced to zero from the initial levels). Simulated multi-model median values indicated that input reduction would lead to an increase in the C sink strength (negative net ecosystem C exchange) in intensive grazing systems: -64 ± 74 g C m^{-2} yr^{-1} (animal density reduction) and -81 ± 74 g C m^{-2} yr^{-1} (N and animal density reduction), against the baseline of -30.5 ± 69.5 g C m^{-2} yr^{-1} (LSU [livestock units] ≥ 0.76 ha^{-1} yr^{-1}). Simulations also indicated a strong effect of N fertilizer reduction on N fluxes, e.g. N_2O-N emissions decreased from 0.34 ± 0.22 (baseline) to 0.1 ± 0.05 g N m^{-2} yr^{-1} (no N fertilization). Simulated decline in grazing intensity had only limited impact on the N balance. The simulated pattern of enteric methane emissions was dominated by high model-to-model variability. The reduction in simulated offtake (animal intake + cut biomass) led to a doubling in net primary production per animal (increased by 11.6 ± 8.1 t C LSU^{-1} yr^{-1} across sites). The highest N_2O-N intensities (N_2O-N/offtake) were simulated at mown and extensively grazed arid sites. We show the possibility of using grassland models to determine sound mitigation practices while quantifying the uncertainties associated with the simulated outputs.

© 2018 Elsevier B.V. All rights reserved.

1. Introduction

Finding solutions to emerging ecological and societal challenges (climate change, food security, ecosystem sustainability) requires improved knowledge of the underlying processes affecting carbon nitrogen (C-N) pools and fluxes in agricultural systems (West and Marland, 2002; Giardina et al., 2014; Campbell and Paustian, 2015). Grassland ecosystems have a potentially important role to play in meeting the challenge of climate change because they can act as a source or sink for atmospheric carbon dioxide (CO_2) (Smith et al., 2008; Oates and Jackson, 2014) and are a source of non-CO_2 greenhouse gases (GHG) such as nitrous oxide (N_2O) and methane (CH_4). Importantly, these GHG emissions can be manipulated by management such as the method of grazing and the fertilizer regime (Soussana et al., 2004; Herrero et al., 2016). Several grassland experiments have addressed the role of management on the short-term GHG balance and global warming potential (e.g. Allard et al., 2007; Soussana et al., 2007; Hörtnagl et al., 2018). However, direct measurement of C-N balances should be supplemented by the use of simulation models, to support the implementation of effective practices and policies in agriculture, e.g. to mitigate GHG emissions (Rosenzweig et al., 2014; Elliott et al., 2015; Folberth et al., 2016). Biogeochemical process models address many of the complex interactions of weather, soil, vegetation and management practices (Bondeau et al., 1999; Churkina et al., 1999; Huntzinger et al., 2012; Warszawski et al., 2014; Chang et al., 2015) and can do so over long time intervals that are not feasible with experimentation. Existing modelling studies have focused on the determination of the C source and sink activity of grasslands (Soussana et al., 2010). Grassland models have been shown to provide adequate accuracy in representing observed yield and GHG emissions across a wide range of environments and management intensities (e.g. White et al., 2008; Chang et al., 2013; Graux et al., 2013; Ben Touhami and Bellocchi, 2015; Ma et al., 2015; Senapati et al., 2016; Ehrhardt et al., 2018).

Models can thus be beneficial for decision makers and farmers because they can be used to explore the productivity and environmental performances of specific systems as a consequence of changed management. However, the effect of management on C and N fluxes in agriculturally managed permanent grasslands (not re-sown more frequently than every five years, which is the focus of this study) is often uncertain (Schulze et al., 2009; Ciais et al., 2010), and such uncertainties are reflected in the outputs of the models used to simulate responses to management (Sándor et al., 2017). Grasslands are highly complex ecosystems and their behaviour is affected by multifaceted interactions of management drivers with water and nutrient availability, soil physics, and vegetation dynamics (Rees et al., 2013; Soussana et al., 2013). The dynamic grassland simulation models developed since the 1990s (e.g. Challinor et al., 2013; Snow et al., 2014; Calanca et al., 2016; J.W. Jones et al., 2017) differ greatly in their treatment of key processes, and hence in their response to environmental and management conditions (Brilli et al., 2017). A thorough assessment of the variation in the response, or sensitivity, of different grassland models to management factors can be critical in determining to what extent simulated responses may vary depending on the model used. From a policy perspective, it is critically important to identify the extent to which management interventions influence C-N fluxes (including productivity) prior to promoting policies that alter farming practices. If the impact of a given practice is uncertain, a sensitivity analysis can give information on the reliability of the models when representing C-N fluxes-management relationships under a variety of conditions. It is thus important to examine model behaviour under changed management in order to characterise the types of responses estimated, contrast the responses of different models and consider the reasons for these differences. In particular, hypotheses about the contribution of grassland management to GHG emissions can be tested via simulation models, which allow understanding, diagnosing and forecasting complex interactions (Chen et al., 2008; Seijan et al., 2011; Graux et al., 2012; Sándor et al., 2017, 2018).

Consequently, using five case studies, we tested the sensitivity of eight grassland models to gradients of management intensity that were selected for their potential to mitigate GHG emissions (e.g. Soussana et al., 2014; Abdalla et al., 2017). With the aim of increasing the reliability and confidence in simulated results, a multi-model ensemble approach was adopted to explore patterns of simulated C and N responses against imposed gradients of N fertilization and animal stocking rate (to which grassland models are generally sensitive, after Brilli et al., 2017). For this study, we included a range of well-known grassland models, and used them to simulate biogeochemical and related outputs (productivity and energy measures). The wider ensemble analysis presented in Ehrhardt et al. (2018) forms the baseline for the work presented here, which analyses factors that may explain the major differences observed in model responses. We further explored to what extent multi-model ensembles can be used to help identify farming practices that reduce GHG emissions. While restricting the analysis to a limited set of management options, this study examines a wide range of output variables and thus provides a framework for assessing grassland performance where direct casual links with farming practices are not obvious, and changes in performance are difficult to measure. As a corollary outcome, viewing and interpreting a variety of model outputs lay ground for future model developments.

2. Materials and methods

We refer to a sub-set of the grassland models described in Ehrhardt et al. (2018), in which models were initialized and calibrated using vegetation and soil variables, and surface-to-atmosphere fluxes at four sites worldwide. We used an ensemble of grassland models (Table 1) and compared their sensitivity to changes in management by comparing

Table 1
The biogeochemical models used for testing the impact of grassland management options.

Model/version	Description/references
APSIM-SWIM v7.7 APSIM-SoilWater v7.7	APSIM (The Agricultural Production Systems sIMulator; Holzworth et al., 2014) simulates several systems through the interaction among plants, animals, soil, climate and management. The model allows the analysis of the whole-farm system, including livestock, crop and pasture sequences and rotations. Users can select between two soil water models: the capacitance-based SoilWater (Probert et al., 1998) and SWIM, which is based on Richards' equation (Huth et al., 2012). The pasture model was that of Li et al. (2011).
DayCent v4.5 2006 DayCent v4.5 2010 DayCent v4.5 2013	DayCent is the daily time-step adaptation of the biogeochemical model CENTURY (Parton et al., 1998). It simulates biomass growth, soil C dynamics, N leaching, gaseous emissions (e.g. N_2O, NO, N_2, NH_3, CH_4 and CO_2) and C fluxes (e.g. NPP, NEE) in croplands, grasslands, forests and savannahs, as affected by management practices (such as fertilization, tillage, pruning, cutting and grazing) and specific external disturbances (e.g. fires). Different versions of the model result in different parameter settings and a few variations in the model structure. DayCent v4.5 2006 applies grazing on a daily basis as linear impact on aboveground biomass and root/shoot ratio, with aboveground biomass removed as a percentage of total aboveground biomass. DayCent v4.5 2010 and 2013 apply grazing on a daily basis with aboveground biomass removed as a percentage of total aboveground biomass rather than as continuous grazing. In DayCent models after 2013, water stress effect on biomass production differs from the previous versions.
LPJmL v.3.5.003	LPJmL (Lund-Potsdam-Jena managed Land) explicitly simulates key ecosystem processes such as photosynthesis, plant and soil respiration, C allocation, evapotranspiration and phenology of nine plant functional types representing natural vegetation at the level of biomes (Sitch et al., 2003), and of 12 plant functional types (Bondeau et al., 2007; Rolinski et al., 2018).
PaSim	PaSim (Pasture Simulation model; Riedo et al., 1998; Calanca et al., 2007; Ma et al., 2015) is a process-based, grassland-specific ecosystem model that simulates grassland productivity and GHG emissions to the atmosphere. The model consists of sub-models for vegetation, grazing animals, microclimate, soil biology, soil physics and management.
SPACSYS v5.0	SPACSYS (Wu et al., 2007, 2015) is a multi-layer, field scale, weather-driven and daily-time-step dynamic simulation model. The current version includes a generic plant growth and development, C and N cycling, plus simulation of soil water that includes representation of water flow to field drains as well as downwards through the soil layers, together with a heat transfer component. The model simulates root architecture.

simulated outputs against gradients of management practices. Model anonymity was maintained throughout the process and model results are presented without attributing them to specific models or modelling teams.

We present multi-model medians and box-plots, and focus on long-term averages for the same four grassland sites (G1 to G4) described in Ehrhardt et al. (2018) plus an additional site (G5) for which full calibration was only completed after the initial publication (Table 2). Overall, there is a lack of case studies in Asia, Africa and South America (which would have extended the comprehensiveness of the research), but sites from G1 to G5 are intended to represent situations commonly encountered in temperate grasslands. While the choices made are described in Ehrhardt et al. (2018), in summary it was thanks to international collaborations that we could pool and share experimental data for five grassland sites (one more than in Ehrhardt et al., 2018). These sites provided high-quality, previously published data encompassing climate, soil, agricultural practices, and C and N fluxes.

To analyse the sensitivity of models with respect to changes in grassland management practices, viz. animal stocking density and N fertilization, management scenarios were obtained by adjusting the observed baseline management (business-as-usual) for each site with systematic decrements over a range of values (Table 3). Sensitivity is defined as the proportional change in models outputs that results from a change in a given factor (here management practices).

In our study-sites, two major practices are responsible for C and N fluxes from grasslands: (1) vegetation removal and (2) fertilizer inputs. The harvesting of vegetation was predominantly controlled by grazing animals for the majority of sites. The exception was G5 where the grazing was light and vegetation was predominantly removed by cutting. Accordingly, a reduction in grassland use was assessed by a limitation of livestock density, either alone or together with reduction (down to cessation) of fertilizer N in N-fertilised sites (G3, G4 and G5) (Table 3). The livestock unit (LSU) based on the grazing equivalent of one adult cow was used to compare different animal types (yearling steers, non-lactating sheep, ewes, lambs, heifers and calves).

Impacts of the defined changes in management were calculated on the changes in a set of output variables related to biomass production and C-N fluxes (Table 4). Fluxes of CO_2 included emissions from ecosystem respiration (R_{ECO}), respiration from plants (R_{PLANT}), soil (R_{SOIL}) and grazing animals (R_{ANIMAL}) as well as estimates of the plant production of organic compounds from atmospheric CO_2 (GPP) and other system variables: Net Ecosystem Exchange, NEE = R_{ECO} − GPP; Net Primary Production, NPP = GPP − R_{PLANT}; Net Biome Production, NBP = −NEE + C losses through enteric CH_4 emissions at pasture, forage harvests and milk production at pasture.

Methane released from soil and enteric fermentation in animals was included in the list of non-CO_2 fluxes, along with the gaseous N compounds emitted to the atmosphere: N_2 (N gas), NO_x (N oxides: the sum of N monoxide, NO, and N dioxide, NO_2), N_2O (nitrous oxide) and NH_3 (ammonia). Nitrogen lost by nitrate (NO_3) leaching was also examined. The biological information included productivity measures such as the plant biomass produced above – and below-ground (AB and BB), two outputs of agronomic interest – the plant biomass consumed by grazing animals (Intake) or otherwise harvested (HAB), and their sum (Offtake) – and the energy that ultimately is utilised by grazing animals ($ME_{Offtake}$: offtake metabolisable energy).

To estimate the amount of plant biomass available for feeding animals, the annual NPP values were normalized by animal stocking rates. We also expressed C and N fluxes relative to the overall productivity of the system, so that we could express the intensity of GHG emissions on the basis of productivity (i.e. g of emitted C or N per g C of harvested or per g C ingested dry matter, DM). This approach is similar to the concept of 'yield-scaled emission' or emissions intensity as defined by Van Groenigen et al. (2010) and has important policy significance and delivers results that are relevant to stakeholders (Venterea et al., 2011; Valin et al., 2013). For this purpose, three additional variables were analysed, representing the ratios of CO_2-C, N_2O-N and CH_4-C emissions to the total amount of C biomass (Offtake) consumed by animals (Intake) and harvested as fodder (HAB): $Int_{CO2\text{-}C}$ = −NEE/Offtake, $Int_{N2O\text{-}N}$ = N_2O-N/Offtake, $Int_{CH4\text{-}C}$ = CH_4-C/Offtake.

Following Sándor et al. (2016), we report the proportional change, named effect size of mean annual output variables from a change in each factor relative to the baseline management at each site.

The N_2O-N emission factor (EF) for fertilizer was calculated as percent ratio of the total yearly N_2O-N emissions over the amount of the annual N fertilizer. This simplified version of the N_2O-N emission factor calculation does not take into account of background emissions because not all the models allowed for a consistent estimation of this component. For this reason, following the IPCC (2006) guidelines, 1 kg N_2O-N ha^{-1} yr^{-1} background emission was subtracted from the simulated values.

The present study was based on yearly aggregated model outputs. R software (R Core Team, 2016) was used for statistical computing and visualization. Accounting for the different global warming potential (GWP) of CO_2, CH_4 and N_2O, total GHG balances were achieved by

Table 2
Selected grassland sites for the modelling exercise.

General description	Grassland sites				
Site code	G1	G2	G3	G4	G5
Country	United States	New Zealand	France	United Kingdom	Switzerland
Location	Mandan	Flockhouse	Laqueuille	Easter Bush	Chamau
Climate[a]	Dfb (humid continental)	Cfb (oceanic)	Cfb (oceanic)	Cfb (oceanic)	Cfb (oceanic)
Latitude	46.77	−40.20	45.64	55.52	47.20
Longitude	−100.89	175.30	2.74	−3.33	8.40
Elevation a.s.l. (m)	591	30	1040	190	393
Simulation period	2003–2006	1997–2008	2003–2012	2002–2010	2010–2013
Mean annual minimal air temperature (°C)[b]	0.0 ± 1.0	9.1 ± 0.5	4.0 ± 0.6	4.6 ± 0.9	4.4 ± 0.6
Mean annual maximal air temperature (°C)[b]	11.9 ± 1.3	17.6 ± 0.6	11.0 ± 0.8	11.4 ± 0.8	14.7 ± 0.8
Mean annual cumulated precipitation (mm)[b]	411 ± 128	896 ± 107	1047 ± 144	961 ± 142	1084 ± 143
Management					
Type	Grazed	Grazed	Grazed	Grazed/mown	Grazed/mown
Animal type	Yearling steers	Non-lactating sheep	Heifers	Ewes, lambs, heifers and calves	Sheep
Mean annual number of grazing days[c]	107	22	163	162	14
Stocking rate (LSU ha^{-1} yr^{-1})[c]	0.08	1.29	1.34	0.76	0.10
Vegetation type	C3 grasses	C3 grasses, legumes, forbs, C4 grasses	C3 grasses, legumes, forbs	C3 grasses	C3 grasses, legumes
Mean annual number of cutting events[c]	0	0	0	0.9	6.3
Total annual N fertilization (kg N ha^{-1} yr^{-1})[c]	0	0	210	220	230
Soil properties					
Soil type[d]	Calcic Siltic Chernozem	Mollic Umbrisol	Loamic Andosol	Eutric Cambisol	Gleysol
Maximum depth of the soil profile (m)	4	0.9	0.9	1.0	1.0
Number of documented layers	6	4	5	5	4
Soil texture: ~sand (%)[e]	29.7	93.1	24.6	22.9	57.2
~silt (%)[e]	51.0	3.8	55.5	19.0	28.9
~clay (%)[e]	19.5	3.1	21.8	58.1	14.0
Bulk density (g cm^{-3})[e]	1.17	1.20	0.67	1.45	1.34
References	Liebig et al. (2006, 2010, 2013)	Newton et al. (2010, 2014)	Allard et al. (2007) and Klumpp et al. (2011)	Skiba et al. (2013) and S.K. Jones et al. (2017)	Imer et al. (2013) and Merbold et al. (2014)

[a] Köppen-Geiger climate classification (Kottek et al., 2006).
[b] Mean minimum and maximum air temperatures, and precipitation totals calculated over 30 years (1980–2009) using AgMERRA (https://data.giss.nasa.gov/impacts/agmipcf/agmerra) meteorological datasets (Ruane et al., 2015).
[c] Mean values over the simulation period. Grazing at G2 site was on a rotational basis, i.e. animals were brought in at intervals for short periods at a high stocking rate, while at all other sites grazing was by set-stocking, i.e. animals were maintained continuously on the pasture at a low stocking rate.
[d] World Reference Base for Soil Resources (FAO, 2014).
[e] Mean values across multiple layers.

converting CH_4 and N_2O emissions rates to CO_2 equivalents (CO_2e) using the 100-year Global Warming Potential (GWP_{100}) as established in national GHG inventories, i.e.: 1 kg N_2O = 298 kg CO_2e, 1 kg CH_4 = 25 kg CO_2e (IPCC, 2006; https://www.epa.gov/ghgemissions/understanding-global-warming-potentials). Mass factors were also applied to model outputs, the latter being expressed in C and N units: 1 kg CH_4-C = 1.33 kg CH_4, 1 kg N_2O-N = 1.57 kg N_2O, 1 kg CO_2-C = 3.67 kg CO_2.

Table 3
Design of management options (where 100% indicates the baseline - business-as-usual - management scenario).

Action[a]	Sites	Description
Reduction of livestock density (LD)	G1, G2, G3, G4, G5	The livestock density in the pasture was decreased in five steps of 10% (from 100% down to 50% of the livestock density indicated by the default standard management) Abbreviations[b]: LD_{90}, LD_{80}, LD_{70}, LD_{60}, LD_{50}
Reduction of livestock density (LD) and nitrogen (N) fertilizer	G3, G4, G5	The amount of mineral or slurry N added to the pasture was decreased in five steps of 20% (from 100% to 0% of the N amount indicated by the default standard management), while the livestock density in the pasture is decreased in five steps of 10% (from 100% down to 50% of the livestock density indicated by the default standard management) Abbreviations[b]: $LD_{90}N_{80}$, $LD_{80}N_{60}$, $LD_{70}N_{40}$, $LD_{60}N_{20}$, $LD_{50}N_0$

[a] When animal density was decreased, cutting events (if present) were left unaltered. When present, supplementary feeding was proportionally reduced along with the animal density.
[b] Percent livestock density (90, 80, 70, 60, 50) or N fertilizer (80, 60, 40, 20, 0) against baseline (business-as-usual). Without assessing all possible LD x N combinations, we focussed on reducing overall levels of management intensity through reductions in N inputs and grazing levels (according to most agri-environment schemes for grassland; Atkinson et al., 2005).

Table 4
Model outputs (annual cumulative) generated by each model (✓: available; NA: not available) and assessed in the study. The identities of models were kept anonymous by using the same model codes as in Ehrhardt et al. (2018).

Variable/models		M05	M06	M07	M08	M16	M22	M24	M28
CO_2 fluxes	GPP (gross primary production): g C m^{-2}	✓	✓	✓	✓	✓	✓	✓	✓
	NPP (net primary production): g C m^{-2}	✓	✓	✓	✓	✓	✓	✓	✓
	NEE (net ecosystem exchange): g C m^{-2}	✓	✓	✓	✓	✓	✓	✓	✓
	NBP (net biome production): g C m^{-2}	NA	✓	NA	NA	✓	NA	✓	✓
	R_{ECO} (ecosystem respiration): g C m^{-2}	✓	✓	✓	✓	✓	✓	✓	✓
	R_{PLANT} (plant respiration): g C m^{-2}	✓	✓	✓	✓	✓	✓	✓	✓
	R_{SOIL} (soil respiration): g C m^{-2}	✓	✓	NA	✓	✓	✓	✓	✓
	R_{ANIMAL} (animal respiration): g C m^{-2}	NA	✓	NA	NA	✓	NA	✓	NA
Non CO_2 fluxes[a]	CH_4 emissions (methane)[b]: g C m^{-2}	NA	✓	✓	✓	✓	NA	✓	NA
	N_2O (nitrous oxide) emissions: g N m^{-2}	✓	✓	✓	✓	✓	NA	✓	✓
	NH_3 (ammonia) emissions: g N m^{-2}	NA	✓	NA	✓	✓	NA	✓	NA
	NO_x (nitrogen oxides) emissions: g N m^{-2}	NA	NA	✓	✓	NA	NA	NA	NA
	N_2 (nitrogen gas) emissions: g N m^{-2}	NA	✓	✓	✓	NA	NA	✓	✓
	N (nitrogen) leaching: g N m^{-2}	NA	✓	NA	✓	✓	NA	✓	✓
Productivity and energy	Aboveground biomass (AB): g DM m^{-2}	NA	✓	✓	✓	✓	✓	✓	✓
	Belowground biomass (BB): g DM m^{-2}	NA	✓	✓	✓	✓	NA	✓	✓
	Harvested biomass (HAB): g DM m^{-2}	NA	✓	✓	✓	✓	NA	✓	✓
	Animal intake (intake): g DM m^{-2}	NA	✓	NA	✓	✓	✓	✓	✓
	Metabolisable energy of Offtake (grazing plus harvesting) ($ME_{Offtake}$): MJ kg^{-1} DM	NA	✓	NA	NA	NA	NA	✓	NA

[a] Fluxes are expressed in units of C (CH_4-C) and N (N_2O-N, etc.).
[b] CH_4 emissions include emissions from both animals (enteric) and their manure. The latter were estimated for M16, and the former were estimated for M06, M08 and M24; all estimations were based on Clark et al. (2003).

3. Results and discussion

Simulated results are presented and discussed separately, with selected graphs, for the following groups of variables: CO_2 fluxes, non-CO_2 fluxes, productivity and energy, and emission intensities. Additional results are provided in the Supplementary material (Figs. A to S).

3.1. CO_2 fluxes

In the baseline scenario, GPP showed a wide range of variations in multi-model medians (137.2–1732.4 g C m^{-2}), while animal respiration (R_{ANIMAL}) was the output with the least divergent results among the models (0.0–211.6 g C m^{-2}; Fig. 1). For the R_{ANIMAL}, model differences tended to be smaller at lower input levels, especially when animal density was reduced without reductions in N fertilization (0.0–143.9 g C m^{-2}). For plant respiration (R_{PLANT}), an increase of model variability was associated with the reduced influence of the livestock. It is also interesting to note that net ecosystem exchange (NEE) values showed large variability among models with intermediate intensification levels, e.g. 70% reduction of LD and 40% reduction of N. The greatest variability was simulated under the mowing-dominated G5 site (Figs. A and B in Supplementary material), but the model variability was also high under mowing and grazing combined management at G4 and under intensively grazed (by heifers) management at the G3 site. Reduction in N fertilization tended to decrease NEE variablity at the G3 and G4 sites. In G5, the analysis of proportional changes indicated - with the combined reduction of N fertilization and animal density - a clear linear decrease in NEE compared to the baseline (Fig. C in Supplementary material). Since NEE is defined as the difference between ecosystem respiration (R_{ECO}) and gross primary production (GPP), the variability of its basic components have an effect on the spread of NEE values. The ensemble uncertainty of GPP and R_{ECO} were highest at G2, G3 and G4 sites, associated with the highest animal densities (Table 2). This suggests that the intensification of grazing management tends to increase the variation of GPP and R_{ECO} estimates between models, with a smaller uncertainty envelope at the G1 and G5 sites, where altered animal density variation is very low (0.1–0.04 LSU ha^{-1} yr^{-1}).

The five grassland sites showed different dynamics in C fluxes with respect to the simulated management options, with NEE varying between −231.3 and +189.2 g C m^{-2} yr^{-1}, considering all sites and simulation years. These results suggest higher NEE (−19.0 ± 75.9 g and −47.6 ± 89.8 C m^{-2} yr^{-1} for baseline and LD_{50}, respectively), or lower C uptake, than Soussana et al. (2007) concluded from nine European grassland sites equipped with eddy-covariance flux measurements, which showed an average net sink of atmospheric CO_2 with NEE of −240 ± 70 g C m^{-2} yr^{-1} (which is in the range −486.3 to 24.8 g C m^{-2} yr^{-1}, or −1783 to −91 g CO_2 m^{-2} yr^{-1}, provided by Hörtnagl et al., 2018 for managed grasslands in Central Europe). The site-by-site analysis (Fig. C in Supplementary material) indicated, except at the G5 site, that C uptake was the dominant process. At the G5 site, an NEE of <0 only occurred with $LD_{70}N_{40}$ management options. At this site, the greatest model uncertainty in NEE values occurred with



Fig. 1. Changes in CO_2 fluxes (g C m^{-2}) calculated over multiple years at five sites, for ten altered management scenarios and the baseline (as in Table 3; LD: livestock density; N: nitrogen). For each management level, triangles show the multi-model (as in Table 4) mean, black lines show multi-model median. Boxes delimit the 25th and 75th percentiles. Whiskers are 10th and 90th percentiles. Points indicate outliers.

the $LD_{70}N_{40}$ management option (Fig. 1), related to differently simulated CO_2 release and uptake processes. According to the median values, the G5 site reached the highest amount of C sequestration (~113.7 g C m^{-2} yr^{-1}) at $LD_{50}N_0$. The general observation from the modelled sites of increasingly negative NEE in response to increasing N inputs is consistent with experimental observations that report increases in the flow of C to the soil in response to long-term fertilizer N use (Ammann et al., 2007; Skiba et al., 2013). For the period during which the C is sequestered, it is removed from the atmosphere and does not contribute to global warming. This effect is simulated at both grazed (G1, G2 and G3) and grazing dominated (G4) sites, for all scenarios. Owing to the large quantity of harvested aboveground biomass (332.1 ± 62.3 g DM m^{-2} yr^{-1} on average for the baseline scenario), the G5 site seems to release C from the soil. However, the grazing removal (395.8 ± 38 g DM m^{-2} yr^{-1} on average for the baseline scenario of G3 site) may drive less radical changes in the C balance. Overall, these simulation results are consistent with eddy-covariance measurement data (Senapati et al., 2014; Koncz et al., 2017), in which mown treatments were observed to release C, while grazed treatments acted as net C sinks. For instance, in Senapati et al. (2014) mown treatments had lower annual net C storage (22.7 ± 32.3 g C m^{-2} yr^{-1}, net sink), related to hay removal, than grazed plots (140.9 ± 69.9 g C m^{-2} yr^{-1}, net sink) - though the observed site (Lusignan, France) was recently converted from cropland to grassland, so would be expected to be increasing in soil C (Senapati et al., 2014).

Considering all the sites, the estimated average C exchange (net biome productivity, NBP, Fig. 1) ranged between −176.9 (sink) and +140.4 g C m^{-2} yr^{-1} (source), with its extremes at $LD_{50}N_0$ and LD_{60} management options, respectively. This high variability was caused by different management systems at G4 (grazed and mown) site (Figs. A and B in Supplementary material), while the extensification combined with N reduction tended to increase C storage in some cases, e.g. at G5 site. Owing to the high organic C exports (from haycut and/or intensive cattle grazing: 1.34 and 1.21 LSU ha^{-1} yr^{-1}), which could be greater than C imports from manure and slurry, the soil processes would be dominated by C emissions at the G2 site (intensive scenarios). The N mitigation reduced the net biome production at G3 site even further.

Ecosystem respiration (R_{ECO}), together with R_{PLANT} and R_{SOIL}, showed a linear decrease as LD and N levels simultaneously decreased, but tended to increase with a reduction in animal density only (Fig. 1). Animal respiration (R_{ANIMAL}) tended to decrease as animal density decreased, though the multi-model median line (Fig. 1) was associated with some uncertainty at the baseline and $LD_{70}N_{40}$ options (i.e. 30% less animals and 60% less N fertilizer). Site-by-site analyses showed (Figs. A and B in Supplementary material) that the greatest simulated R_{ECO} occurred with G3 and G4 grazing systems. Reduction in N fertilizer tended to decrease R_{ECO}, however the variability of soil respiration (R_{SOIL}) increased the uncertainty at G4, particularly under sheep-heifer grazing. Based on model simulations, the main losses of CO_2 at the G4 site were caused by R_{SOIL} and plant respiration (R_{PLANT}). Simulated yearly R_{ANIMAL} values, and their proportional changes (Fig. D in Supplementary material) with respect to the baseline management showed a distinct emission decrease with extensification (which is not the case with R_{PLANT} and R_{SOIL}, Figs. A and B in Supplementary



Fig. 2. Comparison of the combined effect of livestock density reduction (LD from 90% down to 50% of initial density) and the N fertilization reduction (N from 80% down to 0% of initial amount) at G3, G4 and G5 sites for gross primary production (GPP), ecosystem respiration (R_{ECO}) and net ecosystem exchange (NEE) using the multi-model median.

material), at a quasilinear rate (by $y = -0.56 + 0.05x$, R^2~1 for LD changes and $y = -0.62 + 0.05x$, $R^2 = 0.99$ for N and LD combined options at G3 site). The data also suggest that there were substantial differences among models in the estimated effects of altered management. These differences were amplified when N fertilization was decreased jointly with animal density (e.g. option $LD_{70}N_{40}$ at G4 and G5; Fig. D in Supplementary material), which suggests that interactions between the two factors may make a sizeable contribution to this variability in the response of different models.

Some relationships between model outputs and management inputs were apparent from an analysis of data at each site. GPP decreased strongly with stocking rates while R_{ANIMAL} increased, with the exception of G2 site (Fig. E in Supplementary material). The simulated outputs at this site may reflect different and non-linear responses of alternative models (Fig. F in Supplementary material). Often, the offtake increased and NPP decreased with management intensification, albeit with large differences between models. With M06, the highest Offtake was simulated when animal density decreased by 30%, while this happened at 10% lower animal density with M24. M08 simulated the highest NPP value at 30% less livestock density, while with M28 the highest NPP values were simulated at the most intensive management condition.

The influence of N fertilization was investigated at G3, G4 and G5 sites (Fig. 2) by comparing the combined effect of livestock density reduction and N fertilization reduction. In terms of GPP (Fig. 2) and NPP (Fig. L in Supplementary material), the simulations showed a considerable decrease in GPP with >60% less applied N fertilizer at site G4. The R_{ECO} (Fig. 3, middle panel) values also decreased with extensification, where the N reduction had a greater effect at the G3 and G4 sites. Animal respiration was driven by livestock density (Fig. G in Supplementary material), while soil and plant respiration were mainly influenced by N inputs (lower R_{PLANT} and R_{SOIL} with lower N inputs). In terms of NEE (Fig. 2, lower panel) and NBP (Fig. G in Supplementary material) the trend was less obvious, owing to differences in management and site conditions.

The effectiveness of the different management strategies such as fertilizer amounts, different animal stocking rates, grazing alone or combined with mowing, was also influenced by site-specific soil (type and depth) and weather conditions (ie. Precipitation). If we are to distinguish between environmental from management effects, then precipitation patterns must be taken into account as it can also have an influence on the results of the CO_2 fluxes (e.g. Polley et al., 2010). In our simulation study, the amount of precipitation showed a positive correlation with R_{SOIL} and all the other investigated CO_2 fluxes (Fig. H in Supplementary material) apart from NEE and NBP. The respiration outputs demonstrated higher sensitivity to N fertilization, than to percent livestock density changes. NEE values suggested greater respiration in very arid years (such as some years at G1 site, where annual mean precipitation was 271 ± 141 mm), where the animal density reduction did not reduce the amount of CO_2 emissions. A recent review highlighted the particular sensitivity of warm and dry climates to change in stocking density where increased livestock density was associated with significantly lower rates of C sequestration (Abdalla et al.,



Fig. 3. Changes in non-CO_2 fluxes (g C m^{-2}, g N m^{-2}, log_{10} g N m^{-2}) calculated over multiple years at five sites, for ten altered management scenarios and the baseline (as in Table 3). For each management level, triangles show the multi-model (as in Table 4) mean, black lines show multi-model median. Boxes delimit the 25th and 75th percentiles. Whiskers are 10th and 90th percentiles. Hollow circles indicate outliers (LD from 90% down to 50% of initial density, N fertilization from 80% to 0% of initial amount).

2018). Gilmanov et al. (2007) pointed out that organic and semi-arid grassland sites have the potential to become C sources. With decreasing stocking rate, NEE tended to increase above 800 mm annual precipitation. In humid and very humid years, the soil may be saturated and anaerobic, and organic C decomposition can be slowed or stopped under these conditions (yet anaerobic decomposition of partly decomposed organic matter may emit methane, e.g. Bannert et al., 2012). The variability of NEE decreased with N reduction, e.g. the most extensive treatment showed the smallest difference from zero (Fig. H in Supplementary material).

3.2. Non-CO_2 fluxes

A recent IPCC (2014) report and other analyses (Schulze et al., 2009; Tubiello et al., 2014; Gerber et al., 2015; Hörtnagl et al., 2018) highlight the importance of the reduction of non-CO_2 GHGs, as an important element of agricultural mitigation strategies. Particularly methane and nitrous oxide cause longer-term warming effects than CO_2. There were clear trends in model responses, indicating decreases in N_2O-N, NH_3-N, NO_x-N, N_2 and NO_3-N leaching with reductions in N fertilizer, while there were no clear trends in the response to animal livestock reduction (Fig. 3, all models and sites confounded). Only NO_3-N leaching showed a distinctly linear decrease with simultaneous decreases of N fertilizer and livestock density, suggesting a close dependence of this output on N fertilization input. Methane (CH_4-C) emissions tended to decrease with decreasing livestock density and also with reductions in N fertilization.

For N_2O emissions, relative effect size analysis (Fig. I in Supplementary material) and simulated emissions (Fig. 4 and Fig. J in Supplementary material, respectively) revealed linear decreasing trends as both livestock density and N fertilizer were reduced (sites G3, G4 and G5, Fig. 4, bottom). The greatest mitigation of N_2O-N emission was obtained by reductions in N fertilizer at the G4 and G5 sites, where the initial 0.51 and 0.63 g N_2O-N m^{-2} were reduced to 0.11 and 0.16 g N_2O-N m^{-2}, respectively. In terms of total GHG emissions, using the 100-year Global Warming Potential (GWP_{100}), the N mitigation from baseline to zero would reduce the multi-model median simulated N_2O emissions by 135.7, 187.1 and 219.9 g CO_2e m^{-2} yr^{-1} at G3, G4 and G5 sites, respectively. This corresponds to ~16–25% of C sink potential, reported by Soussana et al. (2007) across nine European grassland sites, but a larger percentage (~40–65%) of the sink potentials determined in this study.

The reduction of N fertilizer logically decreases the N_2O-N emissions, as reported here (Fig. J in Supplementary material) and by experimental studies (Cardenas et al., 2010; Bell et al., 2016; Hörtnagl et al., 2018). Our results at the G3 site showed the same trend (Table 5), when the N_2O-N emissions are compared to the applied N fertilizer amounts, the estimated (simplified) N_2O-N emission factors (percent ratios of the total yearly N_2O-N emissions over the amount of annually applied N fertilizer, both in kg N ha^{-1}). Our simulated results (varying between 1.0 and 3.5% across sites and treatments) are not far from the IPCC (2006) default EF for fertilizer N value, which is 1%. At G4 and G5 sites, the EF values tended to increase as grassland management received less N fertilizer (Table 5), which suggest some non-linear reduction of N_2O under reduced fertilizer supply, which can be explained by a decrease of plant N uptake with decreasing N fertilizer rate (e.g. Lü et al., 2014). Negative relationships between N use efficiency and soil N availability were observed in a variety of ecosystems, including grasslands (e.g. Yuan et al., 2006). Decreased N uptake from the soil and less efficient use of the N assimilated by plants leave more N available for microbes in the soil (which is the most important factor for N_2O-N emissions). Thus, the most intensive systems (G3, G4 and G5) had the highest CO_2e emission rates while the N_2O-N emission factors varied between the managements options. There was no trend in median values of simulated N_2O emissions and LD levels, with the exception of G1 and G4 (Fig. I in Supplementary material). At the G1 site, a slight decreasing trend was noticeable with decreasing grazing intensity, with increasingly diverging results among models as more extensive management was introduced.

Overall, the different N fluxes (Fig. 4 and Fig. K in Supplementary material) tended to decrease with reduced N fertilization, mainly after a 60% reduction in the amount of N applied in both grazed and combined (mown and grazing) systems.



Fig. 4. Relationship between nitrous oxide emissions (multi-year averages of seven models) given in N_2O-N and CO_2e forms, and increasing animal stocking rate at all sites comparing livestock density reductions (top graph) and livestock density and N fertilization reduction (bottom graph), as in Table 3.

Regarding CH_4-C emissions (Fig. 5), the G3 site had much higher model uncertainty than other sites, mostly due to high estimates from M06 and M24. The simulated multi-model median values were the highest, with lower livestock density. For example, the multi model multi-year average baseline simulation was 4.6 g C m^{-2}, which was reduced to 3.6 g C m^{-2} with stocking rate reduction alone, and to 3.4 g C m^{-2} in combination with N reduction at the G3 site (Fig. 5). The main agricultural source of CH_4 at the G3 site was the intensive heifer grazing system. Other sites were less affected because either they had grazing sheep (G2 and G5) or were less intensively grazed (low cattle density in G1, combined sheep and cattle grazing in G4).

Site-specific circumstances, mainly soil properties (particularly soil N availability), and precipitation patterns, have considerable effect on the N balance of the grassland sites. Our modelled outputs show that soils tended to release more N_2O-N and CH_4-C (Fig. L in Supplementary material) in humid and very humid years, while the net N emissions were lower during drier years. The highest N_2O-N emissions and also N_2 and NO_x emissions (Fig. L in Supplementary material) were simulated when annual precipitation was around 1000 mm, owing to the more available N, because high rainfall rates increase the rate of N transport to deeper soil layers and increase nitrate leaching (Fig. L in Supplementary material). Also, there would be more anaerobic microsites leading to greater rates of denitrification in waterlogged soils (Smith et al., 1998). Besides, higher NPP values (Fig. H in Supplementary material) were associated with elevated precipitation, thus higher organic N inputs to soil may contribute to the larger N_2O-N emissions. This indicates that the intensity of N losses tends to be associated with annual precipitation levels, and N losses can be effectively mitigated by reducing livestock density and/or N fertilization rates. Our simulations indicate reductions in N fertilization as the most effective option for mitigating non-CO_2 fluxes, mainly in humid areas, an observation that is consistent with a number of previous studies (Bouwman et al., 2002; Rees et al., 2013) but further studies are required considering the high variability of model responses.

Table 5
N_2O-N emission factors: multi-model median at the three N fertilised sites (as in Table 2). Grey cells indicate the lowest values.

Management options	G3	G4	G5
Baseline	1.3%	1.8%	2.3%
$LD_{90}N_{80}$	1.4%	1.2%	2.0%
$LD_{80}N_{60}$	1.6%	2.0%	1.9%
$LD_{70}N_{40}$	1.4%	2.2%	2.5%
$LD_{60}N_{20}$	1.0%	1.7%	3.5%
$LD_{50}N_0$	–	–	–

3.3. Productivity and energy outputs

Some decreasing trends with management extensification can be observed in the box-plots of Fig. 6, e.g. for simulated Offtake and Intake, while aboveground and belowground biomass increased with lower stocking rates, combining simulation results from five sites. In terms of harvested aboveground biomass, a drop was simulated with no N fertilization (e.g. baseline simulation: 421.7 ± 118.6 g DM m^{-2}, $LD_{50}N_0$ management: 200.9 ± 78.8 g DM m^{-2} across the multi-year site averages).

In fertilised sites, where LD levels have been assessed alone and in combination with N fertilisation levels, Fig. 6 and Fig. M in the Supplementary material indicate that, overall, reducing N fertilizer rate will have more effect on aboveground biomass and biomass offtake than reducing livestock density, as data points relative to sites G3, G4 and G5 tend to stay below the 1:1 line. The effect of N fertilizer reduction starts becoming visible at G4 with 60% N reduction, while only with no N fertilization is this effect visible at G5. However, livestock density reduction has a greater effect on animal intake, belowground biomass and $ME_{Offtake}$ (Fig. M in Supplementary material). Animal intake decreases considerably with extensification (Fig. 6 and Fig. N in Supplementary material).

The annual NPP values, normalized by animal stocking rates, are shown in Fig. 7 for each management option (Fig. 7). The G5 site (mowing dominated) was excluded from this analysis owing to the very low stocking rate practised at this site, thus a relationship was established of animal intake (not offtake) with NPP over stocking rate ratio. There



Fig. 5. Relationship between methane emissions (multi-year averages of five models), expressed as CH_4-C and CO_2e, and increasing animal stocking rate at all sites, comparing livestock density reduction (top graph) and livestock and N fertilization reduction (bottom graph) as given in Table 3.

was a decreasing trend in animal intake with extensification when animals have access to more biomass per head. This trend is also supported by the relationship between the minimum required amount of biomass per animal and the productivity of the sites under different management options. Using 1.5 LSU ha^{-1} yr^{-1} as an overall reference estimate of potential ecological carrying capacity (e.g. UK Rural Payments Agency, 2003), which is equal to 1.5 adult cattle on 1 ha pasture field, we see that around ~46% of the total biomass produced each year is consumed by animals in the most intensive grazing systems (baseline) of G2 and G3 sites (Fig. 7). With extensification, this ratio is reduced to ~23% at LD_{50} (average of G2 and G3 sites).

Overall, grassland productivity increases with annual precipitation levels, though uncertainties can be large (e.g. seasonal waterlogging spells and heat waves may have negative effects on grassland productivity also with rainfall >1000 mm yr^{-1}), indicating higher sensitivity to animal density reduction (which has some positive effects) than to



Fig. 6. Changes in productivity and energy outputs (g DM m^{-2}, MJ kg^{-1} DM) calculated over multiple years at five sites, for ten altered management scenarios and the baseline (as in Table 3). For each management level, triangles show the multi-model (as in Table 4) mean, black lines show multi-model median. Boxes delimit the 25th and 75th percentiles. Whiskers are 10th and 90th percentiles. Points indicate outliers.



Fig. 7. Relationship between animal intake (Intake) (multi-year averages of eight models, expressed with two units) and the ratio of net primary production (NPP) (multi-year averages of eight models) over animal stocking rate for different livestock densities (as in Table 3).

N fertilization reduction (with even greater negative effects) of $ME_{Offtake}$, Intake, Offtake, AB and BB (Fig. O in Supplementary material).

3.4. CO_2, CH_4 and N_2O intensities

Extensification, mostly through N input reduction, increased the variability of results in the case of methane intensity. Overall, N_2O-N intensity reduces with greater extensification (Fig. P in Supplementary material), mainly driven by N fertilizer reduction, while CO_2-C intensity shows different patterns at each site. In G4, in particular, CO_2-C intensity is >1 when animal density was reduced by 60%. Multi-model median simulations suggest that reducing N fertilization does not affect methane emissions. However, at the G3 site (grazed by heifers), which shows the greatest intensities (Fig. Q in Supplementary material), CH_4-C emissions increase with reducing animal density (with cattle having a larger area to graze, and thus more biomass available for feeding).

For N_2O-N intensity (Fig. R in Supplementary material), model variability increased with reductions in animal density at the G1 and G3 sites (cattle grazing systems). Under sheep grazing (G2, G4, G5 sites), different models did not differ much in their output when reducing animal density. The intensity of C (in the form of CO_2-C and CH_4-C) and N (in the form of N_2O-N) emissions with respect to biomass offtake did not change with extensification (Fig. P in Supplementary material). The plot of CH_4-C intensity values against NPP-stocking rate ratios (Fig. 8) show the extensification at G3 site increased the simulation uncertainty. For sheep grazing systems, methane emissions did not vary greatly with management options. However, CO_2-C and N_2O-N intensities, and their simulation uncertainties, increased with extensification, when animals had more available biomass.

In relation to annual precipitation levels, CO_2-C, CH_4-C and N_2O-N intensities (Fig. R in Supplementary material) showed different patterns for arid and humid conditions. In the case of CO_2-C intensity, C sequestration was moving around its equilibrium at humid conditions, while for drier years it showed different patterns. N fertilization reduction may increase C fixation, with its maximum at 30% less animal density and 60% less N fertilization.

Overall, the carbon sink increased with extensification (baseline: ~ −70, LD_{50}: ~−175 and $LD_{50}N_0$: ~−329 CO_2eq) (Fig. 9), but N_2O and CH_4 emissions decreased. Livestock density reduction showed greater effect on CH_4-CO_2eq reduction (baseline: ~108, LD_{50}: ~84 and $LD_{50}N_0$: ~75 CO_2eq), while N fertilization reduction considerably reduced the N_2O emission from ~160 and ~152 CO_2eq (baseline and LD50, respectively) to ~49 CO_2eq with no N fertilization.

4. Summary and conclusions

This is the first multi-model study to simulate the effect on C and N fluxes of reduced grazing intensity and N fertilizer inputs in multiple grassland systems across the globe. By mobilizing a multi-model approach, it has provided an improved understanding of GHG flux dynamics in pastures. This study confirms that grasslands (which have the advantage of potentially acting as a C and N sink compared to many croplands) can be exploited for GHG mitigation in beef and dairy production, because C and N sequestrations can, under some circumstances, offset GHG emissions.

Simulated C fluxes indicated that there may be some shift towards a C sink (NEE < 0) with decreasing inputs, though it depends on complex, multifaceted processes of C fixation (GPP) and release (R_{ECO}) occurring in the ecosystem. This is especially true for G3 and G4 sites, while grasslands managed with low animal densities may not support C sequestration under arid conditions or in the presence of high organic C exports from mowing. Simulated N outputs showed the considerable effect of N fertilizer reduction on C and N emissions, while changes in animal density only slightly affected the N balance. Both simulated CH_4-C and N emissions (including leaching) were, as expected, highly sensitive to precipitation levels, with higher values being seen under humid conditions (annual precipitation >1000 mm). This indicates the importance of considering climate patterns when determining budgets of C and N under varying management options. With the most intensively grazed systems, ~35–40% of the simulated net primary production was grazed by animals, with this ratio decreasing to ~13% with decreasing stocking rates. The greatest enteric CH_4-C intensities were estimated for intensive grazing systems, while the highest estimates of N_2O-N intensities were found for mown and extensively grazed arid systems. Considering the dynamic behaviour of grassland systems, the amounts of C and N sequestered or released are not the same each year. However, uncertainties in the year-to-year variations are not critical in this context as our focus was on capturing major trends and levels rather than modelling exact annual or seasonal fluxes.

While suggesting the possibility of using models to determine sound mitigation practices, the present study also showed limitations. Our findings are based purely on simulated data and lack evaluation against measured outputs (experimental trials that have appeared in the published literature give us, at a minimum, a hint at what comprehensive assessment of multi-model ensemble would look like). Although the models used in this study are only a subset of the available grassland models, we think that the various model types and variants (and related parameterizations) evaluated here are reasonably representative of current approaches. Another study limitation is that grassland practices other than grazing density and N fertilization were not assessed. For instance, an option that has not been accounted for in this study is increasing the proportion of legume species in the sward which can allow for reduced use of N fertilizer, and has the potential to mitigate GHG emissions (e.g. Lanigan et al., 2013; Fuchs et al., 2018). For an analysis of the mitigation potential of legumes, we refer readers to a parallel study underway as part of the Model4Pastures project (https://www.faccejpi.



Fig. 8. Relationship between CO_2-C, N_2O-N and CH_4-C intensity outputs (multi-year averages of eight, seven and five models, respectively) and the ratio of net primary production (NPP) output and animal stocking rate for different livestock densities (as in Table 3).



Fig. 9. Ensemble modelling of carbon sink (C sink), nitrous oxide emissions (N_2O) and methane emissions (CH_4) in CO_2eq form using multi-site averages of the multi-model median of eight models under grassland mitigation options (as in Table 3).

com/Research-Themes-and-Achievements/GHG-Mitigation/multi-partners-call/Models4Pastures). Other options, such as organic N fertilization, nitrification inhibitors or supplemental feeding, which are common practices in grassland management, have been left out given that state-of-the-art models are not unambiguously sensitive to such management interventions (Brilli et al., 2017). These difficulties, and those associated with model-to-model variability, suggest that some development work would be sensible given the importance of grasslands in supporting the broader GHG emissions reduction agenda. Despite their limitations, biogeochemical models (which evolve with the progress of research) are today a valuable tool for evaluating alternative options for mitigation of GHG emissions through grassland management. It is still rare for results in support to management decisions to be reported by an assessment of uncertainty. Our results show the potential for associating quantification of uncertainties with the results of grassland modelling under alternative management.

Acknowledgement

This work was undertaken by the CN-MIP project of the Joint Programming Initiative 'FACCE' (https://www.faccejpi.com) under the auspices of the Global Research Alliance for Agricultural Greenhouse Gases - Integrative Research Group (http://globalresearchalliance.org/research/integrative). The project, coordinated by the French National Institute for Agricultural Research (INRA) (ANR-13-JFAC-0001), received funding by the 'FACCE' Multi-partner Call on Agricultural Greenhouse Gas Research through its national financing bodies. The input of PS and NF contributes to projects with support from UK-NERC (U-GRASS: NE/M016900/1) and FACCE-JPI via Defra. VS and PN were funded by the New Zealand Government to support the objectives of the Livestock Research Group of the Global Research Alliance on Agricultural Greenhouse Gases. FE acknowledges support through a grant from the French Environment and Energy Management Agency (ADEME, n° 12-60-C0023). KF and LM acknowledge funds received from the Swiss National Science Foundation (40FA40_154245/1 grant agreement) and from the Doc Mobility grant. Exchanges between French and Italian authors were supported by the Galileo programme (CLIMSOC: 39625XE for France, G18-631 for Italy).

We acknowledge Stephanie K. Jones, Kairsty Topp (Scotland's Rural College, EH9 3JG Edinburgh, United Kingdom) and Ute Skiba (Centre for Ecology and Hydrology Edinburgh, United Kingdom), who contributed to collecting the data used in this study for the G4 site. Lianhai Wu (Rothamsted Research, Sustainable Soil and Grassland Systems Department, United Kingdom) and Rich Conant (NREL, Colorado State University, Fort Collins, CO, USA) are also acknowledged for their contribution to the modelling work performed with SPACSYS and DayCent v4.5 2013, respectively.

Appendix A. Supplementary data

Supplementary data to this article can be found online at https://doi.org/10.1016/j.scitotenv.2018.06.020.

References

Abdalla, M., Richards, M., Pogson, M., Smith, J.U., Smith, P., 2017. Estimating the effect of nitrogen fertilizer on the greenhouse gas balance of soils in Wales under current and future climate. Reg. Environ. Chang. 16, 2357–2368.

Abdalla, M., Hastings, A., Chadwick, D.R., Jones, D.L., Evans, C.D., Jones, M.B., Rees, R.M., Smith, P., 2018. Critical review of the impacts of grazing intensity on soil organic carbon storage and other soil quality indicators in extensively managed grasslands. Agric. Ecosyst. Environ. 253, 62–81.

Allard, V., Soussana, J.-F., Falcimagne, R., Berbigier, P., Bonnefond, J.M., Ceschia, E., D'hour, P., Hénault, C., Laville, P., Martin, C., Pinarès-Patino, C., 2007. The role of grazing management for the net biome productivity and greenhouse gas budget (CO_2, N_2O and CH_4) of semi-natural grassland. Agric. Ecosyst. Environ. 12, 47–58.

Ammann, C., Flechard, C.R., Leifeld, J., Neftel, A., Fuhrer, J., 2007. The carbon budget of newly established temperate grassland depends on management intensity. Agric. Ecosyst. Environ. 121, 5–20.

Atkinson, P.W., Fuller, R.J., Vickery, J.A., Conway, G.J., Tallowin, J.R.B., Smith, R.E.N., Haysom, K.A., Ings, T.C., Asteraki, E.J., Brown, V.K., 2005. Influence of agricultural management, sward structure and food resources on grassland field use by birds in lowland England. J. Appl. Ecol. 42, 932–942.

Bannert, A., Bogen, C., Esperschuetz, J., 2012. Anaerobic oxidation of methane in grassland soils used for cattle husbandry. Biogeosciences 9, 3891–3899.

Bell, M.J., Cloy, J.M., Topp, C.F.E., Ball, B.C., Bagnall, A., Rees, R.M., Chadwick, D.R., 2016. Quantifying N_2O emissions from intensive grassland production: the role of synthetic fertiliser type, application rate, timing, and nitrification inhibitors. J. Agric. Sci. 154, 812–827.

Ben Touhami, H., Bellocchi, G., 2015. Bayesian calibration of the Pasture Simulation model (PaSim) to simulate European grasslands under water stress. Eco. Inform. 30, 356–364.

Bondeau, A., Kicklighter, D.W., Kaduk, J., the participants of the Potsdam NdP Model Intercomparison, 1999. Comparing global models of terrestrial net primary productivity (NPP): importance of vegetation structure on seasonal NPP estimates. Glob. Chang. Biol. 5, 35–45.

Bondeau, A., Smith, P.C., Zaehle, S., Schaphoff, S., Lucht, W., Cramer, W., Gerten, D., Lotze-Campen, H., Müller, C., Reichstein, M., Smith, B., 2007. Modelling the role of agriculture for the 20th century global terrestrial carbon balance. Glob. Chang. Biol. 13, 679–706.

Bouwman, A.F., Boumans, L.J.M., Batjes, N.H., 2002. Emissions of N_2O and NO from fertilized fields: summary of available measurement data. Glob. Biogeochem. Cycles 16, 1058.

Brilli, L., Bechini, L., Bindi, M., Carozzi, M., Cavalli, D., Conant, R., Dorich, C.D., Doro, L., Ehrhardt, F., Farina, R., Ferrise, R., Fitton, N., Francaviglia, R., Grace, P., Iocola, I., Klumpp, K., Léonard, J., Martin, R., Massad, R.S., Recous, S., Seddaiu, G., Sharp, J., Smith, P., Smith, W.N., Soussana, J.-F., Bellocchi, G., 2017. Review and analysis of strengths and weaknesses of agro-ecosystem models for simulating C and N fluxes. Sci. Total Environ. 598, 445–470.

Calanca, P., Vuichard, N., Campbell, C., Viovy, N., Cozic, A., Fuhrer, J., Soussana, J.-F., 2007. Simulating the fluxes of CO_2 and N_2O in European grasslands with the Pasture Simulation model (PaSim). Agric. Ecosyst. Environ. 121, 164–174.

Calanca, P., Deléglise, C., Martin, R., Carrère, P., Mosimann, E., 2016. Testing the ability of a simple grassland model to simulate the seasonal effects of drought on herbage growth. Field Crop Res. 187, 12–23.

Campbell, E.E., Paustian, K., 2015. Current developments in soil organic matter modeling and the expansion of model applications: a review. Environ. Res. Lett. 10, 123004.

Cardenas, L.M., Thorman, R., Ashlee, N., Butler, M., Chadwick, D., Chambers, B., Cuttle, S., Donovan, N., Kingston, H., Lane, S., Dhanoa, M.S., Scholefield, D., 2010. Quantifying annual N_2O emission fluxes from grazed grassland under a range of inorganic fertiliser nitrogen inputs. Agric. Ecosyst. Environ. 136, 218–226.

Challinor, A.J., Smith, M.S., Thornton, P., 2013. Use of agro-climate ensembles for quantifying uncertainty and informing adaptation. Agric. For. Meteorol. 170, 2–7.

Chang, J., Viovy, N., Vuichard, N., Ciais, P., Wang, T., Cozic, A., Lardy, R., Graux, A.-I., Klumpp, K., Martin, R., Soussana, J.-F., 2013. Incorporating grassland management in ORCHIDEE: model description and evaluation at eddy-covariance sites in Europe. Geosci. Model Dev. 6, 2165–2181.

Chang, J., Ciais, P., Viovy, N., Vuichard, N., Herrero, M., Havlik, P., Wang, X., Sultan, B., Soussana, J.-F., 2015. Effect of climate change, CO_2 trends, nitrogen addition, and land-cover and management intensity changes on the carbon balance of European grasslands. Glob. Chang. Biol. 22, 338–350.

Chen, D.L., Li, Y., Grace, P., Mosier, A.R., 2008. N_2O emissions from agricultural lands: a synthesis of simulation approaches. Plant Soil 309, 169–189.

Churkina, G., Running, S.W., Schloss, A.L., the participants of the Potsdam NdP Model Intercomparison, 1999. Comparing global models of terrestrial net primary productivity (NPP): the importance of water availability. Glob. Chang. Biol. 5, 46–55.

Ciais, P., Soussana, J.-F., Vuichard, N., Luyssaert, S., Don, A., Janssens, I.A., Piao, S.L., Dechow, R., Lathière, J., Maignan, F., Wattenbach, M., Smith, P., Ammann, C., Freibauer, A., Schulze, E.D., the CARBOEUROPE Synthesis Team, 2010. The greenhouse gas balance of European grasslands. Biogeosci. Discuss. 7, 5997–6050.

Clark, H., Brookes, I., Walcroft, A., 2003. Enteric Methane Emissions From New Zealand Ruminants 1990–2001 Calculated Using an IPCC Tier 2 Approach. Report Prepared for the Ministry of Agriculture and Forestry by AgResearch and Massey University, Ministry of Agriculture and Forestry, Wellington, New Zealand.

Ehrhardt, F., Soussana, J.-F., Bellocchi, G., Grace, P., Mcauliffe, R., Recous, S., Sandor, R., Smith, P., Snow, V., Migliorati, M.D., Basso, B., Bhatia, A., Brilli, L., Doltra, J., Dorich, C.D., Doro, L., Fitton, N., Giacomini, S.J., Grant, B., Harrison, M.T., Jones, S.K., Kirschbaum, M.U., Klumpp, K., Laville, P., Leonard, J., Liebig, M.A., Lieffering, M., Martin, R., Massad, R., Meier, E., Merbold, L., Moore, A.D., Myrgiotis, V., Newton, P., Pattey, E., Rolinski, S., Sharp, J., Smith, W.N., Wu, L., Zhang, Q., 2018. Assessing uncertainties in crop and pasture ensemble model simulations of productivity and N_2O emissions. Glob. Chang. Biol. 24, e603-e616.

Elliott, J., Müller, C., Deryng, D., Chryssanthacopoulos, J., Boote, K.J., Büchner, M., Foster, I., Glotter, M., Heinke, J., Iizumi, T., Izaurralde, R.C., Mueller, N.D., Ray, D.K., Rosenzweig, C., Ruane, A.C., Sheffield, J., 2015. The Global Gridded Crop Model Intercomparison: data and modeling protocols for phase 1 (v1.0). Geosci. Model Dev. 8, 261–277.

FAO, 2014. World reference base for soil resources 2014. World Soil Resources Report. 106 (Rome. 120 pp.).

Folberth, C., Skalský, R., Moltchanova, E., Balkovič, J., Azevedo, L.B., Obersteiner, M., van der Velde, M., 2016. Uncertainty in soil data can outweigh climate impact signals in global crop yield simulations. Nat. Commun. 7, 11872.

Fuchs, K., Hörtnagl, L., Buchmann, N., Eugster, W., Snow, V., Merbold, L., 2018. Management matters: testing a mitigation strategy for nitrous oxide emissions on intensively managed grassland. Biogeosci. Discuss. https://doi.org/10.5194/bg-2018-192.

Gerber, P.J., Mottet, A., Opio, C.I., Falcucci, A., Teillard, F., 2015. Environmental impacts of beef production: review of challenges and perspectives for durability. Meat Sci. 109, 2–12.

Giardina, C.P., Litton, C.M., Crow, S.E., Asner, G.P., 2014. Warming-related increases in soil CO_2 efflux are explained by increased below-ground carbon flux. Nat. Clim. Chang. 4, 822–827.

Gilmanov, T., Soussana, J.-F., Aires, L., Allard, V., Ammann, C., Balzarolo, M., Barcza, Z., Bemhofer, C., Campbell, C.L., Cernusca, A., Cescatti, A., Clifton-Brown, J., Dirks, B.O.M., Dore, S., Eugster, W., Fuhrer, J., Gimeno, C., Gruenwald, T., Haszpra, L., Hensen, A., Ibrom, A., Jacobs, A.F.G., Jones, M.B., Lanigan, G., Laurila, T., Lohila, A., Manca, G., Marcolla, B., Nagy, Z., Pilegaard, K., Pinter, K., Pio, C., Raschi, A., Rogiers, N., Sanz, M.J., Stefani, P., Sutton, M., Tuba, Z., Valentini, R., Williams, M.L., Wohlfahrt, G., 2007. Partitioning of the tower-based net CO_2 exchange in European grasslands into gross primary productivity and ecosystem respiration components using light response functions analysis. Agric. Ecosyst. Environ. 121, 93–120.

Graux, A.I., Lardy, R., Bellocchi, G., Soussana, J.-F., 2012. Global warming potential of French grassland-based dairy livestock systems under climate change. Reg. Environ. Chang. 12, 751–763.

Graux, A.-I., Bellocchi, G., Lardy, R., Soussana, J.-F., 2013. Ensemble modelling of climate change risks and opportunities for managed grasslands in France. Agric. For. Meteorol. 170, 114–131.

Herrero, M., Henderson, B., Havlík, P., Thornton, P.K., Conant, R.T., Smith, P., Wirsenius, S., Hristov, A.N., Gerber, P., Gill, M., Butterbach-Bahl, K., Valin, H., Garnett, T., Stehfest, E.,

2016. Greenhouse gas mitigation potentials in the livestock sector. Nat. Clim. Chang. 6, 452–461.

Holzworth, D.P., Huth, N.I., Devoil, P.G., Zurcher, E.J., Herrmann, N.I., Mclean, G., Chenu, K., van Oosterom, E.J., Snow, V., Murphy, C., Moore, A.D., Brown, H., Whish, J.P.M., Verrall, S., Fainges, J., Bell, L.W., Peake, A.S., Poulton, P.L., Hochman, Z., Thorburn, P.J., Gaydon, D.S., Dalgliesh, N.P., Rodriguez, D., Cox, H., Chapman, S., Doherty, A., Teixeira, E., Sharp, J., Cichota, R., Vogeler, I., Li, F.Y., Wang, E., Hammer, G.L., Robertson, M.J., Dimes, J.P., Whitbread, A.M., Hunt, J., van Rees, H., McClelland, T., Carberry, P.S., Hargreaves, J.N.G., Macleod, N., McDonald, C., Harsdorf, J., Wedgwood, S., Keating, B.A., 2014. APSIM - evolution towards a new generation of agricultural systems simulation. Environ. Model. Softw. 62, 327–350.

Hörtnagl, L., Barthel, M., Buchmann, N., Eugster, W., Butterbach-Bahl, K., Díaz-Pinés, E., Zeeman, M., Klumpp, K., Kiese, R., Bahn, M., Hammerle, A., Lu, H., Ladreiter-Knauss, T., Burri, S., Merbold, L., 2018. Greenhouse gas fluxes over managed grasslands in Central Europe. Glob. Chang. Biol. https://doi.org/10.1111/gcb.14079.

Huntzinger, D.N., Post, W.M., Wei, Y., Michalak, A.M., West, T.O., Jacobson, A.R., Baker, I.T., Chen, J.M., Davis, K.J., Hayes, D.J., Hoffman, F.M., Jain, A.K., Liu, S., Mcguire, A.D., Neilson, R.P., Potter, C., Poulter, B., Price, D., Raczka, B.M., Tian, H.Q., Thornton, P., Tomelleri, E., Viovy, N., Xiao, J., Yuan, W., Zeng, N., Zhao, M., Cook, R., 2012. North American Carbon Program (NACP) regional interim synthesis: terrestrial biospheric model intercomparison. Ecol. Model. 232, 144–157.

Huth, N.I., Bristow, K.L., Verburg, K., 2012. SWIM3: Model use, calibration, and validation. Trans. ASABE 55, 1303–1313.

Imer, D., Merbold, L., Eugster, W., Buchmann, N., 2013. Temporal and spatial variations of soil CO_2, CH_4 and N_2O fluxes at three differently managed grasslands. Biogeosciences 10, 5931–5945.

IPCC, 2006. N_2O emissions from managed soils, and CO_2 emissions from lime and urea application. In: De Klein, C., Novoa, R.S.A., Ogle, S., Smith, K.A., Rochette, P., Wirth, T.C., McConkey, B.G., Mosier, A., Rypdal, K. (Eds.), Guidelines for National Greenhouse Gas Inventories. vol. 4. IGES, Japan:pp. 1–54 (Available at https://www.ipcc-nggip.iges.or.jp/public/2006gl/pdf/4_Volume4/V4_11_Ch11_N2O&CO2.pdf (access on 18 May 2018)).

IPCC, 2014. In: Core Writing Team, Pachauri, R.K., Meyer, L.A. (Eds.), Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. IPCC, Geneva, Switzerland (151 pp.).

Jones, J.W., Antle, J.M., Basso, B.O., et al., 2017. Brief history of agricultural systems modelling. Agric. Syst. 155, 240–254.

Jones, S.K., Helfter, C., Anderson, M., Coyle, M., Campbell, C., Famulari, D., Di Marco, C., van Dijk, N., Tang, Y.S., Topp, C.F.E., Kiese, R., Kindler, R., Siemens, J., Schrumpf, M., Kaiser, K., Nemitz, E., Levy, P.E., Rees, R.M., Sutton, M.A., Skiba, U.M., 2017. The nitrogen, carbon and greenhouse gas budget of a grazed, cut and fertilised temperate grassland. Biogeosciences 14, 2069–2088.

Klumpp, K., Tallec, T., Guix, N., Soussana, J.-F., 2011. Long-term impacts of agricultural practices and climatic variability on carbon storage in a permanent pasture. Glob. Chang. Biol. 17, 3534–3545.

Koncz, P., Pintér, K., Balogh, J., Papp, M., Hidy, D., Csintalan, Zs., Molnár, E., Szaniszló, A., Kampfi, Gy, Horváth, L., Nagy, Z., 2017. Extensive grazing in contrast to mowing is climate-friendly based on the farm-scale greenhouse gas balance. Agric. Ecosyst. Environ. 240, 121–134.

Kottek, M., Grieser, J., Beck, C., Rudolf, B., Rubel, F., 2006. World map of the Köppen-Geiger climate classification updated. Meteorol. Z. 15, 259–263.

Lanigan, G., Finn, J., Lüscher, A., 2013. Mitigation & adaptation options: grass-legume mixtures. AnimalChange E-Newsletter http://www.animalchange.eu/pdf/newsletter/nov2013.pdf.

Li, F.Y., Snow, V.O., Holzworth, D.P., 2011. Modelling the seasonal and geographical pattern of pasture production in New Zealand. N. Z. J. Agric. Res. 54, 331–352.

Liebig, M.A., Gross, J.R., Kronberg, S.L., Hanson, J.D., Frank, A.B., Phillips, R.L., 2006. Soil response to long-term grazing in the northern Great Plains of North America. Agric. Ecosyst. Environ. 115, 270–276.

Liebig, M.A., Gross, J.R., Kronberg, S.L., Phillips, R.L., Hanson, J.D., 2010. Grazing management contributions to net global warming potential: a long-term evaluation in the Northern Great Plains. J. Environ. Qual. 39, 799–809.

Liebig, M.A., Kronberg, S.L., Hendrickson, J.R., Dong, X., Gross, J.R., 2013. Carbon dioxide efflux from long-term grazing management systems in a semiarid region. Agric. Ecosyst. Environ. 164, 137–144.

Lü, X.-T., Dijkstra, F.A., Kong, D.-L., Wang, Z.-W., Han, X.-G., 2014. Plant nitrogen uptake drives responses of productivity to nitrogen and water addition in a grassland. Sci. Rep. 4, 4817. https://doi.org/10.1038/srep04817.

Ma, S., Lardy, R., Graux, A.-I., Ben Touhami, H., Klumpp, K., Martin, R., Bellocchi, G., 2015. Regional-scale analysis of carbon and water cycles on managed grassland systems. Environ. Model. Softw. 72, 356–371.

Merbold, L., Eugster, W., Stieger, J., Zahniser, M., Nelson, D., Buchmann, N., 2014. Greenhouse gas budget (CO_2, CH_4 and N_2O) of intensively managed grassland following restoration. Glob. Chang. Biol. 20, 1913–1928.

Newton, P.C.D., Lieffering, M., Bowatte, W.M.S.D., Brock, S.C., Hunt, C.L., Theobald, P.W., Ross, D.J., 2010. The rate of progression and stability of progressive nitrogen limitation at elevated atmospheric CO_2 in a grazed grassland over 11 years of Free Air CO_2 enrichment. Plant Soil 336, 433–441.

Newton, P.C.D., Lieffering, M., Parsons, A.J., Brock, S.C., Theobald, P.W., Hunt, C.L., Theobald, P.W., Ross, D.J., 2014. Selective grazing modifies previously anticipated responses of plant community composition to elevated CO_2 in a temperate grassland. Glob. Chang. Biol. 20, 158–169.

Oates, L.G., Jackson, R.D., 2014. Livestock management strategy affects net ecosystem carbon balance of subhumid pasture. Rangel. Ecol. Manag. 67, 19–29.

Parton, W.J., Hartman, M., Ojima, D., Schimel, D., 1998. DAYCENT and its land surface submodel: description and testing. Glob. Planet. Chang. 19, 35–48.

Polley, H.W., Emmerich, W., Bradford, J.A., Sims, P.L., Johnson, D.A., Saliendra, N.Z., Svejcar, T., Angell, R., Frank, A.B., Phillips, R.L., Snyder, K.A., Morgan, J.A., Sanabria, J., Mielnick, P.C., Dugas, W.A., 2010. Precipitation regulates the response of net ecosystem CO_2 exchange to environmental variation on United States rangelands. Rangel. Ecol. Manag. 63, 176–186.

Probert, M.E., Dimes, J.P., Keating, B.A., Dalal, R.C., Strong, W.M., 1998. APSIM's water and nitrogen modules and simulation of the dynamics of water and nitrogen in fallow systems. Agric. Syst. 56, 1–28.

R Core Team, 2016. A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria http://www.R-project.org.

Rees, R.M., Augustin, J., Alberti, G., Ball, B.C., Boeckx, P., Cantarel, A., Castaldi, S., Chirinda, N., Chojnicki, B., Giebels, M., Gordon, H., Grosz, B., Horvath, L., Juszczak, R., Klemedtsson, A.K., Klemedtsson, L., Medinets, S., Machon, A., Mapanda, F., Nyamangara, J., Olesen, J.E., Reay, D.S., Sanchez, L., Cobena, A.S., Smith, K.A., Sowerby, A., Sommer, M., Soussana, J.-F., Stenberg, M., Topp, C.F.E., Van Cleemput, O., Vallejo, A., Watson, C.A., Wuta, M., 2013. Nitrous oxide emissions from European agriculture - an analysis of variability and drivers of emissions from field experiments. Biogeosciences 10, 2671–2682.

Riedo, M., Grub, A., Rosset, M., Fuhrer, J., 1998. A Pasture Simulation model for dry matter production, and fluxes of carbon, nitrogen, water and energy. Ecol. Model. 105, 141–183.

Rolinski, S., Müller, C., Heinke, J., Weindl, I., Biewald, A., Bodirsky, B.L., Bondeau, A., Boons-Prins, E.R., Bouwman, A.F., Leffelaar, P.A., te Roller, J.A., Schaphoff, S., Thonicke, K., 2018. Modeling vegetation and carbon dynamics of managed grasslands at the global scale with LPJmL 3.6. Geosci. Model Dev. 11, 429–451.

Rosenzweig, C., Elliott, J., Deryng, D., Ruane, A.C., Müller, C., Arneth, A., Boote, K.J., Folberth, C., Glotter, M., Khabarov, N., Neumann, K., Piontek, F., Pugh, T.A., Schmid, E., Stehfest, E., Yang, H., Jones, J.W., 2014. Assessing agricultural risks of climate change in the 21st century in a global gridded crop model intercomparison. Proc. Natl. Acad. Sci. 111, 3268–3273.

Ruane, A.C., Goldberg, R., Chryssanthacopoulos, J., 2015. Climate forcing datasets for agricultural modeling: merged products for gap-filling and historical climate series estimation. Agric. For. Meteorol. 200, 233–248.

Sándor, R., Barcza, Z., Hidy, D., Lellei-Kovács, E., Ma, S., Bellocchi, G., 2016. Modelling of grassland fluxes in Europe: evaluation of two biogeochemical models. Agric. Ecosyst. Environ. 215, 1–19.

Sándor, R., Barcza, Z., Acutis, M., Doro, L., Hidy, D., Köchy, M., Minet, J., Lellei-Kovács, E., Ma, S., Perego, A., Rolinski, S., Ruget, F., Sanna, M., Seddaiu, G., Wu, L., Bellocchi, G., 2017. Multi-model simulation of soil temperature, soil water content and biomass in Euro-Mediterranean grasslands: uncertainties and ensemble performance. Eur. J. Agron. 88, 22–40.

Sándor, R., Picon-Cochard, C., Martin, R., Louault, F., Klumpp, K., Borras, D., Bellocchi, G., 2018. Plant acclimation to temperature: developments in the Pasture Simulation model. Field Crop Res. https://doi.org/10.1016/j.fcr.2017.05.030.

Schulze, E.D., Luyssaert, S., Ciais, P., Freibauer, A., Janssens, I.A., Soussana, J.-F., Smith, P., Grace, J., Levin, I., Thiruchittampalam, B., Heimann, M., Dolman, A.J., Valentini, R., Bousquet, P., Peylin, P., Peters, W., Roedenbeck, C., Etiope, G., Vuichard, N., Wattenbach, M., Nabuurs, G.J., Poussi, Z., Nieschulze, J., Gash, J.H., CarboEurope Team, 2009. Importance of methane and nitrous oxide for Europe's terrestrial greenhouse-gas balance. Nat. Geosci. 2, 842–850.

Seijan, V., Lal, R., Lakritz, J., Ezeji, T., 2011. Measurement and prediction of enteric methane emission. Int. J. Biometeorol. 55, 1–16.

Senapati, N., Chabbi, A., Gastal, F., Smith, P., Mascher, N., Loubet, B., 2014. Net carbon storage measured in a mowed and grazed temperate sown grassland shows potential for carbon sequestration under grazed systems. Carbon Manage. 5, 131–144.

Senapati, N., Jansson, P.E., Smith, P., Chabbi, A., 2016. Modelling heat, water and carbon fluxes in mown grassland under multi-objective and multi-criteria constraints. Environ. Model. Softw. 80, 201–224.

Sitch, S., Smith, B., Prentice, I.C., Arneth, A., Bondeau, A., Cramer, W., Kaplan, J.O., Levis, S., Lucht, W., Sykes, M.T., Thonicke, K., Venevsky, S., 2003. Evaluation of ecosystem dynamics, plant geography and terrestrial carbon cycling in the LPJ dynamic global vegetation model. Glob. Chang. Biol. 9, 161–185.

Skiba, U., Jones, S.K., Drewer, J., Helfter, C., Anderson, M., Dinsmore, K., McKenzie, R., Nemitz, E., Sutton, M.A., 2013. Comparison of soil greenhouse gas fluxes from extensive and intensive grazing in a temperate maritime climate. Biogeosciences 10, 1231–1241.

Smith, K.A., Thomson, P.E., Clayton, H., McTaggart, I.P., Conen, F., 1998. Effects of temperature, water content and nitrogen fertilisation on emissions of nitrous oxide by soils. Atmos. Environ. 32, 3301–3309.

Smith, P., Martino, D., Cai, Z., Gwary, D., Janzen, H.H., Kumar, P., McCarl, B., Ogle, S., O'Mara, F., Rice, C., Scholes, R.J., Sirotenko, O., Howden, M., McAllister, T., Pan, G., Romanenkov, V., Schneider, U., Towprayoon, S., Wattenbach, M., Smith, J.U., 2008. Greenhouse gas mitigation in agriculture. Philos. Trans. R. Soc., B 363, 789–813.

Snow, V., Rotz, C.A., Moore, A.D., Martin-Clouaire, R., Johnson, I.R., Hutchings, N.J., Eckard, R.J., 2014. The challenges - and some solutions - to process-based modelling of grazed agricultural systems. Environ. Model. Softw. 62, 420–436.

Soussana, J.-F., Pilegaard, K., Ambus, P., Berbigier, P., Ceschia, E., Clifton-Brown, J., Czobel, S., de Groot, T., Fuhrer, J., Horvath, L., Hensen, A., Jones, M., Kasper, G., Martin, C., Milford, C., Nagy, Z., Neftel, A., Raschi, A., Rees, R.M., Skiba, U., Stefani, P., Saletes, S., Sutton, M.A., Tuba, Z., Weidinger, T., 2004. Annual greenhouse gas balance of European grasslands - first results from the GreenGrass project. International Conference Greenhouse Gas Emissions From Agriculture-mitigation Options and Strategies, 10–12 February, Leipzig, Germany, pp. 25–30.

Soussana, J.-F., Allard, V., Pilegaard, K., Ambus, P., Amman, C., Campbell, C., Ceschia, E., Clifton-Brown, J., Czobel, S., Domingues, R., Flechard, C., Fuhrer, J., Hensen, A., Horvath, L., Jones, M., Kasper, G., Martin, C., Nagy, Z., Neftel, A., Raschi, A., Baronti,

S., Rees, R., Skiba, U., Stefani, P., Manca, G., Sutton, M., Tuba, Z., Valentini, R., 2007. Full accounting of the greenhouse gas (CO_2, N_2O, CH_4) budget of nine European grassland sites. Agric. Ecosyst. Environ. 121, 121–134.

Soussana, J.-F., Tallec, T., Blanfort, V., 2010. Mitigating the greenhouse gas balance of ruminant production systems through carbon sequestration in grasslands. Animal 4, 334–350.

Soussana, J.-F., Barioni, L.G., Ben-Ari, T., Conant, R., Gerber, P., Havlik, P., Ickowicz, A., Howden, M., 2013. Managing grassland systems in a changing climate: the search for practical solutions. 22nd International Grasslands Congress, 15–19 September, Sydney, Australia (18 pp.).

Soussana, J.-F., Klumpp, K., Ehrhardt, F., 2014. The role of grassland in mitigating climate change. Proceedings of EGF at 50: The Future of European Grasslands. Grassland Science in Europe 19, pp. 75–87.

Tubiello, F., Salvatore, M., Cóndor Golec, R., Ferrara, A., Rossi, S., Biancalani, R., Federici, S., Jacobs, H., Flammini, A., 2014. Agriculture, Forestry and Other Land Use Emissions by Sources and Removals by Sinks 1999–2011 Analysis, Working Paper Series ESS/14-02. FAO Statistical Division, Rome, Italy Available at. http://www.fao.org/docrep/019/i3671e/i3671e.pdf, Accessed date: 15 May 2018.

UK Rural Payments Agency, 2003. Cattle scheme literature: Notes for guidance 2003. The Rural Payments Agency of the UK Department for Environment, Food and Rural Affairs through. https://www.gov.uk/government/organisations/rural-payments-agency.

Valin, H., Havlik, P., Mosnier, A., Herrero, M., Schmid, E., Obersteiner, M., 2013. Agricultural productivity and greenhouse gas emissions: trade-offs or synergies between mitigation and food security? Environ. Res. Lett. 8, 035019.

Van Groenigen, J.W., Velthof, G.L., Oenema, O., Van Groenigen, K.J., Van Kessel, C., 2010. Towards an agronomic assessment of N_2O emissions: a case study for arable crops. Eur. J. Soil Sci. 61, 903–913.

Venterea, R.T., Bijesh, M., Dolan, M.S., 2011. Fertilizer source and tillage effects on yield-scaled nitrous oxide emissions in a corn cropping system. J. Environ. Qual. 40, 1521–1531.

Warszawski, L., Frieler, K., Huber, V., Piontek, F., Serdeczny, O., Schewe, J., 2014. The Inter-Sectoral Impact Model Intercomparison Project (ISI-MIP): project framework. Proc. Natl. Acad. Sci. U. S. A. 111, 3228–3232.

West, T.O., Marland, G., 2002. A synthesis of carbon sequestration, carbon emissions, and net carbon flux in agriculture: comparing tillage practices in the United States. Agric. Ecosyst. Environ. 91, 217–232.

White, T.A., Johnson, I.R., Snow, V.O., 2008. Comparison of outputs of a biophysical simulation model for pasture growth and composition with measured data under dryland and irrigated conditions in New Zealand. Grass Forage Sci. 63, 339–349.

Wu, L., Mcgechan, M.B., Mcroberts, N., Baddeley, J.A., Watson, C.A., 2007. SPACSYS: integration of a 3D root architecture component to carbon, nitrogen and water cycling-model description. Ecol. Model. 200, 343–359.

Wu, L., Rees, R.M., Tarsitano, D., Zhang, X., Jones, S.K., Whitmore, A.P., 2015. Simulation of nitrous oxide emissions at field scale using the SPACSYS model. Sci. Total Environ. 530–531, 76–86.

Yuan, Z.Y., Li, L.H., Han, X.G., Chen, S.P., Wang, Z.W., Chen, Q.S., Bai, W.M., 2006. Nitrogen response efficiency increased monotonically with decreasing soil resource availability: a case study from a semiarid grassland in northern China. Oecologia 148, 564–572.