



**HAL**  
open science

## The interplay between total mercury, methylmercury and dissolved organic matter in fluvial systems: A latitudinal study across Europe

Andrea Garcia Bravo, Dolly Kothawala, Katrin Attermeyer, Emmanuel Tessier, Pascal Bodmer, José L.J. Ledesma, Joachim Audet, Joan Pere Casas-Ruiz, Nuria Catalan, Sophie Cauvy-Fraunié, et al.

### ► To cite this version:

Andrea Garcia Bravo, Dolly Kothawala, Katrin Attermeyer, Emmanuel Tessier, Pascal Bodmer, et al. The interplay between total mercury, methylmercury and dissolved organic matter in fluvial systems: A latitudinal study across Europe. *Water Research*, 2018, 144, pp.172-182. 10.1016/j.watres.2018.06.064 . hal-01982331


**HAL Id: hal-01982331**

**<https://hal.science/hal-01982331>**

Submitted on 3 Nov 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



**Manchester  
Metropolitan  
University**

---

Bravo, AG and Kothawala, DN and Attermeyer, K and Tessier, E and Bodmer, P and Ledesma, JLJ and Audet, J and Casas-Ruiz, JP and Catalán, N and Cauvy-Fraunié, S and Colls, M and Deininger, A and Evtimova, VV and Fonvielle, JA and Fuß, T and Gilbert, P and Herrero Ortega, S and Liu, L and Mendoza-Lera, C and Monteiro, J and Mor, JR and Nagler, M and Niedrist, GH and Nydahl, AC and Pastor, A and Pegg, J and Gutmann Roberts, C and Pilotto, F and Portela, AP and González-Quijano, CR and Romero, F and Rulík, M and Amouroux, D (2018) The interplay between total mercury, methylmercury and dissolved organic matter in fluvial systems: A latitudinal study across Europe. *Water Research*, 144. pp. 172-182. ISSN 0043-1354 (In Press)

---

**Downloaded from:** <http://e-space.mmu.ac.uk/621418/>

**Version:** Accepted Version

**Publisher:** Elsevier

**DOI:** <https://doi.org/10.1016/j.watres.2018.06.064>

Please cite the published version

<https://e-space.mmu.ac.uk>

# The interplay between total mercury, methylmercury and dissolved organic matter in fluvial systems: A latitudinal study across Europe

Andrea G. Bravo<sup>1,\*</sup>, Dolly N. Kothawala<sup>2,+</sup>, Katrin Attermeyer<sup>2</sup>, Emmanuel Tessier<sup>3</sup>, Pascal Bodmer<sup>4,5</sup>, José L. J. Ledesma<sup>6</sup>, Joachim Audet<sup>6</sup>, Joan Pere Casas-Ruiz<sup>7</sup>, Núria Catalán<sup>7</sup>, Sophie Cauvy-Fraunié<sup>8</sup>, Miriam Colls<sup>7</sup>, Anne Deininger<sup>9</sup>, Vesela V. Evtimova<sup>10</sup>, Jérémy A. Fonvielle<sup>11</sup>, Thomas Fuß<sup>12,13</sup>, Peter Gilbert<sup>14</sup>, Sonia Herrero Ortega<sup>11</sup>, Liu Liu<sup>5</sup>, Clara Mendoza-Lera<sup>8</sup>, Juliana Monteiro<sup>15</sup>, Jordi-René Mor<sup>7,16</sup>, Magdalena Nagler<sup>17</sup>, Georg H. Niedrist<sup>18</sup>, Anna C. Nydahl<sup>2</sup>, Ada Pastor<sup>7</sup>, Josephine Pegg<sup>19,20</sup>, Catherine Gutmann Roberts<sup>19</sup>, Francesca Pilotto<sup>9</sup>, Ana Paula Portela<sup>15</sup>, Clara Romero González-Quijano<sup>12</sup>, Ferran Romero<sup>7</sup>, Martin Rulík<sup>21</sup>, David Amouroux<sup>3,\*</sup>

<sup>1</sup> Department of Environmental Chemistry, Institute of Environmental Assessment and Water Research (IDAEA), Spanish National Research Council (CSIC), Barcelona, Spain

<sup>2</sup> Limnology/Department of Ecology and Genetics, Uppsala University, Uppsala, Sweden

<sup>3</sup> CNRS/ UNIV PAU & PAYS ADOUR, Institut des Sciences Analytiques et de Physico-Chimie pour l'Environnement et les Matériaux, UMR5254, MIRA, Pau, France

<sup>4</sup> Institute for Environmental Sciences, University of Koblenz-Landau, Landau, Germany

<sup>5</sup> Chemical Analytics and Biogeochemistry, Leibniz-Institute of Freshwater Ecology and Inland Fisheries, Berlin, Germany

<sup>6</sup> Department of Aquatic Sciences and Assessment, Swedish University of Agricultural Sciences, Uppsala, Sweden

<sup>7</sup> Catalan Institute for Water Research (ICRA), Girona, Spain

<sup>8</sup> IRSTEA, UR RIVERLY, 69616 Villeurbanne Cedex, France

<sup>9</sup> Department of Ecology and Environmental Science, Umeå University, Umeå, Sweden

<sup>10</sup> Department of Aquatic Ecosystems, Institute of Biodiversity and Ecosystem Research, Bulgarian Academy of Sciences, Sofia, Bulgaria

<sup>11</sup> Experimental Limnology, Leibniz-Institute of Freshwater Ecology and Inland Fisheries (IGB), Stechlin, Germany

<sup>12</sup> Ecohydrology, Leibniz-Institute of Freshwater Ecology and Inland Fisheries (IGB), Berlin, Germany

<sup>13</sup> WasserCluster Biological Station Lunz, Lunz am See, Austria

<sup>14</sup> The Environmental Research Institute, University of Highlands and Islands, Thurso, Scotland, UK,

<sup>15</sup> Research Centre in Biodiversity and Genetic Resources (CIBIO), University of Porto, Porto, Portugal

<sup>16</sup> Department of Evolutionary Biology, Ecology and Environmental Sciences, Faculty of Biology, University of Barcelona (UB), Barcelona, Spain

<sup>17</sup> Microbial Resource Management, Institute of Microbiology, University of Innsbruck, Innsbruck, Austria

<sup>18</sup> River and Conservation Research, Institute of Ecology, University of Innsbruck, Innsbruck, Austria

<sup>19</sup> Department of Life and Environmental Sciences, Bournemouth University, UK

<sup>20</sup> University Centre Sparsholt, Winchester, UK

<sup>21</sup> Department of Ecology and Environmental Sciences, Palacky University in Olomouc, Olomouc, Czech Republic

+ These authors contributed equally to this work.

Corresponding authors: [Andrea G Bravo \(jandriugarcia@gmail.com\)](mailto:jandriugarcia@gmail.com) & [David Amouroux \(david.amouroux@univ-pau.fr\)](mailto:david.amouroux@univ-pau.fr)

**Keywords: Mercury, methylmercury, streams, rivers, freshwaters, organic matter, EuroRun, fluorescence**

**Abstract**

54 Large-scale studies are needed to identify the drivers of total mercury (THg) and  
55 monomethyl-mercury (MeHg) concentrations in aquatic ecosystems. Studies attempting to  
56 link dissolved organic matter (DOM) to levels of THg or MeHg are few and geographically  
57 constrained. Additionally, stream and river systems have been understudied as compared to  
58 lakes. Hence, the aim of the study was to examine the influence of DOM concentration and  
59 composition, morphological descriptors, land uses and water chemistry on THg and MeHg  
60 concentrations and the percentage of THg as MeHg (%MeHg) in 29 streams across Europe  
61 spanning from 41°N to 64 °N. THg concentrations (0.06 – 2.78 ng L<sup>-1</sup>) were highest in  
62 streams characterized by DOM with a high terrestrial soil signature and low nutrient content.  
63 MeHg concentrations (78 – 160 pg L<sup>-1</sup>) varied non-systematically across systems.  
64 Relationships between DOM bulk characteristics and THg and MeHg suggest that while soil  
65 derived DOM inputs control THg concentrations, autochthonous DOM (aquatically produced)  
66 and the availability of electron acceptors for Hg methylating microorganisms (e.g. sulfate)  
67 drive %MeHg and potentially MeHg concentration. Overall, these results highlight the large  
68 spatial variability in THg and MeHg concentrations at European scale, and underscore the  
69 importance of DOM composition on Hg cycling in fluvial systems.

70

## 71 **1. INTRODUCTION**

72 Mercury (Hg) is a hazardous substance with the potential to produce significant  
73 adverse neurological and other health effects in severely exposed humans, with particular  
74 concerns on unborn children and infants (Mason et al., 2012). Anthropogenic emissions and  
75 long-range transport of atmospheric-Hg have led to Hg concentrations above background  
76 levels in surface reservoirs (atmosphere, oceans or terrestrial) on a global scale (Amos et al.,  
77 2013). Aquatic food webs are particularly vulnerable to increases in Hg, since its methylated  
78 form (monomethyl-mercury, MeHg) accumulates in organisms and biomagnifies in aquatic  
79 food webs (Morel et al., 1998), even at low concentrations. As a consequence, millions of  
80 people are readily exposed to harmful concentrations of MeHg via fish consumption (Mason  
81 et al., 2012).

82 In order to prevent human exposure to Hg, a global legally binding agreement, the  
83 Minamata Convention, was signed by 128 countries and entered into force in September  
84 2017. Prior to the Minamata Convention, the European Water Framework Directive (WFD,  
85 2008) classified Hg as one of 30 priority hazardous substances and set the maximum  
86 allowable concentrations of environmental quality standards to 70 ng L<sup>-1</sup>. However, in aquatic  
87 systems, without any point anthropogenic or geological sources, total-Hg (THg)  
88 concentration typically ranges from less than 0.05 to 12 ng L<sup>-1</sup> (Dennis et al., 2005; Driscoll et  
89 al., 2007; Wiener et al., 2003). Because the maximum allowable concentrations of  
90 environmental quality standards were much higher than naturally relevant concentration  
91 levels of THg, national monitoring programs did not have the incentive to detect THg at  
92 ambient concentrations. There is even less information about MeHg background levels,  
93 which generally are between one and two orders of magnitude lower as compared to THg  
94 levels (Driscoll et al., 2007). In this context, streams and rivers are of special concern  
95 because very few studies have reported THg and MeHg concentrations at a wide  
96 geographical scale (Balogh et al., 2002; Brigham et al., 2009; Eklöf et al., 2015; Lescord et  
97 al., 2018; Marvin-DiPasquale et al., 2009; Tsui et al., 2009). While rivers and streams have,  
98 however, been hypothesized to be important contributors to the global biogeochemical Hg  
99 budget (Amos et al., 2014; Kocman et al., 2017; Schartup et al., 2015a; Sunderland and  
100 Mason, 2007), quantitative information on Hg levels is largely lacking, especially in the  
101 temperate zone. Filling this gap in knowledge is an important prerequisite in the ongoing  
102 quest to reduce uncertainties associated with the global biogeochemical cycling of Hg and for  
103 the development of remediation and risk management strategies within the context of the  
104 Minamata Convention.

105 Freshwaters receive Hg from a variety of sources including atmospheric deposition  
106 (Fitzgerald et al., 1998), release from industrial discharge (Bravo et al., 2015, 2014), and  
107 lateral export from terrestrial sources (e.g. soils and wetlands) (Brigham et al., 2009; Eklöf et  
108 al., 2012). Diffuse inputs from land management activities and remobilization of Hg  
109 previously accumulated in terrestrial ecosystems can be an important source comparable in

110 magnitude to direct Hg anthropogenic inputs (e.g. industrial releases or mining activities) to  
111 surface waters (Kocman et al., 2017). Long-term accumulation of Hg in terrestrial systems  
112 has been explained by the strong binding between atmospherically deposited Hg and sulfur  
113 sites within organic matter (OM) (Skylberg et al., 2006; Xia et al., 1999). In boreal  
114 ecosystems, characterized by OM-rich soils, the strong correlation between THg and  
115 dissolved organic carbon (DOC) found in several studies (Brigham et al., 2009; Driscoll et al.,  
116 1995; Eklöf et al., 2012; Tsui et al., 2009) denotes that the mobilization of OM is an important  
117 vector for Hg transport from catchment soils to surface waters (Eklöf et al., 2012; Grigal,  
118 2002). Besides stream water DOC, the proportion of wetland area within the catchment and  
119 the discharge have been used to predict THg concentrations in surface waters (Brigham et  
120 al., 2009; Eklöf et al., 2015; Tsui et al., 2009). In contrast, the relationship between DOC and  
121 MeHg remains poorly understood (Brigham et al., 2009; Tsui et al., 2009). The variation of  
122 MeHg concentrations in aquatic ecosystems is indeed more complex, as MeHg is potentially  
123 derived from numerous sources, not only terrestrial sources. In boreal streams, the wetland  
124 proportion of the catchment has also been linked to MeHg freshwater concentrations  
125 (Brigham et al., 2009). Wetlands, characterized by oxygen-depletion and low to intermediate  
126 nutrient levels, are known to be important sites for MeHg production and a relevant source of  
127 MeHg to inland waters (Schaefer et al., 2014; Tjerngren et al., 2012). Besides the  
128 contribution of MeHg produced in wetlands to stream waters, in-stream MeHg formation  
129 might also be an additional key source of MeHg (Balogh et al., 2002; Eklöf et al., 2015; Tsui  
130 et al., 2009), especially in catchments lacking extensive wetland coverage. The methylation  
131 of inorganic divalent Hg (IHg) to MeHg is primarily mediated by anaerobic micro-organisms  
132 (Gilmour et al., 2013) and is controlled by temperature, availability of electron acceptors (e.g.  
133  $\text{SO}_4^{2-}$  and  $\text{Fe}^{3+}$ ) (Bravo et al., 2015) and the molecular composition of OM, which influences  
134 IHg-speciation (Graham et al., 2013; Jonsson et al., 2014), IHg-availability (Chiasson-Gould  
135 et al., 2014; Mazrui et al., 2016; Schaefer and Morel, 2009; Schartup et al., 2015b) and  
136 microbial activity (Bravo et al., 2017). Specifically, IHg methylation processes are enhanced  
137 by the presence of algal OM (Bravo et al., 2017; Schartup et al., 2013) and fresh humic OM

138 leached from organic rich soils (Herrero Ortega et al., 2018). Indeed, elevated MeHg  
139 concentrations have been linked to periods of high algal production (Gascon Diez et al.,  
140 2016; Soerensen et al., 2017) and leaf fall (Balogh et al., 2002). However most studies  
141 investigating the role of OM as a transport vector for Hg and/or a key factor controlling MeHg  
142 formation have been carried out in the boreal biome. Consequently, there is still a high  
143 amount of uncertainty in identifying factors governing THg and MeHg levels in stream waters,  
144 particularly at a broader geographical scale.

145         The molecular composition of dissolved organic matter (DOM) is variable and can be  
146 an important factor in predicting THg and potentially, MeHg (Bravo et al., 2017; Herrero  
147 Ortega et al., 2018; Lescord et al., 2018; Noh et al., 2017; Schartup et al., 2015b, 2013). A  
148 common fingerprinting approach used to characterize the bulk DOM composition in inland  
149 waters is fluorescence spectroscopy (Fellman et al., 2010). The increasingly widespread use  
150 of fluorescence can be attributed to its wide accessibility, ease of use, and low costs. Three  
151 commonly used indexes (fluorescence index (McKnight et al., 2001), freshness index  
152 (Parlanti et al., 2000) and humification index (Zsolnay et al., 1999)) are particularly useful for  
153 distinguishing between DOM derived from different sources (McKnight et al., 2001), and  
154 degrees of freshness and humification (Zsolnay et al., 1999). Likewise, commonly known re-  
155 occurring peaks distributed across the fluorescence excitation-emission matrix (EEM) can  
156 provide insights into sample specific DOM characteristics. Fluorescence has indeed  
157 demonstrated the potential to reveal key DOM molecular characteristics associated with THg  
158 and MeHg concentrations (Herrero Ortega et al., 2018; Lescord et al., 2018; Noh et al., 2017;  
159 Schartup et al., 2015b).

160         The aim of this study was to identify the relationships between THg and MeHg  
161 concentrations and DOM composition (sources and quality) across a diversity of European  
162 streams. Unraveling the factors determining THg and MeHg background levels in the aquatic  
163 environment at a broad geographical scale is crucial for a better understanding of Hg cycling.  
164 For this purpose, we measured THg and MeHg concentrations, DOM concentration and

165 composition, anions and cations in filtered and unfiltered stream water at 29 sites with  
166 contrasting catchment characteristic across a wide latitudinal gradient across Europe.

167

## 168 **2. MATERIAL AND METHODS**

### 169 **2.1 Organizational Framework**

170 Water sampling for THg and MeHg concentration measurements was organized  
171 within the framework of the project called EuroRun: *Assessing CO<sub>2</sub> fluxes from European*  
172 *running waters* (available from <https://freshproject-eurorun.jimdo.com/>). EuroRun represents  
173 the first Collaborative European Freshwater Science Project for Young Researchers aiming  
174 to create synergistic interactions among young and early career scientists across Europe.  
175 Within the context of the collaborative EuroRun project, the EuroRun\_Mercury initiative  
176 included 12 teams distributed in 8 countries, which sampled 29 stream sites (Fig. 1). With  
177 56% of participants being female, EuroRun\_Mercury was gender balanced. The teams  
178 represented different regions across Europe. For a specific region, each team selected  
179 several sites with contrasting dominant land uses (i.e. forest or agriculture) and a wide range  
180 of hydrological and watershed characteristics (Table S1).

181

### 182 **2.2 Sampling**

183 The sampling procedure was explained and demonstrated to one lead representative  
184 from each team, who participated in a workshop performed in Sweden in September 2016.  
185 All the material needed to collect water samples along with a detailed written protocol was  
186 provided to participants during the workshop (see SI). The sampling took place during 3<sup>rd</sup> –  
187 14<sup>th</sup> October 2016 and teams from Austria (AUT), Bulgaria (BGR), the Czech Republic  
188 (CZE), Germany (DEU), Spain (ESP), France (FRA), Great Britain (GBR), Portugal (PRT)  
189 and Sweden (SWE) collected water samples for this study (Fig. 1). The 29 sampled sites  
190 were located across a wide latitudinal gradient (from 41°N to 64 °N) and altitude range (from  
191 1 to 1560 m above sea level), and spanned a wide range of hydrological (discharge from 0.1  
192 to 56 m<sup>3</sup> s<sup>-1</sup>) and catchment characteristics (Table S1). Each team provided hydrological



193 (discharge, stream wetted width and stream order) and catchment characteristics data (land  
194 use in % and catchment area).

195 A total of 4 water samples were collected at each site. Amber borosilicate bottles  
196 (VWR, 250 mL) were used to store water and analyze THg and MeHg species in filtered and  
197 unfiltered fractions. A previous study determined the cleaning procedures suitable for THg  
198 and MeHg analyses with these bottles (Cavalheiro et al., 2016). In the field, borosilicate  
199 bottles were rinsed three times with filtered (0.45  $\mu\text{m}$ , Sterivex-HV) or unfiltered water. For  
200 dissolved (<sub>D</sub>) THg<sub>D</sub> and MeHg<sub>D</sub> analyses about 250 mL of water were collected directly into  
201 the amber borosilicate bottle, filtered (0.45  $\mu\text{m}$ , Sterivex-HV) and acidified (1% v/v final  
202 concentration; HCl Ultrex II, J. T. Baker) on site. A subsample of filtered water was placed  
203 into 40 mL pre-muffled EPA borosilicate glass vials (VWR) with PTFE caps for analysis of  
204 DOC concentration, DOM optical properties and concentration of anions and cations.  
205 Unfiltered water samples were collected for determination of the total fraction (<sub>T</sub>) of THg<sub>T</sub>,  
206 MeHg<sub>T</sub> (acidified) and total organic carbon (TOC) concentration. Samples were kept at 4°C  
207 and shipped to the laboratory by courier within 48 h. A subsample (10 mL) of unfiltered water  
208 was filtered in the laboratory, and stored frozen for later determination of anions and cations.  
209 Because four of the vials broke during transport, these samples could not be determined  
210 (marked as not determined in Table 1 and S2).

211 Results for unfiltered water samples correspond to the total fraction (THg<sub>T</sub>, MeHg<sub>T</sub>  
212 and TOC), whereas the results for filtered water samples represent the dissolved fraction  
213 (THg<sub>D</sub>, MeHg<sub>D</sub>, DOC, DOM optical properties, anions and cations).

214

## 215 **2.3 Chemical Analyses**

### 216 **2.3.1 Analysis of Inorganic-Hg and Methylmercury**

217 Inorganic-Hg (IHg) and MeHg (monomethylmercury) concentrations (Table 1) were  
218 measured using species-specific isotope dilution and capillary gas chromatography (Trace  
219 GC Ultra, Thermo Fisher, Waltham, MA, USA equipped with a TriPlus RSH auto-sampler)  
220 hyphenated to inductively coupled plasma mass spectrometer (Thermo X Series 2)

221 (Cavalheiro et al., 2016; Rodriguez Martín-Doimeadios et al., 2003). The detection limits of  
222 this method are 0.02 ng L<sup>-1</sup> for IHg and 0.005 ng L<sup>-1</sup> for MeHg. The measurement error  
223 (calculated by analyzing each sample three times) was less than 15% and 10% for IHg and  
224 MeHg concentrations, respectively. Based on previous studies (Bravo et al., 2014;  
225 Monperrus et al., 2004; Rodriguez Martín-Doimeadios et al., 2003), THg was calculated as  
226 the sum of IHg and MeHg.

### 227 **2.3.2 TOC, DOC and DOM optical properties**

228 TOC and DOC concentrations were determined on unfiltered and filtered samples,  
229 respectively. Prior to analysis, samples were acidified to a pH of 2 with 0.1 M HCl, and  
230 analyzed using a Shimadzu TOC-V<sub>CPH</sub> total carbon analyzer. Optical properties were  
231 determined on filtered water using an Aqualog spectrofluorometer (Horiba), which measures  
232 absorbance and fluorescence simultaneously. The fluorescence spectrophotometer has a  
233 routine protocol before running samples that involves i) instrument validation (ensuring the  
234 signal : noise ratio <20,000 counts ii) ensuring the excitation lamp is calibrated to a given  
235 wavelength (maximum intensity at 467 nm), iii) the emission energy is calibrated using pure  
236 water at an emission of 397 nm and an excitation of 350 nm. Fluorescence spectra were  
237 collected across excitation wavelengths (Ex<sub>λ</sub>) 250 to 445 nm at 5 nm increments and across  
238 emission wavelengths (Em<sub>λ</sub>) ranging from 300–600 nm. Bandpass and resolution were 5 nm.  
239 All spectra were measured in a 1 cm quartz cuvette and automatically blank-subtracted,  
240 corrected for instrument biases and inner filter effects using the FluorEssence™ software  
241 (Horiba). Samples were measured for fluorescence at a 2 second integration time, with the  
242 exception of five more optical dense samples from Great Britain and Sweden, which were  
243 measured at 0.5 seconds. All intensities were normalized to 1 second integration time post-  
244 analysis and the excitation-emission matrixes (EEM) were expressed in fluorescence units  
245 (FU).

246 Peak picking was used to define the relative intensity of four common peaks across  
247 the EEM including the Peak A (λ<sub>ex</sub>: 260 nm, λ<sub>em</sub>: 400–460 nm), Peak C (λ<sub>ex</sub>: 320–360 nm, λ<sub>em</sub>:  
248 420–460 nm), and Peak M (λ<sub>ex</sub>: 290–310 nm, λ<sub>em</sub>: 370–410 nm) associated with soil

249 terrestrial DOM, as well as a protein-like peak T ( $\lambda_{\text{ex}}$ : 275 nm,  $\lambda_{\text{em}}$ : 340–350 nm) that is  
250 considered to be of algal/bacterial origin (i.e. autochthonous to the aquatic environment)  
251 (Fellman et al., 2010; Huguet et al., 2009). The summed area of total fluorescence (TF)  
252 under the EEMs was used to normalize each of the four peaks, which were expressed as a  
253 percent of TF (i.e. %Peak A = Peak A / TF x 100%). We also calculated three common  
254 fluorescence indexes: i) the fluorescence index (FI, calculated as the ratio between emission  
255 at wavelengths of 470 and 520 nm, at an excitation of 370 nm), an indicator of source being  
256 more microbial ( $\approx 1.8$ ) or terrestrial ( $\approx 1.2$ ) (Cory et al., 2007; McKnight et al., 2001), ii) the  
257 freshness index (calculated as the ratio of emission at 380 nm, divided by the emission  
258 maxima between 420 and 435 nm, at an excitation of 310 nm) which reflects the age of OM,  
259 with higher values representing more recently produced OM (Parlanti et al., 2000) and iii) the  
260 humification index (HIX, ratio of peak areas across an emission spectra [435–480 nm / 300–  
261 345 nm]), indicating the degree of humification (Zsolnay et al., 1999). The absorbance at 254  
262 nm ( $A_{254}$ ) was used to calculate carbon specific absorbance ( $\text{SUVA}_{254}$ ), a commonly used  
263 metric of the degree of aromaticity, as  $A_{254}/\text{DOC}$  concentration normalized to a 1 m path  
264 length ( $\text{L mg C}^{-1} \text{ m}^{-1}$ ) (Weishaar et al., 2003).

### 265 **2.3.3 Anions and Cations**

266 Inorganic analyses were conducted on a Metrohm IC system (883 Basic IC Plus and  
267 919 Autosampler Plus). Anions ( $\text{F}^-$ ,  $\text{Cl}^-$ ,  $\text{NO}_3^-$  and  $\text{SO}_4^{2-}$ ) were separated on a Metrosep A  
268 Supp 5 analytical column (150 x 4.0 mm) fitted with a Metrosep A Supp 4/5 guard column at  
269  $0.7 \text{ mL min}^{-1}$  using a carbonate eluent (3.2 mM  $\text{Na}_2\text{CO}_3$  + 1.0 mM  $\text{NaHCO}_3$ ). A Metrosep C4  
270 column (250 x 2.0 mm) with a Metrosep C4 guard column was used for separation of the  
271 cations ( $\text{Na}^+$ ,  $\text{NH}_4^+$ ,  $\text{K}^+$ ,  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$ ) with an eluent of 1.7 mM nitric acid and 0.7 mM dipicolinic  
272 acid at a flow rate of  $0.2 \text{ mL min}^{-1}$ .

### 273 **2.4 Statistical Analyses**

274 Kolmogorov–Smirnov test and quantile–quantile plots were used to evaluate the  
275 normality of the data. The distribution of THg concentrations were slightly skewed due to the  
276 high concentrations measured at sites in Great Britain. The center and dispersion of THg

277 concentrations were summarized by the median and the inter-quartile range (IQR). DOM  
278 optical parameters, MeHg and %MeHg met normality after being log-transformed. Parametric  
279 tests were used to analyze linear correlations (Pearson) between DOM optical parameters,  
280 THg and MeHg concentrations, and %MeHg and to compare Hg concentrations in the total  
281 and dissolved fractions (t-test). All statistical tests were carried out in R 3.2.4 ([http://www.R-](http://www.R-project.org/)  
282 [project.org/](http://www.R-project.org/)) (R Core Team, 2016).

283 Principle component analysis (PCA) was used as an initial diagnostic tool to examine  
284 key relationships and identify strongly correlated variables including optical measurements  
285 (i.e. %Peak A, %Peak C, %Peak M and %Peak T, SUVA<sub>254</sub>, fluorescence index, freshness  
286 index, humification index), TOC, sulfate, nitrate and calcium concentrations, land use and  
287 stream order. This complete information was available for 21 of the 29 studied sites.  
288 Accordingly, the samples with incomplete information (DEU1\_1, DEU1\_2, GBR2\_2, FRA1\_2,  
289 PRT1\_1, CZE1\_1, GBR2\_1 and ESP2\_3) were excluded from the PCA analysis. All data  
290 were log-transformed, scaled and centered prior to PCA analysis. PCA analysis were carried  
291 out using the *prcomp* function of the package *stats* in R 3.2.4 (R Core Team, 2016).

292 Partial least squares (PLS) regression models were subsequently applied to predict  
293 THg concentration (Y response variable) from DOM optical properties, water chemistry and  
294 catchment characteristics (X predictor variables, Table S5 and Table S6). A second PLS,  
295 excluding the sites with high THg concentrations (Great Britain, GBR1\_1, GBR1\_2), was also  
296 tested (Supplementary information, Table S7). The PLS model performance was examined  
297 based on the cumulative goodness of fit ( $R^2Y$ , explained variation), and the cumulative  
298 goodness of prediction (or  $Q^2$ , predicted variation). The variable influence on projections (VIP  
299 scores) provided a means of assigning the importance of X variables on the overall PLS  
300 model. X-predictor variables with an influence score  $\geq 1$  were considered highly influential,  
301 between 0.8 and 1.0 moderately influential, and  $< 0.8$  less influential predictors (Eriksson et  
302 al., 1999). The PLS models were performed using the *pls* function of the *pls* package in R  
303 (Mevik and Wehrens, 2007). Graphics were built in R 3.2.4 (R Core Team, 2016) and  
304 modified to fit the journal requirements with Inkscape 0.92 (<https://inkscape.org/es/>).

305 THg and MeHg loads of both total and dissolved fractions (Table S3) were calculated  
306 with the stream discharge value provided by the participants (Table S1) and the  
307 concentrations measured in this study (Table 1).

### 308 **3. RESULTS**

#### 309 **3.1 Mercury Levels (THg, IHg, MeHg and %MeHg)**

310 THg concentrations were consistently low among sites (Table 1). Median values for  
311 total (THg<sub>T</sub>) and dissolved fractions (THg<sub>D</sub>) were 0.46 ng L<sup>-1</sup> (interquartile range, IQR: 0.28 –  
312 0.8) and 0.24 ng L<sup>-1</sup> (IQR: 0.28 – 0.49), respectively. Streams in Great Britain (GBR) and  
313 Austria (AUT) had the highest THg<sub>T</sub> and THg<sub>D</sub> concentrations (Fig. 2a, Table 1). THg<sub>T</sub>  
314 concentrations were generally higher than THg<sub>D</sub> (p = 0.004), although both were significantly  
315 correlated (R<sup>2</sup>=0.62, p = 0.04; Table S3). IHg<sub>T</sub> was also slightly higher than IHg<sub>D</sub> (Fig. 2b).  
316 MeHg<sub>T</sub> concentrations ranged between 7.8 and 159 pg L<sup>-1</sup> and were similar (p = 0.07) to  
317 MeHg<sub>D</sub> (6.6 – 134 pg L<sup>-1</sup>; Fig. 2c). The highest MeHg concentrations, for total and dissolved  
318 fractions, were found at sites from Sweden (SWE2\_1), Great Britain (GBR1\_1) and Spain  
319 (ESP1\_3). The %MeHg was very similar in both the total (%MeHg<sub>T</sub>) and dissolved (%MeHg<sub>D</sub>)  
320 fractions (Fig. 2d) but varied among sites, ranging from 1 to 43%. The highest %MeHg<sub>T</sub> was  
321 found at a Spanish site (ESP1\_3). The median of THg<sub>T</sub> and MeHg<sub>T</sub> loads were 3.9 kg year<sup>-1</sup>  
322 (IQR: 0.6 – 6.8) and 0.3 kg year<sup>-1</sup> (IQR: 0.1 – 0.5), respectively.

323 Significant correlations were found between the total and dissolved fraction of THg,  
324 IHg and MeHg (Table S3). As total and dissolved fractions were correlated, we used the total  
325 fraction (THg<sub>T</sub>, MeHg<sub>T</sub>, %MeHg<sub>T</sub>) for all the following statistical analyses (PCA and PLS).

#### 326 **3.2 Water Chemistry**

327 DOC and TOC concentrations ranged between 0.9 and 18.5 mg L<sup>-1</sup> and 0.9 and 22.2  
328 mg L<sup>-1</sup>, respectively. DOC and TOC were significantly correlated (R<sup>2</sup> = 0.99, p < 0.01). The  
329 highest concentrations were found in Great Britain (GBR1\_1, GBR1\_2) and Sweden  
330 (SWE1\_1) and the lowest in Spain (ESP1\_2, ESP2\_1) and Austria (AUT2\_1, AUT2\_2,  
331 AUT2\_3; Table S2). The concentrations of sulfate, a well-known electron acceptor for Hg  
332 methylating microorganisms, varied largely and ranged from 1.3 (AUT1\_1) to 731 (ESP2\_3)

333 mg L<sup>-1</sup>. About 50% of the studied sites presented sulfate concentrations lower than 12.6 mg  
334 L<sup>-1</sup> (IQR: 6.8 – 66 mg L<sup>-1</sup>). Spanish sites presented the highest sulfate concentrations (Table  
335 S2). Nitrate concentration varied between 0.04 (GBR1\_1) and 23 (ESP2\_2) mg L<sup>-1</sup>. The  
336 median value for nitrate concentration was 1.9 mg L<sup>-1</sup> (IQR: 0.7 – 2.7).

### 337 **3.3 DOM Optical Characteristics**

338 Very distinct fluorescence signatures were observed between study sites, with four  
339 illustrative EEMs representing contrasting signatures displayed in Figure 3. Sites from Great  
340 Britain showed a strong optical signature representing terrestrially derived DOM, as indicated  
341 by the high intensity of Peaks A and C, while in contrast, sites from Spain (ESP) and the  
342 Czech Republic (CZE), with high Peak T intensities, represented DOM optical signatures of  
343 waters typically enriched in autochthonous DOM (Fig. 3, Table S2). The wide range of bulk  
344 DOM optical characteristics captured in this dataset can be attributed to the broad  
345 geographical scale of this study.

346 Analyzed water samples also captured a wide range of DOM characteristics based on  
347 each of the measured optical indexes. The SUVA<sub>254</sub>, which is associated with the presence  
348 of aromatic (Weishaar et al., 2003) and high molecular mass compounds (Chowdhury, 2013)  
349 typically spans from <1 to a maximum of 6 L mg C<sup>-1</sup> m<sup>-1</sup> (Weishaar et al., 2003), and was  
350 found to range from 2.4 (CZE1\_2) to 5.2 (GBR1\_1, GBR1\_2) L mg C<sup>-1</sup> m<sup>-1</sup>. HIX values  
351 spanned from 2.2 to 25 (Table S2). The highest HIX values were measured in Austria  
352 (AUT1\_2) and Great Britain (GBR1\_1, GBR1\_2). Some of the lowest values were observed  
353 in the Czech Republic (CZE1\_2) and Austria (AUT2\_2). While high HIX values (> 10) have  
354 been linked to the presence of complex molecules and strongly humified organic material,  
355 mainly of terrestrial origin, low values (<4) have been associated with autochthonous OM  
356 (Huguet et al., 2009). The fluorescence index (FI) covered almost the full range of this index,  
357 from a value of 1.3, representing terrestrial sources (degraded plant and soil OM; lower  
358 values), in GBR1\_2, to a value of 1.8, indicating microbial sources (including extracellular  
359 release and leachate from bacteria and algae) in ESP2\_2, ESP3\_3 (Cory et al., 2007;  
360 McKnight et al., 2001). The freshness index, typically associated with the contribution of

361 recently produced microbial-derived DOM (Parlanti et al., 2000), also spanned a wide range  
362 from 0.34 to 0.91 and was positively correlated to the FI and negatively correlated to the HIX  
363 and SUVA<sub>254</sub> (Fig. 4).

364 The wide distribution of DOM composition between sites indicated differences in the  
365 source, processing pathways and biological reactivity of the DOM, as shown by the PCA  
366 analysis (Fig. 5). The first principal component (PC1) explained 41% of the total variance  
367 across 21 European sites (Fig. 5a). Positive loadings on PC1 were associated with high  
368 TOC, A<sub>254</sub> and HIX, indicating a terrestrial DOM signature. In contrast, negative loadings of  
369 PC1 were found for agricultural land use, NO<sub>3</sub><sup>-</sup>, SO<sub>4</sub><sup>2-</sup>, Ca<sup>2+</sup>, fluorescence Peak T and Peak  
370 M, freshness index and FI, likely associated with fresh and recently produced algal and  
371 microbial DOM (Ohno, 2002). Running waters from Great Britain (GBR1\_1; GBR1\_2),  
372 Sweden (SWE1\_1, SWE2\_1) and Austria (AUT1\_1, AUT1\_2) were characterized by low  
373 anions, cations and high concentrations of terrestrial humidified DOM (positive scores in PC1  
374 Fig. 5b). In contrast, negative scores (Fig. 5b) of running waters from Spain (ESP) and the  
375 Czech Republic indicated an enrichment in nutrients and fresh autochthonous OM. Also,  
376 negative loadings for the percentage of urban and agricultural areas in the studied  
377 catchments suggest that anthropogenic activities might enrich running waters in nutrients  
378 and enhance primary production (Fig. 5a). The second component (PC2) explained 14% of  
379 the variance. Positive loadings on PC2 were found for the fluorescence Peak A and the ratio  
380 of the Peaks A to C (A/C). Both Peaks A and C have been linked to the presence of humic  
381 substances (Fellman et al., 2010). However, an increase in the relative abundance of the  
382 peak A compared to Peak C (A/C peak ratio) has been related to DOM processing, as Peak  
383 C is lost preferentially to Peak A under both photo- and biodegradation processes  
384 (Kothawala et al., 2012). Therefore, positive loadings of PC2 indicated terrestrial DOM that  
385 had been processed (Fig. 5a). On the opposite end, negative loadings on PC2 were  
386 associated with aromatic compounds (SUVA<sub>254</sub>) (Weishaar et al., 2003) and terrestrial DOM  
387 (Peak C) yet not strongly degraded (Kothawala et al., 2012).

### 388 **3.4 Predictors of THg, MeHg and %MeHg Concentrations**

389 A PLS model (PLS model I, Fig. 6) identified key factors driving THg<sub>T</sub> concentrations  
390 in running waters. From the 29 available sites, only 21 sites were included in the PLS model  
391 which contained a complete set of required variables. To interpret the PLS loading plots, the  
392 distance and positioning of Y variables (THg, MeHg or %MeHg) relative to X variables  
393 indicated how well, or poorly, they related to each other. The greater the distance a variable  
394 (X or Y) lies from the origin, the greater its overall influence. Variables situated close together  
395 on the PLS plot are positively correlated, while variables opposite to each other are  
396 negatively correlated. The proportion of variance in the X variables (i.e. DOM composition,  
397 anions, cations and environmental factors) explained by the model is indicated by the R<sup>2</sup>X.  
398 The R<sup>2</sup>Y value indicates the proportion of variance in THg explained by the model. With two  
399 predictive components (R<sup>2</sup>X, 50%), the variance of THg concentrations was well explained  
400 (R<sup>2</sup>Y, 83.2%) and relatively well predicted (Q<sup>2</sup>, 60%) by the PLS model I (Fig. 6, Table S5,  
401 Table S6). PLS model I indicates that while TOC and A<sub>254</sub> presented the highest positive  
402 correlation with THg, freshness index, FI and agriculture were negatively correlated. When  
403 not including sites highly influenced by terrestrial DOM inputs, GBR1\_1 and GBR1\_2, a  
404 second PLS (PLS model II), with two predictive components (R<sup>2</sup>X, 40%), explained the  
405 variance of THg<sub>T</sub> concentrations relatively well (R<sup>2</sup>Y, 82%) but could not predict them (Q<sup>2</sup>, -  
406 2%) (Table S7).

407 It was not possible to predict a significant proportion of the variability in MeHg  
408 concentrations using a PLS model with the measured variables. Nevertheless, the  
409 concentration of MeHg was significantly correlated to %MeHg (Fig. 4). Additionally, the  
410 %MeHg was significantly and positively correlated to Peak M (%), FI and sulfate  
411 concentration and negatively related to HIX and THg<sub>T</sub> (Fig. 4).

412

## 413 **4. Discussion**

### 414 **4.1 Hg levels in European running waters**

415 Our results provide new information on THg and MeHg concentrations in streams at a  
416 European scale. THg concentrations across the studied European running waters were in the



417 same range as those measured in pristine streams of northern Sweden (Eklöf et al., 2012),  
418 USA (Oregon, Wisconsin and Florida, Brigham et al., (2009); California, Tsui et al., (2009)  
419 and Minnesota, Balogh et al., (2002)) and France (Cavalheiro et al., 2016) and were lower  
420 than those reported for rivers typically contaminated by point sources (Baptista-Salazar et al.,  
421 2017; Bravo et al., 2014). Therefore, none of the study sites appeared to be affected by point  
422 sources of Hg.

423 Fluvial systems are an important transport pathway and ultimately deliver Hg  
424 originating from catchment soils to the marine environment. At a broader scale, we attempted  
425 to provide a rough estimate discharge of THg<sub>T</sub> and MeHg<sub>T</sub> of European rivers considering  
426 median values obtained in this study and the total European river discharge (3100 km<sup>3</sup>)  
427 (Shiklomanov and Rodda, 2004). Without considering point source of contamination, such as  
428 chlor-alkali or mining activities, known to increase stream Hg concentrations, the estimated  
429 European river discharge of THg<sub>T</sub> and MeHg<sub>T</sub> would be 1426 kg year<sup>-1</sup> (IQR: 868 – 2494)  
430 and 97 kg year<sup>-1</sup> (IQR: 60 – 254), respectively. Particles are an important vector of Hg  
431 transport in streams (Schuster et al., 2011) and can influence hydrologically mediated  
432 processes such as sedimentation and/or remobilization (e.g heavy rain events) of Hg  
433 previously deposited within river systems, which can largely affect THg<sub>T</sub> and MeHg<sub>T</sub> loads  
434 (Baptista-Salazar et al., 2017; Moreno-Brush et al., 2016; Sunderland and Mason, 2007).  
435 Measured THg and MeHg concentrations were notably higher in the total fraction compared  
436 to the dissolved fraction, highlighting the role of particles in the transport and fate of Hg in the  
437 studied fluvial systems. As precise hydrological data were unavailable here, and our  
438 calculations are based on a single sampling campaign, and the discharge of THg and MeHg  
439 only provides an initial rough estimate to be considered with care. In order to improve upon  
440 current estimates of European Hg discharge into the oceans (Kocman et al., 2017), greater  
441 monitoring of THg and MeHg is required across fluvial systems, particularly beyond the  
442 boreal ecozone.

#### 443 **4.2 Terrestrial DOM: An Important Source of THg in European Running Waters**

444 Using a multivariate approach comparing 21 streams spanning a wide geographical  
445 area across Europe, we were able to discriminate between the relative influence of bulk  
446 DOM optical characteristics and environmental variables on THg and MeHg concentrations.  
447 PC1 of the PLS model showed positive correlations between the highly influential predictors  
448 TOC and  $A_{254}$  with THg concentrations in the studied sites and suggests that terrestrial DOM  
449 is an important driver of THg in streams (Fig. 6). Sites characterized by humic-rich soils, such  
450 as those in Sweden and Great Britain (Jones et al., 2005), presented the highest THg<sub>T</sub> and  
451 THg<sub>D</sub> concentrations. The positive correlation between terrestrial DOM and THg  
452 concentration in boreal rivers and streams has been previously reported in the literature  
453 (Brigham et al., 2009; Eklöf et al., 2012). As atmospheric Hg deposition during the industrial  
454 era have led to the accumulation of Hg in organic rich soils (Johansson and Tyler, 2001), the  
455 remobilization of Hg-enriched soil OM is an important process controlling THg concentrations  
456 in streams (Baptista-Salazar et al., 2017; Brigham et al., 2009; Driscoll et al., 1995; Eklöf et  
457 al., 2012; Kirk and Louis, 2009; Tsui et al., 2009) and lakes (Bravo et al., 2017).

458 It is noteworthy to highlight the lack of a significant correlation between TOC and  
459 THg<sub>T</sub> concentration ( $p = 0.6$ ) when the sites with the highest TOC and THg<sub>T</sub> concentrations  
460 (SWE2\_1, GBR1\_1 and GBR1\_2) were excluded from the statistical analyses. Also, a poor  
461 predictability was obtained in a PLS model (PLS model II) built for THg<sub>T</sub> excluding the sites  
462 highly influenced by terrestrial DOM (GBR1\_1 and GBR1\_2, Table S7). Therefore, TOC, and  
463 DOM components associated with terrestrial DOM, failed to predict THg<sub>T</sub> concentrations in  
464 southern parts of Europe (FRA, ESP, CZE, AUT, BGR and DEU). This suggests that other  
465 Hg sources are relevant for the studied southern streams. Direct inputs of Hg deposited by  
466 wet and dry atmospheric depositional processes (Brigham et al., 2009; Domagalski et al.,  
467 2016; Lee et al., 2001) could contribute to THg<sub>T</sub> concentrations in streams with low inputs of  
468 terrestrial DOM. Besides atmospheric deposition, catchment characteristics and changes in  
469 runoff may be relevant in determining THg<sub>T</sub> in streams and rivers (Baptista-Salazar et al.,  
470 2017; Domagalski et al., 2016). Therefore, our results confirm the role of terrestrial OM as an  
471 important source of Hg for streams but highlight the uncertainties on the drivers of THg

472 concentrations at broader global scale when considering systems with low loads of terrestrial  
473 OM.

#### 474 **4.3 Autochthonous DOM and Nutrients: Important Drivers of MeHg Formation in** 475 **European Running Waters**

476 In this study, MeHg concentrations measured in running waters were in the same range as  
477 those previously reported for Sweden (Skylberg et al., 2003), USA (California, Tsui et al.,  
478 (2009); Florida, Brigham et al., (2009)) and France (Cavalheiro et al., 2016). Similarly, the  
479 %MeHg measured in samples collected in Sweden and Great Britain were similar to those  
480 previously reported in the literature (Brigham et al., 2009; Skylberg et al., 2003). In contrast  
481 to THg, MeHg concentrations were not correlated to TOC,  $A_{254}$  or HIX (Fig. 4), suggesting  
482 that terrestrial sources are not the main vector for MeHg in the studied streams. MeHg  
483 concentrations have been related to the concentration of autochthonous OM in reservoirs,  
484 lakes and in the Baltic sea (Gascon Diez et al., 2016; Noh et al., 2017; Soerensen et al.,  
485 2017).

486 The %MeHg has been considered a proxy of net MeHg formation (Drott et al., 2008).  
487 In the studied European streams studied here, the positive correlation between %MeHg to  
488 Peak M and the FI indicated that autochthonous OM is important for MeHg formation. Peak  
489 M, as well as Peak A and Peak C, have been generally described as humic-like peaks  
490 (Fellman et al., 2010). Peaks A and C have been linked to DOM derived primarily from  
491 vascular plant sources, aromatic and with a higher-molecular-weight fraction of the DOM  
492 pool (Coble, 1996). In contrast, Peak M represents compounds less aromatic and of lower  
493 molecular weight as compared to peaks A and C (Fellman et al., 2010). In a previous study  
494 performed in freshwaters, strong correlations between Peak M and Peak T (protein-like)  
495 were reported, but weaker relationships were observed between Peak M and Peaks A and C  
496 (Kothawala et al., 2014). Indeed, Peak M was originally used as a marker for autochthonous  
497 DOM from planktonic production (Fellman et al., 2010). A positive correlation of the %MeHg  
498 with Peak M and the FI thus suggests that net MeHg production is linked to fresh  
499 autochthonous OM. In contrast, the negative correlation between %MeHg and HIX suggests

500 that high concentrations of highly humified terrestrial DOM hampered Hg methylation  
501 processes most likely by decreasing Hg bioavailability (Chiasson-Gould et al., 2014;  
502 Soerensen et al., 2017) and/or bacterial activity (Bravo et al., 2017). High Hg methylation  
503 rates have been associated with algal derived OM (Bravo et al., 2017; Gascon Diez et al.,  
504 2016; Herrero Ortega et al., 2018; Mazrui et al., 2016). Our study provides further evidence  
505 of the potential role of algal and/or microbial DOM in the *in situ* formation of MeHg in  
506 streams.

507         The methylation of IHg to MeHg in aquatic systems has been primarily attributed to  
508 the activity of sulfate-reducing bacteria, iron-reducing bacteria (FeRB), methanogens and  
509 Firmicutes (Gilmour et al., 2013) in oxygen limiting conditions of sediments (Bravo et al.,  
510 2017), hypolimnion (Eckley and Hintelmann, 2006) and settling particles (Gascon Diez et al.,  
511 2016). As sulfate can be used as an electron acceptor by Hg methylating microorganisms  
512 (Pak and Bartha, 1998), the positive significant correlation found between the %MeHg and  
513 the sulfate concentration, suggests an important role of SRB in MeHg formation in running  
514 waters and provides further evidence for the relevance of *in situ* MeHg formation.

515         Stream sulfate mainly originates from atmospheric deposition and/or geogenic  
516 sources (Ledesma et al., 2018). Because sulfate release is accompanied by cation leaching  
517 (Prechtel et al., 2001), high concentrations of both sulfate and cations (e.g. calcium, Table  
518 S2) in Spanish sites indicate an effect of geological substrate of the river basin on the stream  
519 water chemistry. Although it was not the focus of this work, these results underline the  
520 relevance of geology and catchment characteristics on stream water properties and  
521 ultimately also for Hg cycling.

522

## 523 **Conclusion**

524         Our findings indicated that streams enriched in terrestrial DOM, as typical of Northern  
525 study sites, revealed the highest THg concentrations, suggesting the source was from  
526 organic rich catchment soils. In contrast, Southern European streams were enriched in  
527 nutrients and autochthonous DOM and although inputs of THg and terrestrial DOM from the

528 catchment were lower, we propose that MeHg levels were regulated by. We propose that the  
529 elevated MeHg concentrations in some of these sites could be due to a higher *in situ* MeHg  
530 formation, likely related to enhanced microbial activity, which is boosted by the availability of  
531 autochthonous algal/microbial derived DOM and sulfate. Our results show the interplay  
532 between DOM components and THg and MeHg concentrations in streams and provide new  
533 insights into Hg cycling in aquatic systems at a broad geographical European scale.

534

## 535 **References**

- 536 Amos, H.M., Jacob, D.J., Kocman, D., Horowitz, H.M., Zhang, Y., Dutkiewicz, S., Horvat, M., Corbitt,  
537 E.S., Krabbenhoft, D.P., Sunderland, E.M., 2014. Global biogeochemical implications of mercury  
538 discharges from rivers and sediment burial. *Environ. Sci. Technol.* 48, 9514–9522.  
539 doi:10.1021/es502134t
- 540 Amos, H.M., Jacob, D.J., Streets, D.G., Sunderland, E.M., 2013. Legacy impacts of all-time  
541 anthropogenic emissions on the global mercury cycle. *Global Biogeochem. Cycles* 27, 410–421.  
542 doi:10.1002/gbc.20040
- 543 Balogh, S.J., Huang, Y., Offerman, H.J., Meyer, M.L., Johnson, D.K., 2002. Episodes of elevated  
544 methylmercury concentrations in prairie streams. *Environ. Sci. Technol.* 36, 1665–1670.  
545 doi:10.1021/es011265w
- 546 Baptista-Salazar, C., Richard, J.H., Horf, M., Rejc, M., Gosar, M., Biester, H., 2017. Grain-size  
547 dependence of mercury speciation in river suspended matter, sediments and soils in a mercury  
548 mining area at varying hydrological conditions. *Appl. Geochemistry* 81, 132–142.  
549 doi:10.1016/j.apgeochem.2017.04.006
- 550 Bravo, A.G., Bouchet, S., Guédron, S., Amouroux, D., Dominik, J., Zopfi, J., 2015. High methylmercury  
551 production under ferruginous conditions in sediments impacted by sewage treatment plant  
552 discharges. *Water Res.* 80, 245–255. doi:10.1016/j.watres.2015.04.039
- 553 Bravo, A.G., Bouchet, S., Tolu, J., Björn, E., Mateos-Rivera, A., Bertilsson, S., 2017. Molecular  
554 composition of organic matter controls methylmercury formation in boreal lakes. *Nat. Commun.*  
555 8, 14255. doi:10.1038/ncomms14255

556 Bravo, A.G., Cosio, C., Amouroux, D., Zopfi, J., Chevalley, P.-A., Spangenberg, J.E., Ungureanu, V.-  
557 G., Dominik, J., 2014. Extremely elevated methyl mercury levels in water, sediment and  
558 organisms in a Romanian reservoir affected by release of mercury from a chlor-alkali plant.  
559 *Water Res.* 49, 391–405.

560 Brigham, M.E., Wentz, D. a., Aiken, G.R., Krabbenhoft, D.P., 2009. Mercury cycling in stream  
561 ecosystems. 1. Water column chemistry and transport. *Environ. Sci. Technol.* 43, 2720–2725.  
562 doi:10.1021/es802694n

563 Cavalheiro, J., Sola, C., Baldanza, J., Tessier, E., Lestremau, F., Botta, F., Preud'homme, H.,  
564 Monperrus, M., Amouroux, D., 2016. Assessment of background concentrations of  
565 organometallic compounds (methylmercury, ethyllead and butyl- and phenyltin) in French aquatic  
566 environments. *Water Res.* 94, 32–41. doi:10.1016/j.watres.2016.02.010

567 Chiasson-Gould, S.A., Blais, J.M., Poulain, A.J., 2014. Dissolved organic matter kinetically controls  
568 mercury bioavailability to bacteria. *Environ. Sci. Technol.* 48, 3153–3161.  
569 doi:10.1021/es4038484

570 Chowdhury, S., 2013. Trihalomethanes in drinking water: Effect of natural organic matter distribution.  
571 *Water* 39, 1–8.

572 Coble, P.G., 1996. Characterization of marine and terrestrial DOM in seawater using excitation-  
573 emission matrix spectroscopy. *Mar. Chem.* 51, 325–346. doi:10.1016/0304-4203(95)00062-3

574 Cory, R.M., McKnight, D.M., Chin, Y.P., Miller, P., Jaros, C.L., 2007. Chemical characteristics of fulvic  
575 acids from Arctic surface waters: Microbial contributions and photochemical transformations. *J.*  
576 *Geophys. Res. Biogeosciences* 112, 1–14. doi:10.1029/2006JG000343

577 Dennis, I.F., Clair, T.A., Driscoll, C.T., Kamman, N., Chalmers, A., Shanley, J., Norton, S.A., Kahl, S.,  
578 2005. Distribution patterns of mercury in Lakes and Rivers of northeastern North America.  
579 *Ecotoxicology* 14, 113–123. doi:10.1007/s10646-004-6263-0

580 Domagalski, J., Majewski, M.S., Alpers, C.N., Eckley, C.S., Eagles-Smith, C.A., Schenk, L., Wherry,  
581 S., 2016. Comparison of mercury mass loading in streams to atmospheric deposition in  
582 watersheds of Western North America: Evidence for non-atmospheric mercury sources. *Sci.*  
583 *Total Environ.* 568, 638–650. doi:10.1016/j.scitotenv.2016.02.112

584 Driscoll, C.T., Blette, V., Yan, C., Schofield, C.L., Munson, R., Holsapple, J., 1995. The role of  
585 dissolved organic carbon in the chemistry and bioavailability of mercury in remote Adirondack  
586 lakes. *Water, Air, Soil Pollut.* 80, 499–508. doi:10.1007/BF01189700

587 Driscoll, C.T., Han, Y.-J., Chen, C.Y., Evers, D.C., Lambert, K.F., Holsen, T.M., Kamman, N.C.,  
588 Munson, R.K., 2007. Mercury contamination in forest and freshwater ecosystems in the  
589 northeastern United States. *Bioscience* 57, 17. doi:10.1641/B570106

590 Drott, A., Lambertsson, L., Björn, E., Skyllberg, U., 2008. Do potential methylation rates reflect  
591 accumulated methyl mercury in contaminated sediments? *Environ. Sci. Technol.* 42, 153–158.

592 Eckley, C.S., Hintelmann, H., 2006. Determination of mercury methylation potentials in the water  
593 column of lakes across Canada. *Sci. Total Environ.* 368, 111–125.  
594 doi:10.1016/j.scitotenv.2005.09.042

595 Eklöf, K., Fölster, J., Sonesten, L., Bishop, K., 2012. Spatial and temporal variation of THg  
596 concentrations in run-off water from 19 boreal catchments, 2000-2010. *Environ. Pollut.* 164, 102–  
597 109. doi:10.1016/j.envpol.2012.01.024

598 Eklöf, K., Kraus, A., Futter, M., Schelker, J., Meili, M., Boyer, E.W., Bishop, K., 2015. Parsimonious  
599 model for simulating total mercury and methylmercury in boreal streams based on riparian flow  
600 paths and seasonality. *Environ. Sci. Technol.* 49, 7851–7859. doi:10.1021/acs.est.5b00852

601 Eriksson, L., Johansson, E., Wold, S., 1999. *Introduction to Multi- and Megavariate Data Analysis*  
602 *Using Projection Methods (PCA & PLS)*. Umetrics.

603 Fellman, J.B., Hood, E., Spencer, R.G.M., 2010. Fluorescence spectroscopy opens new windows into  
604 dissolved organic matter dynamics in freshwater ecosystems: A review. *Limnol. Oceanogr.* 55,  
605 2452–2462. doi:10.4319/lo.2010.55.6.2452

606 Fitzgerald, W.F., Engstrom, D.R., Mason, R.P., Nater, E.A., 1998. The case for atmospheric mercury  
607 contamination in remote Areas. *Environ. Sci. Technol.* 32, 1–7.

608 Gascon Diez, E., Loizeau, J.-L., Cosio, C., Bouchet, S., Adatte, T., Amouroux, D., Bravo, A.G., 2016.  
609 Role of settling particles on mercury methylation in the oxic water column of freshwater systems.  
610 *Environ. Sci. Technol.* 50, 11672–11679. doi:10.1021/acs.est.6b03260

611 Gilmour, C.C., Podar, M., Bullock, A.L., Graham, A.M., Brown, S.D., Somenahally, A.C., Johs, A.,

612 Hurt, R.A., Bailey, K.L., Elias, D.A., 2013. Mercury methylation by novel microorganisms from  
613 new environments. *Environ. Sci. Technol.* 47, 11810–11820. doi:10.1021/es403075t

614 Graham, A.M., Aiken, G.R., Gilmour, C.C., 2013. Effect of dissolved organic matter source and  
615 character on microbial Hg methylation in Hg-S-DOM solutions. *Environ. Sci. Technol.* 47, 5746–  
616 5754. doi:10.1021/es400414a

617 Grigal, D.F., 2002. Inputs and outputs of mercury from terrestrial watersheds: a review. *Environ. Rev.*  
618 10, 1–39. doi:10.1139/a01-013

619 Herrero Ortega, S., Catalán, N., Björn, E., Gröntoft, H., Hilmarsson, T.G., Bertilsson, S., Wu, P.,  
620 Bishop, K., Levanoni, O., Bravo, A.G., 2018. High methylmercury formation in ponds fueled by  
621 fresh humic and algal derived organic matter. *Limnol. Oceanogr.* 63, S44–S53.  
622 doi:10.1002/lno.10722

623 Huguet, A., Vacher, L., Relexans, S., Saubusse, S., Froidefond, J.M., Parlanti, E., 2009. Properties of  
624 fluorescent dissolved organic matter in the Gironde Estuary. *Org. Geochem.* 40, 706–719.  
625 doi:10.1016/j.orggeochem.2009.03.002

626 Johansson, K., Tyler, G., 2001. Impact of atmospheric long range transport of lead, mercury and  
627 cadmium on the Swedish forest environment. *Water, Air Soil Pollut. Focus* 1, 279–297.

628 Jones, R.J.A., Hiederer, R., Rusco, E., Montanarella, L., 2005. Estimating organic carbon in the soils  
629 of Europe for policy support. *Eur. J. Soil Sci.* 56, 655–671. doi:10.1111/j.1365-  
630 2389.2005.00728.x

631 Jonsson, S., Skjällberg, U., Nilsson, M.B., Lundberg, E., Andersson, A., Björn, E., 2014. Differentiated  
632 availability of geochemical mercury pools controls methylmercury levels in estuarine sediment  
633 and biota. *Nat. Commun.* 5, 4624. doi:10.1038/ncomms5624

634 Kirk, J.L., Louis, V.L.S., 2009. Multiyear total and methyl mercury exports from two major sub-arctic  
635 rivers draining into Hudson Bay, Canada. *Environ. Sci. Technol.* 43, 2254–2261.  
636 doi:10.1021/es803138z

637 Kocman, D., Wilson, S.J., Amos, H.M., Telmer, K.H., Steenhuisen, F., Sunderland, E.M., Mason, R.P.,  
638 Outridge, P., Horvat, M., 2017. Toward an assessment of the global inventory of present-day  
639 mercury releases to freshwater environments. *Int. J. Environ. Res. Public Health* 14.



640 doi:10.3390/ijerph14020138

641 Kothawala, D., Stedmon, C., Müller, R., Weyhenmeyer, G., Köhler, S., Tranvik, L.J., 2014. Controls of  
642 dissolved organic matter quality : evidence from a large-scale boreal lake survey. *Glob. Chang.*  
643 *Biol.* 20, 1101–1114. doi:10.1111/gcb.12488

644 Kothawala, D.N., von Wachenfeldt, E., Koehler, B., Tranvik, L.J., 2012. Selective loss and  
645 preservation of lake water dissolved organic matter fluorescence during long-term dark  
646 incubations. *Sci. Total Environ.* 433, 238–246. doi:10.1016/j.scitotenv.2012.06.029

647 Ledesma, J.L.J., Futter, M.N., Blackburn, M., Lidman, F., Grabs, T., Sponseller, R.A., Laudon, H.,  
648 Bishop, K.H., Köhler, S.J., 2018. Towards an Improved Conceptualization of Riparian Zones in  
649 Boreal Forest Headwaters. *Ecosystems* 21, 297–315. doi:10.1007/s10021-017-0149-5

650 Lee, D.S., Nemitz, E., Fowler, D., Kingdon, R.D., 2001. Modelling atmospheric mercury transport and  
651 deposition across Europe and the UK 35, 5455–5466.

652 Lescord, G.L., Emilson, E.J.S., Johnston, T.A., Branfireun, B.A., Gunn, J.M., 2018. Optical Properties  
653 of Dissolved Organic Matter and Their Relation to Mercury Concentrations in Water and Biota  
654 Across a Remote Freshwater Drainage Basin. *Environ. Sci. Technol.* acs.est.7b05348.  
655 doi:10.1021/acs.est.7b05348

656 Marvin-DiPasquale, M., Lutz, M.A., Brigham, M.E., Krabbenhoft, P., Aiken, G.R., Orem, W.H., Hall,  
657 B.D., Lutz, M.A., Brigham, M.E., Krabbenhoft, D.P., 2009. Production and Bed Sediment. 2. Pore  
658 Water Partitioning Mercury Cycling in Stream Methylmercury Production and Bed Sediment -  
659 Pore Water Partitioning. *Environ. Sci. Technol.* 43, 2726–2732. doi:10.1021/es802698v

660 Mason, R.P., Choi, A.L., Fitzgerald, W.F., Hammerschmidt, C.R., Lamborg, C.H., Soerensen, A.L.,  
661 Sunderland, E.M., 2012. Mercury biogeochemical cycling in the ocean and policy implications.  
662 *Environ. Res.* 119, 101–117. doi:10.1016/j.envres.2012.03.013

663 Mazrui, N.M., Jonsson, S., Thota, S., Zhao, J., Mason, R.P., 2016. Enhanced availability of mercury  
664 bound to dissolved organic matter for methylation in marine sediments. *Geochim. Cosmochim.*  
665 *Acta* 194, 153–162. doi:10.1016/j.gca.2016.08.019

666 McKnight, D.M., Boyer, E.W., Westerhoff, P.K., Doran, P.T., Kulbe, T., Andersen, D.T., 2001.  
667 Spectrofluorometric characterization of dissolved organic matter for indication of precursor

668 organic material and aromaticity. *Limnol. Oceanogr.* 46, 38–48. doi:10.4319/lo.2001.46.1.0038

669 Mevik, B.-H., Wehrens, R., 2007. The pls Package: Principal Component and Partial Least Squares  
670 Regression in R. *J. Stat. Software, Artic.* 18.

671 Monperrus, M., Krupp, E., Amouroux, D., Donard, O.F., Rodríguez Martín-Doimeadios, R., 2004.  
672 Potential and limits of speciated isotope-dilution analysis for metrology and assessing  
673 environmental reactivity. *TrAC Trends Anal. Chem.* 23, 261–272. doi:10.1016/S0165-  
674 9936(04)00313-9

675 Morel, F.M.M., Kraepiel, A.M.L., Amyot, M., 1998. The chemical cycle and bioaccumulation of  
676 mercury. *Annu. Rev. Ecol. Syst.* 29, 543–566. doi:10.1146/annurev.ecolsys.29.1.543

677 Moreno-Brush, M., Rydberg, J., Gamboa, N., Storch, I., Biester, H., 2016. Is mercury from small-scale  
678 gold mining prevalent in the southeastern Peruvian Amazon? *Environ. Pollut.* 218, 150–159.  
679 doi:10.1016/j.envpol.2016.08.038

680 Noh, S., Kim, J., Hur, J., Hong, Y., Han, S., 2017. Potential contributions of dissolved organic matter to  
681 monomethylmercury distributions in temperate reservoirs as revealed by fluorescence  
682 spectroscopy. *Environ. Sci. Pollut. Res.* doi:10.1007/s11356-017-0913-2

683 Ohno, T., 2002. Fluorescence inner-filtering correction for determining the humification index of  
684 dissolved organic matter. *Environ. Sci. Technol.* 36, 742–746. doi:10.1021/es0155276

685 Pak, K., Bartha, R., 1998. Mercury methylation and demethylation in anoxic lake sediments and by  
686 strictly anaerobic bacteria. *Appl. Environ. Microbiol.* 64, 1013–1017.

687 Parlanti, E., Wörz, K., Geoffroy, L., Lamotte, M., 2000. Dissolved organic matter fluorescence  
688 spectroscopy as a tool to estimate biological activity in a coastal zone submitted to  
689 anthropogenic inputs. *Org. Geochem.* 31, 1765–1781. doi:10.1016/S0146-6380(00)00124-8

690 Prechtel, A., Alewell, C., Armbruster, M., Bittersohl, J., Cullen, J.M., Evans, C.D., Helliwell, R.,  
691 Kopáček, J., Marchetto, A., Matzner, E., Meesenburg, H., Moldan, F., Moritz, K., Veselý, J.,  
692 Wright, R.F., 2001. Response of sulphur dynamics in European catchments to decreasing  
693 sulphate deposition. *Hydrol. Earth Syst. Sci.* 5, 311–326. doi:10.5194/hess-5-311-2001

694 R Core Team, 2016. R: A Language and Environment for Statistical Computing.

695 Rodriguez Martín-Doimeadios, R.C., Monperrus, M., Krupp, E., Amouroux, D., Donard, O.F.X., 2003.  
696 Using speciated isotope dilution with GC-inductively coupled plasma MS to determine and  
697 unravel the artificial formation of monomethylmercury in certified reference sediments. *Anal.*  
698 *Chem.* 75, 3202–3211. doi:10.1021/ac026411a

699 Schaefer, J.K., Kronberg, R.-M., Morel, F.M.M., Skyllberg, U., 2014. Detection of a key Hg methylation  
700 gene, *hgcA*, in wetland soils. *Environ. Microbiol. Rep.* 6, 441–447. doi:10.1111/1758-2229.12136

701 Schaefer, J.K., Morel, F.M.M., 2009. High methylation rates of mercury bound to cysteine by  
702 *Geobacter sulfurreducens*. *Nat. Geosci.* 2, 123–126. doi:10.1038/ngeo412

703 Schartup, A.T., Balcom, P.H., Soerensen, A.L., Gosnell, K.J., Calder, R.S.D., Mason, R.P.,  
704 Sunderland, E.M., 2015a. Freshwater discharges drive high levels of methylmercury in Arctic  
705 marine biota. *Proc. Natl. Acad. Sci.* 112, 11789–11794. doi:10.1073/pnas.1505541112

706 Schartup, A.T., Mason, R.P., Balcom, P.H., Hollweg, T.A., Chen, C.Y., 2013. Methylmercury  
707 production in estuarine sediments: Role of organic matter. *Environ. Sci. Technol.* 47, 695–700.

708 Schartup, A.T., Ndu, U.C., Balcom, P.H., Mason, R.P., Sunderland, E.M., 2015b. Contrasting effects  
709 of marine and terrestrially derived dissolved organic matter on mercury speciation and  
710 bioavailability in seawater. *Environ. Sci. Technol.* 5965–5972.

711 Schuster, P.F., Striegl, R.G., Aiken, G.R., Krabbenhoft, D.P., Dewild, J.F., Butler, K., Kamark, B.,  
712 Dornblaser, M., 2011. Mercury Export from the Yukon River Basin and Potential Response to a  
713 Changing Climate 9262–9267.

714 Shiklomanov, I.A., Rodda, J.C., 2004. *World Water Resources at the Beginning of the Twenty-First*  
715 *Century*, Cambridge University Press (UK).

716 Skyllberg, U., Bloom, P.R., Qian, J., Lin, C.-M., Bleam, W.F., 2006. Complexation of mercury(II) in soil  
717 organic matter: EXAFS evidence for linear two-coordination with reduced sulfur groups. *Environ.*  
718 *Sci. Technol.* 40, 4174–80.

719 Skyllberg, U., Qian, J., Frech, W., Xia, K., Bleam, W.F., 2003. Distribution of mercury, methyl mercury  
720 and organic sulphur species in soil, soil solution and stream of a boreal forest catchment.  
721 *Biogeochemistry* 64, 53–76. doi:10.1023/A:1024904502633

722 Soerensen, A.L., Schartup, A.T., Skrobonja, A., Björn, E., 2017. Organic matter drives high interannual

723 variability in methylmercury concentrations in a subarctic coastal sea. *Environ. Pollut.* 229, 531–  
724 538. doi:10.1016/j.envpol.2017.06.008

725 Sunderland, E.M., Mason, R.P., 2007. Human impacts on open ocean mercury concentrations. *Global*  
726 *Biogeochem. Cycles* 21. doi:10.1029/2006GB002876

727 Tjerngren, I., Karlsson, T., Björn, E., Skjellberg, U., 2012. Potential Hg methylation and MeHg  
728 demethylation rates related to the nutrient status of different boreal wetlands. *Biogeochemistry*  
729 108, 335–350. doi:10.1007/s10533-011-9603-1

730 Tsui, M.T.K., Finlay, J.C., Nater, E.A., 2009. Mercury bioaccumulation in a stream network. *Environ.*  
731 *Sci. Technol.* 43, 7016–7022. doi:10.1021/es901525w

732 Weishaar, J.L., Aiken, G.R., Bergamaschi, B. a, Fram, M.S., Fujii, R., Mopper, K., 2003. Evaluation of  
733 specific ultraviolet absorbance as an indicator of the chemical composition and reactivity of  
734 dissolved organic carbon. *Environ. Sci. Technol.* 37, 4702–4708.

735 WFD, 2008. Directive 2008/105/EC of 16 December 2008 on environmental quality standards in the  
736 field of water policy, amending and subsequently repealing Council Directives 82/176/EEC,  
737 83/513/EEC, 84/156/EEC, 84/491/ECC, 86/280/ECC and amending Directive 2000/60/EC, The  
738 European Parliament and the Council of the European Union. *Official Journal of the European*  
739 *Union.* doi:http://eur-lex.europa.eu/legal-content/EN/TXT/?uri=celex:32008L0105

740 Wiener, J.G., Krabbenhoft, D.P., Heinz, G.H., Scheuhammer, A.M., 2003. Ecotoxicology of mercury,  
741 in: Hoffman, D.J., Rattner, B.A., Jr., G.A.B., Jr., J.C. (Eds.), *Handbook of Ecotoxicology*, Second  
742 Edition. Lewis Publishers, Boca Raton, FL, pp. 409–463. doi:10.1201/9781420032505.ch16

743 Xia, K., Skjellberg, U., Bleam, W.F., Bloom, P.R., Nater, E.A., Helmke, P.A., 1999. X-ray absorption  
744 spectroscopic evidence for the complexation of Hg(II) by reduced sulfur in soil humic substances.  
745 *Env. Sci Technol* 33, 257–261.

746 Zsolnay, A., Baigar, E., Jimenez, M., Steinweg, B., Saccomandi, F., 1999. Differentiating with  
747 fluorescence spectroscopy the sources of dissolved organic matter in soils subjected to drying.  
748 *Chemosphere* 38, 45–50. doi:10.1016/S0045-6535(98)00166-0

749 **Acknowledgements**

750 This research was partially funded the Generalitat de Catalunya (Beatriu de Pinos BP-00385-  
751 2016) grants to AGB. DNK would like to acknowledge the Swedish National Science  
752 Foundation, Vetenskapsrådet (VR), for a starting grant. We thank the initiators of the first  
753 Collaborative European Freshwater Science Project for Young Researchers, the European  
754 Federation of Freshwater Sciences (EFFS) board, the European Fresh and Young  
755 Researchers (EFYR) and the representatives of the Fresh Blood for Fresh Water (FBFW)  
756 meetings. We also thank the seven national freshwater societies financing this project,  
757 namely the Deutsche Gesellschaft für Limnologie e.V. (DGL; Germany), Swiss Society for  
758 Hydrology and Limnology (Switzerland), Iberian Association for Limnology (AIL; Spain and  
759 Portugal), Italian Association of Oceanography and Limnology (Italy), Freshwater Biological  
760 Association (FBA; United Kingdom), French Limnological Association (AFL; France), Group  
761 of Austrian Members of SIL (SIL-Austria), as well as the Leibniz-Institute of Freshwater  
762 Ecology and Inland Fisheries for additional funds. CML was funded by the French Agency for  
763 Biodiversity (ONEMA-AFB, Action 13, “Colmatage, échange snappe-rivière et processus  
764 biogéochimiques). NC held a “Juan de la Cierva” postdoctoral grant (FJCI-2014-23064).  
765 Furthermore, we would like to thank Christian Noss, Elvira de Eyto, Brian C. Doyle, Adam  
766 Bednařík, Tea Basic, Georgina Busst, David Fletcher, Danny Sheath, Laura Barral-Fraga,  
767 Anna Freixa, Xisca Timoner, Marcus Klaus, Wiebke Schulz, Nina Pansch, Nikolay Simov and  
768 Lyubomir A. Kenderov for field assistance, Christoffer Bergvall for laboratory assistance and  
769 Viktor Rosenberg for the GIS map. David Kocman is acknowledged for the discussion on Hg  
770 loads in rivers.

771

#### 772 **Authors Contribution:**

773 AGB conceived the study. KA and PB coordinated the sampling and the teams involved in  
774 the study. AGB and ET performed GC–ICP–MS analyses of Hg with guidance from DA. DNK  
775 conducted fluorescence, spectrometry and DOC and TOC analyses. KA measured cations  
776 and anions. AGB and DNK conducted all statistical analyses. AGB, DNK and KA built the  
777 figures. AGB wrote the manuscript with significant assistance and comments from DNK, DA,

778 KA and PB. The rest of the co-authors carried out the sampling and contributed to the writing  
779 of the MS.

780

781

**Table 1.** Concentrations of inorganic-Hg (IHg), methylmercury (MeHg), total-Hg (THg = IHg+ MeHg) and the percentage of MeHg as THg (%MeHg) in the dissolved (filtered 0.45  $\mu\text{m}$ : <sub>D</sub>) and total (unfiltered: <sub>T</sub>) fraction of the studied streams.

Country	Stream ID	IHg <sub>D</sub>	MeHg <sub>D</sub>	THg <sub>D</sub>	MeHg <sub>D</sub>	IHg <sub>T</sub>	MeHg <sub>T</sub>	THg <sub>T</sub>	MeHg <sub>T</sub>
		ng L <sup>-1</sup>	pg L <sup>-1</sup>	ng L <sup>-1</sup>	%	ng L <sup>-1</sup>	pg L <sup>-1</sup>	ng L <sup>-1</sup>	%
Austria	AUT1_1	1.04	12	1.05	1.2	0.96	16	0.97	1.6
	AUT1_2	0.8	19	0.82	2.3	0.93	19	0.95	2.0
	AUT2_1	0.14	38	0.18	21.1	0.52	54	0.57	9.5
	AUT2_2	0.06	7	0.06	10.4	0.05	8	0.06	13.0
	AUT2_3	0.2	9	0.21	4.4	0.34	26	0.37	7.0
Bulgaria	BGR1_1	0.19	24	0.21	11.4	0.26	27	0.28	9.8
	BGR1_2	0.2	39	0.24	16.4	0.23	31	0.26	12.0
Czech Rep	CZE1_1	0.45	53	0.51	10.4	nd	nd	nd	nd
	CZE1_2	0.24	7	0.25	2.8	0.64	112	0.75	15.0
Germany	DEU1_1	0.17	67	0.23	28.7	0.3	12	0.31	3.9
	DEU1_2	0.14	36	0.17	20.8	0.34	60	0.4	15.0
	DEU2_1	0.14	10	0.15	6.7	0.4	56	0.46	12.2
	DEU2_2	0.11	26	0.13	20.1	0.55	52	0.6	8.7
Spain	ESP1_1	0.3	67	0.37	18.2	0.37	89	0.46	19.3
	ESP1_2	0.22	16	0.24	6.7	0.13	14	0.14	9.7
	ESP1_3	0.16	122	0.28	43.0	0.21	159	0.37	43.1
	ESP2_1	0.14	38	0.17	21.8	0.1	31	0.13	23.5
	ESP2_2	0.53	52	0.58	9.1	0.86	76	0.94	8.1
	ESP2_3	0.5	53	0.56	9.6	0.14	65	0.2	32.5
France	FRA1_1	0.47	15	0.48	3.1	0.49	15	0.51	2.9
	FRA1_2	0.39	17	0.41	4.2	0.86	20	0.88	2.2
Great Britain	GBR1_1	2.17	134	2.3	5.8	2.66	124	2.78	4.5
	GBR1_2	nd	nd	nd	nd	2.31	24	2.33	1.0
	GBR2_1	0.19	45	0.24	19.2	0.33	88	0.42	20.9
	GBR2_2	0.14	17	0.16	11.1	0.27	15	0.28	5.4
Portugal	PRT1_1	0.25	20	0.27	7.5	nd	nd	nd	nd
Sweden	SWE1_1	nd	nd	nd	nd	0.44	89	0.53	16.8
	SWE2_1	0.48	135	0.61	21.9	0.7	153	0.86	17.8
	SWE2_2	0.13	26	0.16	16.2	0.15	30	0.18	16.4

**Figure 1.** Distribution of the 29 sampling sites across nine European countries (see exact coordinates in Table 1).

**Figure 2.** Box and whisker plots of the distribution of a) total Hg (THg) in the unfiltered (total) fraction THg<sub>T</sub> and the filtered (dissolved) fraction THg<sub>D</sub> (n = 27), b) inorganic Hg (IHg) in the total IHg<sub>T</sub> and dissolved fractions IHg<sub>D</sub> (n = 27), c) monomethyl-Hg (MeHg) concentrations (n = 27) in the total (MeHg<sub>T</sub>) and dissolved (MeHg<sub>D</sub>) fractions and d) the %MeHg (n = 27). Boxes represent 25<sup>th</sup> (Q1) and 75<sup>th</sup> (Q3) percentile. The median is indicated within the box. The lower whiskers represent the lowest value within 1.5 IQR of the Q1, and the highest within 1.5 IQR of Q3.

**Figure 3.** Excitation-emission matrixes from sites presenting contrasted DOM characteristics a) and b) show samples with an eminently terrestrially derived DOM rich in humic peaks A and C. Plots c) and d) display samples with a marked microbial or algal signature as indicated by the high intensities in the peak T region. Intensities are shown in fluorescence units.

**Figure 4.** Significant ( $p < 0.05$ ,  $n=21$ ) correlations between THg<sub>T</sub>, MeHg<sub>T</sub>, the %MeHg<sub>T</sub>, DOM optical properties and stream characteristics. The ellipses have their eccentricity parametrically scaled to the Pearson correlation value (narrower ellipses represent higher correlation values). The orientation of the ellipses indicate negative (red) or positive (blue) correlations (scale on the right).

**Figure 5.** Principal Component Analysis (PCA) of European streams (n = 21) showing the explained variability arising from principle components 1 (PC1) and 2 (PC2) based on bulk optical characteristics of dissolved organic matter (DOM) and environmental variables including stream order and catchment cover, as presented in (a) the loadings plot, and (b) the corresponding scores plot displaying the 21 study sites.

**Figure 6.** Partial least squares loadings plot predicting the variability of total mercury (THg) concentration (Y variable, red) with several DOM optical properties and geographical and water chemistry characteristics as predictors (X variables). Fluorescence peaks are expressed as percentage of total fluorescence. Variables in blue were identified to be highly influential predictors of THg (with a variable influence on projection of  $\geq 1.0$ )


**Graphical Abstract.** Conceptual framework of (a) streams characterized by high inputs of terrestrial DOM and high concentrations of THg and (b) streams enriched in microbial/algal DOM depicting high MeHg formation (%MeHg).





Figure 1  
[Click here to download high resolution image](#)



Figure 2



**Figure 3**

**Figure 4**






Figure 6



**Electronic Supplementary Material (for online publication only)**

**[Click here to download Electronic Supplementary Material \(for online publication only\): EuroRun\\_Hg\\_SI\\_v4.docx](#)**

**Method Details (MethodsX)**

[Click here to download Method Details \(MethodsX\): MethodsX-Bravo et al\\_v3.docx.zip](#)