



HAL
open science

Colloidal stability of polyphenols in young red wine by Acacia gum: the major implication of arabinogalactan-proteins rich in proteins

Michael Nigen, Rafael Apolinar Valiente, Pascale Williams, Thierry Doco,
Nerea Iturmendi, Virginie Moine, Isabelle Jaouen, Christian Sanchez

► To cite this version:

Michael Nigen, Rafael Apolinar Valiente, Pascale Williams, Thierry Doco, Nerea Iturmendi, et al.. Colloidal stability of polyphenols in young red wine by Acacia gum: the major implication of arabinogalactan-proteins rich in proteins. 16. Food Colloids Conference, 2016, Wageningen, Netherlands. 2016, Food Colloids Conference. hal-01984265

HAL Id: hal-01984265

<https://hal.science/hal-01984265v1>

Submitted on 16 Jan 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Colloidal stability of polyphenols in young red wine by Acacia gum: The major implication of arabinogalactan-proteins rich in proteins

Michaël NIGEN¹, Rafael APOLINAR-VALIENTE¹, Pascale WILLIAMS², Thierry DOCO², Néréa ITURMENDI³, Virginie MOINE³, Isabelle JAOUEN⁴ and Christian SANCHEZ¹

¹ UMR IATE, UM-INRA-CIRAD-SupAgro, 2 place Pierre Viala, 34060 Montpellier, France

² UMR SPO, UM-INRA-SupAgro, 2 place Pierre Viala, 34060 Montpellier, France

³ BIOLAFFORT, 126 quai de ka Souys, 33100 Bordeaux, France

⁴ ALLAND & ROBERT, 9 rue Saintonge, 75003 Paris, France

Introduction

Acacia gum (GA) is a dried exudate obtained from *Acacia senegal* and *Acacia seyal* trees. GA macromolecules are highly branched heteropolysaccharides belonging to the arabinogalactan-protein (AGP) family. GA can be defined as a continuum of molecular species differing especially by their molar mass, charge density and hydrophobicity index. GA are composed of galactose, arabinose, rhamnose, glucuronic acid and a small amount of protein (0.5-3%). GA is widely used in food and non-food industries for its functional properties. In oenology, GA is used as additive to ensure the colloidal stability of young red wine precluding or minimizing the precipitation of polyphenols. In this study, we investigated the efficiency of GA and its molecular fractions towards polyphenols instabilities.

Materials & Methods

Acacia senegal gum was fractionated by Hydrophobic Interaction Chromatography (HIC) with the recovery of three fractions. They were named HIC-F1 (85-94% of GA), HIC-F2 (6-18% of GA) and HIC-F3 (1-3% of GA), and classified in that order according to a growing hydrophobic index. The stability of polyphenols in hydro-alcoholic matrix at pH 3.5 and 10° C was investigated in the presence of GA and its HIC fractions. The critical concentrations (C_{crit}) of GA and HIC fractions needed to stabilize the polyphenols were determined.

Results

Table 1. Biochemical composition of GA and its HIC fractions.

	Galactose (%)	Arabinose (%)	Rhamnose (%)	Glucuronic Acid (%)	Amino acids (mg.g ⁻¹)
GA	33.9	33.9	14.7	16.6	21.5
HIC-F1	36.9	29.7	14.1	18.1	5.6
HIC-F2	35.9	33.4	13.0	17.2	52.5
HIC-F3	33.0	37.1	12.9	16.4	105.9

- Sugar composition of GA and its three HIC fractions is similar.

- HIC fractions differ by their amino-acids content: HIC-F3 and HIC-F2 fractions contain ~20 and ~10 times more amino acids than HIC-F1 fraction.




Fig 1. Molar mass (M_w) distribution of GA and its HIC fractions.

- HIC-F1 contains mainly low M_w macromolecules ($M_w < 7.5 \times 10^5$ g.mol⁻¹).
- HIC-F2 and HIC-F3 contain mainly high M_w macromolecules ($M_w > 7.5 \times 10^5$ g.mol⁻¹).

Table 2. Critical concentration (C_{crit}) of GA and its HIC fractions towards polyphenols instability.

Samples	Critical Concentration (g.L ⁻¹)
GA	0.245 ± 0.006
HIC-F1	0.996 ± 0.090
HIC-F2	0.059 ± 0.002
HIC-F3	0.027 ± 0.003

- HIC fractions differ by their efficiency towards the polyphenols instability: HIC-F3 and HIC-F2 fractions are ~37 and ~17 times more efficient than HIC-F1 fraction.




Fig 2. Stability of polyphenols in presence of GA and its HIC fraction.

Conclusions

GA and its molecular HIC fractions prevent the polyphenols instability in hydro-alcoholic solutions. The HIC fractions can be classified according to their efficiency towards polyphenols instability as the following: HIC-F3 > HIC-F2 >> HIC-F1.

The polyphenols are preferentially stabilized by the high molar mass macromolecules rich in proteins (amino-acids) of *Acacia senegal* gum. The colloidal stabilizing properties of these macromolecules could be attributed to their remarkable physico-chemical properties such as their charge density and their hydrophobic behavior.