



HAL
open science

On Pillai's problem with Pell numbers and powers of 2

Mohand Ouamar Hernane, Florian Luca, Salah Eddine Rihane, Alain Togbé

► **To cite this version:**

Mohand Ouamar Hernane, Florian Luca, Salah Eddine Rihane, Alain Togbé. On Pillai's problem with Pell numbers and powers of 2. Hardy-Ramanujan Journal, 2019, Atelier Digit_Hum, pp.22 - 31. 10.46298/hrj.2019.5102 . hal-01984492

HAL Id: hal-01984492

<https://hal.science/hal-01984492>

Submitted on 17 Jan 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

On Pillai's problem with Pell numbers and powers of 2

Mohand Ouamar Hernane, Florian Luca, Salah Eddine Rihane and Alain Togbé

To the memory of S. Srinivasan

Abstract. In this paper, we find all integers c having at least two representations as a difference between a Pell number and a power of 2.

Keywords. Pell numbers, Linear form in logarithms, reduction method.

2010 Mathematics Subject Classification. primary 11B39; secondary 11J86.

1. Introduction

It is well-known that the sequence $\{P_n\}_{n \geq 0}$ of Pell numbers is defined by

$$P_0 = 0, \quad P_1 = 1, \quad P_{n+2} = 2P_{n+1} + P_n, \quad n \geq 0.$$

The first Pell numbers are

$$0, 1, 2, 5, 12, 29, 70, 169, 408, 985, 2378, 5741, 13860, \dots$$

In this paper, we are interested in the Diophantine equation

$$P_n - 2^m = c \tag{1.1}$$

for a fixed c and variable m and n . In particular, we are interested in those integers c admitting at least two representations as a difference between a Pell number and a power of 2. This is a variation of the equation

$$a^x - b^y = c, \tag{1.2}$$

in non-negative integers (x, y) where a, b, c are given fixed positive integers. The history of equation (1.2) is very rich and goes back to 1935 when Herschfeld [He35], [He36] studied the particular case $(a, b) = (2, 3)$. Extending Herschfeld's work, Pillai [Pi36], [Pi37] proved that if a, b are coprime positive integers then there exists $c_0(a, b)$ such that if $c > c_0(a, b)$ is an integer, then equation (1.2) has at most one positive integer solution (x, y) . Since then, variations of equation (1.2) has been intensively studied. Some recent results related to equation (1.1) are obtained by the third author and his collaborators in which they replaced Pell numbers P_n by the Fibonacci numbers F_n (see [DdLuR17]), Tribonacci numbers (see [BrLuYa17]), and k -generalized Fibonacci numbers (see [DdGLu17]).

The aim of this paper is to prove the following result.

Theorem 1.1. *The only integers c having at least two representations of the form $P_n - 2^m$ are $c \in \{-4, -3, -2, -1, 0, 1, 4\}$. Furthermore, all the representations of the above integers as $P_n - 2^m$ with integers $n \geq 0$ and $m \geq 0$ are given by*

$$\begin{aligned} -4 &= P_4 - 2^4 = P_0 - 2^2; \\ -3 &= P_5 - 2^5 = P_3 - 2^3 = P_1 - 2^1; \\ -2 &= P_2 - 2^2 = P_0 - 2^1; \\ -1 &= P_1 - 2^1 = P_0 - 2^0; \\ 0 &= P_2 - 2^1 = P_1 - 2^0; \\ 1 &= P_3 - 2^2 = P_2 - 2^0; \\ 4 &= P_4 - 2^3 = P_3 - 2^0. \end{aligned} \tag{1.3}$$

We organize this paper as follows. In Section 2, we recall some results useful for the proof of Theorem 1.1. The proof of Theorem 1.1 is done in the last section.

2. Auxiliary results

The next tools are related to the transcendental approach to solve Diophantine equations. For a non-zero algebraic number γ of degree d over \mathbb{Q} , whose minimal polynomial over \mathbb{Z} is $a \prod_{j=1}^d (X - \gamma^{(j)})$, we denote by

$$h(\gamma) = \frac{1}{d} \left(\log |a| + \sum_{j=1}^d \log \max \left(1, |\gamma^{(j)}| \right) \right)$$

the usual absolute logarithmic height of γ .

Lemma 2.1. *Let $\gamma_1, \dots, \gamma_s$ be a real algebraic numbers and let b_1, \dots, b_s be nonzero rational integer numbers. Let D be the degree of the number field $\mathbb{Q}(\gamma_1, \dots, \gamma_s)$ over \mathbb{Q} and let A_j be a positive real number satisfying*

$$A_j = \max\{Dh(\gamma_j), |\log \gamma_j|, 0.16\} \text{ for } j = 1, \dots, s.$$

Assume that

$$B \geq \max\{|b_1|, \dots, |b_s|\}.$$

If $\gamma_1^{b_1} \cdots \gamma_s^{b_s} \neq 1$, then

$$|\gamma_1^{b_1} \cdots \gamma_s^{b_s} - 1| \geq \exp(-C(s, D)(1 + \log B)A_1 \cdots A_s),$$

where $C(s, D) := 1.4 \cdot 30^{s+3} \cdot s^{4.5} \cdot D^2(1 + \log D)$.

Lemma 2.2. *Assume that τ and μ are real numbers and M is a positive integer. Let p/q be the convergent of the continued fraction of the irrational τ such that $q > 6M$, and let A, B, μ be some real numbers with $A > 0$ and $B > 1$. Let $\varepsilon = \|\mu q\| - M \cdot \|\tau q\|$, where $\|\cdot\|$ denotes the distance from the nearest integer. If $\varepsilon > 0$, then there is no solution of the inequality*

$$0 < m\tau - n + \mu < AB^{-k}$$

in positive integers m, n and k with

$$m \leq M \quad \text{and} \quad k \geq \frac{\log(Aq/\varepsilon)}{\log B}.$$

3. Proof of Theorem 1.1

Assume that there exist positive integers n, m, n_1, m_1 such that $(n, m) \neq (n_1, m_1)$, and

$$P_n - 2^m = P_{n_1} - 2^{m_1}.$$

Because of the symmetry, we can assume that $m \geq m_1$. If $m = m_1$, then $P_n = P_{n_1}$, so $(n, m) = (n_1, m_1)$, contradicting our assumption. Thus, $m > m_1$. Since

$$P_n - P_{n_1} = 2^m - 2^{m_1}, \tag{3.1}$$

and the right-hand side is positive, we get that the left-hand side is also positive and so $n > n_1$. Thus, $n \geq 2$ and $n_1 \geq 1$. We use the Binet formula

$$P_k = \frac{\alpha^k - \beta^k}{2\sqrt{2}} \quad \text{for all } k \geq 0,$$

where $(\alpha, \beta) := (1 + \sqrt{2}, 1 - \sqrt{2})$ are the roots of the characteristic equation of the Pell sequence. It is well-known that

$$\alpha^{k-2} \leq P_k \leq \alpha^{k-1} \quad \text{for all } k \geq 1.$$

Using the equation (3.1), we get

$$\alpha^{n-4} \leq P_{n-2} \leq P_n - P_{n_1} = 2^m - 2^{m_1} < 2^m, \quad (3.2a)$$

$$\alpha^{n-1} \geq P_n > P_n - P_{n_1} = 2^m - 2^{m_1} \geq 2^{m-1}, \quad (3.2b)$$

therefore

$$1 + \left(\frac{\log 2}{\log \alpha} \right) (m - 1) < n < \left(\frac{\log 2}{\log \alpha} \right) m + 4. \quad (3.3)$$

If $n < 150$, then $m \leq 200$. We ran a computer program for $2 \leq n_1 < n \leq 150$ and $1 \leq m_1 < m < 200$ and found only the solutions from list (1.3). From now, we assume that $n \geq 150$.

Note that the inequality (3.3) implies that $\frac{m}{2} < n$. So, to solve equation (3.1), we need an upper bound for n .

3.A. Bounding n

Using the Binet formula in the Diophantine equation (3.1), we get

$$\begin{aligned} \left| \frac{\alpha^n}{2\sqrt{2}} - 2^m \right| &= \left| \frac{\beta^n}{2\sqrt{2}} + \frac{\alpha^{n_1} - \beta^{n_1}}{2\sqrt{2}} - 2^{m_1} \right| \leq \frac{\alpha^{n_1} + 2}{2\sqrt{2}} + 2^{m_1} \\ &\leq \frac{\alpha^{n_1}}{\sqrt{2}} + 2^{m_1} < 2 \max\{\alpha^{n_1}, 2^{m_1}\}. \end{aligned}$$

Multiplying by 2^{-m} , using the relation (3.2a) and using the fact that $2 < \alpha$, we get

$$\left| (\sqrt{2})^{-1} \alpha^n 2^{-m-1} - 1 \right| < 2 \max \left\{ \frac{\alpha^{n_1}}{2^m}, 2^{m_1-m} \right\} < \max\{\alpha^{n_1-n+5}, 2^{m_1-m+1}\}. \quad (3.4)$$

For the left-hand side, we apply Theorem 2.1 with the data

$$s = 3, \quad \gamma_1 = \sqrt{2}, \quad \gamma_2 = \alpha, \quad \gamma_3 = 2, \quad b_1 = -1, \quad b_2 = n, \quad b_3 = -m - 1.$$

Throughout we work with $\mathbb{K} := \mathbb{Q}(\sqrt{2})$ with $D = 2$. Since $\max\{1, n, m + 1\} \leq 2n$ we take $B := 2n$. Furthermore, we take $A_1 := 2h(\gamma_1) = \log 2$, $A_2 := 2h(\gamma_2) = \log \alpha$, $A_3 := 2h(\gamma_3) = 2 \log 2$. Put

$$\Lambda = (\sqrt{2})^{-1} \alpha^n 2^{-m-1} - 1.$$

If $\Lambda = 0$, then $\alpha^{2n} \in \mathbb{Q}$, which is false. Thus, $\Lambda \neq 0$. Then, by Theorem 2.1, the left-hand side of (3.4) is bounded as

$$\log |\Lambda| > -1.4 \cdot 30^6 \cdot 3^{4.5} \cdot 2^2 (1 + \log 2)(1 + \log 2n)(\log 2)(\log \alpha)(2 \log 2).$$

Comparing with (3.4), we get

$$\min\{(n - n_1 - 5) \log \alpha, (m - m_1 - 1) \log 2\} < 8.3 \times 10^{11} (1 + \log 2n),$$

which gives

$$\min\{(n - n_1) \log \alpha, (m - m_1) \log 2\} < 8.4 \times 10^{11} (1 + \log 2n).$$

Now the argument splits into two cases.

Case 1. $\min\{(n - n_1) \log \alpha, (m - m_1) \log 2\} = (n - n_1) \log \alpha$.

In this case, we rewrite (3.1) as

$$\left| \left(\frac{\alpha^{n-n_1} - 1}{2\sqrt{2}} \right) \alpha^{n_1} - 2^m \right| = \left| \frac{\beta^n - \beta^{n_1}}{2\sqrt{2}} - 2^{m_1} \right| < 2^{m_1} + 1 \leq 2^{m_1+1},$$

which implies

$$\left| \left(\frac{\alpha^{n-n_1} - 1}{2\sqrt{2}} \right) \alpha^{n_1} 2^{-m} - 1 \right| < 2^{m_1-m+1}. \quad (3.5)$$

We put

$$\Lambda_1 = \left(\frac{\alpha^{n-n_1} - 1}{2\sqrt{2}} \right) \alpha^{n_1} 2^{-m} - 1.$$

Clearly, $\Lambda_1 \neq 0$, for if $\Lambda_1 = 0$, then

$$(\alpha^{n-n_1} - 1)\alpha^{n_1} = \sqrt{2} \times 2^m.$$

Conjugating this relation in \mathbb{K} , we get that

$$(\alpha^{n-n_1} - 1)\alpha^{n_1} = -(\beta^{n-n_1} - 1)\beta^{n_1}. \quad (3.6)$$

The absolute value of the left-hand side is at least $\alpha^n - \alpha^{n_1} \geq \alpha^{n-2} \geq \alpha^{188}$, while the absolute value of the right-hand side is at most $(|\beta|^{n-n_1} + 1)|\beta|^{n_1} < 2$, which is a contradiction.

We apply Theorem 2.1 by taking $s = 3$, and

$$\gamma_1 = \frac{\alpha^{n-n_1} - 1}{2\sqrt{2}}, \quad \gamma_2 = \alpha, \quad \gamma_3 = 2, \quad b_1 = 1, \quad b_2 = n_1, \quad b_3 = -m.$$

The minimal polynomial of γ_1 divides

$$8x^2 - 8P_{n-n_1}x - ((-1)^{n-n_1} + 1 - Q_{n-n_1}),$$

where $\{Q_k\}_{k \geq 0}$ is the Pell-Lucas companion sequence of the Pell sequence given by $Q_0 = 2$, $Q_1 = 1$, $Q_{k+2} = 2Q_{k+1} + Q_k$ for all $k \geq 0$, for which the Binet formula of its general term is

$$Q_k = \alpha^k + \beta^k \quad \text{for all } k \geq 0.$$

Thus, we obtain

$$\begin{aligned} h(\gamma_1) &\leq \frac{1}{2} \left(\log 8 + \log \left(\frac{\alpha^{n-n_1} + 1}{2\sqrt{2}} \right) \right) < \frac{1}{2} \log(4\sqrt{2}\alpha^{n-n_1}) \\ &< \frac{1}{2}(n - n_1 + 2) \log \alpha < 4.2 \times 10^{11}(1 + \log 2n). \end{aligned} \quad (3.7)$$

So, we can take $A_1 := 8.4 \times 10^{11}(1 + \log 2n)$. Further, as before, we can take $A_2 := \log \alpha$ and $A_3 := 2 \log 2$. Finally, since $\max\{1, n_1, m\} \leq 2n$, we can take $B := 2n$. We then get that

$$\log |\Lambda_1| > -1.4 \cdot 30^6 \cdot 3^{4.5} \cdot 2^2(1 + \log 2)(1 + \log 2n) \times (8.4 \times 10^{11}(1 + \log 2n))(\log \alpha)(2 \log 2).$$

Thus,

$$\log |\Lambda_1| > -1.1 \cdot 10^{24}(1 + \log 2n)^2.$$

Comparing this with (3.5), we get that

$$(m - m_1) \log 2 < 1.1 \cdot 10^{24}(1 + \log 2n)^2.$$

Case 2. $\min\{(n - n_1) \log \alpha, (m - m_1) \log 2\} = (m - m_1) \log 2$.

In this case, we rewrite (3.1) as

$$\left| \frac{\alpha^n}{2\sqrt{2}} - 2^{m_1}(2^{m-m_1} - 1) \right| = \left| \frac{\beta^n + \alpha^{n_1} - \beta^{n_1}}{2\sqrt{2}} \right| < \frac{\alpha^{n_1} + 2}{2\sqrt{2}} < \alpha^{n_1},$$

so

$$\begin{aligned} |(\sqrt{2}(2^{m-m_1} - 1))^{-1} \alpha^n 2^{-m_1-1} - 1| &< \frac{\alpha^{n_1}}{2^m - 2^{m_1}} \leq \frac{2\alpha^{n_1}}{2^m} \\ &\leq 2\alpha^{n_1-n+4} < \alpha^{n_1-n+5}. \end{aligned} \quad (3.8)$$

Let

$$\Lambda_2 = (\sqrt{2}(2^{m-m_1} - 1))^{-1} \alpha^n 2^{-m_1-1} - 1.$$

Clearly, $\Lambda_2 \neq 0$, for if $\Lambda_2 = 0$ implies $\alpha^{2n} \in \mathbb{Q}$, which is not possible.

We apply again Theorem 2.1. In this application, we take again $s = 3$, and

$$\gamma_1 = \sqrt{2}(2^{m-m_1} - 1), \quad \gamma_2 = \alpha, \quad \gamma_3 = 2, \quad b_1 = -1, \quad b_2 = n, \quad b_3 = -m_1 - 1.$$

The minimal polynomial of γ_1 is $x^2 - 2(2^{m-m_1} - 1)^2$. Thus

$$h(\gamma_1) = \log \left(\sqrt{2}(2^{m-m_1} - 1) \right) < (m - m_1 + 1) \log 2 < 8.5 \times 10^{11}(1 + \log 2n).$$

So, we can take $A_1 := 1.7 \times 10^{12}(1 + \log 2n)$. Further, as before, we can take $A_2 := \log \alpha$ and $A_3 := 2 \log 2$. Finally, since $\max\{1, n, m_1 + 1\} \leq 2n$, we can take $B := 2n$.

We then get that

$$\log |\Lambda_2| > -1.4 \cdot 30^6 \cdot 3^{4.5} \cdot 2^2(1 + \log 2)(1 + \log 2n) \times (1.7 \times 10^{12}(1 + \log 2n))(\log \alpha)(2 \log 2).$$

Thus,

$$\log |\Lambda_1| > -2.1 \cdot 10^{24}(1 + \log 2n)^2.$$

Comparing this with (3.8), we get that

$$(n - n_1) \log \alpha < 2.2 \cdot 10^{24}(1 + \log 2n)^2.$$

Thus, in both Case 1 and Case 2, we have

$$\begin{aligned} \min\{(n - n_1) \log \alpha, (m - m_1) \log 2\} &< 8.4 \cdot 10^{11}(1 + \log 2n) \\ \max\{(n - n_1) \log \alpha, (m - m_1) \log 2\} &< 2.2 \cdot 10^{24}(1 + \log 2n)^2. \end{aligned} \quad (3.9)$$

We now finally rewrite equation (3.1) as

$$\left| \frac{(\alpha^{n-n_1} - 1)}{2\sqrt{2}} \alpha^{n_1} - 2^{m_1}(2^{m-m_1} - 1) \right| = \left| \frac{\beta^n - \beta^{n_1}}{2\sqrt{2}} \right| < |\beta|^{n_1} = \frac{1}{\alpha^{n_1}}.$$

Dividing both sides by $2^m - 2^{m_1}$, we get

$$\begin{aligned} \left| \left(\frac{\alpha^{n-n_1} - 1}{\sqrt{2}(2^{m-m_1} - 1)} \right) \alpha^{n_1} 2^{-m_1-1} - 1 \right| &< \frac{1}{\alpha^{n_1}(2^m - 2^{m_1})} \leq \frac{2}{\alpha^{n_1} 2^m} \\ &\leq 2\alpha^{4-n-n_1} \leq \alpha^{4-n}, \end{aligned} \quad (3.10)$$

because $\alpha^{n_1} \geq \alpha > 2$. To find a lower-bound on the left-hand side, we use again Theorem 2.1 with $s = 3$, and

$$\gamma_1 = \frac{\alpha^{n-n_1} - 1}{\sqrt{2}(2^{m-m_1} - 1)}, \quad \gamma_2 = \alpha, \quad \gamma_3 = 2, \quad b_1 = 1, \quad b_2 = n_1, \quad b_3 = -m_1 - 1.$$

Using $h(x/y) = h(x) + h(y)$ for any two nonzero algebraic numbers x and y , we have

$$\begin{aligned} h(\gamma_1) &\leq h\left(\frac{\alpha^{n-n_1} - 1}{\sqrt{2}}\right) + h(2^{m-m_1} - 1) < \frac{1}{2}(n - n_1 + 2) \log \alpha + (m - m_1) \log 2 \\ &< 2.5 \cdot 10^{24}(1 + \log 2n)^2, \end{aligned}$$

where in the above chain of inequalities, we used the argument from (3.7) as well as the bound (3.9). So, we can take $A_1 := 5 \cdot 10^{24}(1 + \log 2n)^2$ and certainly $A_2 := \log \alpha$ and $A_3 := 2 \log 2$. We need to show that if we put

$$\Lambda_3 = \left(\frac{\alpha^{n-n_1} - 1}{\sqrt{2}(2^{m-m_1} - 1)}\right) \alpha^{n_1} 2^{-m_1-1} - 1,$$

then $\Lambda_3 \neq 0$. But $\Lambda_3 = 0$ leads to

$$(\alpha^{n-n_1} - 1) \alpha^{n_1} = \sqrt{2}(2^m - 2^{m_1}),$$

which upon conjugation in \mathbb{K} leads to (3.6), which is impossible. Thus, $\Lambda_3 \neq 0$. Theorem 2.1 gives

$$\log |\Lambda_3| > -1.4 \cdot 30^6 \cdot 3^{4.5} \cdot 2^2(1 + \log 2)(1 + \log 2n) \times (5 \times 10^{24}(1 + \log 2n)^2)(\log \alpha)(2 \log 2),$$

which together with (3.10) gives

$$(n - 4) < 6 \cdot 10^{36}(1 + \log 2n)^3,$$

leading to $n < 7 \cdot 10^{42}$.

3.B. Reducing n

We now need to reduce the above bound for n and to do so we make use of Lemma 2.2 several times. To begin with, we return to (3.4) and put

$$\Gamma := n \log \alpha - m \log 2 - \log(2\sqrt{2}).$$

For technical reasons we assume that $\min\{n - n_1, m - m_1\} \geq 20$. We go back to the inequalities for Λ , Λ_1 , Λ_2 . Since we assume that $\min\{n - n_1, m - m_1\} \geq 20$ we get $|e^\Gamma - 1| = |\Lambda| < \frac{1}{4}$. Hence, $|\Lambda| < \frac{1}{2}$ and since the inequality $|x| < 2|e^x - 1|$ holds for all $x \in (-\frac{1}{2}, \frac{1}{2})$, we get

$$|\Gamma| < 2 \max\{\alpha^{n_1-n+5}, 2^{m_1-m+1}\} \leq \max\{\alpha^{n_1-n+6}, 2^{m_1-m+2}\}.$$

Assume $\Gamma > 0$. We then have the inequality

$$\begin{aligned} 0 < n \left(\frac{\log \alpha}{\log 2}\right) - m - \frac{3}{2} &< \max\left\{\frac{\alpha^6}{\log 2} \alpha^{-(n-n_1)}, \frac{4}{\log 2} 2^{-(m-m_1)}\right\} \\ &< \max\{286 \alpha^{-(n-n_1)}, 6 \cdot 2^{-(m-m_1)}\}. \end{aligned}$$

We apply Lemma 2.2 with

$$\tau = \frac{\log \alpha}{\log 2}, \quad \mu = -\frac{3}{2}, \quad (A, B) = (286, \alpha) \text{ or } (6, 2).$$

Let $\tau = [a_0, a_1, \dots] = [1, 3, 1, 2, 6, 1, 2, 11, 2, 2, 1, \dots]$ be the continued fraction of τ . We choose $M := 7 \cdot 10^{42}$ and consider the 89-th convergent

$$\frac{p}{q} = \frac{p_{89}}{q_{89}} = \frac{348317200801236358620935022888502708006954949997}{273930475375768068118103206105181460963472890299}.$$

It satisfies $q = q_{89} > 6M$. Further, it yields $\varepsilon > 0.49$, and therefore either

$$n - n_1 \leq \frac{\log(286q/\varepsilon)}{\log \alpha} < 132, \quad \text{or} \quad m - m_1 \leq \frac{\log(6q/\varepsilon)}{\log 2} < 126.$$

In the case of $\Gamma < 0$, we consider the following inequality:

$$\begin{aligned} m \left(\frac{\log 2}{\log \alpha} \right) - n + \frac{\log(2\sqrt{2})}{\log \alpha} &< \max \left\{ \frac{\alpha^6}{\log \alpha} \alpha^{-(n-n_1)}, \frac{4}{\log \alpha} 2^{-(m-m_1)} \right\} \\ &< \max \{ 225 \alpha^{-(n-n_1)}, 5 \cdot 2^{-(m-m_1)} \}, \end{aligned}$$

instead and apply Lemma 2.2 with

$$\tau = \frac{\log 2}{\log \alpha}, \quad \mu = \frac{\log(2\sqrt{2})}{\log \alpha}, \quad (A, B) = (225, \alpha) \quad \text{or} \quad (5, 2).$$

Let $\tau = [a_0, a_1, \dots] = [0, 1, 3, 1, 2, 6, 1, 2, 11, 2, 2, 1, \dots]$ be the continued fraction of τ (note that the current τ is just the reciprocal of the previous τ). Again, we choose $M = 4.2 \cdot 10^{43}$, and in this case we consider the 90-th convergent

$$\frac{p}{q} = \frac{p_{90}}{q_{90}} = \frac{116053085442077720965142600370109382756030794793}{91268753856671834783420178148248093030868562298},$$

which satisfies $q = q_{90} > 6M$. This yields again $\varepsilon > 0.49$, and therefore either

$$n - n_1 \leq \frac{\log(225q/\varepsilon)}{\log \alpha} < 132, \quad \text{or} \quad m - m_1 \leq \frac{\log(5q/\varepsilon)}{\log 2} < 126.$$

These bounds agree with the bounds obtained in the case that $\Gamma > 0$. As a conclusion, we have either $n - n_1 \leq 131$ or $m - m_1 \leq 125$ whenever $\Gamma \neq 0$.

Now, we have to distinguish between the cases $n - n_1 \leq 131$ and $m - m_1 \leq 125$. First, let assume that $n - n_1 \leq 131$. In this case, we consider inequality (3.5) and assume that $m - m_1 \geq 20$. We put

$$\Gamma_1 = n_1 \log \alpha - m \log 2 + \log \left(\frac{\alpha^{n-n_1} - 1}{2\sqrt{2}} \right).$$

Then inequality (3.5) implies that

$$|\Gamma_1| < \frac{4}{2^{m-m_1}}.$$

If we further assume that $\Gamma_1 > 0$, we then get

$$0 < n_1 \left(\frac{\log \alpha}{\log 2} \right) - m + \frac{\log((\alpha^{n-n_1} - 1)/(2\sqrt{2}))}{\log 2} < \frac{4}{(\log 2)2^{m-m_1}} < \frac{6}{2^{m-m_1}}.$$

Again we apply Lemma 2.2 with the same τ as in the case when $\Gamma > 0$. We use the 89-th convergent $p/q = p_{89}/q_{89}$ of τ as before. But in this case we choose $(A, B) := (6, 2)$ and use

$$\mu_k = \frac{\log((\alpha^k - 1)/(2\sqrt{2}))}{\log 2},$$

instead of μ for each possible value of $k := n - n_1 \in [1, 2, \dots, 131]$. We have problems at $k \in \{1, 4\}$. We discard these values for now and will treat them later. For the remaining values of k , we get $\varepsilon > 0.0004$. Hence, by Lemma 2.2, we get

$$m - m_1 < \frac{\log(6q/0.0004)}{\log 2} < 172.$$

Thus, $n - n_1 \leq 131$ implies $m - m_1 \leq 171$, unless $n - n_1 \in \{1, 4\}$. A similar conclusion is reached if $\Gamma_1 < 0$ with the same two exceptions for $n - n_1 \in \{1, 4\}$. The reason we have a problem at $k \in \{1, 4\}$ is because

$$\frac{\alpha - 1}{2\sqrt{2}} = \frac{1}{2} \quad \text{and} \quad \frac{\alpha^4 - 1}{2\sqrt{2}} = 2\alpha^2.$$

So, $\Gamma_1 = n_1 \log \alpha - (m + 1) \log 2$, or $(n_1 + 2) \log \alpha - (m - 1) \log 2$ when $k = 1, 4$ respectively. Thus we get that

$$\left| \tau - \frac{m+1}{n_1} \right| < \frac{4}{2^{m-m_1} n_1}, \quad \text{or} \quad \left| \tau - \frac{m-1}{n_1+2} \right| < \frac{4}{2^{m-m_1} (n_1+2)},$$

respectively. Assume that $m - m_1 > 150$. Then $2^{m-m_1} > 8 \times (8 \times 10^{42}) > 8 \times (n_1 + 2)$, therefore

$$\frac{4}{2^{m-m_1} n_1} < \frac{1}{2n_1^2} \quad \text{and} \quad \frac{4}{2^{m-m_1} (n_1+2)} < \frac{1}{2(n_1+2)^2}.$$

By a criterion of Legendre, it follows that $(m+1)/n_1$ or $(m-1)/(n_1+2)$ are convergents of τ , respectively. So, say one of $(m+1)/n_1$ or $(m-1)/(n_1+2)$ is of the form p_k/q_k for some $k = 0, 1, \dots, 84$. Here, we use that $q_{84} > 8 \times 10^{42} > n_1 + 2$. Then

$$\frac{1}{(a_k + 2)q_k^2} < \left| \tau - \frac{p_k}{q_k} \right|.$$

Since $\max\{a_k : k = 0, \dots, 84\} = 100$, we get

$$\frac{1}{102q_k^2} < \frac{4}{2^{m-m_1}q_k} \quad \text{and} \quad q_k \text{ divides one of } \{n_1, n_1 + 2\}.$$

Thus,

$$2^{m-m_1} \leq 4 \times 102(n_1 + 2) < 4 \times 102 \times 8 \times 10^{42}$$

giving $m - m_1 \leq 152$.

Now let us turn to the case that $m - m_1 \leq 125$ and let us consider inequality (3.8). We put

$$\Gamma_2 = n \log \alpha - m_1 \log 2 + \log(1/(2\sqrt{2}(2^{m-m_1} - 1))),$$

and we assume that $n - n_1 \geq 20$. We then have

$$|\Gamma_2| < \frac{2\alpha^5}{\alpha^{n-n_1}}.$$

Assuming $\Gamma_2 > 0$, we get

$$0 < n \left(\frac{\log \alpha}{\log 2} \right) - m_1 + \frac{\log((1/(2\sqrt{2}(2^{m-m_1} - 1))))}{\log \alpha} < \frac{2\alpha^5}{(\log 2)\alpha^{n-n_1}} < \frac{237}{\alpha^{n-n_1}}.$$

We apply again Lemma 2.2 with the same τ , q , M , $(A, B) := (237, \alpha)$ and

$$\mu_k = \frac{\log((1/(2\sqrt{2}(2^k - 1))))}{\log 2} \quad \text{for } k = 1, 2, \dots, 125.$$

We get $\varepsilon > 0.003$, therefore

$$n - n_1 < \frac{\log(237q/0.003)}{\log \alpha} < 137.$$

A similar conclusion is reached when $\Gamma_2 < 0$. To conclude, we first get that either $n - n_1 \leq 131$ or $m - m_1 \leq 125$. If $n - n_1 \leq 131$, then $m - m_1 \leq 171$, and if $m - m_1 \leq 125$ then $n - n_1 \leq 136$. In conclusion, we always have $n - n_1 < 137$ and $m - m_1 < 172$.

Finally we go to (3.10). We put

$$\Gamma_3 = n_1 \log \alpha - m_1 \log 2 + \log \left(\frac{\alpha^{n-n_1} - 1}{2\sqrt{2}(2^{m-m_1} - 1)} \right).$$

Since $n \geq 200$, inequality (3.10) implies that

$$|\Gamma_3| < \frac{2}{\alpha^{n-4}} = \frac{2\alpha^4}{\alpha^n}.$$

Assume that $\Gamma_3 > 0$. Then

$$0 < n_1 \left(\frac{\log \alpha}{\log 2} \right) - m_1 + \frac{\log((\alpha^k - 1)/(2\sqrt{2}(2^l - 1)))}{\log 2} < \frac{2\alpha^4}{(\log 2)\alpha^n} < \frac{99}{\alpha^n},$$

where $(k, l) := (n - n_1, m - m_1)$. We apply again Lemma 2.2 with the same $\tau, M, q, (A, B) := (99, \alpha)$ and

$$\mu_{k,l} = \frac{\log((\alpha^k - 1)/(2\sqrt{2}(2^l - 1)))}{\log 2} \quad \text{for } 1 \leq k \leq 137, 1 \leq l \leq 171.$$

We have problem at $(k, l) = (1, 1), (4, 1)$ (as for the case of Γ_1) and additionally for $(k, l) = (8, 2)$ since

$$\frac{\alpha^8 - 1}{2\sqrt{2}(2^2 - 1)} = 4\alpha^4;$$

We discard the cases $(k, l) = (4, 1), (12, 1), (8, 2)$ for the time being. For the remaining ones, we get $\varepsilon > 0.0002$, so we obtain

$$n \leq \frac{\log(99q/0.0002)}{\log \alpha} < 139.$$

A similar conclusion is reached when $\Gamma_3 < 0$. Hence, $n < 150$. Now we look at the cases $(k, l) = (1, 1), (4, 1), (8, 2)$. The cases $(k, l) = (1, 1), (4, 1)$ can be treated as we did before when we showed that $n - n_1 \leq 137$ implies $m - m_1 \leq 171$. The case when $(k, l) = (8, 2)$ can be dealt with similarly as well. Namely, it gives

$$|(n_1 + 4)\tau - (m_1 + 2)| < \frac{99}{\alpha^n}.$$

Hence,

$$\left| \tau - \frac{m_1 + 2}{n_1 + 4} \right| < \frac{99}{(n_1 + 4)\alpha}. \quad (3.11)$$

Since $n \geq 150$, we have $\alpha^n > 2 \times 99 \times (8 \times 10^{42}) > 2 \times 99 \times (n_1 + 4)$, which shows that the right-hand side of inequality (3.11) is at most $2/(n_1 + 4)^2$. By Legendre's criterion, we get that $(m_1 + 2)/(n_1 + 4) = p_k/q_k$ for some $k \in [0, 84]$. We then get by an argument similar to a previous one that

$$\alpha^n \leq 99 \times 102 \times (8 \times 10^{42}),$$

giving $n \leq 123$. So, the conclusion is that $n < 150$ holds also in the case of the pair $(k, l) = (8, 2)$. However, this contradicts our working assumption that $n \geq 150$.

Theorem 1.1 is therefore proved.

Acknowledgements

We thank the referee for comments which improved the quality of this paper. F. L. was supported in part by grant CPRR160325161141 and an A-rated scientist award both from the NRF of South Africa and by grant no. 17-02804S of the Czech Granting Agency. F.L. worked on this paper during visits at the Mathematics Department to the Purdue University Northwest, USA and the Max Planck Institute for Mathematics, Bonn, Germany in February and March of 2018, respectively. He thanks these institutions for the generous support and hospitality.

References

- [BaDa69] A. Baker and H. Davenport, The equations $3x^2 - 2 = y^2$ and $8x^2 - 7 = z^2$, *Quart. J of Math. Ser.(2)* **20** (1969), 129–137.
- [BrLuYa17] J. J. Bravo, F. Luca, and K. Yazán, On a problem of Pillai with Tribonacci numbers and powers of 2, *Bull. Korean Math. Soc.* **54.3** (2017), 1069–1080.
- [ChPiZi17] K. C. Chim, I. Pink and V. Ziegler, On a variant of Pillai's problem, *Int. J. Number Theory* **13.7** (2017), 1711–1717.
- [ChPiZi16] K. C. Chim, I. Pink and V. Ziegler, On a variant of Pillai's problem II, *J. Number Theory* **183** (2018), 269–290.
- [DdLuR17] M. Ddamulira, F. Luca and M. Rakotomalala, On a problem of Pillai with Fibonacci numbers and powers of 2, *Proc. Math. Sci.* **127.3** (2017), 411–421.
- [DdGLu17] M. Ddamulira, C. A. Gómze and F. Luca, On a problem of Pillai with k -generalized Fibonacci numbers and powers of 2, (Preprint), 24 pages.
- [DuPe98] A. Dujella and A. Pethö, A generalization of a theorem of Baker and Davenport, *Quart. J. Math. Oxford Ser.* **49.3** (1998), 291–306.
- [He35] A. Herschfeld, The equation $2^x - 3^y = d$, *Bull. Amer. Math. Soc.* **41** (1935), 631.
- [He36] A. Herschfeld, The equation $2^x - 3^y = d$, *Bull. Amer. Math. Soc.* **42** (1936), 231–234.
- [Ma00] E. M. Matveev, An explicit lower bound for a homogeneous rational linear form in logarithms of algebraic numbers II, *Izv. Math.* **64.6** (2000), 1217–1269.
- [Mi04] P. Mihăilescu, Primary cyclotomic units and a proof of Catalan's conjecture, *J. Reine Angew. Math.* **572** (2004), 167–195.
- [Pi36] S. S. Pillai, On $a^x + b^y = c$, *J. Indian Math. Soc. (N.S.)* **2** (1936), 119–122.
- [Pi37] S. S. Pillai, A correction to the paper On $a^x + b^y = c$, *J. Indian Math. Soc. (N.S.)* **2** (1937), 215.

Mohand Ouamar Hernane

Université des Sciences et de la Technologie Houari-Boumediène (USTHB)
 Faculté de Mathématiques, Laboratoire d'Algèbre et Théorie des Nombres, BP 32
 16111 Bab-Ezzouar Alger, Algérie
e-mail: mhernane@usthb.dz

Florian Luca

School of Mathematics, University of the Witwatersrand
 Private Bag 3, Wits 2050
 Johannesburg, South Africa
 AND
 Max Planck Institute for Mathematics, Vivatsgasse 7
 53111 Bonn, Germany
 AND
 Department of Mathematics, Faculty of Sciences
 University of Ostrava, 30 Dubna 22
 701 03 Ostrava 1, Czech Republic *e-mail*: Florian.Luca@wits.ac.za

Salah Eddine Rihane

Université des Sciences et de la Technologie Houari-Boumediène (USTHB)
 Faculté de Mathématiques, Laboratoire d'Algèbre et Théorie des Nombres, BP 32
 16111 Bab-Ezzouar Alger, Algérie
e-mail: salahrihane@hotmail.fr

Alain Togbé

Department of Mathematics, Statistics, and Computer Science
 Purdue University Northwest, 1401 S, U.S. 421
 Westville IN 46391, USA
e-mail: atogbe@pnw.edu