

# Explicit $H^1$ -Estimate for the solution of the Lamé system with mixed boundary conditions.

Djamal AIT-AKLI<sup>1</sup>, Abdelkader MERAKEB<sup>2</sup>

February 1, 2020

## Abstract

In this paper we consider the Lamé system on a polygonal convex domain with mixed boundary conditions of Dirichlet-Neumann type. An explicit  $L^2$  norm estimate for the gradient of the solution of this problem is established. This leads to an explicit bound of the  $H^1$  norm of this solution. Note that the obtained upper-bound is not optimal.

**keywords:** Lamé system; Korn's inequality; Poincaré's inequality; inequality of trace; explicit estimates.

**AMS subject classification:** 35J57, 74B05

<sup>1 2</sup>

## 1 Introduction

The static equilibrium of a deformable structure occupying a domain  $\Omega$  subset of  $\mathbb{R}^2$  is governed by the Lamé linear elasto-static system, see [5]. In this paper, we restrict the study to a convex domain  $\Omega$  whose boundary has a polygonal shape that posses  $m + 1$  edges with  $m \geq 2$ . We denote  $\Gamma = \cup \Gamma_i$  its boundary and  $d(\Omega)$  its diameter. This system is given by

$$\begin{cases} Lu = f & \text{a.e in } \Omega, \\ \sigma \cdot \vec{n}_i = g_i & \text{on } (\Gamma - \Gamma_0) \cap \Gamma_i, \quad 1 \leq i \leq m \\ u = 0 & \text{on } \Gamma_0. \end{cases} \quad (1)$$

We need to assume that the components  $\Gamma_i$  of the boundary  $\Gamma$  fulfill a condition similar to  $(H_2)$  of Theorem 2.3 stated in the paper [9]. Actually, for our purpose, a stronger condition is needed and it is formulated in (5) below. The vector function  $u = (u^1, u^2)$  satisfying the system (8) describes a displacement in the plane. In this model, we impose a homogeneous Dirichlet

---

<sup>1</sup>L2CSP, UMMTO.15000, Tizi-Ouzou, ALGERIA. (d.aitakli@yahoo.com)

<sup>2</sup>L2CSP, UMMTO. 15000, Tizi-Ouzou, ALGERIA.(merakeb\_kader@yahoo.fr)

condition on  $\Gamma_0$  and a Neumann condition on the rest of the boundary. The equality on the boundary is understood in the sense of the trace. We denote  $L$  the Lamé operator defined by:

$$Lu := -\operatorname{div} \sigma(u) = -\operatorname{div}[2\mu\varepsilon(u) + \lambda \operatorname{Tr} \varepsilon(u) Id] \quad (2)$$

We assume the data functions  $f$  and  $g$  at the right hand side to satisfy  $f \in [L^2(\Omega)]^2$  and  $g \in [H^{\frac{1}{2}}(\Gamma - \Gamma_0)]^2$ . The vector  $\vec{n}_i$  represents the outside normal to  $\Gamma_i$ . We write  $\mu$  and  $\lambda$  the Lamé's coefficients. We place ourselves in the isotropic framework, the deformation tensor  $\varepsilon$  is defined by

$$\varepsilon(u) = \frac{1}{2}(\nabla u + \nabla^t u), \quad (3)$$

The weak form of problem (1) is (see [2], [5]):

find  $u \in V$ ;  $\forall v \in V$

$$\int_{\Omega} 2\mu\varepsilon(u)\varepsilon(v) + \lambda \operatorname{div} u \operatorname{div} v \, dx = \int_{\Omega} f v \, dx + \int_{\Gamma - \Gamma_0} g v \, d\sigma(x) \quad (4)$$

where

$$V = \{v \in [H^1(\Omega)]^2; \quad v = 0 \quad \text{on} \quad \Gamma_0\}$$

The problem of existence and uniqueness in  $V$  of the solution of (4) is classic, (see [2]).

If we denote  $\theta$  the interior angle between the edges  $\Gamma_j$  and  $\Gamma_k$  such that  $\bar{\Gamma}_j \cap \bar{\Gamma}_k \neq \emptyset$  and if we denote  $\gamma$  the interior angle between the Neumann part of the boundary  $\Gamma_N$  and the Dirichlet part of the boundary  $\Gamma_D$  such that  $\Gamma_N \cap \Gamma_D \neq \emptyset$ , then we impose

$$\theta \leq \pi, \quad \gamma \leq \pi. \quad (5)$$

The reason behind this assumption on the boundary is to get a better regularity of the solution of the weak problem (4). Precisely in that case we have, following [9],  $u \in [H^{\frac{3}{2}+\nu}(\Omega)]^2$  for some positive  $\nu$ , which implies in particular, using the appropriate Sobolev embedding, see [1], that  $u \in [C^{0, \frac{1}{2}+\nu}(\bar{\Omega})]^2$  i.e.  $u$  is  $(\frac{1}{2} + \nu)$ -holder continuous. Let us denote

$$\|\varepsilon(u)\|_{0,\Omega} := \left( \int_{\Omega} \varepsilon(u)\varepsilon(u) \, dx \right)^{\frac{1}{2}}, \quad \|\nabla u\|_{0,\Omega} := \left( \int_{\Omega} |\nabla u^1|^2 + |\nabla u^2|^2 \, dx \right)^{\frac{1}{2}}.$$

By using the second Korn inequality, see [7], the trace and the Poincaré's inequalities, one easily gets from (4) the following estimate

$$\|\nabla u\|_{0,\Omega} \leq \frac{1}{c_k} \frac{1}{2\mu} (c_p \|f\|_{0,\Omega} + c_{p,t} \|g\|_{\frac{1}{2}, \Gamma - \Gamma_0}), \quad (6)$$

where  $c_{p,t}$  is a constant that depends of Poincaré constant and the constant of trace inequality.  $c_k$  is the constant of the Korn's inequality. Note that the value of the constant  $c_k$  and  $c_{p,t}$  appearing in (6) are unknown and can not be explicitly lower-bounded in the general case. We propose to determine explicitly these constants. The main result of this work is stated in the following theorem:

**Theorem 1.** *The unique weak solution  $u$  of (4) on the polygonal domain  $\Omega$  admits the explicit upper bound*

$$\|\nabla u\|_{0,\Omega} \leq \frac{1}{\sqrt{\mu}} [1 + d(\Omega)\|f\|_{0,\Omega} + 8(|\Gamma| + \|g\|_{0,\Gamma}^2)], \quad (7)$$

where  $d(\Omega)$  represent the diameter of  $\Omega$  and  $|\Gamma|$  the measure of the boundary of  $\Omega$ .

Estimate (7) is similar to estimate (6) in the sense that the constants appearing there are the same. Before demonstrating this theorem, it is useful to go through some remarks and results. These auxiliary results are needed in order to adequately get a decomposition of the solution of the main problem, which belongs to  $H^1(\Omega)$ , into functions that are still in  $H^{\frac{1}{2}}(\Gamma)$ . Denote  $x_i$ , for  $1 \leq i \leq m$ , the vertex of the polygon that connects  $\Gamma_{i-1}$  with  $\Gamma_i$  and  $x_0$  the one that connects  $\Gamma_m$  to  $\Gamma_0$ . Define the auxiliary function  $u_\epsilon \in H^1(\Omega)$  as the unique solution to the following Dirichlet problem

$$\begin{cases} Lu_\epsilon = f & \text{a.e in } \Omega, \\ u_\epsilon = u_\epsilon^d & \text{on } \Gamma. \end{cases} \quad (8)$$

Where  $u_\epsilon^d$  is the trace on the boundary  $\Gamma$  of the function

$$\phi_\epsilon(x)u(x) \quad (9)$$

such that if  $\epsilon < \frac{|\Gamma_i|}{2} \forall i$ ,  $0 \leq i \leq m$  then  $\phi_\epsilon$  is defined by

$$\begin{cases} \phi_\epsilon(x) & = 0, & \|x - x_i\| \leq \epsilon^2, & 0 \leq i \leq m; \\ \phi_\epsilon(x) & = \exp\left[-\frac{\epsilon^{\frac{1}{2}}(\epsilon - \|x - x_i\|)}{\|x - x_i\| - \epsilon^2}\right], & \epsilon^2 < \|x - x_i\| < \epsilon, & 0 \leq i \leq m; \\ \phi_\epsilon(x) & = 1, & \epsilon \leq \|x - x_i\|, & 0 \leq i \leq m, \end{cases}$$

let us denote

$$D_{i,\epsilon} := \{x \in \mathbb{R}^2 \text{ such that } \|x - x_i\| < \epsilon^2\}.$$

We easily see that  $\phi_\epsilon \in C^0(\overline{\Omega})$ , consequently, there will be no jump when passing to the distributional derivative and thus  $\nabla u_\epsilon \in L^2(\Omega)$  i.e.  $u_\epsilon \in H^1(\Omega)$ . It is shown, using Lebesgue's dominated convergence theorem for instance, that  $\|\phi_\epsilon - 1\|_{0,\Gamma_i} \rightarrow 0$  i.e. we have convergence in  $L^2$  along the edge  $\Gamma_i$ . The functions  $\phi_\epsilon$  are identically zero on a small neighborhood of the respective vertices of the polygon.

In the sequel, we denote  $u_\epsilon$  the vector-valued function  $u_\epsilon = (u_\epsilon^1, u_\epsilon^2)$ .

## 2 Weak problem for $u_\epsilon$ and approximation results

First of all, we construct the weak problem verified by the approximating function  $u_\epsilon$ . With the approximating displacement  $u_\epsilon \in V$  is associated the approximating stress tensor

$$\sigma_\epsilon := 2\mu\varepsilon(u_\epsilon) + \lambda \operatorname{Tr} \varepsilon(u_\epsilon)I, \quad (10)$$

since  $Lu_\epsilon = \operatorname{div} \sigma_\epsilon = f$ , then  $\sigma_\epsilon \in [H(\operatorname{div})(\Omega)]^{2 \times 2}$ . For a fixed  $\epsilon$ , by density of the regular functions in the space  $H(\operatorname{div})(\Omega)$ , there exists  $\sigma_\epsilon^n \in [C^\infty(\bar{\Omega})]^{2 \times 2}$  such that  $\sigma_\epsilon^n \rightarrow \sigma_\epsilon$  in  $[H(\operatorname{div})(\Omega)]^{2 \times 2}$ . This means

$$\|\sigma_\epsilon^n - \sigma_\epsilon\|_{\operatorname{div}, \Omega} := \|\operatorname{div} \sigma_\epsilon^n - \operatorname{div} \sigma_\epsilon\|_{0, \Omega} + \|\sigma_\epsilon^n - \sigma_\epsilon\|_{0, \Omega} \rightarrow 0 \quad (11)$$

when  $n \rightarrow \infty$ . We pose  $\operatorname{div} \sigma_\epsilon^n = f^n$ , then integrating by part against a test function  $v \in [C^\infty(\bar{\Omega})]^2 \cap V$  yields the following

$$\int_{\Omega} \sigma_\epsilon^n \nabla v = \int_{\Omega} f^n v + \int_{\Gamma} \sigma_\epsilon^n \cdot \vec{n} v d\sigma.$$

Passing to the limit in  $n$  using (11), we find  $\forall v \in [C^\infty(\bar{\Omega})]^2 \cap V$

$$\int_{\Omega} \sigma_\epsilon \nabla v = \int_{\Omega} f v + \langle \sigma_\epsilon \cdot \vec{n}, v \rangle_{[H^{\frac{1}{2}}]^\prime(\Gamma - \Gamma_0) \times [H^{\frac{1}{2}}]^\prime(\Gamma - \Gamma_0)},$$

where  $\sigma_\epsilon \cdot \vec{n} =: g^\epsilon \in [H^{\frac{1}{2}}(\Gamma - \Gamma_0)]^\prime$  is the image of the normal component  $\sigma_\epsilon$  by the trace operator on  $\Gamma$ . Since, following the main result in [4],  $[C^\infty(\bar{\Omega})]^2 \cap V$  is a dense subset of  $V \subset H^1(\Omega)$ , then, according to the definition (3) and the expression (10), the function  $u_\epsilon$  satisfy

$\forall v \in V$ ,

$$\int_{\Omega} 2\mu\varepsilon(u_\epsilon)\varepsilon(v) + \int_{\Omega} \lambda \operatorname{div} u_\epsilon \operatorname{div} v = \int_{\Omega} f v + \langle g^\epsilon, v \rangle_{[H^{\frac{1}{2}}]^\prime(\Gamma - \Gamma_0) \times [H^{\frac{1}{2}}]^\prime(\Gamma - \Gamma_0)}; \quad (12)$$

this is the weak problem satisfied by the approximating function  $u_\epsilon$ .

Let us recall, (see [8]), that the  $H^{\frac{1}{2}}$ -norm in one dimension on  $\Gamma_i$  is defined by:

$$\|u\|_{\frac{1}{2}, \Gamma_i} := (\|u\|_{0, \Gamma_i}^2 + \int_{\Gamma_i} \int_{\Gamma_i} \frac{|u(x) - u(y)|^2}{\|x - y\|^2} dx dy)^{\frac{1}{2}}.$$

**Remark 2.** For any sufficiently small  $\epsilon > 0$ , it is possible to cover  $\Omega$  with a collection of open sets  $(W_j^\epsilon)_j$  such that for each  $j$ ,  $W_j^\epsilon \cap \Gamma$  is either empty or equals one of the following subsets: for some  $0 \leq i \leq m - 1$

$$1) \quad \Gamma_i^{1, \epsilon} := \{x \in \Gamma_i; \quad 0 < \|x - x_i\| < 2\epsilon\};$$

- 2)  $\Gamma_i^{2,\epsilon} := \{x \in \Gamma_i; \quad 0 < \|x - x_{i+1}\| < 2\epsilon\};$   
 3)  $\Gamma_i^{3,\epsilon} := \{x \in \Gamma_i; \quad \|x - x_i\| > \frac{3}{2}\epsilon \quad \text{and} \quad \|x - x_{i+1}\| > \frac{3}{2}\epsilon\};$   
 4)  $\Gamma_i^{4,\epsilon} = \{x \in \Gamma_i \cup \Gamma_{i+1}; \quad \|x - x_{i+1}\| < \epsilon^2\};$

and for  $i = m$

- 1)  $\Gamma_m^{1,\epsilon} := \{x \in \Gamma_m; \quad 0 < \|x - x_m\| < 2\epsilon\};$   
 2)  $\Gamma_m^{2,\epsilon} := \{x \in \Gamma_m; \quad 0 < \|x - x_0\| < 2\epsilon\};$   
 3)  $\Gamma_m^{3,\epsilon} := \{x \in \Gamma_m; \quad \|x - x_m\| > \frac{3}{2}\epsilon \quad \text{and} \quad \|x - x_0\| > \frac{3}{2}\epsilon\};$   
 4)  $\Gamma_m^{4,\epsilon} = \{x \in \Gamma_m \cup \Gamma_0; \quad \|x - x_0\| < \epsilon^2\}$

let  $(\vartheta_j^\epsilon)_j$ , with  $\text{supp } \vartheta_j^\epsilon \subset W_j^\epsilon$ , a  $C^1$ -partition of unity with respect to this cover; since  $\vartheta_j^\epsilon \in H^{\frac{1}{2}}(\Gamma)$  then

$$\begin{aligned}
 \|\phi_\epsilon - 1\|_{\frac{1}{2},\Gamma} &= \|(\phi_\epsilon - 1) \sum_j \vartheta_j^\epsilon\|_{\frac{1}{2},\Gamma} \\
 &\leq \sum_{i=0}^m \|(\phi_\epsilon - 1) \sum_{j, \text{supp } \vartheta_j^\epsilon \subset \Gamma_i} \vartheta_j^\epsilon\|_{\frac{1}{2},\Gamma_i} + \sum_{i=0}^m \|(\phi_\epsilon - 1)\|_{\frac{1}{2},\Gamma_i^{4,\epsilon}} \\
 &\leq \sum_{i=0}^m \|\phi_\epsilon - 1\|_{\frac{1}{2},\Gamma_i} + (m+1) \|(\phi_\epsilon - 1)\|_{\frac{1}{2},\Gamma_0^{4,\epsilon}} \\
 &= \sum_{i=0}^m \|\phi_\epsilon - 1\|_{\frac{1}{2},\Gamma_i} + (m+1) \|(\phi_\epsilon - 1)\|_{0,\Gamma_0^{4,\epsilon}} \tag{13}
 \end{aligned}$$

So, using the definition and symmetry of  $\phi_\epsilon$ , we get for all  $0 \leq i \leq m$

$$\begin{aligned}
 \|\phi_\epsilon - 1\|_{\frac{1}{2},\Gamma_i} &= \|(\phi_\epsilon - 1) \sum_{j, \text{supp } \vartheta_j^\epsilon \subset \Gamma_i} \vartheta_j^\epsilon\|_{\frac{1}{2},\Gamma_i} \\
 &\leq \|(\phi_\epsilon - 1) \sum_{j, \text{supp } \vartheta_j^\epsilon \subset \Gamma_i^{1,\epsilon}} \vartheta_j^\epsilon\|_{\frac{1}{2},\Gamma_i} + \|(\phi_\epsilon - 1) \sum_{j, \text{supp } \vartheta_j^\epsilon \subset \Gamma_i^{2,\epsilon}} \vartheta_j^\epsilon\|_{\frac{1}{2},\Gamma_i} \\
 &\quad + \|(\phi_\epsilon - 1) \sum_{j, \text{supp } \vartheta_j^\epsilon \subset \Gamma_i^{3,\epsilon}} \vartheta_j^\epsilon\|_{\frac{1}{2},\Gamma_i} + 2 \|(\phi_\epsilon - 1)\|_{\frac{1}{2},\Gamma_i^{4,\epsilon}} \\
 &\leq \sum_{j=1}^3 \|\phi_\epsilon - 1\|_{\frac{1}{2},\Gamma_i^{j,\epsilon}} + 2 \|\phi_\epsilon - 1\|_{\frac{1}{2},\Gamma_i^{4,\epsilon}} \\
 &= 2 \|\phi_\epsilon - 1\|_{\frac{1}{2},\Gamma_i^{1,\epsilon}} + 2 \|\phi_\epsilon - 1\|_{\frac{1}{2},\Gamma_i^{4,\epsilon}}
 \end{aligned}$$

thus we have

$$\begin{aligned}
 \|\phi_\epsilon - 1\|_{\frac{1}{2},\Gamma_i} &\leq 2 \|\phi_\epsilon - 1\|_{0,\Gamma_i^{1,\epsilon}} + 2 \left( \int_{\Gamma_i^{1,\epsilon}} \int_{\Gamma_i^{1,\epsilon}} \frac{|\phi_\epsilon(x) - \phi_\epsilon(y)|^2}{\|x - y\|^2} dx dy \right)^{\frac{1}{2}} \\
 &\quad + 2 \|(\phi_\epsilon - 1)\|_{0,\Gamma_i^{4,\epsilon}}. \tag{14}
 \end{aligned}$$

**Lemma 3.** *The functions  $\phi_\epsilon$  admit the following limits*

$$\text{for all } 0 \leq i \leq m \quad \|\phi_\epsilon - 1\|_{\frac{1}{2}, \Gamma_i} \rightarrow 0 \quad \text{as } \epsilon \rightarrow 0$$

*Proof.* If we choose the vertex point  $x_i$  as the origin of the  $\mathbb{R}^2$ -orthonormal coordinate system such that  $\Gamma_i$  is supported by the positive half  $x$ -axis then the abscisses of  $x \in \Gamma_i^{1, \epsilon} \equiv ]0, 2\epsilon[$  satisfy

$$\|x - x_i\| = |x| = x.$$

the  $H^{\frac{1}{2}}$ -semi-norm of  $\phi_\epsilon$  on  $\Gamma_i$  writes

$$|\phi_\epsilon - 1|_{\frac{1}{2}, \Gamma_i}^2 := \int_{\Gamma_i} \int_{\Gamma_i} \frac{|\phi_\epsilon(x) - \phi_\epsilon(y)|^2}{\|x - y\|^2} dx dy. \quad (15)$$

it yields by using the definition of  $\phi_\epsilon$

$$|\phi_\epsilon - 1|_{\frac{1}{2}, \Gamma_i}^2 \leq 2 \int_0^\epsilon \int_0^\epsilon \frac{|\phi_\epsilon(x) - \phi_\epsilon(y)|^2}{|x - y|^2} dx dy. \quad (16)$$

Consider the decomposition of (16) into four partial double integrals

$$1) \int_0^{\epsilon^2} \int_0^{\epsilon^2} \frac{|\phi_\epsilon(x) - \phi_\epsilon(y)|^2}{|x - y|^2} dx dy = 0,$$

this is obvious.

$$2) \int_{\epsilon^2}^\epsilon \int_{\epsilon^2}^\epsilon \frac{|\phi_\epsilon(x) - \phi_\epsilon(y)|^2}{|x - y|^2} dx dy \leq \int_{\epsilon^2}^\epsilon \int_{\epsilon^2}^\epsilon \frac{|\exp[-\frac{\epsilon^{\frac{1}{2}}(\epsilon-x)}{(x-\epsilon^2)}] - \exp[-\frac{\epsilon^{\frac{1}{2}}(\epsilon-y)}{(y-\epsilon^2)}]|^2}{|x - y|^2} dx dy.$$

The function  $F(x) := \exp[-\frac{\epsilon^{\frac{1}{2}}(\epsilon-x)}{(x-\epsilon^2)}]$  is  $C^1(] \epsilon^2, \epsilon[)$  and thus lipschitz. We have, using the fact that  $x \rightarrow F'(x)$  is increasing on  $[\epsilon^2, \frac{\epsilon}{2}]$ , that

$$|F'(x)| \leq \frac{\epsilon^{\frac{1}{2}}(1-\epsilon)}{\epsilon(\frac{1}{2}-\epsilon)^2} \exp(\frac{-\epsilon^{\frac{1}{2}}}{\frac{1}{2}-\epsilon}) =: L_1,$$

$\forall x \in [\epsilon^2, \frac{\epsilon}{2}]$ . On the other hand

$$|F'(x)| \leq \frac{\epsilon^{\frac{1}{2}}}{\epsilon(1-\epsilon)} \exp(\frac{2}{1-\epsilon}) =: L_2,$$

$\forall x \in [\frac{\epsilon}{2}, \epsilon]$ . Therefore we conclude that

$$|F'(x)| \leq L := \max(L_1, L_2) \leq L_1 + L_2$$

for all  $x \in [\epsilon^2, \epsilon]$ . This yields

$$\begin{aligned} & \left( \int_{\epsilon^2}^{\epsilon} \int_{\epsilon^2}^{\epsilon} \frac{|\phi_{\epsilon}(x) - \phi_{\epsilon}(y)|^2}{|x - y|^2} dx dy \right)^{\frac{1}{2}} \\ & \leq \left( \int_{\epsilon^2}^{\epsilon} \int_{\epsilon^2}^{\epsilon} \frac{|\frac{\epsilon^{\frac{1}{2}}}{\epsilon(1-\epsilon)} \exp(\frac{2}{1-\epsilon}) + \frac{\epsilon^{\frac{1}{2}(1-\epsilon)}}{\epsilon(\frac{1}{2}-\epsilon)^2} \exp(\frac{-\frac{\epsilon^{\frac{1}{2}}}{\frac{1}{2}-\epsilon})|^2 |x - y|^2}{|x - y|^2} dx dy \right)^{\frac{1}{2}} \\ & \leq \left( \int_{\epsilon^2}^{\epsilon} \int_{\epsilon^2}^{\epsilon} \left| \frac{\epsilon^{\frac{1}{2}}}{\epsilon(1-\epsilon)} \exp(\frac{2}{1-\epsilon}) \right|^2 dx dy \right)^{\frac{1}{2}} + \left( \int_{\epsilon^2}^{\epsilon} \int_{\epsilon^2}^{\epsilon} \left| \frac{\epsilon^{\frac{1}{2}(1-\epsilon)}}{\epsilon(\frac{1}{2}-\epsilon)^2} \exp(\frac{-\frac{\epsilon^{\frac{1}{2}}}{\frac{1}{2}-\epsilon}) \right|^2 dx dy \right)^{\frac{1}{2}} \\ & \rightarrow 0 \quad \text{as } \epsilon \rightarrow 0. \end{aligned}$$

$$\begin{aligned} 3) & \int_0^{\epsilon^2} \int_{\epsilon^2}^{\epsilon} \frac{|\phi_{\epsilon}(x) - \phi_{\epsilon}(y)|^2}{|x - y|^2} dx dy \\ & \leq \int_0^{\epsilon^2} \int_{\epsilon^2}^{\frac{\epsilon}{2}} \frac{1}{|x - \epsilon^2|^2} \exp^2 \left[ -\frac{\epsilon^{\frac{1}{2}}(\epsilon - x)}{(x - \epsilon^2)} \right] dx dy + \int_0^{\epsilon^2} \int_{\frac{\epsilon}{2}}^{\epsilon^2} \frac{1}{|x - \epsilon^2|^2} \exp^2 \left[ -\frac{\epsilon^{\frac{1}{2}}(\epsilon - x)}{(x - \epsilon^2)} \right] dx dy \\ & \leq \epsilon^2 \int_{\epsilon^2}^{\frac{\epsilon}{2}} \frac{1}{|x - \epsilon^2|^2} \exp^2 \left[ -\frac{\epsilon^{\frac{1}{2}+1}(1 - \epsilon)}{2(x - \epsilon^2)} \right] dx + \epsilon^2 \int_{\frac{\epsilon}{2}}^{\epsilon} \frac{1}{|\frac{\epsilon}{2} - \epsilon^2|^2} \exp^2 \left[ \frac{\epsilon^{\frac{1}{2}}(2\epsilon)}{(\frac{\epsilon}{2} - \epsilon^2)} \right] dx \\ & \leq \frac{\epsilon^{\frac{1}{2}} \exp(-\frac{\epsilon^{\frac{3}{2}}(1-\epsilon)}{x})}{1 - \epsilon} \Big|_{x=0}^{x=\epsilon} + \frac{1}{(\frac{1}{2} - \epsilon)^2} \frac{\epsilon}{2} \exp^2 \left[ \frac{2\epsilon^{\frac{1}{2}}}{(\frac{1}{2} - \epsilon)} \right] \\ & \rightarrow 0 \quad \text{as } \epsilon \rightarrow 0, \end{aligned}$$

here we used the properties of the exponential function and elementary majorizations.

$$4) \int_{\epsilon^2}^{\epsilon} \int_0^{\epsilon^2} \frac{|\phi_{\epsilon}(x) - \phi_{\epsilon}(y)|^2}{|x - y|^2} dx dy = \int_{\epsilon^2}^{\epsilon} \int_0^{\epsilon^2} \frac{|\exp[-\frac{\epsilon^{\frac{1}{2}}(y-\epsilon)}{(y-\epsilon^2)}]|^2}{|x - y|^2} dx \rightarrow 0,$$

proceed in the same way as for 3).

Combining these integrals on one hand and using the facts:

$$\|\phi_{\epsilon} - 1\|_{0,\Gamma} \rightarrow 0 \quad \text{and} \quad 2\|(\phi_{\epsilon} - 1)\|_{0,\Gamma_i^{4,\epsilon}} \rightarrow 0$$

for all  $0 \leq i \leq m$  on the other hand yield, using (14), the result of lemma 3. Consequently, using (13), it yields also

$$\|\phi_{\epsilon} - 1\|_{\frac{1}{2},\Gamma} \rightarrow 0.$$

□

Since  $u$  is  $(\frac{1}{2} + \iota)$ -holder continuous and thus uniformly continuous on  $\Omega$ , the result of lemma (3) implies

$$\|u_{\epsilon} - u\|_{\frac{1}{2},\Gamma} \leq \|u(\phi_{\epsilon} - 1)\|_{\frac{1}{2},\Gamma} \leq \|u\|_{\infty,\Gamma} \|\phi_{\epsilon} - 1\|_{\frac{1}{2},\Gamma} \rightarrow 0.$$

One proves the following approximation lemma:

**Lemma 4.** *The function  $u_\epsilon$  defined by (8) and the distribution  $g^\epsilon$  appearing in problem (12) satisfy respectively the following limits*

$$a) \|\nabla u_\epsilon - \nabla u\|_{0,\Omega} \rightarrow 0, \quad b) \|g^\epsilon - g\|_{[H^{\frac{1}{2}}]', \Gamma - \Gamma_0} \rightarrow 0$$

as  $\epsilon \rightarrow 0$ .

*Proof.* a) The RHS,  $f \in L^2(\Omega)$ , of problem (8) being fixed, consider the linear operator  $G$  that associate to each  $u^d \in H^{\frac{1}{2}}(\Gamma)$  the corresponding unique solution  $u$  of problem (8).

$$G : (H^{\frac{1}{2}}(\Gamma), \|\cdot\|_{\frac{1}{2},\Gamma}) \rightarrow A \subset V, \\ u^d := u|_\Gamma \rightarrow K(u^d) = u.$$

Where  $(A, \|\cdot\|_{H^1(\Omega)})$  denote the range of  $H^{\frac{1}{2}}(\Gamma)$  under  $G$ . The inverse operator  $G^{-1}$  identifies with the trace operator which is obviously well defined and bijective for  $u \in A$ . Using the trace inequality on  $\Gamma$ , there exists  $c > 0$  such that  $\forall u \in A$

$$\|u\|_{\frac{1}{2},\Gamma} \leq c \|\nabla u\|_{0,\Omega}$$

this implies the continuity of the linear bijective operator  $G^{-1}$ . According to the Banach isomorphism theorem the operator  $G$  is continuous, this means that there exists  $c_{-1} > 0$  such that for all  $u \in A$  we have

$$\|\nabla u\|_{0,\Omega} \leq c_{-1} \|u\|_{\frac{1}{2},\Gamma}$$

thus, using the limit proved in lemma (3)

$$\|\nabla u - \nabla u_\epsilon\|_{0,\Omega} \leq c_{-1} \|u - u_\epsilon\|_{\frac{1}{2},\Gamma} \rightarrow 0, \quad (17)$$

this proves a).

b) We do the same reasoning as for a). Given  $g \in [H^{\frac{1}{2}}(\Gamma - \Gamma_0)]'$ , let  $w \in V$  be the unique solution of

$$\int_{\Omega} 2\mu \varepsilon(w) \varepsilon(v) + \lambda \operatorname{div} w \operatorname{div} v \, dx = \langle g, v \rangle_{[H^{\frac{1}{2}}(\Gamma - \Gamma_0)]', [H^{\frac{1}{2}}(\Gamma - \Gamma_0)]} \quad , \quad (18)$$

for all  $v \in V$ . Choosing  $v = w$ , there exists  $c' > 0$  such that

$$\|\nabla w\|_{0,\Omega} \leq c' \|g\|_{[H^{\frac{1}{2}}(\Gamma - \Gamma_0)]'}. \quad (19)$$

Let  $K$  be the operator that associate to each data  $g \in [H^{\frac{1}{2}}(\Gamma - \Gamma_0)]'$  the solution function  $w$  of the corresponding problem (18):

$$K : [H^{\frac{1}{2}}(\Gamma - \Gamma_0)]' \rightarrow D \subset V \\ g \rightarrow K(g) = w.$$



Where  $(D, \|\cdot\|_{H^1(\Omega)})$  denote the range of  $[H^{\frac{1}{2}}]'(\Gamma - \Gamma_0)$  under  $K$ . Following existence and uniqueness result for problem (18),  $K$  is well defined, furthermore it is linear and invertible. An equivalent formulation of (19) is: there exists a constant  $c' > 0$  such that  $\forall g \in [H^{\frac{1}{2}}(\Gamma - \Gamma_0)]'$ , we have

$$\|K(g)\|_{H^1} \leq c' \|g\|_{[H^{\frac{1}{2}}]'(\Gamma - \Gamma_0)}$$

i.e.  $K$  is continuous. Then, according to Banach's isomorphism theorem, we deduce that  $\exists c'_{-1} > 0$  such that

$$\|g\|_{[H^{\frac{1}{2}}(\Gamma - \Gamma_0)]'} \leq c'_{-1} \|\nabla w\|_{0,\Omega}. \quad (20)$$

Rewriting (4) with  $g \in H^{\frac{1}{2}}(\Gamma - \Gamma_0) \equiv D \subset [H^{\frac{1}{2}}(\Gamma - \Gamma_0)]'$  then subtracting (4) and (12) member-to-member, one find that  $u - u_\epsilon$  satisfy:  $\forall v \in V$ ,

$$\int_{\Omega} 2\mu(\varepsilon(u_\epsilon) - \varepsilon(u))\varepsilon(v) + \lambda \operatorname{div}(u_\epsilon - u) \operatorname{div} v \, dx = \langle g^\epsilon - g, v \rangle_{[H^{\frac{1}{2}}(\Gamma - \Gamma_0)]', [H^{\frac{1}{2}}(\Gamma - \Gamma_0)]}.$$

Applying (20) to  $w = u_\epsilon - u$  we get:

$$\|g^\epsilon - g\|_{[H^{\frac{1}{2}}(\Gamma - \Gamma_0)]'} \leq c'_{-1} \|\nabla u_\epsilon - \nabla u\|_{0,\Omega}.$$

Considering (17), we infer  $b$ ).  $\square$

**Remark 5.** A consequence of the previous lemma: for arbitrary small  $\beta > 0$ , there exists  $\epsilon_1 > 0$  such that  $\forall \epsilon, 0 < \epsilon < \epsilon_1$  we have

$$i) \quad \|g^\epsilon - g\|_{[H^{\frac{1}{2}}(\Gamma - \Gamma_0)]'} \leq \beta, \quad ii) \quad \|\nabla u_\epsilon - \nabla u\|_{0,\Omega} \leq \beta$$

iii) Since by assumption  $g \in H^{\frac{1}{2}}(\Gamma - \Gamma_0)$  then, by using the continuity of the canonical embedding

$$I : H^{\frac{1}{2}}(\Gamma - \Gamma_0) \rightarrow L^2(\Gamma - \Gamma_0),$$

we have

$$\|g\|_{0,\Gamma - \Gamma_0} \leq \|g\|_{\frac{1}{2},\Gamma - \Gamma_0}.$$

### 3 Auxiliary lemmas

Let  $\epsilon_1$  be such as defined in remark (5). For the rest of the paper, we fix  $\epsilon$ ,  $0 < \epsilon < \epsilon_1$ .

Before presenting a proof of the main result, we intend to state two auxiliary lemmas. These lemmas are established using, principally, Theorem 2.1 proved in [10]. In order to apply the later result, (assumption 1.2) in

that same paper need to be met. Our polygonal domain satisfy very well that assumption.

These lemmas convey the essential idea in the demonstration of the main result. The idea is to carry out a particular decomposition, alluded to in the introduction, of the approximating function to the solution  $u_\epsilon$  of problem (8). This key idea is illustrated in the following example. Consider  $v \in$  to be the solution of problem silmilar to (8)

$$\begin{cases} Lv = f & \text{a.e in } \Omega, \\ v = v^d & \text{on } \cup_i \Gamma_i. \end{cases}$$

Let's decompose this problem into two other problems

$$S_1^i \begin{cases} Lv_1^i = f & \text{a.e in } \Omega, \\ v_1^i = v^d & \text{on } \Gamma_i. \\ v_1^i = 0 & \text{on } \cup_i \Gamma_i - \Gamma_i. \end{cases} \quad S_2^i \begin{cases} Lv_2^i = f & \text{a.e in } \Omega, \\ v_2^i = 0 & \text{on } \Gamma_i. \\ v_2^i = v^d & \text{on } \cup_i \Gamma_i - \Gamma_i. \end{cases}$$

Provided that the problems are well posed, the functions  $v_1^i$  and  $v_2^i$  are well defined. On the other hand, it is easy to see that  $v_1^i = v$  on  $\Gamma_i$  does not generally imply  $\sigma(v_1^i) \cdot n = \sigma(v)$  on  $\Gamma_i$ . So the idea is to find a decomposition that makes such a property to be fulfilled. i.e. such that  $v_1^i$  can be approximated by a function  $v_1^{i,n}$  such that  $\sigma(v_1^{i,n}) \cdot n$  is an approximation of  $\sigma(v_1^i)$  on  $\Gamma_i$  in the sense of trace. The adequate use of Theorem 2.1 proved in [10] can make this decomposition possible.

Using the principal result in ([4]), we can find  $(u_\epsilon^p)_p \in C^\infty(\bar{\Omega})$  such that for each  $p$ ,  $u_\epsilon^p$  vanishes on the same part of  $\Gamma$  where does  $u_\epsilon$  and such that

$$\|\nabla u_\epsilon^p - \nabla u_\epsilon\|_{0,\Omega} \rightarrow 0 \quad \text{as } p \rightarrow \infty \quad (21)$$

If we denote

$$f^p := L(u_\epsilon^p) \quad g_p^\epsilon := \sigma(u_\epsilon^p) \cdot \vec{n} \quad (22)$$

then, a particular consequence of (21) is: given an arbitrary small  $\beta > 0$ , we can fix  $p$  such that

$$\|\nabla u_\epsilon^p - \nabla u_\epsilon\|_{0,\Omega} \leq \frac{1}{4\sqrt{\mu}}, \quad (23)$$

and, by using the Banach isomorphism theorem with homogeneous Dirichlet condition on  $\Gamma_0$ , we get

$$(i) \|f^p - f\|_{0,\Omega} \leq \frac{1}{4\sqrt{\mu}} \quad (ii) \|g_p^\epsilon - g^\epsilon\|_{[H^{\frac{1}{2}}(\Gamma - \Gamma_0)]'} \leq \beta. \quad (24)$$

We fix, for the rest of the paper,  $p$  for which (23) and (24) are fulfilled.

**Lemma 6.** *Let  $\delta_0$  be an arbitrary small number. There exists two functions  $u_1$  and  $u_2$  such that*

$$i) \quad u_\epsilon^p = u_1 + u_2.$$

$$ii) \quad \begin{cases} L(u_1) = f_1 & \text{in } \Omega \\ u_1 = 0 & \text{on } \Gamma - \Gamma_0 \\ u_1 = 0 & \text{on } \Gamma_0 \end{cases} \quad \begin{cases} L(u_2) = f_2 & \text{in } \Omega \\ u_2 = u_\epsilon^p & \text{on } \Gamma - \Gamma_0, \\ u_2 = 0 & \text{on } \Gamma_0. \end{cases}$$

$$iii) \quad \sigma(u_1) \cdot \vec{n} = 0 \quad \text{on } \Gamma \text{ and thus, } \sigma(u_2) \cdot \vec{n} = g_p^\epsilon \quad \text{on } \Gamma.$$

$$iv) \quad f_2 := f^p - f_1 \text{ satisfy } \|f^p - f_1\|_{0,\Omega} \leq \frac{\delta_0}{2m}.$$

*Proof.* Let  $\delta_0 > 0$  be an arbitrary small number. Consider the decomposition

$$u_\epsilon^p := w + z$$

where  $w$  and  $z$  are respectively the solutions of the following problems

$$\begin{cases} L(w) = f^p & \text{in } \Omega \\ w = 0 & \text{on } \Gamma - \Gamma_0 \\ w = 0 & \text{on } \Gamma_0 \end{cases} \quad \begin{cases} L(z) = 0 & \text{in } \Omega \\ z = u_\epsilon^p & \text{on } \Gamma - \Gamma_0 \\ z = 0 & \text{on } \Gamma_0; \end{cases}$$

observe that  $\sigma(w) \cdot \vec{n}$  does not necessary equal zero and thus  $\sigma(z) \cdot \vec{n}$  does not necessary equal  $g$ . Nevertheless, by density, there is functions  $w_n \in C_0^\infty(\Omega)$  such that

$$\|\nabla w_n - \nabla w\|_{0,\Omega} \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

One should note that, following the definition of compactly supported functions in  $\Omega$ ,  $\sigma(w_n) \cdot \vec{n} = 0$  on  $\partial\Omega$  for all  $n$ .

Using the same argument as in the proof of lemma (4), and for the fixed homogeneous Dirichlet boundary condition, there exists  $C_0 > 0$  depending merely on the domain  $\Omega$  such that

$$\|L(w_n) - f^p\|_{0,\Omega} \leq C_0 \|\nabla w_n - \nabla w\|_{0,\Omega} \rightarrow 0 \text{ as } n \rightarrow \infty, \quad (25)$$

thus, there exists an element  $w^*$  in  $\{w_n\}_{n \in \mathbb{N}}$  that satisfy the estimate

$$\|L(w^*) - f^p\|_{0,\Omega} \leq \frac{\delta_0}{m}. \quad (26)$$

Decompose  $w := w^* + w_2$  where  $w_2$  is a solutions of

$$\begin{cases} L(w_2) = f^p - L(w^*) & \text{in } \Omega \\ w_2 = 0 & \text{on } \Gamma - \Gamma_0 \\ w_2 = 0 & \text{on } \Gamma. \end{cases}$$

One should remark that

$$\sigma(w_2) \cdot \vec{n} + \sigma(z_1) \cdot \vec{n} = g_p^\epsilon,$$

indeed,  $\sigma(w_2) \cdot \vec{n} + \sigma(z_1) \cdot \vec{n} = \sigma(u) \cdot \vec{n} - \sigma(w^*) \cdot \vec{n} = g_p^\epsilon - 0$ . Pose

$$u_1 := w^* \quad , \quad u_2 := w_2 + z_1,$$

and

$$f_1 := L(w^*) \quad , \quad f_2 := f^p - L(w^*),$$

this concludes the lemma.  $\square$

It should be pointed out that we repeatedly apply the essential argument in lemma (4), namely the Banach isomorphism theorem, in order to obtain estimates for the inverse of some operators.

**Lemma 7.** *Let  $u_1, u_2$  be such as defined in lemma (6). Given  $\delta_1, \delta_2 > 0$  arbitrary small numbers, there exists  $m$  functions  $u_{2,i}, 1 \leq i \leq m$ , such that*

$$u_{2,i} = 0 \quad \text{on } \Gamma - \Gamma_i,$$

and such that  $g_i := \sigma(u_{2,i}) \cdot \vec{n}$  satisfy

$$\|g_i - g_p^\epsilon\|_{0,\Gamma_i} \leq \frac{\delta_1}{m}. \quad (27)$$

moreover,  $u'_2 := \sum_{i=1}^m u_{2,i}$  satisfy

$$\|\nabla(u_1 + u'_2) - \nabla u_\epsilon^p\|_{0,\Omega} \leq \delta_2. \quad (28)$$

The proof is done in two main steps. In the first one, we construct the functions  $u_{2,i}$  and in the second step we prove that these functions actually satisfy estimate (27) and (28).

*Proof.* Let  $\delta_1, \delta_2$  be arbitrary small numbers. The functions  $u_{2,i}$  will be defined by introducing the intermediate functions  $X_i, Y_i, \tilde{Y}_i^n$  and  $V_i$  respectively in subsections a), b), c) and d). As we define these functions, we demonstrate some estimates that will be used to derive (27) and (28) in the second step.

**Step 1:** Fix  $i, 1 \leq i \leq m$ . Consider the decomposition

$$u_2 = X_i + \sum_{j \neq i} X_j = X_i + Z_i ,$$

where  $X_i$  and  $Z_i$  are the solutions of

$$(P_1) \begin{cases} L(X_i) & = 0 & \text{in } \Omega \\ X_i & = u_\epsilon^p & \text{on } \Gamma_i \\ X_i & = 0 & \text{on } \Gamma - \Gamma_i \cup \Gamma_0 \\ X_i & = 0 & \text{on } \Gamma_0 \end{cases} \quad (P'_1) \begin{cases} L(Z_i) & = f_2 & \text{in } \Omega \\ Z_i & = 0 & \text{on } \Gamma_i \\ Z_i & = u_\epsilon^p & \text{on } \Gamma - \Gamma_i \cup \Gamma_0 \\ Z_i & = 0 & \text{on } \Gamma_0. \end{cases}$$

One should notice that these two problems are well posed, this can be seen by considering the regularity of  $u_\epsilon^p$  on the boundary and the fact that  $u_\epsilon^p$  vanish identically on a neighborhood of the the vertices. Since  $\Omega$  is a Lipschitz domain, it possesses the  $W^{1,2}$ -Sobolev extension property.

a) According to (theorem 2.1, [10]) see Index A at the end, there exists  $X_i^n \in C_0^\infty(\mathbb{R}^2)$  such that we have

$$(h1) \quad \|\nabla X_i^n - \nabla X_i\|_{0,\Omega} \leq \frac{\delta_3}{m} \quad (h2) \quad \text{supp}(X_i^n) \cap (\Gamma - \Gamma_i) = \emptyset,$$

where  $\delta_3 > 0$  is an adequately chosen real number. A consequence of (h2) is:

$$(h3) \quad \sigma(X_i^n) \cdot \vec{n} = 0 \text{ on } \Gamma - \Gamma_i, \quad X_i^n = 0 \text{ on } \Gamma - \Gamma_i.$$

Furthermore, with the homogeneous Dirichlet condition on  $\Gamma - \Gamma_i$  being fixed in the problem  $P_1$  and in the problem solved by  $X_i^n$  then, using the Banach isomorphism theorem, there exists  $C_1 > 0$  such that

$$\|\sigma(X_i^n) \cdot \vec{n} - \sigma(X_i) \cdot \vec{n}\|_{0,\Gamma_i} + \|L(X_i^n) - 0\|_{0,\Omega} \leq C_1 \|\nabla X_i^n - \nabla X_i\|_{0,\Omega} \leq \frac{C_1 \delta_3}{m}.$$

We choose  $\delta_3$  in assumption (h1) such that

$$(c1) : \quad \frac{C_1 \delta_3}{m} < \frac{\delta_0}{4m}$$

where  $\delta_0$  is a positive real number whose choice will be precised later (see condition (c4) below). Fix  $n_1$  such that  $F_i^{n_1} := L(X_i^{n_1})$  satisfy

$$\|F_i^{n_1} - 0\|_{0,\Omega} \leq \frac{\delta_0}{4m}. \quad (29)$$

Denote  $\tilde{X}_i^{n_1} := X_i - X_i^{n_1}$ , it does satisfy

$$X_i^{n_1} + \tilde{X}_i^{n_1} + Z_i = u_\epsilon^p \quad (30)$$

i.e.

$$\left\{ \begin{array}{l} L(X_i^{n_1}) = F_i^n \quad \text{in } \Omega \\ X_i^{n_1} = x_i^{n_1} \quad \text{on } \Gamma_i \\ X_i^{n_1} = 0 \quad \text{on } \Gamma - \Gamma_i \end{array} \right. \quad \left\{ \begin{array}{l} L(\tilde{X}_i^{n_1}) = -F_i^n \quad \text{in } \Omega \\ \tilde{X}_i^{n_1} = u_\epsilon^p - x_i^{n_1} \quad \text{on } \Gamma_i \\ \tilde{X}_i^{n_1} = 0 \quad \text{on } \Gamma - \Gamma_i, \end{array} \right.$$

for the same reasons as for  $(P_1)$  and  $(P_2)$ , these problems are well posed. Considering  $(h3)$  and  $(30)$ , the function  $\tilde{X}_i^{n_1} + Z_i$  satisfy

$$(P_3) \left\{ \begin{array}{l} L(\tilde{X}_i^{n_1} + Z_i) = -F_i^{n_1} + f_2 \quad \text{in } \Omega \\ \tilde{X}_i^{n_1} + Z_i = u_\epsilon^p - x_i^{n_1} \quad \text{on } \Gamma_i \\ \sigma(\tilde{X}_i^{n_1} + Z_i) \cdot \vec{n} = g_p^\epsilon \quad \text{on } \Gamma - \Gamma_i \cup \Gamma_0 \\ \tilde{X}_i^{n_1} + Z_i = 0 \quad \text{on } \Gamma_0. \end{array} \right.$$

b) Let  $Y_i$  be the function defined by

$$(P_4) \left\{ \begin{array}{l} L(Y_i) = -F_i^{n_1} + f_2 \quad \text{in } \Omega \\ Y_i = 0 \quad \text{on } \Gamma_i \\ \sigma(Y_i) \cdot \vec{n} = g_p^\epsilon \quad \text{on } \Gamma - \Gamma_i \cup \Gamma_0 \\ Y_i = 0 \quad \text{on } \Gamma_0, \end{array} \right.$$

Thanks to assumption (5), this problem is well posed. There exists a trace constant  $c_t$  such that, using  $(h1)$ , we have

$$\|(u_\epsilon^p - x_i^{n_1})\|_{0,\Gamma_i} \leq c_t \times \|\nabla X_i - \nabla X_i^{n_1}\|_{0,\Omega} \leq \frac{c_t \times \delta_3}{m}; \quad (31)$$

for the fixed  $-F_i^{n_1} + f_2 \in L^2(\Omega)$ ,  $g_p^\epsilon \in L^2(\Gamma - \Gamma_i \cup \Gamma_0)$  and the homogeneous Dirichlet condition on  $\Gamma_0$  in problems  $(P_3)$  and  $(P_4)$ , and by using an argument similar to that in part (a) of the proof of lemma (4) (Banach isomorphism theorem), there exists a constant  $C_3 > 0$  such that

$$\begin{aligned} \|\nabla Y_i - \nabla(\tilde{X}_i^{n_1} + Z_i)\|_{0,\Omega} &\leq C_3 \|Y_i - (u_\epsilon^p - x_i^{n_1})\|_{0,\Gamma_i} \\ &= C_3 \|0 - (u_\epsilon^p - x_i^{n_1})\|_{0,\Gamma_i} \\ &\leq \frac{c_t \times C_3 \delta_3}{m}, \end{aligned} \quad (32)$$

and by using the trace inequality again we have

$$\begin{aligned} \|Y_i - (\tilde{X}_i^{n_1} + Z_i)\|_{0,\Gamma - \Gamma_i} &\leq c_t \|\nabla Y_i - \nabla(\tilde{X}_i^{n_1} + Z_i)\|_{0,\Omega} \\ &\leq c_t^2 \times C_3 \frac{\delta_3}{m}. \end{aligned} \quad (33)$$

Considering (30) and (h3), we easily see that  $\tilde{X}_i^{n_1} + Z_i = u_\epsilon^p$  on  $\Gamma - \Gamma_i$ ; and thus, estimates (32), (33) become respectively

$$\|\nabla(X_i^{n_1} + Y_i) - \nabla u_\epsilon^p\|_{0,\Omega} \leq \frac{c_t \times C_3 \delta_3}{m}. \quad (34)$$

$$\|Y_i - u_\epsilon^p\|_{0,\Gamma - \Gamma_i} \leq \frac{c_t^2 \times C_3 \delta_3}{m}. \quad (35)$$

On the other hand, since  $Y_i = 0$  on  $\Gamma_i$  then, using the result (Theorem 2.1, [10]), there exists  $Y_i^n \in C_0^\infty(\mathbb{R}^2)$  such that

$$(h4) \quad \|\nabla Y_i^n - \nabla Y_i\|_{0,\Omega} \leq \frac{\delta'_1}{m} \quad \text{and} \quad \text{supp}(Y_i^n) \cap \Gamma_i = \emptyset,$$

where  $\delta'_1 > 0$  is an adequately chosen real number, thus

$$\sigma(Y_i^n) \cdot \vec{n} = 0 \quad \text{on} \quad \Gamma_i \quad \text{and} \quad Y_i^n = 0 \quad \text{on} \quad \Gamma_i \quad (36)$$

for the homogeneous Dirichlet condition on  $\Gamma_i \cup \Gamma_0$  for problem  $(P_4)$  and the problem solved by  $Y_i^n$ , and using Banach isomorphism theorem, there exists  $C'_1 > 0$  such that

$$\|L(Y_i^n) - (-F_i^{n_1} + f_2)\| \leq C'_1 \|\nabla Y_i^n - \nabla Y_i\|_{0,\Omega}.$$

We choose  $\delta'_1 > 0$  in assumption (h4) such that

$$(c2) : \quad \frac{C'_1 \delta'_1}{m} < \frac{\delta_0}{4m},$$

where  $\delta_0$  is as defined in lemma (6) i.e. need to satisfy condition (c4) below. Fix  $n_2 > n_1$  such that

$$h_i^{n_2} := L(Y_i^{n_2}) \quad (37)$$

satisfy

$$\|h_i^{n_2} - (-F_i^{n_1} + f_2)\|_{0,\Omega} \leq \frac{\delta_0}{4m};$$

consequently, considering (iv) of lemma (6) and (29), we have

$$\|h_i^{n_2} - 0\| \leq \frac{\delta_0}{m}. \quad (38)$$

c) Denote  $\tilde{Y}_i^{n_2} := Y_i - Y_i^{n_2}$ , we have

$$(P_5) \begin{cases} L(Y_i^{n_2}) & = h_i^{n_2} & \text{in } \Omega \\ Y_i^{n_2} & = 0 & \text{on } \Gamma_i \\ Y_i^{n_2} & = y_i^{n_2} & \text{on } \Gamma - \Gamma_i \cup \Gamma_0 \\ Y_i^{n_2} & = 0 & \text{on } \Gamma_0 \end{cases} \quad (P_6) \begin{cases} L(\tilde{Y}_i^{n_2}) & = -h_i^{n_2} - F_i^{n_1} + f_2 & \text{in } \Omega \\ \tilde{Y}_i^{n_2} & = 0 & \text{on } \Gamma_i \\ \tilde{Y}_i^{n_2} & = Y_i - y_i^{n_2} & \text{on } \Gamma - \Gamma_i \cup \Gamma_0 \\ \tilde{Y}_i^{n_2} & = 0 & \text{on } \Gamma_0, \end{cases}$$

estimate (34) rewrites

$$\|\nabla(\tilde{Y}_i^{n_2} + Y_i^{n_2} + X_i^{n_1}) - \nabla u_\epsilon^p\|_{0,\Omega} \leq \frac{c_t \times C_3 \delta_3}{m}, \quad (39)$$

denote

$$g' := \sigma(\tilde{Y}_i^{n_2}) \cdot \vec{n} \in L^2(\Gamma - \Gamma_0).$$

d) Let  $V_i$  be the function defined by

$$(P_7) \begin{cases} L(V_i) & = -h_i^{n_2} - F_i^{n_1} + f_2 & \text{in } \Omega \\ \sigma(V_i) \cdot \vec{n} & = g' & \text{on } \Gamma_i \\ V_i & = 0 & \text{on } \Gamma - \Gamma_i \cup \Gamma_0 \\ V_i & = 0 & \text{on } \Gamma_0. \end{cases}$$

Set

$$u_{2,i} := X_i^{n_1} + V_i,$$

**Step 2:** The functions  $u_{2,i}$  are the unique solutions of

$$\begin{cases} L(u_{2,i}) & = -h_i^{n_2} + f_2 & \text{in } \Omega \\ u_{2,i} & = X_i^{n_1} + V_i & \text{on } \Gamma_i \\ u_{2,i} & = 0 & \text{on } \Gamma - \Gamma_i \cup \Gamma_0 \\ u_{2,i} & = 0 & \text{on } \Gamma_0. \end{cases}$$

Using the estimates established in step 1, we are ready to prove (27) and (28).

**Proof of estimate (27):** Since  $\sigma(V_i) \cdot \vec{n} = \sigma(\tilde{Y}_i^{n_2}) \cdot \vec{n}$  on  $\Gamma_i$ , then we have for  $g_i := \sigma(V_i + X_i^{n_1}) \cdot \vec{n}$

$$\begin{aligned} \|g_i - g_p^\epsilon\|_{0,\Gamma_i} &= \|\sigma(V_i + X_i^{n_1}) \cdot \vec{n} - g_p^\epsilon\|_{0,\Gamma_i} \\ &= \|\sigma(\tilde{Y}_i^{n_2} + X_i^{n_1}) \cdot \vec{n} - \sigma(u_\epsilon^p) \cdot \vec{n}\|_{0,\Gamma_i} \end{aligned}$$

Using (36)

$$\begin{aligned} \|g_i - g_p^\epsilon\|_{0,\Gamma_i} &= \|\sigma(\tilde{Y}_i^{n_2} + Y_i^{n_2} + X_i^{n_1}) \cdot \vec{n} - \sigma(u_\epsilon^p) \cdot \vec{n}\|_{0,\Gamma_i} \\ &\leq \|\sigma(\tilde{Y}_i^{n_2} + Y_i^{n_2} + X_i^{n_1}) \cdot \vec{n} - \sigma(u_\epsilon^p) \cdot \vec{n}\|_{0,\Gamma - \Gamma_0}; \end{aligned}$$



for the fixed homogeneous Dirichlet condition on  $\Gamma_0$ , and according to the Banach isomorphism theorem, there exists a constant  $C_2 > 0$  depending merely on  $\Omega$  such that, using (39) and since we have  $Y_i = Y_i^{n_2} + \tilde{Y}_i^{n_2}$

$$\begin{aligned} \|g_i - g_p^\epsilon\|_{0,\Gamma_i} &\leq \|\sigma(Y_i + X_i^{n_1}) \cdot \vec{n} - \sigma(u_\epsilon^p) \cdot \vec{n}\|_{0,\Gamma-\Gamma_0} \\ &\leq C_2 \|\nabla(Y_i + X_i^{n_1}) - \nabla u_\epsilon^p\|_{0,\Omega} \\ &\leq \frac{c_t \times C_3 C_2 \delta_3}{m}. \end{aligned}$$

If  $\delta_3$  also fulfill condition

$$(c3) : \quad \frac{c_t \times C_3 C_2 \delta_3}{m} < \frac{\delta_1}{m},$$

where  $\delta_1$  is the arbitrary small number assumed at the beginning of the proof then, estimate (27) is proved.

In order to prove estimate (28), we need to estimate the trace of  $X_i^{n_1} + V_i$  on  $\Gamma_i$ .

1) Firstly remark, by using (h3), that

$$V_i + X_i^{n_1} = 0 \quad \text{on } \Gamma - \Gamma_i.$$

2) For the fixed  $-(h_i^{n_2} + F_i^{n_1} - f_2) \in L^2(\Omega)$ ,  $g' \in L^2(\Gamma_i)$  and the homogeneous Dirichlet condition on  $\Gamma_0$  in problems  $(P_6)$  and  $(P_7)$ , the Banach isomorphism theorem ensures the existence of a constant  $C_4 > 0$  depending merely on  $\Omega$  such that

$$\|\nabla V_i - \nabla \tilde{Y}_i^{n_2}\|_{0,\Omega} \leq C_4 \|V_i - \tilde{Y}_i^{n_2}\|_{0,\Gamma-\Gamma_i} = C_4 \|0 - (Y_i - y_i^{n_2})\|_{0,\Gamma-\Gamma_i},$$

hence, considering (h4) and using trace inequality, we have

$$C_4 \|0 - (Y_i - y_i^{n_2})\|_{0,\Gamma-\Gamma_i} \leq c_t C_4 \|\nabla Y_i - \nabla Y_i^{n_2}\|_{0,\Omega} \leq c_t C_4 \frac{\delta_1'}{m},$$

thus,

$$\|\nabla V_i - \nabla \tilde{Y}_i^{n_2}\|_{0,\Omega} \leq c_t C_4 \frac{\delta_1'}{m}. \quad (40)$$

Using trace inequality and (40) we obtain

$$\|V_i - 0\|_{0,\Gamma_i} \leq \|V_i - \tilde{Y}_i^{n_2}\|_{0,\Gamma} \leq c_t \|\nabla V_i - \nabla \tilde{Y}_i^{n_2}\|_{0,\Omega} \leq c_t^2 \times C_4 \frac{\delta_1'}{m}. \quad (41)$$

Estimates (31) and (41) imply

$$\begin{aligned}
\|V_i + X_i^{n_1} - u_\epsilon^p\|_{0,\Gamma_i} &\leq \|V_i - 0\|_{0,\Gamma_i} + \|X_i^{n_1} - u_\epsilon^p\|_{0,\Gamma_i} \\
&= \|V_i - 0\|_{0,\Gamma_i} + \|x_i^{n_1} - u_\epsilon^p\|_{0,\Gamma_i} \\
&\leq c_t^2 \times C_4 \frac{\delta_1'}{m} + \frac{c_t \delta_3}{m}.
\end{aligned} \tag{42}$$

**Proof of estimate (28):** Write  $u_2' := \sum_i u_{2,i} = \zeta' + \eta'$  where  $\zeta'$  and  $\eta'$  are respectively the solutions of

$$(P_8) \begin{cases} L(\zeta') = -\sum_{i=1}^m (h_i^{n_2} - f_2) & \text{in } \Omega \\ \zeta' = 0 & \text{on } \Gamma_i \\ \zeta' = 0 & \text{on } \Gamma - \Gamma_i \end{cases} \quad \begin{cases} L(\eta') = 0 & \text{in } \Omega \\ \eta' = \sum_{i=1}^m u_{2,i} & \text{on } \Gamma - \Gamma_0 \\ \eta' = 0 & \text{on } \Gamma_0, \end{cases}$$

and write  $u_2 = \zeta + \eta$  where  $\zeta$  and  $\eta$  are the solutions of

$$(P_9) \begin{cases} L(\zeta) = f_2 & \text{in } \Omega \\ \zeta = 0 & \text{on } \Gamma_i \\ \zeta = 0 & \text{on } \Gamma - \Gamma_i \end{cases} \quad \begin{cases} L(\eta) = 0 & \text{in } \Omega \\ \eta = u_\epsilon^p & \text{on } \Gamma - \Gamma_0 \\ \eta = 0 & \text{on } \Gamma_0. \end{cases}$$

If we denote  $c_j k$  the constant in the Korn inequality then the weak form of the problem solved by  $\zeta - \zeta'$  yields

$$\begin{aligned}
\|\nabla \zeta - \nabla \zeta'\|_{0,\Omega} &\leq c_k d(\Omega) \|f_2 + \sum_i (h_i^{n_2} - f_2)\|_{0,\Omega} \\
&\leq d(\Omega) (\|f_2\|_{0,\Omega} + \sum_i \|h_i^{n_2}\|_{0,\Omega} + \sum_i \|f_2\|_{0,\Omega})
\end{aligned}$$

using (iv) of lemma (6) and (38), we have

$$\|\nabla \zeta - \nabla \zeta'\|_{0,\Omega} \leq d(\Omega) \left( \frac{\delta_0}{2m} + \delta_0 + \frac{\delta_0}{2} \right). \tag{43}$$

On the other hand, the Banach isomorphism theorem ensures, for the null source term and the Dirichlet homogeneous condition on  $\Gamma_0$  in problems (P<sub>8</sub>) and (P<sub>9</sub>), that there exists a constant  $C_5 > 0$  such that

$$\begin{aligned}
\|\nabla \eta - \nabla \eta'\|_{0,\Omega} &\leq C_5 \|\eta - \eta'\|_{0,\Gamma - \Gamma_0} = C_5 \|u_\epsilon^p - \sum_{i=1}^m u_{2,i}\|_{0,\Gamma - \Gamma_0} \\
&\leq C_5 \sum_{i=1}^m \|u_\epsilon^p - \sum_{i=1}^m u_{2,i}\|_{0,\Gamma_i} \\
&= C_5 \sum_{i=1}^m \|u_\epsilon^p - u_{2,i}\|_{0,\Gamma_i}.
\end{aligned}$$

Applying estimate (42) with  $u_{2,i} = V_i + X_i^{n_1}$

$$\|\nabla\eta - \nabla\eta'\|_{0,\Omega} \leq C_5(c_t^2 \times C_4\delta_1' + c_t\delta_3) \quad (44)$$

Combining (43) and (44), we obtain using triangular inequality

$$\begin{aligned} \|\nabla u_2 - \nabla u_2'\|_{0,\Omega} &\leq \|\nabla\zeta - \nabla\zeta'\| + \|\nabla\eta - \nabla\eta'\|_{0,\Omega} \\ &\leq d(\Omega)\left(\frac{\delta_0}{2m} + \delta_0 + \frac{\delta_0}{2}\right) + C_5(c_t^2 \times C_4\delta_1' + c_t\delta_3), \end{aligned}$$

Finally, if  $\delta_0$ ,  $\delta_3$  and  $\delta_1'$  are subject to fulfill the condition

$$(c4) : \quad d(\Omega)\left(\frac{\delta_0}{2m} + \delta_0 + \frac{\delta_0}{2}\right) + C_5(c_t^2 \times C_4\delta_1' + c_t\delta_3) < \delta_2.$$

where  $\delta_2$  is the arbitrary small number assumed at the beginning of the proof then,

$$\|\nabla u_\epsilon^p - \nabla(u_1 + u_2')\|_{0,\Omega} \leq \delta_2$$

This proves (45). □

**Remark 8.** 1) Let  $g_p^\epsilon$  such as defined in (22), we can easily show that

$$\|g_p^\epsilon\|_{0,\Gamma-\Gamma_0}^2 \leq \|g\|_{0,\Gamma-\Gamma_0}^2 + \delta_6^2, \quad (45)$$

where  $\delta_6$  is an arbitrary small positive real number. Indeed, write using triangular inequality

$$\begin{aligned} | \|g_p^\epsilon\|_{0,\Gamma-\Gamma_0}^2 - \int_{\Gamma-\Gamma_0} g g_p^\epsilon \, d\sigma | &= | \langle g_p^\epsilon, g_p^\epsilon \rangle_{[H^{\frac{1}{2}}(\Gamma-\Gamma_0)]', H^{\frac{1}{2}}(\Gamma-\Gamma_0)} \\ &\quad - \langle g, g_p^\epsilon \rangle_{[H^{\frac{1}{2}}(\Gamma-\Gamma_0)]', H^{\frac{1}{2}}(\Gamma-\Gamma_0)} | \\ &\leq | \langle g_p^\epsilon - g, g_p^\epsilon \rangle_{[H^{\frac{1}{2}}(\Gamma-\Gamma_0)]', H^{\frac{1}{2}}(\Gamma-\Gamma_0)} | \\ &\leq \|g_p^\epsilon - g\|_{[H^{\frac{1}{2}}(\Gamma-\Gamma_0)]'} \|g_p^\epsilon\|_{\frac{1}{2},\Gamma-\Gamma_0} \\ &\leq (\|g_p^\epsilon - g\|_{[H^{\frac{1}{2}}(\Gamma-\Gamma_0)]'} + \|g - g\|_{[H^{\frac{1}{2}}(\Gamma-\Gamma_0)]'}) \|g_p^\epsilon\|_{\frac{1}{2},\Gamma-\Gamma_0} \end{aligned}$$

Since, following estimate (ii) of (24),  $g_p^\epsilon$  converges in  $[H^{\frac{1}{2}}(\Gamma - \Gamma_0)]'$  i.e. weakly in  $H^{\frac{1}{2}}(\Gamma - \Gamma_0)$  hence, there exists  $C' > 0$  such that

$$\|g_p^\epsilon\|_{\frac{1}{2},\Gamma-\Gamma_0} \leq C'$$

for all  $p \in \mathbb{N}$ . Thus,

$$\|g_p^\epsilon\|_{0,\Gamma-\Gamma_0}^2 - \int_{\Gamma-\Gamma_0} gg_p^\epsilon d\sigma \leq C' (\|g_p^\epsilon - g^\epsilon\|_{[H^{\frac{1}{2}}(\Gamma-\Gamma_0)]'} + \|g^\epsilon - g\|_{[H^{\frac{1}{2}}(\Gamma-\Gamma_0)]'}) ,$$

using (i) of remark (5) and estimates (ii) of (24) and for an adequate choice of  $\beta$ , the parameters  $p$  and  $\epsilon$  can be chosen such that

$$\|g_p^\epsilon\|_{0,\Gamma-\Gamma_0}^2 - \int_{\Gamma-\Gamma_0} gg_p^\epsilon d\sigma \leq 2\beta C'$$

and thus, Cauchy-Schwarz yields

$$\|g_p^\epsilon\|_{0,\Gamma-\Gamma_0}^2 \leq \int_{\Gamma-\Gamma_0} gg_p^\epsilon + 2\beta C' \leq \|g\|_{0,\Gamma-\Gamma_0} \|g_p^\epsilon\|_{0,\Gamma-\Gamma_0} + 2\beta C' ,$$

Young inequality on the other hand gives us

$$\|g_p^\epsilon\|_{0,\Gamma-\Gamma_0}^2 \leq \frac{1}{2} \|g\|_{0,\Gamma-\Gamma_0}^2 + \frac{1}{2} \|g_p^\epsilon\|_{0,\Gamma-\Gamma_0}^2 + 2\beta C'$$

and hence, there exists  $p$  such that  $g_p^\epsilon$  satisfy

$$\|g_p^\epsilon\|_{0,\Gamma-\Gamma_0}^2 \leq \|g\|_{0,\Gamma-\Gamma_0}^2 + 4\beta C' ,$$

pose

$$\delta_6^2 := 4\beta C'$$

since, following (i) of remark (5) and (24),  $\beta$  is arbitrary small then the same holds with  $\delta_6$ .

2) on the other hand, if  $u_{2,i}$  and  $h_i^{n_2}$  are respectively such as defined in lemma (7) and (37) then

$$\begin{aligned} \|L(u_{2,i})\|_{0,\Omega} &= \|h_i^{n_2} + f_2\|_{0,\Omega} \leq \|h_i^{n_2}\|_{0,\Omega} + \|f_2\|_{0,\Omega} \\ &\leq \frac{\delta_0}{m} + \frac{\delta_0}{2m} =: \tau, \end{aligned} \quad (46)$$

where  $\delta_0$  is arbitrary small and satisfy (c4) and thus,  $\tau$  is also arbitrary small.

### 3.1 Extension of the functions $u_{2,i}$

We introduce some useful lemmas, which will play important roles in the proof of theorem (1).

Since we are looking for explicit estimates, we should use Poincaré, trace and Korn's inequalities relatively to suitable geometric configurations i.e. for which they are explicitly formulated. The configuration that best fits our polygonal convex domain  $\Omega$  is the half-plane  $\mathbb{R}^{2+}$  containing the convex polygonal domain  $\Omega$  for Korn's inequality, the square  $S_d$  with edge's length equal to  $d(\Omega)$  for the Poincaré inequality and with edge  $\Gamma_i$  for the trace inequality. These squares are taken to be subsets of the half-plane containing  $\Omega$ . Thus we determine these constants thanks to results available for this type of domains. The need to consider the functions in and outside the domain  $\Omega$  suggests to extend by zero the functions  $u_{2,i}$  outside the convex domain  $\Omega$ . The definition of the functions  $u_{2,i}$  is adapted to make such an extension.

We consider for  $i$ ,  $1 \leq i \leq m$ , the extension by zero of  $u_{2,i}$  from the convex domain  $\Omega$  to the half-plane  $\mathbb{R}^{2+}$  containing  $\Omega$  such that  $\Gamma_i \subset \partial\mathbb{R}^{2+}$ .

$$\tilde{u}_{2,i} = \begin{cases} u_{2,i}, & a.e. \quad x \in \overline{\Omega}, \\ 0, & x \in \mathbb{R}^{2+} - \overline{\Omega}. \end{cases} \quad (47)$$

We have, obviously, the following

$$\|\partial_{x_i} \tilde{u}_{2,i}\|_{0,\mathbb{R}^{2+}} = \|\partial_{x_i} \tilde{u}_{2,i}\|_{0,\Omega} = \|\partial_{x_i} u_{2,i}\|_{0,\Omega}. \quad (48)$$

The inequalities are established for the extended  $H^1$  regular functions defined on a square containing the convex polygonal domain  $\Omega$ .

### 3.2 Explicit constant in the Poincaré inequality

We show in the following lemma that the function  $u_{2,i} \in V_i$  satisfy the Poincaré inequality for which we determine explicitly the constant.

**Lemma 9.** *For all  $i$ ,  $0 \leq i \leq m$ , the function  $u_{2,i}$  satisfy:*

$$\|u_{2,i}\|_{0,\Omega} \leq d(\Omega) \|\nabla u_{2,i}\|_{0,\Omega}, \quad (49)$$

*the constant  $d(\Omega)$  means the diameter of  $\Omega$ .*

*Proof.* We establish Poincaré inequality for one of the two components  $u_{2,i}^l$ ,  $l = 1, 2$ , the same estimate hold with the other. Note  $abcd$  the square  $S_d$ , with edge's length equal to  $d(\Omega)$  and subset of the half-plane containing  $\Omega$ , such that  $a = (a_1, a_2)$ ,  $b = (b_1, b_2)$ ,  $c = (c_1, c_2)$  et  $d = (d_1, d_2)$  and such that  $\Gamma_i \subset S_i := [c, d]$ ; so  $\tilde{u}_{2,i}^l = 0$  on  $\partial S_d - \Gamma_i$ .

Since  $\tilde{u}_{2,i}^l$  is absolutely continuous on the lines parallel to the coordinate axis, then applying the fundamental theorem of calculus to  $\tilde{u}_{1,i}^{n,l}$  on

$S_d$  for  $l = 1, 2$ , we have for all  $(x_1, x_2) \in [a_1, d_1] \times [a_2, b_2]$

$$\tilde{u}_{2,i}^l(x_1, x_2) = \int_{a_1}^{x_1} \partial_{x_1} \tilde{u}_{2,i}^l(s, x_2) ds + \tilde{u}_{2,i}^l(a_1, x_2).$$

Since  $(a_1, x_2) \in \partial S_d - \Gamma_i$ , then  $\forall (x_1, x_2) \in [a_1, d_1] \times [a_2, b_2]$

$$\tilde{u}_{2,i}^l(x_1, x_2) = \int_{a_1}^{x_1} \partial_{x_1} \tilde{u}_{2,i}^l(s, x_2) ds.$$

Using Cauchy-Schwarz inequality  $\forall (x_1, x_2) \in [a_1, d_1] \times [a_2, b_2]$

$$|\tilde{u}_{2,i}^l(x_1, x_2)| \leq |x_1 - a|^{1/2} \left( \int_{a_1}^{x_1} |\partial_{x_1} \tilde{u}_{2,i}^l(s, x_2)|^2 ds \right)^{1/2}.$$

Taking the square of the two hand sides of this inequality and using the fact  $|x_1 - a| \leq d(\Omega)$ :  $\forall (x_1, x_2) \in [a_1, d_1] \times [a_2, b_2]$  yields

$$|\tilde{u}_{2,i}^l(x_1, x_2)|^2 \leq |x_1 - a| \int_{a_1}^{x_1} |\partial_{x_1} \tilde{u}_{2,i}^l(s, x_2)|^2 ds \leq d(\Omega) \int_{a_1}^{d_1} |\partial_{x_1} \tilde{u}_{2,i}^l(s, x_2)|^2 ds.$$

Integrating on  $S_d$  with respect to the variables  $x_1$  and  $x_2$ :

$$\begin{aligned} \|\tilde{u}_{2,i}^l\|_{0,S_d}^2 &= \int_{a_2}^{b_2} \int_{a_1}^{d_1} |\tilde{u}_{2,i}^l(x_1, x_2)|^2 dx dx \leq d(\Omega) \int_{a_2}^{b_2} \int_{a_1}^{d_1} \int_{a_1}^{d_1} |\partial_{x_1} \tilde{u}_{2,i}^l(s, x_2)|^2 ds dx dx \\ &\leq d^2(\Omega) \int \int_{S_d} |\partial_x \tilde{u}_{2,i}^l(s, x_2)|^2 ds dx. \end{aligned}$$

According to definition (47) and by considering (48) we get

$$\|u_{2,i}^l\|_{0,\Omega}^2 \leq d^2(\Omega) \|\nabla u_{2,i}^l\|_{0,\Omega}^2.$$

We infer that

$$\begin{aligned} \|u_{2,i}\|_{0,\Omega}^2 &= \|u_{2,i}^1\|_{0,\Omega}^2 + \|u_{2,i}^2\|_{0,\Omega}^2 \leq d^2(\Omega) (\|\nabla u_{2,i}^1\|_{0,\Omega}^2 + \|\nabla u_{2,i}^2\|_{0,\Omega}^2) \\ &= d^2(\Omega) \|\nabla u_{2,i}\|_{0,\Omega}^2. \end{aligned}$$

□

### 3.3 Explicit bound for the trace of $u_{2,i}$

Using mainly the inequality of Poincaré stated in lemma (9) and the trace inequality for  $u_{2,i}$  on the edge of a parallelogram (see lemma 4.2 in [3]), one establishes an explicit bound for the trace of the function  $u_{2,i}$  on  $\Gamma_i$ .

**Lemma 10.** *For all  $i$ , the function  $u_{2,i}$  defined in lemma (7) satisfy:*

$$\|u_{2,i}\|_{0,\Gamma_i} \leq 16|\Gamma_i| \|g_i\|_{0,\Gamma_i} + \delta_9, \quad (50)$$

where  $\delta_9$  is an arbitrary small real positive number.

*Proof.* Let  $u_{2,i}$  be defined on the square  $S_{\Gamma_i}$  with edge  $\Gamma_i$  such that  $S_{\Gamma_i}$  is a subset of the half-plane  $\mathbb{R}^{2+}$  containing  $\Omega$ . Let us write

$$u_{2,i} = \alpha + \varrho,$$

this decomposition is made similarly to that presented in lemma (7) i.e. such that on one hand

$$\|L(\alpha)\|_{0,S_{\Gamma_i}} \quad \text{and} \quad \|\sigma(\alpha) \cdot \vec{n} - g_i\|_{0,\Gamma_i} \quad \text{are small enough} \quad (51)$$

and on the other hand,  $\alpha, \varrho$  vanish respectively on  $\partial S_{\Gamma_i} - \Gamma_i, \Gamma_i$ . We prove (50) for  $u_{2,i}$  considered as the trace on  $\Gamma_i$  of the function  $\alpha$ . Decompose  $\alpha := \alpha_1 + \alpha_2$  in the following way

$$\begin{cases} L(\alpha_1) & = L(\alpha) & \text{in } \Omega \\ \sigma(\alpha_1) \cdot \vec{n} & = 0 & \text{on } \Gamma_i \\ \alpha_1 & = 0 & \text{on } \partial S_{\Gamma_i} - \Gamma_i \end{cases} \quad \begin{cases} L(\alpha_2) & = 0 & \text{in } \Omega \\ \sigma(\alpha_2) \cdot \vec{n} & = \sigma(\alpha) \cdot \vec{n} & \text{on } \Gamma_i, \\ \alpha_2 & = 0 & \text{on } \partial S_{\Gamma_i} - \Gamma_i. \end{cases}$$

Applying the inequality of trace on the boundary of the the square  $S_{\Gamma_i}$ , which is a parallelogram (see lemma 4.2 in [3]), for the components of  $u_{2,i}$  on  $\Gamma_i$ , it yields for both  $k = 1, 2$

$$\|\alpha_k\|_{0,\Gamma_i}^2 \leq 2 \frac{|\Gamma_i|}{|\Gamma_i|^2} \|\alpha_k\|_{0,S_{\Gamma_i}}^2 + 2 \frac{|\Gamma_i|^2}{|\Gamma_i|} \|\nabla \alpha_k\|_{0,S_{\Gamma_i}}^2,$$

simplifying,

$$\|\alpha_k\|_{0,\Gamma_i}^2 \leq 2 \frac{1}{|\Gamma_i|} \|\alpha_k\|_{0,S_{\Gamma_i}}^2 + 2|\Gamma_i| \|\nabla \alpha_k\|_{0,S_{\Gamma_i}}^2,$$

using estimate (49) for the case where  $\Omega$  is the square  $S_{\Gamma_i}$ , with  $d(S_{\Gamma_i}) = \sqrt{2}|\Gamma_i|$ , yields

$$\begin{aligned} \|\alpha_k\|_{0,\Gamma_i}^2 &\leq 2 \frac{2|\Gamma_i|^2}{4|\Gamma_i|} \|\nabla \alpha_k\|_{0,S_{\Gamma_i}}^2 + 2|\Gamma_i| \|\nabla \alpha_k\|_{0,S_{\Gamma_i}}^2 \\ &\leq 3|\Gamma_i| \|\nabla \alpha_k\|_{0,S_{\Gamma_i}}^2, \end{aligned}$$

applying Korn's inequality for the extended function  $\tilde{\alpha}$  defined on the half-plane  $\mathbb{R}^{2+}$  containing  $S_{\Gamma_i}$ , for which case Korn constant equals  $\frac{1}{2}$ , we obtain

$$\|\alpha_k\|_{0,\Gamma_i}^2 \leq 12|\Gamma_i| \|\varepsilon(\alpha_k)\|_{0,S_{\Gamma_i}}^2. \quad (52)$$

Since  $\alpha_1$  and  $\alpha_2$  satisfy

$$\|\varepsilon(\alpha_1)\|_{0,S_{\Gamma_i}}^2 = \int_{S_{\Gamma_i}} L(\alpha)\alpha_1 \, dx \quad \text{and} \quad \|\varepsilon(\alpha_2)\|_{0,S_{\Gamma_i}}^2 = \int_{\Gamma_i} \sigma(\alpha) \cdot \vec{n} \, \alpha_2 \, d\sigma,$$

then, using Cauchy-schwarz, this leads to estimates

$$\|\varepsilon(\alpha_1)\|_{0,S_{\Gamma_i}} \leq 2\sqrt{2}|\Gamma_i| \|L(\alpha)\|_{0,S_{\Gamma_i}} \quad \text{and} \quad \|\varepsilon(\alpha_2)\|_{0,S_{\Gamma_i}} \leq \|\sigma(\alpha) \cdot \vec{n}\|_{0,\Omega} \|\alpha_2\|_{0,\Gamma_i},$$

using these last estimates, (52) becomes

$$\|\alpha_1\|_{0,\Gamma_i}^2 \leq 12 \times 8|\Gamma_i|^2 \|L(\alpha)\|_{0,S_{\Gamma_i}}^2, \quad \|\alpha_2\|_{0,\Gamma_i}^2 \leq 12|\Gamma_i| \|\sigma(\alpha) \cdot \vec{n}\|_{0,\Gamma_i} \|\alpha_2\|_{0,\Gamma_i}.$$

Using (46) and assumption (51), we can find  $\delta_5$  and  $\delta_6$  small enough such that

$$\|\alpha_1\|_{0,\Gamma_i} \leq \delta_5, \quad \|\alpha_2\|_{0,\Gamma_i} \leq 12|\Gamma_i| \|g_i\|_{0,\Gamma_i} + \delta_6,$$

for all  $i$ . Hence, combining these two estimates, we obtain

$$\|u_{2,i}\|_{0,\Gamma_i} \leq \|\alpha_1\|_{0,\Gamma_i} + \|\alpha_2\|_{0,\Gamma_i} \leq \delta_9 + 16|\Gamma_i| \|g_i\|_{0,\Gamma_i}$$

with  $\delta_9 = \delta_5 + \delta_6$  arbitrary small. □

## 4 Proof of theorem 1

We are ready now to present a proof of the main theorem. It uses, principally, lemma (6) and lemma (7).

*Proof. Step 1:* Let  $u_1$  be as defined in lemma (6), it is easy to check that it satisfies

$$2\mu \int_{\Omega} \varepsilon(u_1)\varepsilon(v) \, dx + \lambda \int_{\Omega} \operatorname{div} u_1 \operatorname{div} v \, dx = \int_{\Omega} f_1 v \, dx$$

for all  $v \in H_0^1(\Omega)$ . Choose  $v = u_1$  and use Cauchy-Schwarz inequality

$$2\mu \|\varepsilon(u_1)\|_{0,\Omega}^2 + \lambda \|\operatorname{div} u_1\|_{0,\Omega}^2 \leq \|f_1\|_{0,\Omega} \|u_1\|_{0,\Omega}$$

Using Korn's inequality relatively to the case of homogeneous Dirichlet condition on one hand, and Poincaré inequality on the other hand, yield



$$\frac{\mu}{2} \|\nabla u_1\|_{0,\Omega}^2 \leq \|f_1\|_{0,\Omega} \frac{d(\Omega)}{2} \|\nabla u_1\|_{0,\Omega},$$

using (iv) of lemma (6), we obtain

$$\mu \|\nabla u_1\|_{0,\Omega} \leq d(\Omega) \|f^p\|_{0,\Omega} + d(\Omega) \frac{\delta_0}{m}. \quad (53)$$

Fix  $i$ ,  $1 \leq i \leq m$ . Let  $u_{2,i}$  be such as defined in lemma (7), it is easy to check that  $u_{2,i}$  satisfies

$$2\mu \int_{\Omega} \varepsilon(u_{2,i}) \varepsilon(v) \, dx + \lambda \int_{\Omega} \operatorname{div} u_{2,i} \operatorname{div} v \, dx = \int_{\Omega} L(u_{2,i}) v \, dx + \int_{\Gamma-\Gamma_0} g_i v \, d\sigma$$

for all  $v \in V$ . Choose  $v = u_{2,i}$  and use Cauchy-Schwarz to obtain

$$2\mu \|\varepsilon(u_{2,i})\|_{0,\Omega}^2 + \lambda \|\operatorname{div} u_{2,i}\|_{0,\Omega}^2 \leq \|L(u_{2,i})\|_{0,\Omega} \|u_{2,i}\|_{0,\Omega} + \|g_i\|_{0,\Gamma_i} \|u_{2,i}\|_{0,\Gamma_i},$$

using estimate (49)

$$2\mu \|\varepsilon(u_{2,i})\|_{0,\Omega}^2 \leq \|L(u_{2,i})\|_{0,\Omega} d(\Omega) \|\nabla u_{2,i}\|_{0,\Omega} + \|g_i\|_{0,\Gamma_i} \|u_{2,i}\|_{0,\Gamma_i}.$$

On the other hand, there exists constants  $c_k$ ,  $c_p$  and  $c_t$  such that

$$\|\nabla u_{2,i}\|_{0,\Omega} \leq c_p \|L(u_{2,i})\|_{0,\Omega} + c_t c_t \|g_i\|_{0,\Gamma_i}$$

thus

$$\begin{aligned} 2\mu \|\varepsilon(u_{2,i})\|_{0,\Omega}^2 &\leq d(\Omega) \|L(u_{2,i})\|_{0,\Omega} c_k (c_p \|L(u_{2,i})\|_{0,\Omega} + c_t \|g_i\|_{0,\Gamma_i}) + \|g_i\|_{0,\Gamma_i} \|u_{2,i}\|_{0,\Gamma_i}; \\ &\quad (54) \end{aligned}$$

since, following (27) and (45),  $\|g_i\|_{0,\Omega}$  are uniformly bounded with respect to  $\epsilon$  then, using (46)

$$\begin{aligned} 2\mu \|\varepsilon(u_{2,i})\|_{0,\Omega}^2 &\leq d(\Omega) \tau c_k (c_p \tau + c_t \|g_i\|_{0,\Gamma_i}) + \|g_i\|_{0,\Gamma_i} \|u_{2,i}\|_{0,\Gamma_i} \\ &\leq d(\Omega) \tau c_k (c_p \tau + c_t (\|g\|_{0,\Gamma-\Gamma_0} + \delta_6 + \frac{\delta_1}{m})) + \|g_i\|_{0,\Gamma_i} \|u_{2,i}\|_{0,\Gamma_i}; \\ &\quad (55) \end{aligned}$$

pose

$$\delta_7^2 := d(\Omega) \tau c_k (c_p \tau + c_t (\|g\|_{0,\Gamma-\Gamma_0} + \delta_6 + \frac{\delta_1}{m})),$$

it can be made, thanks to estimate (46), as small as desired. Applying (50), (27) and (45), estimate (55) becomes

$$\begin{aligned}
2\mu \|\varepsilon(u_{2,i})\|_{0,\Omega}^2 &\leq \delta_7^2 + 16|\Gamma_i| \|g_i\|_{0,\Gamma_i}^2 + \delta_9 \|g_i\|_{0,\Gamma_i} \\
&\leq \delta_7^2 + 16|\Gamma_i| \|g_i\|_{0,\Gamma_i}^2 + \delta_9 \left( \frac{\delta_1}{m} + \delta_6 + \|g\|_{0,\Gamma_i} \right) \\
&\leq \frac{\delta_{10}^2}{m^2} + 16|\Gamma_i| \|g_i\|_{0,\Gamma_i}^2,
\end{aligned}$$

where  $\frac{\delta_{10}^2}{m^2} := \delta_7^2 + \delta_9 \left( \frac{\delta_1}{m} + \delta_6 + \|g\|_{0,\Gamma_i} \right)$ , then by simplifying,

$$\sqrt{2\mu} \|\varepsilon(u_{2,i})\|_{0,\Omega} \leq \frac{\delta_{10}}{m} + 4|\Gamma_i|^{\frac{1}{2}} \|g_i\|_{0,\Gamma_i}. \quad (56)$$

One should notice that, by definition of the constants  $\delta_6$ ,  $\delta_7$  and  $\delta_9$ , the constant  $\delta_{10}$  can be made as small as possible.

**Step 2:** Since the deformation  $\varepsilon(u_{2,i})$  is a linear application with respect to the first derivatives of  $u_{2,i}$  then, with the same notations as in (47) and by using (48), we have

$$\|\varepsilon(\tilde{u}_{2,i})\|_{0,\mathbb{R}^{2+}} = \|\varepsilon(\tilde{u}_{2,i})\|_{0,\Omega} = \|\varepsilon(u_{2,i})\|_{0,\Omega_i}.$$

Applying the estimate stated in (corollary 1.2.2, [6]) to  $\tilde{u}_{2,i}$  gives

$$\frac{1}{2} \times \|\nabla u_{2,i}\|_{0,\Omega} = \frac{1}{2} \times \|\nabla \tilde{u}_{2,i}\|_{0,\mathbb{R}^{2+}} \leq \|\varepsilon(\tilde{u}_{2,i})\|_{0,\mathbb{R}^{2+}},$$

thus, (56) becomes

$$\frac{\sqrt{\mu}}{\sqrt{2}} \|\nabla u_{2,i}\|_{0,\Omega} \leq \frac{\delta_{10}}{m} + 4|\Gamma_i|^{\frac{1}{2}} \|g_i\|_{0,\Gamma_i},$$

Young inequality yields

$$\frac{\sqrt{\mu}}{\sqrt{2}} \|\nabla u_{2,i}\|_{0,\Omega} \leq \frac{\delta_{10}}{m} + 2(|\Gamma_i| + \|g_i\|_{0,\Gamma_i}^2).$$

Using approximation result (27), we have

$$\frac{\sqrt{\mu}}{\sqrt{2}} \|\nabla u_{2,i}\|_{0,\Omega} \leq \frac{\delta_{10}}{m} + 4(|\Gamma_i| + \|g_p^\varepsilon\|_{0,\Gamma_i}^2 + \frac{\delta_1^2}{m^2}).$$

Summing over  $i = \overline{1, m}$  we obtain using triangular inequality

$$\frac{\sqrt{\mu}}{\sqrt{2}} \|\nabla u_2'\|_{0,\Omega} \leq \frac{\sqrt{\mu}}{\sqrt{2}} \sum_i \|\nabla u_{2,i}\|_{0,\Omega} \leq \delta_{10} + 4|\Gamma| + 4\|g_p^\varepsilon\|_{0,\Gamma}^2 + 4\delta_1^2. \quad (57)$$

Finally, combining (53) and (57) we get

$$\begin{aligned} & \sqrt{\mu} \|\nabla(u_1 + u'_2)\|_{0,\Omega} \\ & \leq \frac{\delta_0}{m} d(\Omega) + \sqrt{2}\delta_{10} + 4\delta_1^2 + d(\Omega) \|f^p\|_{0,\Omega} + 8(|\Gamma| + \|g_p^\epsilon\|_{0,\Gamma}^2), \end{aligned}$$

considering (28) and posing  $\delta_{11} := \frac{\delta_0}{m} d(\Omega) + \sqrt{2}\delta_{10} + 4\delta_1^2$ ,

$$\sqrt{\mu} \|\nabla u_\epsilon^p\|_{0,\Omega} \leq \delta_{11} + \sqrt{\mu}\delta_2 + d(\Omega) \|f^p\|_{0,\Omega} + 8(|\Gamma| + \|g_p^\epsilon\|_{0,\Gamma}^2). \quad (58)$$

Use (45), estimate (58) becomes

$$\|\nabla u_\epsilon^p\|_{0,\Omega} \leq \frac{1}{\sqrt{\mu}} [8\delta_6^2 + \delta_{11} + \sqrt{\mu}\delta_2 + d(\Omega) \|f^p\|_{0,\Omega} + 8(|\Gamma| + \|g\|_{0,\Gamma}^2)], \quad (59)$$

using (23) and (i) of (24), estimate (59) becomes

$$\|\nabla u_\epsilon\|_{0,\Omega} \leq \frac{1}{\sqrt{\mu}} \left[ \frac{1}{2} + 8\delta_6^2 + \sqrt{\mu}\delta_2 + \delta_{11} + d(\Omega) \|f\|_{0,\Omega} + 8(|\Gamma| + \|g\|_{0,\Gamma}^2) \right].$$

By adequately choosing the positive numbers  $\delta'_j$ 's, we immediately get

$$\|\nabla u_\epsilon\|_{0,\Omega} \leq \frac{1}{\sqrt{\mu}} [1 + d(\Omega) \|f\|_{0,\Omega} + 8(|\Gamma| + \|g\|_{0,\Gamma}^2)]. \quad (60)$$

We conclude the theorem for  $u$  by applying (a) of lemma (4) on one hand, and (iii) of remark (5) on the other hand.  $\square$

Finally; in order to get the explicit  $H^1$  estimate of  $u_\epsilon$ , and so that of  $u$ , we use the poincaré inequality (49) to bound  $\|u\|_{0,\Omega}$  at one hand and the estimate (7) at the other hand.

## Conclusion

In the point of view of numerical analysis, estimate of theorem (1) is interesting. Indeed, error estimates in finite element method of the type

$$\|u - u_h\|_{0,\Omega} \leq Ch \|\nabla u\|_{0,\Omega}$$

involve the quantity  $\|\nabla u\|_{0,\Omega}$ . Assuming that the constant  $C$  can be explicitly computed, then it is possible to explicitly bound  $\|\nabla u\|_{0,\Omega}$  which implies a better estimate of  $\|u\|_{0,\Omega}$ .

Another interesting feature of the estimate (7) that makes it effective is that it does not depend on the characteristic parameters of the polygonal domain  $\Omega$ , namely, the edges's length, their number as well as the measures of the angles. The estimate is therefore indifferently applicable to all polygons. All this allows the possibility to generalize this result, by approximation, to a  $C^1$  class domain.

## A Clarification

Regarding the application of the main result established in [10], we note the following facts. The part of theorem 2.1 in [10] used in the proof of lemma (7) is the equivalence between the following two assertions.

1)  $X_i$  is approximated by smooth functions with support away from  $\Gamma - \Gamma_i$

$$2) \lim_{r \rightarrow 0} \frac{1}{|B(x, r)|} \int_{B(x, r) \cap \Omega} |X_i(y)| \, dy = 0$$

To effectively apply this result, we need to show that the fact  $X_i = 0$  on  $\Gamma - \Gamma_i$  implies 2). Indeed, let  $x \in \Gamma - \Gamma_i$ . We assume for convenience, that  $x$  coincide with the origin  $(0, 0)$  of  $\mathbb{R}^2$ . Since  $\partial\Omega$  is a polygon then, there exists  $c > 0$  independent of  $r$  such that

$$|B(0, r) \cap \Omega| = c|B(0, r)|.$$

On the other hand, the function  $X_i$  solution of problem  $(P_1)$  is continuous on  $\bar{\Omega}$ . One can see this using the following argument: according to the definition of  $w_\epsilon^p$ ,  $X_i \in C^\infty(\partial\Omega)$ . Applying the Whitney extension theorem (see [11]), one can find  $\bar{X}_i \in C^\infty(\bar{\Omega})$  that coincide with  $X_i$  on  $\Gamma$ . Thus,  $X_i - \bar{X}_i \in H^1(\Omega)$  solves the following problem

$$\begin{cases} L(X_i - \bar{X}_i) = -L(\bar{X}_i) & \text{a.e in } \Omega, \\ X_i - \bar{X}_i = 0 & \text{on } \cup_i \Gamma_i. \end{cases}$$

Since  $\Omega$  is convex,  $X_i - \bar{X}_i \in H^2(\Omega)$ . Consequently

$$X_i = (X_i - \bar{X}_i) + \bar{X}_i \in H^2(\Omega)$$

as well. Furthermore, by an adequate Sobolev embedding,  $X_i \in C^0(\bar{\Omega})$ . The Lebesgue differentiation theorem yields:

$$\begin{aligned} \lim_{r \rightarrow 0} \frac{1}{|B(x, r)|} \int_{B(x, r) \cap \Omega} |X_i(y)| \, dy &= \lim_{r \rightarrow 0} \frac{1}{|B(0, r)|} \int_{B(0, r) \cap \Omega} |X_i(y)| \, dy \\ &\leq \lim_{r \rightarrow 0} \frac{1}{|B(0, r) \cap \Omega|} \int_{B(0, r) \cap \Omega} |X_i(y)| \, dy \\ &= |X_i(0)| = 0, \end{aligned}$$

we infer that assertion 2) holds for all  $x \in \Gamma - \Gamma_i$  and thus,  $C_{1,2}$ -almost everywhere. Where  $C_{1,2}$  refers to the 2-capacity of the set  $\Gamma - \Gamma_i$ .

## References

- [1] R. A. Adams. J. J. Fournier. *Sobolev spaces*. 2nd ed., Academic Press, 1975.
- [2] G. Allaire. *Analyse numérique et optimisation. Une introduction à la modélisation et à la simulation numérique*. Editions de l'école polytechnique 2004.
- [3] C. Carstensen, S.A. Sauter, *A posteriori error analysis for elliptic PDEs on domains with complicated structures*, Numerische Mathematik, vol 96, issue 4, pp.691-721, February 2004.
- [4] P. Doktor. A. Zenisek. *The density of infinitely differentiable functions in Sobolev spaces with mixed boundary conditions*. Applications of Mathematics. Volume 51, Issue 5, pp 517-547, October 2006.
- [5] G. Duvaut, J. L. Lions. *Les inéquations en mécanique et en physique*. Dunod, Paris, 1972.
- [6] C. Eck, J. Jarusek, M. Krbec. *Unilateral contact problems: variational methods and existence theorems*. Series: Chapman and Hall/CRC Pure and Applied Mathematics, March 2005.
- [7] C. O. Horgan. *Korn's inequalities and their applications in continuum mechanics*. SIAM Review. Vol. 37, No.4, pp 491-511, December 1995.
- [8] E. Di Nezza. G. Palatucci. E. Valdinoci. *Hitchhiker's guide to the fractional Sobolev spaces*. Bulletin des Sciences Mathématiques. Volume 136, Issue 5 , Pages 521-573, July-August 2012.
- [9] S. Nicaise. *About the Lamé system in a polygonal or polyhedral domain and a coupled problem between the Lamé system and the plate equation. I: regularity of the solutions*. Annali della scuola Normale superiore di Pisa, classe di scienze 4<sup>e</sup> série, tome 19, n<sup>o</sup>3, p.p. 327-361, 1992.
- [10] M. Egert. P. Tolksdorf. *Characterizations of Sobolev Functions that vanish on a part of the boundary*. Discrete and Continuous Dynamical Systems - S, 10(4), 729-743, 2017.
- [11] H. Whitney. *Analytic extensions of differentiable functions defined in closed sets*. Trans. Amer. Math. Soc. 36, 63-89, 1934.