



**HAL**  
open science

## How do food sources drive meiofauna community structure in soft-bottom coastal food webs?

L.H. van Der Heijden, J. Rzeznik-Orignac, R. Asmus, D. Fichet, M. Bréret, P. Kadel, L. Beaugeard, H. Asmus, B. Lebreton

### ► To cite this version:

L.H. van Der Heijden, J. Rzeznik-Orignac, R. Asmus, D. Fichet, M. Bréret, et al.. How do food sources drive meiofauna community structure in soft-bottom coastal food webs?. *Marine Biology*, 2018, 165 (10), pp.166. 10.1007/s00227-018-3419-7 . hal-01994251

**HAL Id: hal-01994251**

**<https://hal.science/hal-01994251>**

Submitted on 18 Feb 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **How do food sources drive meiofauna community structure in soft-bottom coastal food webs?**

2 L.H. van der Heijden<sup>1,2</sup>, J. Rzeznik-Orignac<sup>3</sup>, R. Asmus<sup>2</sup>, D. Fichet<sup>1</sup>, M. Bréret<sup>1</sup>, P. Kadel<sup>2</sup>, L. Beaugéard<sup>1</sup>,  
3 H. Asmus<sup>2</sup>, B. Lebreton<sup>1</sup>

4 <sup>1</sup>*UMR 7266 Littoral, Environnement et Sociétés (CNRS - University of La Rochelle), Institut du littoral et*  
5 *de l'environnement, 2 rue Olympe de Gouges, 17000 La Rochelle, France*

6 <sup>2</sup>*Alfred-Wegener-Institut Helmholtz-Zentrum für Polar- und Meeresforschung, Wattenmeerstation Sylt,*  
7 *Hafenstrasse 43, 25992 List/Sylt, Germany*

8 <sup>3</sup>*Sorbonne Universités, UPMC Univ Paris 06, CNRS, Laboratoire d'Ecogéochimie des Environnements*  
9 *Benthiques (LECOB UMR 8222), Observatoire Océanologique, 66650 Banyuls/mer, France*

10

11 Corresponding author: L.H. van der Heijden

12 Email address: [luuk.van\\_der\\_heijden@univ-lr.fr](mailto:luuk.van_der_heijden@univ-lr.fr)

13 Phone number: +33 (0) 546 507 631

## 14 **Abstract**

15 Meiofauna have a strong role in the functioning of coastal ecosystems but how their food sources  
16 affect their community structure remains unclear, likely due to the influence of abiotic parameters and the  
17 small size of these organisms. The meiofauna and their potential food sources were sampled in several  
18 intertidal habitats (i.e. mudflat, seagrass bed, sandflat) of the Marennes-Oléron Bay, France, and the Sylt-  
19 Rømø Bight, Germany, to assess the relationships between habitat characteristics and meiofauna  
20 community structure. Biomass and quality (carbon/chlorophyll *a* ratios) of food sources were estimated.  
21 Meiofauna community structure based on density and biomass was determined, as well as nematode  
22 diversity and feeding types. Meiofauna density and biomass varied highly within habitats and within  
23 ecosystems, ranging from  $1.7 \cdot 10^5$  ind.  $m^{-2}$  to  $3.4 \cdot 10^6$  ind.  $m^{-2}$  and from  $0.057$  gC  $m^{-2}$  to  $1.541$  gC  $m^{-2}$ ,  
24 respectively. Benthic microalgae and sediment organic matter are important drivers in these food webs as  
25 the density of two important trophic groups of nematodes, non-selective deposit feeders and epigrowth  
26 feeders, varied in accordance with these food sources. No clear relationship was observed between  
27 bacterial biomass and selective deposit feeders (i.e. bacterivores) probably due to the high production rate  
28 of bacteria. Complementary information about production rates of food source as well as data from  
29 trophic markers would provide complimentary information to better understand flows of organic matter in  
30 intertidal habitats, particularly for opportunistic species such as non-selective deposit feeders.

## 31 **Introduction**

32 Soft-bottom intertidal habitats are highly productive (Underwood and Kromkamp 1999; Hemminga  
33 and Duarte 2000), fueling complex food webs and providing food resources for a large diversity of top  
34 consumers like fish and birds (Day et al. 1989; Duarte and Cebrián 1996) via trophic pathways going  
35 through meiofauna (Chardy and Dauvin 1992) and macrofauna (Haubois et al. 2005). Meiofauna, here  
36 defined as metazoans associated with sediments with a size range from 40 to 500  $\mu m$ , are considered as  
37 an important trophic component in the functioning of coastal ecosystems. They have an intermediate  
38 trophic position (Leguerrier et al. 2003; Giere 2009) and high secondary production (from 4 to 29 gC  $m^{-2}$   
39  $year^{-1}$ ) (Escaravage et al. 1989; Chardy and Dauvin 1992; Danovaro et al. 2002) that often exceeds that of  
40 the macrofauna (Giere 2009). Meiofauna are connected to the microorganisms in the food web via the  
41 consumption of microalgae and bacteria (Rzeznik-Orignac et al. 2003; Vafeiadou et al. 2014) and to the  
42 food webs of macroconsumers especially via their consumption by small benthic fish and predatory  
43 invertebrates (Coull 1999). Nematodes are the most dominant group in meiofauna; their food resources  
44 can be very specific, or in the contrary, relatively diverse, and their buccal cavity is very distinct based on  
45 these feeding types. This morphological trait has been used for decades to define feeding types (Wieser  
46 1953; Warwick 1971; Heip et al. 1985) and some more recent methods (i.e. feeding behavior

47 experiments; (Moens and Vincx 1997), trophic markers (Moens et al. 2005)) permitted to revise or  
48 precise the feeding types of some genera (e.g. assort of *Metachromadora* from the omnivores/predators to  
49 the epigrowth feeders).

50 Both, biotic and abiotic parameters, have been shown to affect meiofauna community structure.  
51 Among biotic parameters, food availability in general (Castel et al. 1989; Danovaro 1996), availability of  
52 specific food sources such as bacteria and microphytobenthos (Danovaro and Gambi 2002; Fonseca et al.  
53 2011), the quality of the food (Danovaro et al. 2002) and the deposition of organic material of pelagic  
54 origin (Franco et al. 2008) have been identified as structuring parameters. Abiotic parameters such as  
55 sediment grain-size (Steyaert et al. 2003; Dupuy et al. 2015), habitat structure (Fonseca et al. 2011),  
56 oxygen gradients (Josefson and Widbom 1988), intertidal water level (Steyaert et al. 2001) as well as  
57 anthropogenic impacts (Semprucci et al. 2010) are influencing meiofauna community structure. Studies  
58 on meiobenthic communities in intertidal systems have been carried out mainly on single habitats,  
59 focusing on effects of abiotic and biotic parameters at the habitat scale (Reise 1985; Rzeznik-Orignac et  
60 al. 2003; Lebreton et al. 2012). Factors affecting the community structure of meiofauna at a larger scale  
61 (Moens et al. 2013) in intertidal habitats are still poorly known, as few studies have compared several  
62 habitats and/or different ecosystems.

63 The present study investigates whether meiobenthic communities, especially free-living nematodes  
64 and benthic copepods, differ among habitats with different characteristics (i.e. mudflats, seagrass beds  
65 and sandflats; influence of terrestrial organic matter inputs or not) to define which factors are structuring  
66 these communities. Particular attention is paid to the role of food sources, as they have been highlighted  
67 as important drivers of ecosystem functioning in intertidal habitats (Asmus and Asmus 1985; Underwood  
68 and Kromkamp 1999; McLusky and Elliott 2004). Aim of this study is therefore to define if changes in  
69 biomass and quality of food sources can affect meiofauna community structure. If yes, the objective is to  
70 determine—among all available food sources (i.e. benthic microalgae, sediment organic matter (SOM),  
71 suspended particulate organic matter (SPOM) and bacteria)—what food sources play a role and how  
72 changes in these food sources influence meiofauna community structure. In this aim, a temporal sampling  
73 was done in spring and autumn in several habitats to define the temporal and spatial variations of the  
74 food sources and those of meiofauna community structure.

## 75 **Materials and methods**

### 76 **Study sites**

77 *Marennes-Oléron Bay*

78 The Marennes-Oléron Bay is a semi-enclosed system, located in the middle of the western Atlantic  
79 coast of France, between the Oléron Island and the mainland (Fig. 1). The bay has a surface of 180 km<sup>2</sup>  
80 (Gouleau et al. 2000) and is influenced by both oceanic and continental inputs, mainly from the Charente  
81 River (river input ranges from 15 to 500 m<sup>3</sup> s<sup>-1</sup>) (Ravail et al. 1988). Tides are semi-diurnal, with a period  
82 of emersion of ca. 6 h per tidal cycle and a tidal range from 0.9 to 6.5 m (Kervella 2009). The system is  
83 dominated by intertidal bare mudflats (91 km<sup>2</sup>, i.e. 58%) which have a flat bottom slope (1:1000) and  
84 represent a large tidal area (up to 4.5 km wide) (Gouleau et al. 2000).

85 The Brouage mudflat, located on the eastern part of the bay, is the most extensive mudflat (40 km<sup>2</sup>)  
86 (Fig. 1). The sediment consists mainly of silt and clay particles (95%) with a size smaller than 63 µm  
87 (Pascal et al. 2009). Seagrass beds, consisting of the species *Zostera noltii* (Lebreton et al. 2009), are  
88 located on the western side of the bay, along the coast of the Oléron Island, which is more influenced by  
89 offshore waters than the eastern side (Dechambenoy et al. 1977). The seagrass meadow is stretched out  
90 over 15 km along the upper part of the flat, with an average width of 1.5 km (Guillaumont 1991).


#### 91 *Sylt-Rømø Bight*

92 The Sylt-Rømø Bight is a semi-enclosed basin, located in the north of Germany between the  
93 mainland of Germany and Denmark, the islands of Sylt (Germany) and of Rømø (Denmark) (Fig. 1). The  
94 bight consists of a mosaic of habitats, mainly intertidal and subtidal sandflats, seagrass beds, and mudflats  
95 (Asmus and Asmus 1985). The bight has a coverage of 404 km<sup>2</sup> of which 135 km<sup>2</sup> are intertidal (Asmus  
96 and Asmus 1985) and the tidal gauge is 1.8 m on average (Asmus and Asmus 1985).

97 The *Arenicola* sandflat is the most represented intertidal habitat in the bay with 91 km<sup>2</sup> (67%)  
98 (Asmus and Asmus 2005). However, *Zostera noltii* seagrass beds expanded over the last years, from 15  
99 km<sup>2</sup> in 1995 to 83 km<sup>2</sup> (i.e. 21% of the bay surface) in 2010 (Reise and Kohlus 2008). The intertidal bare  
100 mudflats are poorly represented in the bight, with an area of 4 km<sup>2</sup> (Fig. 1).

#### 101 *Field sampling and laboratory processes*

102 Three intertidal habitats were studied in the two bays: a bare mudflat (mudflat MO) and a seagrass  
103 bed (seagrass MO) in the Marennes-Oléron Bay, and a bare mudflat (mudflat SR), a seagrass bed  
104 (seagrass SR) and a sandflat (sandflat SR) in the Sylt-Rømø Bight (Fig. 1). Based on knowledge already  
105 gained about these habitats in each of these five locations (Leguerrier et al. 2003; Baird et al. 2007;  
106 Pacella et al. 2013; Saint-Béat et al. 2013) one sampling station that was the most representative for the  
107 studied habitat was selected.



108

109 **Figure 1.** Sampling stations of the mudflats, seagrass beds and sandflat in the Marennnes-Oléron Bay (MO) and the  
 110 Sylt-Rømø Bight (SR) along the European coast. Pelagic sampling stations, where samples for suspended particulate  
 111 organic matter were taken, are indicated by roman numbers in both Marennnes-Oléron Bay (I-II) and Sylt-Rømø  
 112 Bight (III-V).

113 *Sediment grain size*

114 Sediment samples ( $n = 3$ ) were collected randomly with cores ( $\varnothing = 4$  cm) and were split into 0-1 cm  
 115 and 1-5 cm depth fractions. Sediment grain size was determined by laser granulometry analyses for the  
 116 sediment from the seagrass bed and mudflats (Mastersize 2000, Malvern Instruments Limited, United  
 117 Kingdom). The freeze-dried sediment of these habitats was rehydrated before analyses in order to avoid  
 118 the use of ultrasound to disaggregate the sediment. Sediment grain size of freeze-dried samples from the  
 119 sandflat of the Sylt-Rømø Bight was determined by dry sieving. Samples were classified based on their  
 120 median sediment grain size after Wentworth (1922).

121 *Benthic primary producers, composite sources and bacteria*

122 For all stations, sampling was carried out in two seasons, in spring (from May 17 to June 17, 2016)  
 123 and in autumn (from November 14 to December 15, 2016), to compare a season with a high primary  
 124 production with a season with a high load of detrital matter. Sampling was carried out during low tide,  
 125 starting about one hour before lowest water level.

126 Samples for sediment organic matter (SOM), detritus, and seagrass leaves and roots (3 replicates)  
127 were taken using cores ( $\varnothing = 19$  cm) which were separated in two layers (a surface layer from 0-1 cm, a  
128 sub-surface layer from 1-5 cm). The collected material was gently sieved on a 500  $\mu\text{m}$  mesh screen (1  
129 mm for sandflat SR samples) after which the upper fraction was sorted in seagrass leaves, roots and  
130 detritus (brown to black colored leaves and roots). All samples were freeze-dried, then weighed ( $\pm 0.001$   
131 g). Carbon to dry weight ratios were determined on each sample: A small amount was ground to a fine  
132 and homogeneous powder using a ball mill (MM 400, Retsch, Germany) and carbon content was  
133 determined using an elemental analyzer (Flash EA 1112, Thermo Scientific, Italy). Subsamples were  
134 collected in the lower fraction (SOM), freeze-dried, and ground using a mortar and pestle. Chlorophyll *a*  
135 and organic carbon concentrations were determined in sub-samples by fluorimetry (TD-700, Turner  
136 Designs, USA) using the method of Lorenzen (1966) and elemental analyses (Flash EA 1112, Thermo  
137 Scientific, Italy), respectively. Chlorophyll *a* values were converted into carbon biomass using the  
138 conversion ratio (47.63) provided by de Jonge (1980) and used as a proxy of fresh material of microalgae  
139 origin (i.e. microphytobenthos and trapped phytoplankton) in sediment, thereafter called “benthic  
140 microalgae”. Carbonates were removed prior to elemental analyses by adding HCl at 2 mol.l<sup>-1</sup> drop-by-  
141 drop on SOM samples until cessation of bubbling. Subsequently samples were dried at 60°C to constant  
142 weight, freeze-dried and ground again.

143 Bacterial biomass (3 replicates) was determined using cores ( $\varnothing = 4$  cm) which were separated in two  
144 layers (0-1 cm and 1-5 cm). These sediment layers were homogenized with a spatula and 2 mL of this  
145 material were mixed with 2 mL formalin (4%). Samples were deep frozen in liquid nitrogen and stored at  
146 -80°C. The number of bacteria was determined following the protocol of Lavergne et al. (2014). Samples  
147 were diluted (1/2000) using sodium pyrophosphate (P-8010, Sigma Aldrich, France) and Tween 80 (P-  
148 1754, Sigma Aldrich, France), homogenized and then incubated at 4 °C during 30 minutes. Samples were  
149 then sonicated at 60 W during 30 seconds, after which 1 mL of the pre-diluted sample was stained with 20  
150  $\mu\text{L}$  of SYBR Green I (Invitrogen-11540746; 1/5000 final concentration) and incubated in the dark at  
151 room temperature for 15 minutes. Afterwards, 2  $\mu\text{L}$  of calibration beads (1  $\mu\text{M}$ ) were added and the  
152 samples were analyzed using a flow cytometer (BD FACSCANTO II, BD Biosciences, USA). Bacterial  
153 densities were converted into carbon using the conversion factor (20 fgC cell<sup>-1</sup>) established by Lee and  
154 Fuhrman (1987).

#### 155 *Suspended particulate organic matter*

156 Sampling of the suspended particulate organic matter (SPOM) was conducted by collecting samples  
157 of surface water during high tide, in the same timeframe as the benthic samples. Water was sampled at  
158 two stations (I and II) in the Marennes-Oléron Bay and three stations (III, IV and V) in the Sylt-Rømø

159 Bight (Fig. 1). Station I, in the north of the Brouage mudflat, represented the water mass covering the  
160 mudflat MO station. Station II, near the inlet of the Marennes-Oléron bay, represented the water mass  
161 influencing the seagrass MO station. Station II is regularly sampled in the framework of the SOMLIT  
162 surveys (data provided by “Service d’Observation en Milieu Littoral, INSU-CNRS, La Rochelle station”).  
163 Water samples collected at stations III and IV represented water masses covering the sandflat SR (III),  
164 and the mudflat SR and seagrass SR stations (IV), respectively. Station V, near the entrance of the Sylt-  
165 Rømø Bight, represented the SPOM originating from the North Sea.

166 SPOM was sampled by collecting 10 to 100 mL of surface seawater which was pre-filtered on a 250-  
167  $\mu\text{m}$  mesh screen to eliminate large particles/organisms. Water samples were then filtered on  
168 precombusted (4 hours, 450°C) Whatman GF/F glass fiber filters (0.7  $\mu\text{m}$  nominal porosity) under  
169 moderate vacuum and these filters were freeze-dried. Chlorophyll *a* content, used as a proxy of  
170 phytoplankton biomass, of the freeze-dried GF/F filters was measured by fluorimetry using the method of  
171 Lorenzen (1966). Phytoplankton biomass was determined by using the conversion factor (56.4)  
172 established by Lü et al. (2009).

### 173 *Meiofauna*

174 Meiofauna was sampled for taxonomic composition, density (ind.  $\text{m}^{-2}$ ) and biomass ( $\text{gC m}^{-2}$ )  
175 estimations at the same time as benthic primary producers and composite sources. Three replicates were  
176 collected randomly in a ten square meters area using cores ( $\varnothing = 4 \text{ cm}$ ). Cores were separated in two layers  
177 (0-1 cm and 1-5 cm) to study the changes of meiofauna communities related to sediment depth. Samples  
178 were sieved with a 500- $\mu\text{m}$  mesh to eliminate macrofauna and large detritus. Meiofauna was extracted  
179 from the sediment using Ludox HS-40 (Sigma Aldrich, France) (de Jonge and Bouwman 1977) and  
180 retrieved with a 40- $\mu\text{m}$  mesh sieve (Higgins and Thiel 1988). From each replicate, 100 nematodes were  
181 haphazardly taken with fine tweezers (#55, Dumont, Switzerland), mounted on slides in anhydrous  
182 glycerol to prevent dehydration (Seinhorst 1959) and examined microscopically. The taxonomic  
183 composition of the nematode community was determined at least to the genera level based on these 100  
184 individuals according to the identification keys of Platt and Warwick (1983, 1988), and Warwick and  
185 Clarke (1998). Density was calculated for each genera as number of individuals per  $\text{m}^2$  (ind.  $\text{m}^{-2}$ ) and  
186 relative abundance as the percent composition of each genera relative to the total number of organisms.  
187 Species diversity indices were calculated on 100 individuals for each replicate at the five habitats for the  
188 two seasons. Community structure of the other meiofauna groups was not determined due to low  
189 densities. Nematodes were allocated into four trophic groups according to Wieser (1953) as follows:  
190 selective deposit-feeders (1A), non-selective deposit-feeders (1B), epigrowth feeders (2A), and  
191 omnivores/predators (2B). Various nematode genera, including at least one of each from the 4 different



192 feeding groups, were selected for determination of individual dry weight and carbon content. Between 10  
193 and 200 nematodes from the same genera were picked for dry weight determination, freeze-dried and  
194 weighed ( $\pm 1 \mu\text{g}$ ) using a microbalance (ME 5, Sartorius, Germany). Organic carbon composition was  
195 determined using an elemental analyzer (Flash EA 1112, Thermo Scientific, Italy). Mean biomass of  
196 species-specific nematodes and benthic copepods were used to convert the density of nematodes into  
197 grams of carbon per  $\text{m}^2$  and dry weight per  $\text{m}^2$  (Supplementary material, Appendix 1).

## 198 **Data treatment and statistical methods**


199 All analyses were conducted using the R software (R Core Team, 2016) and all hypotheses were  
200 tested at  $p < 0.05$  level. Normality of the data was tested using Shapiro-Wilks tests. Homogeneity of  
201 variance was checked using Bartlett and Levene tests. Average hierarchical clusters (data dissimilarity,  
202 Bray-Curtis distance) were realized to discriminate groups within the food sources and within the  
203 meiofauna trophic structure using the *vegan* package (Oksanen et al. 2017) and *pvclust* package (Suzuki  
204 and Shimodaira 2015). ANOVA followed by Tukey *post-hoc* test were used to compare sediment  
205 characteristics, meiofauna densities, biomass and diversity indices: Hill's diversity  $N_1$  (Hill 1973) and  
206 species evenness  $J'$  (Formula in supplementary material, Appendix 2). Comparisons between sediment  
207 layers were conducted on meiofauna densities ( $\text{ind. m}^{-3}$ ) and biomasses ( $\text{gC m}^{-3}$ ) expressed in volume of  
208 sediment. Comparisons between seasons and habitats were done on meiofauna densities, biomasses and  
209 diversity indices from combined layers (expressed per surface area). Environmental variables have been  
210 analyzed through a preliminary Principal Components Analysis (PCA) to visualize the differences  
211 between habitats based on these variables. In a subsequent step copepods, other meiofauna taxa  
212 (ostracods, foraminifers and polychaetes) and feeding types of nematodes were compared with Bray-  
213 Curtis similarity redundancy relationship analyses (RDA) made with the *vegan* package (Oksanen et al.  
214 2017). The relative abundance of copepods, other meiofauna (ostracods, foraminifers and polychaetes)  
215 and different feeding types of nematodes (dependent variables) were linked to environmental variables  
216 (independent variables) that were selected after stepwise forward procedure. Multicollinearity was tested  
217 for the explanatory variables using the *vif.cca* function in the *vegan* package (values over 10 indicated  
218 collinearity; Oksanen et al. 2017). The significance of the model was tested using a two-way ANOVA.

## 219 **Results**

### 220 **Sediment grain size**

221 Sediment of the mudflats and seagrass beds consisted of a mixture of sand (29 to 64%), silt (29 to  
222 58%) and clay (7 to 15%), while the sandflat consisted mainly of sand (98%) (Fig. 2). Sediments were  
223 classified as silt for mudflat MO, mudflat SR (autumn) and seagrass MO (autumn), very fine sand for

224 seagrass SR, mudflat SR (spring) and seagrass MO (spring) and medium sand for sandflat SR. The mud  
 225 content within the different layers of the same habitat was relatively similar, with a variation lower than  
 226 2%. The sediment grain-size of mudflat MO was lower than all other habitats with a median sediment  
 227 grain-size of  $32 \pm 4.3 \mu\text{m}$ , and the grain-size of sandflat SR was higher than all other habitats with a  
 228 median sediment grain-size of  $407 \pm 16.3 \mu\text{m}$  (Supplementary material, Appendix 3).



229  
 230 **Figure 2.** Ternary diagram presenting the sediment composition of the mudflats, seagrass beds and sandflat in the  
 231 Marennes-Oléron Bay (MO) and the Sylt-Rømø Bight (SR) in spring (grey symbols) and autumn (black symbols).  
 232 Separation by size for clay ( $< 3.9 \mu\text{M}$ ), silty ( $3.9 - 63 \mu\text{M}$ ), and sandy ( $63 - 2000 \mu\text{M}$ ) sediment. ( $n = 3$ )

233 **Potential food sources**

234 The total biomass of benthic food sources ranged from 67.4 (sandflat SR, autumn) to  $447.4 \text{ gC m}^{-2}$   
 235 (mudflat SR, spring) (Fig 3; C and E). SOM was dominant in all habitats sources with proportions  
 236 ranging from 29.3 to 94.5%, representing 19.8 (sandflat SR, autumn) to  $290.3 \text{ gC m}^{-2}$  (mudflat MO,  
 237 autumn). Benthic microalgae were the second most dominant food source, with biomass ranging from  $7.1$   
 238  $\text{gC m}^{-2}$  (3.7%, mudflat MO, spring) to  $70.3 \text{ gC m}^{-2}$  (22.6%, seagrass SR, spring). Detrital matter was  
 239 found mainly in the two seagrass beds and mudflat SR with 4.3 (1.0%, mudflat SR, spring) to  $45.5 \text{ gC m}^{-2}$   
 240 ( $16.7\%$ , seagrass bed MO, autumn). In the seagrass beds, leaves and roots were occurring with biomass  
 241 ranging from 1.6 (0.5%, seagrass SR, autumn) to  $15.0 \text{ gC m}^{-2}$  (5.1%, seagrass MO, spring) and 0.3 (0.1%,  
 242 seagrass SR, autumn) to  $24.4 \text{ gC m}^{-2}$  (8.2%, seagrass MO, autumn). Bacteria represented the lowest  
 243 biomass in all habitats with 0.6 (0.9%, sandflat SR, autumn) to  $20.1 \text{ gC m}^{-2}$  (4.5%, mudflat SR, spring).  
 244 Some differences of food source biomass could be observed between the surface and the sub-surface

245 layers, always with higher biomass in the surface layer when they occurred: detrital matter in mudflat SR  
246 in spring and in seagrass MO in autumn, bacteria in mudflat SR in spring, and benthic microalgae in all  
247 habitats in all seasons except for mudflat SR and sandflat SR in autumn (two-way ANOVA with Tukey  
248 *post-hoc* test, highest p-value 0.032).

249 The average hierarchical cluster of dissimilarities highlighted that food source composition in the  
250 sandflat SR was different than in the other habitats (Fig. 3; A), due to the much lower biomass of SOM in  
251 sandflat SR (29.2 gC m<sup>-2</sup>) than in mudflats and seagrass beds (256.6 gC m<sup>-2</sup>; two-way ANOVA with  
252 Tukey *post-hoc* test,  $p < 0.001$ , Fig. 3C and E). Benthic microalgae biomass was much higher in sandflat  
253 SR (54.2 gC m<sup>-2</sup>) than in mudflat SR (27.8 gC m<sup>-2</sup>), mudflat MO (10.5 gC m<sup>-2</sup>) and seagrass MO (22.6 gC  
254 m<sup>-2</sup>, two-way ANOVA with Tukey *post-hoc* test,  $p < 0.001$ ). Food source composition in both mudflats  
255 and both seagrass beds of the two bays appeared to be similar as highlighted by the clustering but some  
256 variations could be pointed out based on two-way ANOVA. Biomass of bacteria in spring was higher in  
257 mudflat SR (13.5 gC m<sup>-2</sup>) compared to mudflat MO (3.9 gC m<sup>-2</sup>), seagrass MO (4.2 gC m<sup>-2</sup>), seagrass SR  
258 (6.9 gC m<sup>-2</sup>) and sandflat SR (1.8 gC m<sup>-2</sup>, two-way ANOVA with Tukey *post-hoc* test,  $p < 0.001$ ). The  
259 presence of detrital matter in mudflat SR versus its absence in mudflat MO discriminated both mudflats as  
260 well. Lastly, biomass of leaves and roots in spring were higher in the Marennes-Oléron Bay (39.4 gC m<sup>-2</sup>)  
261 compared to the Sylt-Rømø Bight (6.4 gC m<sup>-2</sup>, two-way ANOVA with Tukey *post-hoc* test,  $p < 0.001$ ).

262 Some differences occurred at the seasonal scale as well, with higher biomass of primary producers in  
263 spring when a difference occurred. Such a pattern was observed for benthic microalgae in sandflat SR,  
264 seagrass SR and seagrass MO, and for seagrass leaves in seagrass bed MO (two-way ANOVA with  
265 Tukey *post-hoc* test,  $p < 0.001$ ). Bacteria biomass of mudflat SR and seagrass SR were also higher in  
266 spring compared to autumn (two-way ANOVA followed by Tukey *post-hoc* test,  $p < 0.001$ ). About  
267 detrital matter, seasonal changes followed an opposite pattern when they occurred, with an increase in  
268 biomass of detritus from spring to autumn in mudflat SR and seagrass MO (two-way ANOVA with  
269 Tukey *post-hoc* test,  $p < 0.001$ ). SOM biomass was stable between spring and autumn (two-way ANOVA  
270 with Tukey *post-hoc* test, lowest p-value 0.44).

271 Regarding quality of potential food sources in the different habitats, they varied from being partly  
272 detrital in sandflat SR (57 to 63%) to being almost completely made out of detrital matter in mudflat MO  
273 (from 93 to 99%). Carbon: chlorophyll *a* (C:chl *a*) ratios of SOM were higher in mudflat MO (1201.4),  
274 mudflat SR (681.8) and seagrass MO (504.6) compared to seagrass SR (221.4) and sandflat SR (72.3;  
275 two-way ANOVA with Tukey *post-hoc* test,  $p < 0.001$ , Table 1). SOM had much higher C:chl *a* ratios in  
276 the sub-surface layer than in the surface layer (two-way ANOVA with Tukey *post-hoc* test,  $p < 0.001$ ).  
277 Variation in C:chl *a* ratios between spring and autumn were not observed (two-way ANOVA, lowest p-


278 value 0.33, Table 1). Proportions of SOM that were originating from fresh microalgae (i.e.  
 279 microphytobenthos and trapped phytoplankton) ranged from 3.6 (mudflat MO, spring) to 71.1% (sandflat  
 280 SR, autumn, Table 1). These proportions were much higher in sandflat SR (66.6%) and seagrass SR  
 281 (22.0%) than in mudflat SR (7.0%) and mudflat MO (4.0%, two-way ANOVA with Tukey *post-hoc* test,  
 282 Table 1). Quantities of SPOM ranged between 0.32 and 0.54 gC L<sup>-1</sup>. Minor variations between habitats  
 283 observed within SPOM are reported in the supplementary material (Appendix 4).

284 **Table 1.** Carbon: chlorophyll *a* ratios (mean ± standard deviation, *n* = 3) and percentage of carbon  
 285 originating from fresh microalgae (mean ± standard deviation, %, *n* = 3) in sediment organic matter  
 286 (SOM) in the mudflats, seagrass beds and sandflat in the Marennes-Oléron Bay (MO) and the Sylt-Rømø  
 287 Bight (SR) in spring and autumn. Letters displayed after mean values indicated grouping of similar  
 288 samples (two-way ANOVA with Tukey *post-hoc* tests, *p* < 0.05).

289

Habitat	Carbon: chlorophyll <i>a</i> ratio		Carbon of SOM originating from fresh microalgae	
	Spring	Autumn	%	
	Spring	Autumn	Spring	Autumn
<b>Mudflat MO</b>	1314.3 <sup>E</sup> ± 35.3	1088.5 <sup>D</sup> ± 166.2	3.6 <sup>A</sup> ± 0.1	4.4 <sup>A</sup> ± 0.7
<b>Mudflat SR</b>	643.9 <sup>C</sup> ± 60.1	719.7 <sup>C</sup> ± 60.6	7.4 <sup>A</sup> ± 0.7	6.6 <sup>A</sup> ± 0.6
<b>Seagrass MO</b>	389.2 <sup>B</sup> ± 47.5	619.9 <sup>C</sup> ± 47.6	12.4 <sup>AB</sup> ± 1.4	7.7 <sup>A</sup> ± 0.6
<b>Seagrass SR</b>	193.9 <sup>A</sup> ± 14.3	249.0 <sup>AB</sup> ± 29.8	24.6 <sup>C</sup> ± 1.8	19.3 <sup>BC</sup> ± 2.4
<b>Sandflat SR</b>	76.8 <sup>A</sup> ± 2.2	67.8 <sup>A</sup> ± 9.1	62.0 <sup>D</sup> ± 1.7	71.1 <sup>D</sup> ± 9.8

290



291

292 **Figure 3.** Composition of potential benthic food sources for meiofauna and of meiofauna in the mudflats, seagrass beds and sandflat in the Marennes-Oléron Bay (MO) and the  
 293 Sylt-Rømø Bight (SR) in spring and autumn. **A)** Hierarchical average cluster of dissimilarity for the proportional composition of potential food sources. **B)** Hierarchical average  
 294 cluster of dissimilarity for proportional composition of meiofauna (nematodes and benthic copepods). Approximately unbiased (AU) probability values (p-values) by multiscale  
 295 bootstrap resampling are under each node and clusters (AU  $\geq$  95%) are highlighted with grey rectangles. **C)** Proportional composition of potential food sources. **D)**

296 Proportional composition of benthic copepods and trophic groups of nematodes. Trophic groups of genera with  
297 relative abundance > 5% are displayed. **E)** Biomass of potential food sources. **F)** Biomass of benthic copepods and  
298 trophic groups of nematodes. Selective deposit feeders = 1A, non-selective deposit feeders = 1B, epigrowth feeders  
299 = 2A and omnivores/predators = 2B (Wieser 1953). Abbreviations: Chr = *Chromadora*, Cta = *Chromadorita*, Com =  
300 *Comesa*, Dap = *Daptonema*, Ele = *Eleutherolaimus*, Eno = *Enoplolaimus*, Met = *Metachromadora*, Mic =  
301 *Microlaimus*, Mon = *Monoposthia*, Odo = *Odontophora*, Pra = *Praeacanthoichus*, Pty = *Ptycholaimellus*, Sab =  
302 *Sabatieria*, Spha = *Sphaerolaimus*, Spi = *Spirinia*, Ter = *Terschellingia*, The = *Theristus*, Vis = *Viscosia*. ( $n = 3$ )

### 303 **Meiofauna community structure**

304 The meiofauna community was dominated by nematodes with relative abundance ranging from  
305 41.7% (sandflat SR, surface layer, spring) to 100% (mudflat MO, sub-surface layer, spring), except for  
306 the surface layer of sandflat SR where copepods dominated (55.5%, Fig. 4). Four other meiofauna taxa  
307 occurred in the different habitats: copepods (0 to 55.5%), foraminifera (0 to 7.4%), polychaetes (0 to  
308 5.3%), and ostracods (0 to 0.4%). Total meiofauna density was higher in the surface layer than in the sub-  
309 surface layer in all habitats (two-way ANOVA with Tukey *post-hoc* test, highest  $p$ -value 0.033). Mean  
310 meiofaunal density of both layers combined ranged from  $1.7 \cdot 10^5$  ind.  $m^{-2}$  (mudflat SR, autumn) to  
311  $3.4 \cdot 10^6$  ind.  $m^{-2}$  (mudflat MO, autumn, Appendix 5). Meiofauna densities (both layers combined) were  
312 similar between spring and autumn, except in seagrass SR where densities were higher in spring (two-way  
313 ANOVA with Tukey *post-hoc* test, highest  $p$ -value 0.020, Table 2).

314 Total biomass of nematodes (both layers combined) ranged from  $0.053 \text{ gC } m^{-2}$  ( $1.6 \cdot 10^5$  ind.  $m^{-2}$ ,  
315 mudflat SR, autumn) to  $1.541 \text{ gC } m^{-2}$  ( $2.9 \cdot 10^6$  ind.  $m^{-2}$ , mudflat MO, spring, Fig. 3F). Nematode density  
316 and biomass were higher in surface layers than in the sub-surface layers (two-way ANOVA with Tukey  
317 *post-hoc* test, highest  $p$ -value 0.042). Nematodes from the surface layers dominated the contribution to  
318 the total biomass in all habitats, with lowest proportions in seagrass SR (54.0%) and mudflat SR (55.4%)  
319 and higher proportions in sandflat SR (82.3%), mudflat MO (83.3%) and seagrass SR (95.7%). At the  
320 habitat scale, nematode density and biomass were much higher in mudflat MO (density:  $3.0 \cdot 10^6$  ind.  $m^{-2}$ ;  
321 biomass:  $1.257 \text{ gC } m^{-2}$ ) than in all other habitats (two-way ANOVA with Tukey *post-hoc* test,  $p < 0.001$ ,  
322 Table 2, Appendix 5). Nematode biomass in the seagrass beds ranged from 0.146 (seagrass MO, autumn)  
323 to  $0.459 \text{ gC } m^{-2}$  (seagrass SR, spring). Nematode biomass of the mudflat SR and sandflat SR ranged from  
324  $0.054$  (mudflat SR, autumn) to  $0.150 \text{ gC } m^{-2}$  (mudflat SR, spring). Some temporal variations occurred as  
325 well, with higher nematode densities in spring in seagrass SR and higher nematode biomass in spring in  
326 mudflat MO (two-way ANOVA with Tukey *post-hoc* test, highest  $p$ -value 0.005, Table 2).


327 Total biomass of benthic copepods, which were merely present in the surface layers, ranged from  
328  $0.003 \text{ gC } m^{-2}$  ( $1.0 \cdot 10^4$  ind.  $m^{-2}$ , mudflat SR, autumn) to  $0.110 \text{ gC } m^{-2}$  ( $1.7 \cdot 10^5$  ind.  $m^{-2}$ , mudflat MO,

329 spring, Fig. 3F). Copepod biomass of the mudflat MO (0.095 gC m<sup>-2</sup>) was much higher than in the  
 330 mudflat SR (0.006 gC m<sup>-2</sup>), while copepod density was higher in the seagrass MO than in the seagrass SR  
 331 and mudflat SR (two-way ANOVA with Tukey *post-hoc* test,  $p < 0.001$ ). Densities and biomass of  
 332 benthic copepods were similar in spring and autumn (two-way ANOVA with Tukey *post-hoc* test, lowest  
 333  $p$ -value 0.178, Table 2).

334 **Table 2.** Results of the two-way ANOVA for differences in total meiofauna, nematode, and benthic copepod density  
 335 and nematode and benthic copepod biomass across habitats and seasons.

Source	Total meiofauna			Nematodes		Benthic copepods	
	df	F-ratio	<i>P</i> > F	F-ratio	<i>P</i> > F	F-ratio	<i>P</i> > F
<b>Density</b>							
Habitat	4	363.0	<b>&lt; 0.001</b>	385.8	<b>&lt; 0.001</b>	12.7	<b>&lt; 0.001</b>
Season	1	6.3	<b>0.020</b>	9.6	<b>0.005</b>	1.9	0.178
Habitat:season	4	8.8	<b>&lt; 0.001</b>	9.1	<b>&lt; 0.001</b>	2.8	0.052
Error	20						
<b>Biomass</b>							
Habitat	4			216.2	<b>&lt; 0.001</b>	29.01	<b>&lt; 0.001</b>
Season	1			34.5	<b>&lt; 0.001</b>	0.47	0.503
Habitat:season	4			11.7	<b>&lt; 0.001</b>	2.14	0.113
Error	40						

336 df = degree of freedom; F = F statistic; *P* = probability. Values in bold indicate a significant difference ( $\alpha \leq 0.05$ )



337

338 **Figure 4.** Proportional composition of meiofauna in the mudflats, seagrass beds and sandflat in the Marennes-  
 339 Oléron Bay (MO) and the Sylt-Rømø Bight (SR) in spring and autumn and in each sediment layer (0-1cm/1-5cm).  
 340 Densities (%) are indicated when > 7.5%. ( $n = 3$ )

341 **Nematode community structure**

342 Five thousand four hundred nematodes were determined at genus and/or species level. A total of 139  
 343 species, belonging to 65 genera and 20 families were found at the five habitats in both the Marennes-  
 344 Oléron Bay and the Sylt-Rømø Bight. Eight genera represented 70% of the community in the surface  
 345 layer of all habitats combined: *Daptonema* (1B, 18%), *Chromadorita* (2A, 10%), *Metachromadora* (2A,  
 346 10%), *Sabatieria* (1B, 10%), *Ptycholaimellus* (2A, 7%), *Spirinia* (2A, 6%), *Terschellingia* (1A, 5%), and  
 347 *Praeacanthochus* (2A, 4%). Four genera represented 70% of the community in the sub-surface layer of  
 348 all habitats combined: *Terschellingia* (1A, 28%), *Sabatieria* (1B, 25%), *Spirinia* (2A, 13%), and  
 349 *Metachromadora* (2A, 6%). The surface layer had a higher number of genera (from 3 to 9 genera, mean:  
 350 5.8) that represented 75% of the community compared to the sub-surface layer (from 2 to 9 genera, mean:  
 351 3.4) for all habitats. The ten most dominant genera per habitat, season and sediment layer are presented in  
 352 the supplementary material (Appendix 6). Species richness varied from 8 to 28, Hill's diversity from 2.3  
 353 to 19.1, and species evenness from 0.41 to 0.88 (Table 3). Species richness, diversity and evenness  
 354 decreased with depth at all habitats (two-way ANOVA,  $p < 0.001$ , Table 4). Species richness and  
 355 evenness were higher at the seagrass MO than at the mudflat SR and the seagrass SR (two-way ANOVA  
 356 with Tukey *post-hoc* test,  $p < 0.001$ ). Hills diversity was higher at the seagrass MO than at all other  
 357 habitats (two-way ANOVA with Tukey *post-hoc* test,  $p < 0.001$ ). There was no difference in diversity and  
 358 evenness between seasons (two-way ANOVA, lowest p-value 0.126, Table 4).

359 **Table 3.** Number of nematode genera, Hill's diversity ( $N_I$ ), and species evenness ( $J'$ ) in the mudflats,  
 360 seagrass beds and sandflat in the Marennes-Oléron Bay (MO) and the Sylt-Rømø Bight (SR) in spring  
 361 and autumn. ( $n = 3$ )

	Mudflat MO		Mudflat SR		Seagrass SR		Seagrass MO		Sandflat SR											
	Spring	Autumn	Spring	Autumn	Spring	Autumn	Spring	Autumn	Spring	Autumn										
Layer (in cm)	0-1	1-5	0-1	1-5	0-1	1-5	0-1	1-5	0-1	1-5										
Species richness	22	11	21	13	23	12	15	13	16	11	16	12	24	19	23	14	20	12	21	18
Hill's diversity ( $N_I$ )	14	5	12	6	13	4	9	5	7	3	6	4	15	12	14	8	11	6	12	7
Species evenness ( $J'$ )	0.9	0.6	0.8	0.7	0.8	0.6	0.8	0.6	0.7	0.5	0.6	0.6	0.8	0.9	0.8	0.8	0.8	0.7	0.8	0.7

362



363 **Table 4.** Results of the two-way ANOVA for differences in species richness, Hill's diversity ( $N_i$ ), and  
 364 species evenness ( $J'$ ) between habitats, sediment layers and season in the mudflats, seagrass beds and  
 365 sandflat in the Marennes-Oléron Bay and the Sylt-Rømø Bight.

Source	Species richness			Hill's diversity		Species evenness	
	df	F-ratio	$P > F$	F-ratio	$P > F$	F-ratio	$P > F$
Layer	1	88.3	< <b>0.001</b>	149.6	< <b>0.001</b>	55.9	< <b>0.001</b>
Site	4	11.0	< <b>0.001</b>	29.0	< <b>0.001</b>	16.2	< <b>0.001</b>
Season	1	0.1	0.802	2.4	0.126	0.0	1.000
Layer:site	4	1.5	0.214	4.75	<b>0.003</b>	4.5	<b>0.005</b>
Layer:season	1	6.4	<b>0.015</b>	3.3	0.078	0.6	0.452
Site:season	4	3.1	<b>0.026</b>	1.9	0.134	0.3	0.878
Layer:site:season	4	2.3	0.079	2.5	0.062	2.3	0.077
Error	38						

366 df = degree of freedom; F = F statistic;  $P$  = probability. Values in bold indicate a significant difference ( $\alpha \leq 0.05$ )


367 Community compositions, based on proportions of benthic copepods and of trophic groups of  
 368 nematodes, were similar between the two mudflats (besides the very large difference of biomass), as  
 369 highlighted by the average hierarchical cluster analysis ( $p < 0.05$ , Fig. 3B). The sandflat SR and the two  
 370 seagrass beds had similar communities as well. The difference of trophic structures between the mudflats  
 371 and the other habitats was related to the higher proportions of selective deposit-feeders (1A) and of non-  
 372 selective deposit-feeders (1B) in the mudflats. The mudflat MO was dominated by genera belonging to  
 373 non-selective deposit feeders (36%, 0.579 gC m<sup>-2</sup>), epigrowth feeders (34%, 0.525 gC m<sup>-2</sup>), and selective  
 374 deposit feeders (25%, 0.071 gC m<sup>-2</sup>) with *Terschellingia* (1A, 22%, 0.055 gC m<sup>-2</sup>), *Comesa* (1B, 14%,  
 375 0.082 gC m<sup>-2</sup>) and *Ptycholaimellus* (2A, 13%, 0.048 gC m<sup>-2</sup>) being the most common genera (Fig. 3D). At  
 376 the mudflat SR, the non-selective deposit feeders dominated (55%), and the most abundant genera were  
 377 *Sabatieria* (1B, 43%, 0.042 gC m<sup>-2</sup>) and *Terschellingia* (1A, 13%, 0.004 gC m<sup>-2</sup>). The seagrass bed SR  
 378 was dominated by epigrowth feeders with the genera *Spirinia* (2A, 46%, 0.225 gC m<sup>-2</sup>) and *Chromadorita*  
 379 (2A, 16%, 0.026 gC m<sup>-2</sup>). The seagrass bed MO was dominated by various genera belonging to epigrowth  
 380 feeders (31%, 0.165 gC m<sup>-2</sup>) and non-selective deposit feeders (29%, 0.061 gC m<sup>-2</sup>). There, the biomass  
 381 of nematodes was dominated by the epigrowth feeder genera *Metachromadora* (7%, 0.035 gC m<sup>-2</sup>). The  
 382 sandflat SR was dominated by genera belonging to epigrowth feeders (35%, 0.023 gC m<sup>-2</sup>) and non-  
 383 selective deposit feeders (19%, 0.040 gC m<sup>-2</sup>), with the genera *Microlaimus* (2A, 15%, 0.006 gC m<sup>-2</sup>) and  
 384 *Daptonema* (1B, 13%, 0.028 gC m<sup>-2</sup>) being the most common.

### 385 Relationships between environmental parameters and meiofauna trophic structure

386 The PCA on the environmental variables explained 63.2% of the total variation (Axis 1: 39.9%, axis  
387 2: 23.3%, Fig. 5A). Axis 1 highlights a positive relationship between SPOM, benthic microalgae and  
388 sediment grain size. At the opposite, when biomass of benthic microalgae, SPOM and sediment grain size  
389 were low, quantities of SOM were higher, as well as bacteria and detritus biomass, and  
390 carbon:chlorophyll *a* ratios of SOM. Axis 2 demonstrates that biomass of bacteria increased with biomass  
391 of SOM, SPOM, and benthic microalgae. Sandflat SR is defined by larger sediment grain size, higher  
392 biomass of benthic microalgae and of SPOM, as highlighted on axis 1. Axis 1 also discriminates seagrass  
393 SR from mudflat SR, mudflat MO and seagrass MO (autumn), the later ones being characterized with  
394 lower biomass of benthic microalgae and SPOM and/or higher carbon:chlorophyll *a* ratios of SOM, and  
395 higher biomass of SOM and detritus. In spring axis 2 discriminates well the mudflat and seagrass bed in  
396 the Marennes-Oléron Bay compared to the Sylt-Rømø Bight, which could be due to higher biomass of  
397 bacteria (mudflat SR) and/or benthic microalgae (seagrass SR). In terms of temporal variations, axis 1  
398 highlights low variation in environmental variables between spring and autumn except for seagrass MO,  
399 which could be due to a significant increase in biomass of detritus from spring to autumn in this habitat.  
400 Axis 2 highlights a stronger temporal variation (Fig. 5A), especially in seagrass SR, seagrass MO and  
401 mudflat SR. For seagrass SR this could be due to a decrease in bacteria, benthic microalgae, SPOM and  
402 SOM biomass from spring to autumn, as well as an increase in detrital matter, while changes in seagrass  
403 MO may be related to opposite patterns. For mudflat SR variations could be explained by the decrease in  
404 biomass of benthic microalgae and bacteria and/or the increase in detritus from spring to autumn.

405 The redundancy analyses (RDA) represented 51.8% of the total variation and 81.1% of the  
406 constrained variation (Fig. 5B). Fresh seagrass material was excluded from the RDA due to collinearity  
407 after stepwise forward selection of constraints ( $vif.cca > 10$ ). Axis 1 explains 33.0% of the total variation  
408 and 51.8% of the constrained variation. Axis 2 explains 18.8% of the total variation and 29.4% of the  
409 constrained variation. Both axes significantly explain the model, which highlights relationships between  
410 differences of communities and environmental parameters, particularly biomass of benthic microalgae,  
411 SPOM, detritus, SOM, carbon:chlorophyll *a* ratios and sediment grain size. Patterns highlighted by axis 1  
412 about environmental parameters were similar as those highlighted in the PCA: there was a positive  
413 relationship between benthic microalgae and SPOM biomasses, and sediment grain size. At the opposite,  
414 when benthic microalgae, SPOM and detritus biomasses were low, the carbon:chlorophyll *a* ratio was  
415 higher, as well as biomasses of SOM, bacteria and detritus. Differences of meiofauna trophic structure  
416 within habitats are mainly explained by sediment grain-size, carbon:chlorophyll *a* ratio of SOM, and  
417 biomasses of SOM, SPOM, detritus, benthic microalgae, and bacteria (Fig. 5). Selective deposit feeders  
418 and non-selective deposit feeders occurred mainly within habitats where biomasses of SOM, detritus and  
419 bacteria were high (i.e. sub-surface layers of mudflat SR in both seasons, sub-surface layers of seagrass


420 MO in both seasons, and sub-surface layers of mudflat MO in autumn). Epigrowth feeders dominated in  
 421 habitats where benthic microalgae were an important driving factor (i.e. surface layers of mudflat MO in  
 422 both seasons, sub-surface layers of mudflat MO in spring, surface and sub-surface layers of seagrass SR  
 423 in both seasons, and sub-surface layers of sandflat SR in both seasons). Omnivore/predatory nematodes,  
 424 benthic copepods and other meiofauna taxa occurred in habitats with high biomass of benthic microalgae  
 425 and SPOM, as well as large sediment grain-size (i.e. surface and sub-surface layers of sandflat SR in both  
 426 seasons and surface layers of seagrass MO in both seasons, and surface layers of mudflat SR in both  
 427 seasons).




428 **Figure 5. (A)** Principal component analyses for the environmental factors (arrows) plotted for the  
 429 mudflats, seagrass beds and sandflat in the Marennes-Oléron Bay (MO) and the Sylt-Rømø Bight (SR) in  
 430 spring (closed symbols) and autumn (open symbols). **(B)** Redundancy relationship analyses of  
 431 environmental factors (arrows) selected with stepwise forward procedure plotted against the different  
 432 groups of meiofauna (benthic copepods, trophic groups of nematodes and other meiofauna taxa) and the  
 433 surface (grey symbols) and sub-surface layers (black symbols) of the mudflats, seagrass beds and sandflat  
 434 in the Marennes-Oléron Bay (MO) and the Sylt-Rømø Bight (SR) in spring (closed symbols) and autumn  
 435 (open symbols). Abbreviations: OTHER = other meiofauna (i.e. polychaetes, foraminifers, and  
 436 ostracods); SPOM = suspended particulate organic matter; SOM = sediment organic matter; 1A =  
 437 nematodes, selective deposit feeders; 1B = nematodes, non-selective deposit feeders; 2A = nematodes,  
 438 epigrowth feeders; 2B = nematodes, omnivores/predators.

440 A regression using nematode feeding type data from all habitats highlighted that the ratios between  
 441 densities of epigrowth feeders and densities of non-selective deposit feeders increased when benthic  
 442 microalgae biomass increased ( $F[1, 8] = 10.4, p = 0.01, r^2 = 0.61$ , Fig. 6). The relative abundance of  
 443 selective deposit feeders was positively correlated to the ratio between biomass of fresh organic material

444 (i.e. benthic microalgae and seagrass leaves and roots) and biomass of detrital material (i.e. SOM and  
 445 detritus) ( $F[1, 8] = 23.1, p < 0.01, r^2 = 0.74$ , Fig. 7).



446  
 447 **Figure 6** Relationship between the biomass of benthic microalgae and the ratio between densities of epigrowth  
 448 feeders (2A) and densities of non-selective deposit feeders (1B) in the mudflats, seagrass beds and sandflat in the  
 449 Marennes-Oléron Bay (MO) and the Sylt-Rømø Bight (SR) in spring (grey symbols) and autumn (black symbols).  
 450 ( $n = 3$ )



451  
 452 **Figure 7.** Relationship between the ratio of fresh organic material to detrital material and the relative abundance of  
 453 selective deposit feeders (1A) in the mudflats, seagrass beds and sandflats in the Marennes-Oléron Bay (MO) and  
 454 the Sylt-Rømø Bight (SR) in spring (grey symbols) and autumn (black symbols). ( $n = 3$ )

455 **Discussion**

456 **Habitat description**

457 The different habitats compared in this study are characterized by a broad range of abiotic and biotic  
458 factors. This provides a good framework to determine how changes in these parameters, particularly  
459 potential food resources, may influence meiofauna community structure. Biomass of benthic microalgae  
460 was much higher in habitats with a larger sediment grain size (sandflat SR and seagrass SR) as already  
461 observed in the Marennes-Oléron Bay (Guarini et al. 1998), and benthic microalgae therefore represented  
462 a much higher proportion of the SOM in these habitats. This pattern is quite unusual, as habitats with large  
463 sediment grain size generally support less microphytobenthos biomass than habitats with small sediment  
464 grain size (Cammen 1991; de Jong and de Jonge 1995). Estimations of benthic microalgae biomass were  
465 based on chlorophyll *a* measurements and therefore included trapped fresh phytoplankton. Trapped  
466 phytoplankton represents a lower proportion of fresh microalgae material than benthic microphytobenthos,  
467 although some variations can occur depending on seasons and locations (Asmus 1983; Lebreton et al. 2012).  
468 Microphytobenthos community structure and its production may change as well depending on habitats:  
469 muddy sediments are usually dominated by large epipellic diatoms (Thornton et al. 2002; Azovsky et al.  
470 2004) whereas small epipsammic diatoms are the main taxa in sandy habitats (Asmus and Bauerfeind 1994;  
471 Azovsky et al. 2004). Benthic microalgae production can be very high in muddy sediments (i.e. 372 gC m<sup>-2</sup>  
472 yr<sup>-1</sup> in mudflat MO) (Pinckney and Zingmark 1993; Leguerrier et al. 2003) and, under favorable  
473 conditions, epipellic diatoms can double their standing stock on a daily basis (Underwood and Paterson  
474 1993). All these changes of biomass, community structure and production rates likely affect the use of this  
475 food resource by meiofauna.

476 Biomass of SOM was much lower in sandflat SR than in seagrass beds and mudflats. This is likely  
477 related to the high benthic microalgae production in bare mudflats (MacIntyre et al. 1996) and to the smaller  
478 sediment grain size in mudflats and seagrass beds (Bergamaschi et al. 1997). A decrease in sediment grain  
479 size generally leads to an increase in SOM (Bergamaschi et al. 1997) as fine particles enhance adsorption  
480 of organic compounds (Mayer 1994). Decrease of sediment grain size is also associated to lower  
481 hydrodynamics which increases sedimentation of organic matter particles (Frontalini et al. 2014) and of  
482 finer sediment. The presence of seagrass beds, reducing hydrodynamics and enhancing sedimentation as  
483 well (Leduc et al. 2009), also likely increased the load of SOM. As a result, even if SPOM was relatively  
484 similar between habitats, the presence of meadows may influence the biomass and the nature of the organic  
485 matter available to meiofauna between unvegetated and vegetated habitats (Machás and Santos 1999). The  
486 quality of the organic matter available to consumers was also correlated with sediment grain size, with  
487 SOM dominated by fresh organic matter in the sandflat and being of much lower quality in habitats made  
488 of fine sediments (seagrass bed and mudflats), where loads of detrital matter were higher.

489 Loads of detritus were high in seagrass beds, the largest ones being measured in seagrass MO, probably  
490 due to the much higher biomass of seagrass roots in this habitat compared to seagrass SR. The higher load  
491 of detrital matter in mudflat SR compared to mudflat MO is much likely related to the proximity between  
492 the mudflat and the seagrass bed in the Sylt-Rømø Bight, highlighting that seagrass material can be exported  
493 to bare systems. The presence of seagrass in intertidal systems may also indirectly affect the composition  
494 of the detrital organic matter available to meiofauna and meiofauna distribution in the sediment (Marbà et  
495 al. 2006). The complex root structure of seagrass beds indeed plays a role for the amount of fine particles  
496 and detritus associated with the shallow oxic zone in vegetated sediments (Wetzel et al. 2001), as well as  
497 in the release of oxygen and dissolved organic matter by the roots (Pedersen et al. 1998). Nevertheless,  
498 seagrass material is generally poorly used by most of the consumers as fresh or as detrital material  
499 (Danovaro 1996; Vizzini et al. 2002). The presence of seagrass and detritus as well as smaller sediment  
500 grain size also affected the abundance of bacteria within the studied habitats. Much lower biomass of  
501 bacteria was observed in habitats with lower quantities of detrital matter and SOM and in habitats with  
502 coarser sediment (i.e. sandflat), likely due to the reduced surface for bacteria colonization (Dale 1974;  
503 Mayer 1994).

#### 504 **Meiofauna community structure: spatial and temporal variations**

505 Meiofauna density and biomass varied highly within habitats and ecosystems, with lowest and  
506 highest values observed in mudflat SR (density:  $0.17 \cdot 10^6$  ind.  $m^{-2}$ , biomass:  $0.057$  gC  $m^{-2}$ ) and mudflat  
507 MO (density:  $3.36 \cdot 10^6$  ind.  $m^{-2}$ , biomass:  $1.598$  gC  $m^{-2}$ ), respectively. Meiofauna in all habitats were  
508 dominated by nematodes, followed by benthic copepods, as typically observed in intertidal habitats  
509 (Armonies and Hellwig-Armonies 1987; Rzeznik-Orignac et al. 2003; Leduc and Probert 2011). Data  
510 comparisons (Table 5) highlight that density and biomass of nematodes and benthic copepods measured  
511 in this study are in the range of those previously measured but, more generally, that these values can be  
512 highly variable between different soft-bottom habitats as well as within similar habitats (Heip et al. 1985;  
513 Castel et al. 1989; Kotwicki et al. 2005b).

514 In terms of community structure, proportions of benthic copepods increased from habitats with a  
515 lower complexity (mudflats: 6.2%) to habitats with a higher complexity caused by a higher seagrass  
516 biomass (seagrass beds: 15.5%) or by larger sediment grain size (sandflat: 38.4%). Larger proportions of  
517 copepods in the sandflat could be explained by a higher feeding efficiency in sandier sediment, probably  
518 due to a better accessibility to benthic microalgae and a better ability of these grazers to move in sand (De  
519 Troch et al. 2006). Higher proportions of benthic copepods in seagrass bed MO may be related to a more  
520 developed seagrass bed and therefore a more complex habitat, providing a larger diversity of ecological  
521 niches (Ndaro and Olafsson 1999; De Troch et al. 2001). The presence and development stage of seagrass

522 beds also affected nematode diversity as species richness was higher in the most dense seagrass bed  
 523 (Marennes-Oléron) between the two studied ones. Hill's diversity was also higher in seagrass MO than in  
 524 all other habitats, which could be partially linked to habitat complexity as well (Austen et al. 1998;  
 525 Steyaert et al. 2003; Giere 2009). Proportions of omnivore/predatory nematodes increased as well when  
 526 complexity increased due to larger sediment grain size as previously observed in sandflats of several other  
 527 coastal ecosystems (Warwick 1971; Steyaert et al. 2003). This could be explained by the increase of  
 528 foraging efficiency of predators with an increase in sediment grain size (Gallucci et al. 2005).

529 If some patterns could be observed between meiofauna community structure and habitat type,  
 530 direct relationships between biomass of food sources and density and/or biomass of nematodes and  
 531 benthic copepods could not be observed in this study. This is very likely related to the diversity of feeding  
 532 types observed in nematodes (Moens et al. 2005b; Vafeiadou et al. 2014) and, to a lower extent, in  
 533 benthic copepods (Cnudde et al. 2015), confirming the importance of approaches carried out at the scale  
 534 of the feeding type, as this was done for nematodes in this study.

535 Within these different habitats nematode density, biomass and diversity were similar between  
 536 spring and autumn, except for higher nematode density in seagrass SR and higher nematode biomass in  
 537 mudflat MO in spring. No temporal variation was observed for meiofauna community structure. The lack  
 538 of a clear temporal pattern between spring and autumn (Escaravage et al. 1989; Danovaro and Gambi  
 539 2002; Rzeznik-Orignac et al. 2003) could be partly explained by low temporal variation in parameters  
 540 strongly defining meiofauna community structure (i.e. food sources and sediment grain size; Moens et al.  
 541 2013).

542 **Table 5.** Density (ind. m<sup>-2</sup> \* 10<sup>6</sup>) and biomass (in gC m<sup>-2</sup> and/or g dry weight m<sup>-2</sup>) of nematodes and  
 543 benthic copepods in mudflats, seagrass beds (*Zostera* spp.) and sandflats from this study and from the  
 544 available literature about marine soft-bottom intertidal systems. Average conversion factors used to  
 545 estimate biomass in gC m<sup>-2</sup> from densities are: nematodes: 2.28\*10<sup>6</sup>, benthic copepods: 3.49\*10<sup>6</sup>, and  
 546 from biomass (in g dry-weight m<sup>-2</sup>) are: nematodes: 3.26, benthic copepods: 3.40. Converted values are in  
 547 italic.

Habitat type	Location	Density		Biomass (g dry weight m <sup>-2</sup> )		Biomass (gC m <sup>-2</sup> )		Reference
		Nematodes	Copepods	Nematodes	Copepods	Nematodes	Copepods	
Mudflat	Papanui Inlet, New Zealand	2.1–3.2	0.5–1.1	0.163	0.125	<i>0.05</i>	<i>0.04</i>	Leduc and Probert 2011
	Danish Wadden Sea, Denmark	0.22		0.03–0.4		<i>0.01–0.12</i>		Smidt 1951
	Sylt-Rømø Bight, Germany	0.28	0.03	0.233	0.015	0.102	0.006	This study
	Cornwall, U.K.	1.1				<i>0.48</i>		Ellison 1984
	Marennes-Oléron Bay, France	2.1				<i>0.92</i>		Rzeznik-Orignac et al. 2003
	Sylt-Rømø Bight, Germany	2.4	0.07			<i>1.05</i>	<i>0.02</i>	Armonies and Hellwig-Armonies 1987
	Lynher estuary, U.K.	0.8–22.9		1.4–3.4		<i>0.43–1.49</i>		Warwick and Price 1979

	Marenes-Oléron Bay, France					0.2–1.5	< 0.09	Pascal 2008
	Marenes-Oléron Bay, France	3.03	0.13	6.070	0.346	1.257	0.095	This study
	South Carolina, U.S.A.	2.3–4.4				1.01–1.93		Sikora and Sikora 1982
	European estuaries	0.13–14.5				0.06–6.35		Soetaert et al. 1995
	Papanui Inlet, New Zealand	4.9–5.2	< 0.04	0.480	< 0.009	0.15	< 0.003	Leduc and Probert 2011
	Marenes-Oléron Bay, France	0.65	0.21	1.303	0.459	0.294	0.078	This study
	Sylt-Rømø Bight, Germany	0.81	0.04	0.819	0.015	0.351	0.006	This study
<b>Seagrass bed</b>	Sylt-Rømø Bight, Germany	2.0				0.88		Armonies and Hellwig-Armonies 1987
	Arcachon Bay, France	5.5–8.0		2.4–5.3		0.74–2.32		Castel et al. 1989; Escaravage et al. 1989
	Mira estuary, Portugal	1.8–3.3				0.79–1.46		Materatski et al. 2018
	Marenes-Oléron Bay, France					2.75	0.36	Lebreton 2009
	North Sea intertidal sandflats	0.15	0.11			0.07	0.03	Kotwicki et al. 2005a
	Sylt-Rømø Bight, Germany	0.22	0.14	0.303	0.130	0.124	0.050	This study
	Arcachon Bay, France	0.7–1.2		0.8–1.3		0.25–0.57		Castel et al. 1989; Escaravage et al. 1989
<b>Sandflat</b>	Oosterschelde, Netherlands	5.0	0.13	2.0	0.2	0.61	0.06	Smol et al. 1994
	Sylt-Rømø Bight, Germany	2.3	3.5			1.01	1.00	Armonies and Hellwig-Armonies 1987
	European intertidal flats	0.24–4.1	< 0.35			0.11–1.80	< 0.10	Hicks and Coull 1983; Kotwicki et al. 2005b

548

549 **Benthic microalgae and sediment organic matter are important drivers of nematode community**  
550 **structure**

551 Changes in benthic microalgae biomass influenced the community structure of nematodes: relative  
552 abundance of epigrowth feeders, relying on benthic microalgae, were reduced when biomass of this food  
553 source decreased. Such a relationship between biomass of benthic microalgae and density of epigrowth  
554 feeders had already been observed in a subtidal seagrass meadow (Danovaro and Gambi 2002) and during  
555 grazing experiments (Rzeznik-Orignac and Fichet 2012). A relationship was also observed between  
556 biomass of benthic microalgae and relative abundances of benthic copepods and omnivores/predators.  
557 However, this relationship may be partly mediated by sediment grain size as benthic microalgae biomass  
558 (this study) and relative abundance of benthic copepods increase when sediment get coarser (Hicks and  
559 Coull 1983). Nevertheless, the much higher biomass of benthic microalgae in the sandflat compared to  
560 SOM suggests that consumers rely—at least partly—directly (i.e. copepods) or indirectly (i.e.  
561 omnivores/predators) on benthic microalgae in this habitat. Such a dependence to benthic microalgae as a  
562 food source has been demonstrated for benthic copepods (Leduc et al. 2009).

563 At a more global scale (i.e. whole meiofauna, all habitats combined), no general pattern could be  
564 observed between benthic microalgae biomass and those of nematodes and benthic copepods. Some other  
565 parameters should probably be taken into account to better consider relationships between this food  
566 resource and consumers. Benthic microalgae can indeed reach very high production rates and is a food  
567 source of high quality (Cebrián 1999). Intense grazing by meiofauna may affect benthic microalgae as well.



568 The high primary production of benthic microalgae in mudflat MO ( $372 \text{ gC m}^{-2} \text{ yr}^{-1}$ ; Leguerrier et al. 2003)  
569 and high grazing pressure on this food source by the large biomass of nematodes (Blanchard 1991) could  
570 explain the lack of relationship between benthic microalgae and meiofauna biomass within these habitats.  
571 Quality is of high importance as well: Moens et al. (1999b) suggested that meiofauna migrates actively  
572 towards 'optimal' food sources. In addition to epigrowth feeders, several studies demonstrated that deposit  
573 feeders rely on microalgae (Moens and Vincx 1997; Rzeznik-Orignac et al. 2003; Vafeiadou et al. 2014)  
574 likely due to the better quality of microalgae. A relationship between density of nematodes belonging to  
575 these feeding groups and microalgae has been observed (Moens et al. 1999a). In particular, grazing  
576 experiments using  $^{14}\text{C}$  labelled diatoms indicated the dependence of a non-selective deposit feeding  
577 nematode (*Daptonema*), an epigrowth feeding nematode (*Metachromadora*) and benthic copepods on  
578 microalgae (Rzeznik-Orignac and Fichet 2012). Buccal cavities of epigrowth feeders are appropriate for  
579 consumption of benthic microalgae, due to the presence of a dorsal tooth which is used to pierce diatoms  
580 (Wieser 1953; Moens and Vincx 1997). Vafeiadou et al. (2014), using stable isotopes, confirmed that non-  
581 selective deposit feeding nematodes, epigrowth feeding nematodes and benthic copepods can largely rely  
582 on benthic microalgae in sparsely covered seagrass beds. The non-selective deposit feeders can therefore  
583 be considered as opportunistic feeders, which may change feeding strategies depending on the availability  
584 of food sources (Moens and Vincx 1997; Rzeznik-Orignac et al. 2008). As a result, benthic microalgae are  
585 often considered as the principal food source for nematodes and benthic copepods (Moens et al. 2005a;  
586 Rzeznik-Orignac et al. 2008; Maria et al. 2012).

587 Benthic microalgae followed an opposite pattern than detrital material (SOM and detritus), with  
588 decreasing biomass of benthic microalgae when detrital material increased. SOM biomass was stable  
589 between spring and autumn, as well as between surface and sub-surface layers, but large variations were  
590 observed between the different habitats (i.e. much lower biomass in sandflat SR). Relative abundance of  
591 non-selective deposit feeders, closely related to detrital matter based on the RDA, decreased when biomass  
592 of SOM was lowered. Higher SOM quality in the surface sediment layers could lead to higher biomass,  
593 density and diversity of nematodes in the surface layer compared to the sub-surface layer. As a result, the  
594 important role of SOM is very likely related to its very high amount, as well as variations in quality between  
595 the studied systems. A relationship between SOM and density of non-selective deposit feeders was also  
596 recognized in a subtidal seagrass meadow (Danovaro and Gambi 2002), while a grazing experiment  
597 indicated that non-selective and selective deposit feeding nematode species assimilated predominantly  
598 SOM and bacteria (Tietjen and Lee 1977).

599 As a result, SOM and benthic microalgae appear to be important biotic factors driving the nematode  
600 community structure and, *a fortiori*, the composition of meiofauna assemblages as nematodes represent a

601 large proportion of the meiofauna in coastal ecosystems (Armonies and Hellwig-Armonies 1987; Rzeznik-  
602 Orignac et al. 2003; Leduc and Probert 2011).

### 603 **Role of bacteria as a food source?**

604 Selective deposit feeders had much higher relative abundances in habitats with larger loads of detritus  
605 and SOM compared to those dominated by fresh material. In addition, multivariate analyses indicated that  
606 biomass of bacteria were higher when loads of SOM increased. These results highlight a probable  
607 relationship between bacteria biomass and relative abundance of selective deposit feeders, endorsing that  
608 consumers from this trophic group rely on bacteria (Moens and Vincx 1997), which develop on detrital  
609 particles (Day et al. 1989; Danovaro 1996), or dissolved organic matter (Moens and Vincx 1997). The use  
610 of bacteria as food resources has been confirmed by stable isotope studies demonstrating that selective  
611 deposit feeders can rely on sulfide-oxidizing bacteria (Vafeiadou et al. 2014), such specific diets being  
612 related to their small buccal cavity (Wieser 1953; Moens and Vincx 1997). The lack of direct relationship  
613 between abundances of selective deposit feeders and biomass of bacteria, as in Semprucci et al. (2010),  
614 might be due to the influence of additional abiotic parameters (i.e. oxic conditions (Jensen 1983) and  
615 sediment grain size (Steyaert et al. 2003)), the reliance of bacteria on a large diversity of substrates  
616 (Boschker et al. 2000), and high bacterial production which is only partially grazed (< 28%, Pascal et al.  
617 2009).

618 Multivariate analyses demonstrated as well a relationship between relative abundances of non-  
619 selective deposit feeders and biomass of detritus and, to a lower extent, those of bacteria and SOM.  
620 Nevertheless, no direct relationship was observed between relative abundances of non-selective deposit  
621 feeders and bacteria, SOM or detritus biomass, which is likely related to the more opportunistic feeding  
622 behavior of these nematodes (Moens and Vincx 1997; Rzeznik-Orignac et al. 2008). The shape of the buccal  
623 cavity of non-selective deposit feeders allows them to rely on a much larger diversity of food sources:  
624 benthic microalgae, SOM, bacteria and seagrass detritus, as demonstrated by stable isotope studies  
625 (Rzeznik-Orignac et al. 2008; Maria et al. 2012; Vafeiadou et al. 2014) and observations of feeding behavior  
626 (Moens and Vincx 1997). Biomass of fresh seagrass matter, which varied highly between habitats in this  
627 study, could also not be linked to variations in non-selective deposit feeders. However, the lack of relation  
628 here is expected since this resource is poorly used by these nematodes, and, to a larger extent, to meiofauna  
629 in general, due to its low nutritional value and high lignocellulose content (Vizzini et al. 2002). This was  
630 confirmed by Materatski et al. (2015) who observed no changes in nematode density before and after a  
631 seagrass collapse.

### 632 **Effect of sediment depth on meiofauna communities: influence of oxic conditions**

633 Nematode and benthic copepod densities and biomasses decreased from surface to sub-surface layers  
634 in all habitats, which is a common pattern in seagrass beds and unvegetated habitats (Castel et al. 1989;  
635 Lebreton 2009; Leduc and Probert 2011). Benthic microalgae, which have the highest biomass in the  
636 surface layer, might influence the vertical distribution and migration of nematodes and benthic copepods  
637 (Moens et al. 2013), particularly nematode epigrowth feeders which specifically rely on benthic microalgae  
638 and which were very abundant in all habitats. Differences in abiotic parameters, such as oxygen depletion  
639 towards deeper layers, could affect as well the vertical distribution of meiofauna (Steyaert et al. 2003;  
640 Vanaverbeke et al. 2004), as anoxic conditions are much less favorable to nematodes and benthic copepods.  
641 Besides influencing biomass and densities, anoxic conditions affect diversity and community structure as  
642 well. Diversity of nematodes clearly decreased with depth in all the studied habitats, which was observed  
643 in other studies (Steyaert et al. 2003; Leduc and Probert 2011). *Terschellingia* spp. and *Sabatieria* spp. had  
644 a higher relative abundance in sub-surface layers than in surface layers, related to their higher tolerance  
645 regarding hypoxic conditions (Steyaert et al. 2007). *Terschellingia* spp. and *Sabatieria* spp. are also known  
646 to feed on sulfide-oxidizing bacteria that occur in these hypoxic zones (Semprucci and Balsamo 2012;  
647 Vafeiadou et al. 2014).

## 648 **Conclusion**

649 Benthic microalgae and SOM are two important drivers in the functioning of soft-bottom coastal food  
650 webs as changes in these food resources affect the two most dominant nematode trophic groups (i.e. non-  
651 selective deposit feeders, considered as more opportunistic, and epigrowth feeders, more specifically related  
652 to benthic microalgae). This is likely related to the high biomass of these food sources, as well as the quality  
653 and high production rates of benthic microalgae (MacIntyre et al. 1996; Cebrián 1999). Therefore, this  
654 study highlights the existence of some relationships between composition of food sources and the trophic  
655 structure of the meiofauna community, when studied at the scale of trophic groups. On the contrary, no  
656 clear relationship was observed between bacterial biomass and density of selective deposit feeders  
657 (bacterivores; 1A), despite a relationship between SOM biomass and relative abundances of selective  
658 deposit feeders. This lack of relationship may be related to the high production rates of bacteria and the  
659 influence of abiotic factors, especially oxic conditions. Approaches such as trophic markers (e.g. stable  
660 isotope and/or fatty acid analyses) would be very complementary to the determination of feeding types  
661 based on the morphology of the buccal cavity. Such tools would provide a more comprehensive picture of  
662 food sources really assimilated by meiofauna and highlight how the use of food sources by meiofauna may  
663 change depending on availability. Combining trophic marker data with biomass and production data in food  
664 web models would then provide a much more complete overview of the functioning of coastal food webs.

## 665 **Compliance with ethical standards**

666 **Conflict of interest**

667 All authors declare that they have no conflict of interest.

668 **Ethical statement**

669 All applicable international, national and/or institutional guidelines for the sampling, care and use of  
670 organisms were followed.

671 **Acknowledgements**

672 This study forms part of the Ph.D. thesis of L. van der Heijden. This work was financially  
673 supported by the University of La Rochelle through a grant provided by the ‘Ministère de l’Enseignement  
674 Supérieur et de la Recherche’. This work was supported by the CNRS through the CNRS research chair  
675 provided to B. Lebreton. Financial support was provided by the Alfred Wegener Institute Helmholtz  
676 Centre for Polar and Marine Research, as well as the University of La Rochelle (Mobility for doctoral  
677 students) and the German Academic Exchange Service (DAAD: short-term research grant, 2017, no.  
678 57314023). We acknowledge P. Pineau, B. Hussel, S. Horn, M. Burgdorf, M. Paar and N. Lachaussée for  
679 their support during fieldwork, and G. Guillou and C. Labrune for their support in the analyses. We thank  
680 Q. Bernier for his help in sample processing and analysis. Finally, L. van der Heijden thanks the members  
681 of his Ph.D. committee, M. Graeve, N. Niquil and G. Blanchard, for their guidance, as well as two  
682 anonymous reviewers whose comments greatly improved the manuscript.

683 **References**

- 684 Armonies W, Hellwig-Armonies M (1987) Synoptic patterns of meiofaunal and macrofaunal abundances and  
685 specific composition in littoral sediments. *Helgoländer Meeresuntersuchungen* 41:83–111. doi:  
686 10.1007/BF02365101
- 687 Asmus H, Asmus RM (1985) The importance of grazing food chain for energy flow and production in three  
688 intertidal sand bottom communities of the northern Wadden Sea. *Helgoländer Meeresuntersuchungen* 39:273–  
689 301. doi: 10.1007/BF01992775
- 690 Asmus H, Asmus RM (2005) Significance of suspension-feeder systems on different spatial scales. In: Dame RF,  
691 Olenin S (eds) *The comparative roles of suspension-feeders in ecosystems*. Springer Netherlands, pp 199–219
- 692 Asmus RM (1983) Benthische und pelagische Primärproduktion und Nährsalzbilanz. Eine Freilanduntersuchung im  
693 Watt der Nordsee. Ber. Inst. Meereskunde, Univ. Kiel
- 694 Asmus RM, Bauerfeind E (1994) The microphytobenthos of Königshafen—spatial and seasonal distribution on a  
695 sandy tidal flat. *Helgoländer Meeresuntersuchungen* 48:257–276. doi: 10.1007/BF02367040

- 696 Austen MC, Widdicombe S, Villano-Pitacco N (1998) Effects of biological disturbance on diversity and structure of  
697 meiobenthic nematode communities. *Mar Ecol Prog Ser* 174:233–246. doi: 10.3354/meps174233
- 698 Azovsky AI, Chertoproud ES, Saburova MA, Polikarpov IG (2004) Spatio-temporal variability of micro- and  
699 meiobenthic communities in a White Sea intertidal sandflat. *Estuar Coast Shelf Sci* 60:663–671. doi:  
700 10.1016/j.ecss.2004.03.005
- 701 Baird D, Asmus H, Asmus R (2007) Trophic dynamics of eight intertidal communities of the Sylt-Rømø Bight  
702 ecosystem, northern Wadden Sea. *Mar Ecol Prog Ser* 351:25–41. doi: 10.3354/meps07137
- 703 Bergamaschi BA, Tsamakidis E, Keil RG, Eglinton TI, Montluçon DB, Hedges JI (1997) The effect of grain size and  
704 surface area on organic matter, lignin and carbohydrate concentration, and molecular compositions in Peru  
705 Margin sediments. *Geochim Cosmochim Acta* 61:1247–1260. doi: 10.1016/S0016-7037(96)00394-8
- 706 Blanchard GF (1991) Measurement of meiofauna grazing rates on microphytobenthos: is primary production a  
707 limiting factor? *J Exp Mar Bio Ecol* 147:37–46. doi: 10.1016/0022-0981(91)90035-U
- 708 Boschker HTS, Wielemaker A, Schaub BEM, Holmer M (2000) Limited coupling of macrophyte production and  
709 bacterial carbon cycling in the sediments of *Zostera* spp. meadows. *Mar Ecol Prog Ser* 203:181–189. doi:  
710 10.3354/meps203181
- 711 Cammen LM (1991) Annual bacterial production in relation to benthic microalgal production and sediment oxygen  
712 uptake in an intertidal sandflat and an intertidal mudflat. *Mar Ecol Prog Ser* 71:13–25. doi:  
713 10.3354/meps071013
- 714 Castel J, Labourg P-J, Escavavage V, Auby I, Garcia ME (1989) Influence of seagrass beds and oyster parks on the  
715 abundance and biomass patterns of meio- and macrobenthos in tidal flats. *Estuar Coast Shelf Sci* 28:71–85.
- 716 Cebrián J (1999) Patterns in the fate of production in plant communities. *Am Nat* 154:449–468. doi:  
717 10.1086/303244
- 718 Chardy P, Dauvin JC (1992) Carbon flows in a subtidal fine sand community from the western English Channel: a  
719 simulation analysis. *Mar Ecol Prog Ser* 81:147–161. doi: 10.3354/meps081147
- 720 Cnudde C, Werbrouck E, Lepoint G, Vangansbeke D, Moens T, Troch M De (2015) Trophodynamics of intertidal  
721 harpacticoid copepods based on stable isotopes and fatty acid profiling. *Mar Ecol Prog Ser* 524:225–239. doi:  
722 10.3354/meps11161
- 723 Core Team R (2016) R: A language and environment for statistical computing.
- 724 Coull BC (1999) Role of meiofauna in estuarine soft-bottom habitats. *Aust J Ecol* 24:327–343. doi: 10.1046/j.1442-  
725 9993.1999.00979.x
- 726 Dale NG (1974) Bacteria in intertidal sediment: Factors related to their distribution. *Limnol Oceanogr* 19:509–519.

- 727 Danovaro R (1996) Detritus-Bacteria-Meiofauna interactions in a seagrass bed (*Posidonia oceanica*) of the NW  
728 Mediterranean. *Mar Biol* 127:1–13. doi: 10.1007/BF00993638
- 729 Danovaro R, Gambi C (2002) Biodiversity and trophic structure of nematode assemblages in seagrass systems:  
730 evidence for a coupling with changes in food availability. *Mar Biol* 141:667–677. doi: 10.1007/s00227-002-  
731 0857-y
- 732 Danovaro R, Gambi C, Mirto S (2002) Meiofaunal production and energy transfer efficiency in a seagrass *Posidonia*  
733 *oceanica* bed in the western Mediterranean. *Mar Ecol Prog Ser* 234:95–104. doi: 10.3354/meps234095
- 734 Day JWJ, Hall CAS, Kemp MW, Yáñez-Arancibia A (eds) (1989) *Estuarine ecology*. John Wiley & Sons, New  
735 York
- 736 de Jong DJ, de Jonge VN (1995) Dynamics and distribution of microphytobenthic chlorophyll-a in the Western  
737 Scheldt estuary (SW Netherlands). *Hydrobiologia* 311:21–30. doi: 10.1007/BF00008568
- 738 de Jonge VN (1980) Fluctuations in the organic carbon to chlorophyll-a ratios for estuarine benthic diatom  
739 populations. *Mar Ecol Prog Ser* 2:345–353. doi: 10.3354/meps002345
- 740 de Jonge VN, Bouwman LA (1977) A simple density separation technique for quantitative isolation of  
741 microphytobenthos using the colloidal silica Ludox™. *Mar Biol* 42:143–148. doi: 10.1007/BF00391564
- 742 De Troch M, Gurdebeke S, Fiers F, Vincx M (2001) Zonation and structuring factors of meiofauna communities in a  
743 tropical seagrass bed (Gazi Bay, Kenya). *J Sea Res* 45:45–61. doi: 10.1016/S1385-1101(00)00055-1
- 744 De Troch M, Houthoofd L, Chepurinov V, Vanreusel A (2006) Does sediment grain size affect diatom grazing by  
745 harpacticoid copepods? *Mar Environ Res* 61:265–277. doi: 10.1016/j.marenvres.2005.10.004
- 746 Dechambenoy CL, Pointier F, Sirou F, Vouvé J (1977) Apport de la thermographie infrarouge aéroportée à la  
747 connaissance de la dynamique superficielle des estuaires (système Charente-Seudre-Anse de l' Aiguillon).  
748 *Acad des Sci* 284:1269–1272.
- 749 Duarte CM, Cebrián J (1996) The fate of marine autotrophic production. *Limnol Oceanogr* 41:1758–1766. doi:  
750 10.4319/lo.1996.41.8.1758
- 751 Dupuy C, Nguyen Thanh H, Mizrahi D, Jourde J, Bréret M, Agogué H, Beaugeard L, Bocher P (2015) Structure and  
752 functional characteristics of the meiofauna community in highly unstable intertidal mudbanks in Suriname and  
753 French Guiana ( North Atlantic coast of South America ). *Cont Shelf Res* 110:39–47. doi:  
754 10.1016/j.csr.2015.09.019
- 755 Ellison RL (1984) Foraminifera and meiofauna on an intertidal mudflat, Cornwall, England: Populations; respiration  
756 and secondary production; and energy budget. *Hydrobiologia* 109:131–148. doi: 10.1007/BF00011572
- 757 Escaravage V, Garcia ME, Castel J (1989) The distribution of meiofauna and its contribution to detritic pathways in

- 758 tidal flats (Arcachon Bay, France). In: Ros JD (ed) Topics in Marine Biology. pp 551–559
- 759 Fonseca G, Hutchings P, Gallucci F (2011) Meiobenthic communities of seagrass beds (*Zostera capricorni*) and  
760 unvegetated sediments along the coast of New South Wales, Australia. *Estuar Coast Shelf Sci* 91:69–77. doi:  
761 10.1016/j.ecss.2010.10.003
- 762 Franco MA, Soetaert K, Van Oevelen D, Van Gansbeke D, Costa MJ, Vincx M, Vanaverbeke J (2008) Density,  
763 vertical distribution and trophic responses of metazoan meiobenthos to phytoplankton deposition in  
764 contrasting sediment types. *Mar Ecol Prog Ser* 358:51–62. doi: 10.3354/meps07361
- 765 Frontalini F, Semprucci F, du Châtelet EA, Francescangeli F, Margaritelli G, Rettori R, Spagnoli F, Balsamo M,  
766 Coccioni R (2014) Biodiversity trends of the meiofaunal and foraminiferal assemblages of Lake Varano  
767 (southern Italy). *Proc Biol Soc Washingt* 127:7–22. doi: 10.2988/0006-324X-127.1.7
- 768 Gallucci F, Steyaert M, Moens T (2005) Can field distributions of marine predacious nematodes be explained by  
769 sediment constraints on their foraging success? *Mar Ecol Prog Ser* 304:167–178. doi: 10.3354/meps304167
- 770 Giere O (2009) Meiobenthology: the microscopic motile fauna of aquatic sediments, Second. Springer-Verlag,  
771 Berlin Heidelberg
- 772 Gouleau D, Jouanneau JM, Weber O, Sauriau PGG (2000) Short- and long-term sedimentation on Montportail–  
773 Brouage intertidal mudflat, Marennes–Oléron Bay (France). *Cont Shelf Res* 20:1513–1530. doi:  
774 10.1016/S0278-4343(00)00035-2
- 775 Guarini J-M, Blanchard GF, Bacher C, Gros P, Riera P, Richard P, Gouleau D, Prou J, Galois R, Sauriau P-G (1998)  
776 Dynamics of spatial patterns of microphytobenthic biomass: inferences from a geostatistical analysis of two  
777 comprehensive surveys in Marennes-Oléron Bay (France). *Mar Ecol Prog Ser* 166:131–141.
- 778 Guillaumont B (1991) Utilisation de l'imagerie satellitaire pour des comparaisons spatiales et temporelles en zone  
779 intertidale. In: Elliott M, Ducrotoy J-P (eds) Estuaries and coasts: spatial and temporal intercomparisons.  
780 ECSA19 symposium. Olsen & Olsen, Fredensborg, pp 63–68
- 781 Haubois AG, Guarini JM, Richard P, Fichet D, Radenac G, Blanchard GF (2005) Ingestion rate of the deposit-feeder  
782 *Hydrobia ulvae* (Gastropoda) on epipellic diatoms: Effect of cell size and algal biomass. *J Exp Mar Bio Ecol*  
783 317:1–12. doi: 10.1016/j.jembe.2004.11.009
- 784 Heip C, Vincx M, Vranken G (1985) The ecology of marine nematodes. *Ocean Mar Biol Ann Rev* 23:399–489.
- 785 Hemminga M, Duarte CM (2000) Seagrass ecology, First. Cambridge University Press, Cambridge, Massachusetts
- 786 Hicks GRF, Coull BC (1983) The ecology of marine meiobenthic harpacticoid copepods. *Ocean Mar Biol Ann Rev*  
787 21:67–175. doi: 10.1234/2013/999990.
- 788 Higgins RP, Thiel H (1988) Introduction to the study of meiofauna. Smithsonian Institution Press, Washington DC

- 789 Hill MO (1973) Diversity and evenness: a unifying notation and its consequences. *Ecology* 54:427–432. doi:  
790 10.2307/1934352
- 791 Jensen P (1983) Meiofaunal abundance and vertical zonation in a sublittoral soft bottom, with a test of the Haps  
792 corer. *Mar Biol* 74:319–326. doi: 10.1007/BF00403458
- 793 Josefson AB, Widbom B (1988) Differential response of benthic macrofauna and meiofauna to hypoxia in the  
794 Gullmar Fjord basin. *Mar Biol* 100:31–40. doi: 10.1007/BF00392952
- 795 Kervella S (2009) Dynamique des sédiments fins et mixtes des zones intertidales de la baie de Marennes-Oléron.
- 796 Kotwicki L, De Troch M, Urban-Malinga B, Gheskiere T, Weslawski JM (2005a) Horizontal and vertical  
797 distribution of meiofauna on sandy beaches of the North Sea (The Netherlands, Belgium, France). *Helgol Mar*  
798 *Res* 59:255–264. doi: 10.1007/s10152-005-0001-8
- 799 Kotwicki L, Szymelfenig M, De Troch M, Urban-Malinga B, Węśławski JM (2005b) Latitudinal biodiversity  
800 patterns of meiofauna from sandy littoral beaches. *Biodivers Conserv* 14:461–474. doi: 10.1007/10531-004-  
801 6272-6
- 802 Lavergne C, Beaugiard L, Dupuy C, Courties C, Agogué H (2014) An efficient and rapid method for the  
803 enumeration of heterotrophic prokaryotes in coastal sediments by flow cytometry. *J Microbiol Methods*  
804 105:31–38. doi: 10.1016/j.mimet.2014.07.002
- 805 Lebreton B (2009) Analyse de la structure et du fonctionnement du réseau trophique d'un herbier par approche  
806 multi-traceurs: tracement isotopique naturel et profils acides gras.
- 807 Lebreton B, Richard P, Radenac G, Bordes M, Bréret M, Arnaud C, Mornet F, Blanchard GF (2009) Are epiphytes a  
808 significant component of intertidal *Zostera noltii* beds? *Aquat Bot* 91:82–90. doi:  
809 10.1016/j.aquabot.2009.03.003
- 810 Lebreton B, Richard P, Galois R, Radenac G, Brahmia A, Colli G, Grouazel M, André C, Guillou G, Blanchard GF  
811 (2012) Food sources used by sediment meiofauna in an intertidal *Zostera noltii* seagrass bed: A seasonal stable  
812 isotope study. *Mar Biol* 159:1537–1550. doi: 10.1007/s00227-012-1940-7
- 813 Leduc D, Probert PK (2011) Small-scale effect of intertidal seagrass (*Zostera muelleri*) on meiofaunal abundance,  
814 biomass, and nematode community structure. *J Mar Biol Assoc United Kingdom* 91:579–591. doi:  
815 10.1017/S0025315410001645
- 816 Leduc D, Probert PK, Duncan A (2009) A multi-method approach for identifying meiofaunal trophic connections.  
817 *Mar Ecol Prog Ser* 383:95–111. doi: 10.3354/meps07991
- 818 Lee S, Fuhrman JEDA (1987) Relationships between biovolume and biomass of naturally derived marine  
819 bacterioplankton. *Deep Res Part B Oceanogr Lit Rev* 53:1298–1303. doi: 10.1016/0198-0254(87)96080-8



- 820 Leguerrier D, Niquil N, Boileau N, Rzeznik J, Sauriau PG, Le Moine O, Bacher C (2003) Numerical analysis of the  
821 food web of an intertidal mudflat ecosystem on the Atlantic coast of France. *Mar Ecol Prog Ser* 246:17–37.  
822 doi: 10.3354/meps246017
- 823 Lorenzen CJ (1966) A method for the continuous measurement of in vivo chlorophyll concentration. *Deep Res*  
824 *Oceanogr Abstr* 13:223–227. doi: 10.1016/0011-7471(66)91102-8
- 825 Lü S, Wang X, Han B (2009) A field study on the conversion ratio of phytoplankton biomass carbon to chlorophyll-  
826 a in Jiaozhou Bay, China. *Chinese J Oceanol Limnol* 27:793–805. doi: 10.1007/s00343-009-9221-0
- 827 Machás R, Santos R (1999) Sources of organic matter in Ria Formosa revealed by stable isotope analysis. *Acta*  
828 *Oecologica* 20:463–469. doi: 10.1016/S1146-609X(99)00122-8
- 829 MacIntyre HL, Geider RJ, Miller DC (1996) Microphytobenthos: The ecological role of the “Secret garden” of  
830 unvegetated, shallow-water marine habitats. I. Distribution, abundance and primary production. *Estuaries*  
831 19:202. doi: 10.2307/1352225
- 832 Marbà N, Holmer M, Gacia E, Barron, Marb N (2006) Seagrass beds and coastal biogeochemistry. In: Larkum  
833 AWD, Orth RJ, Duarte CM (eds) *Seagrasses: biology, ecology and conservation*. Dordrecht: Springer, pp  
834 159–192
- 835 Maria TF, Vanaverbeke J, Esteves AM, De Troch M, Vanreusel A (2012) The importance of biological interactions  
836 for the vertical distribution of nematodes in a temperate ultra-dissipative sandy beach. *Estuar Coast Shelf Sci*  
837 97:114–126. doi: 10.1016/j.ecss.2011.11.030
- 838 Materatski P, Vafeiadou AM, Ribeiro R, Moens T, Adão H (2015) A comparative analysis of benthic nematode  
839 assemblages from *Zostera noltii* beds before and after a major vegetation collapse. *Estuar Coast Shelf Sci*  
840 167:256–268. doi: 10.1016/j.ecss.2015.07.001
- 841 Materatski P, Ribeiro R, Moreira-Santos M, Sousa JP, Adão H (2018) Nematode biomass and morphometric  
842 attributes as descriptors during a major *Zostera noltii* collapse. *Mar Biol* 165:1–17. doi:  
843 <http://dx.doi.org/10.1007/s00227-018-3283-5>
- 844 Mayer LM (1994) Relationships between mineral surfaces and organic carbon concentrations in soil and sediments.  
845 *Chem Geol* 114:347–363. doi: 10.1016/0009-2541(94)90063-9
- 846 McLusky DS, Elliott M (2004) *The estuarine ecosystem: ecology, threats and management*. OUP, Oxford
- 847 Moens T, Vincx M (1997) Observations on the feeding ecology of estuarine nematodes. *J Mar Biol Assoc United*  
848 *Kingdom* 77:211–227.
- 849 Moens T, Gansbeke D, Vincx M (1999a) Linking estuarine nematodes to their suspected food. A case study from  
850 the Westerschelde Estuary (south-west Netherlands). *J Mar Biol Assoc United Kingdom* 79:1017–1027. doi:  
851 Doi 10.1017/S0025315499001253

- 852 Moens T, Verbeeck L, de Maeyer A, Swings J, Vincx M (1999b) Selective attraction of marine bacterivorous  
853 nematodes to their bacterial food. *Mar Ecol Prog Ser* 176:165–178. doi: 10.3354/meps176165
- 854 Moens T, Dos Santos GAP, Thompson F, Swings J, Fonsêca-Genevois V, Vincx M, De Mesel I (2005a) Do  
855 nematode mucus secretions affect bacterial growth? *Aquat Microb Ecol* 40:77–83. doi: 10.3354/ame040077
- 856 Moens T, Bouillon S, Gallucci F (2005b) Dual stable isotope abundances unravel trophic position of estuarine  
857 nematodes. *J Mar Biol Assoc United Kingdom* 85:1401–1407. doi: 10.1017/S0025315405012580
- 858 Moens T, Braeckman U, Derycke S, Fonseca G, Gallucci F, Gingold R, Guilini K, Ingels J, Leduc D, Vanaverbeke  
859 J, Van Colen C, Vanreusel A, Vincx M (2013) Ecology of free-living marine nematodes. In: Schmidt-Rhaesa  
860 A (ed) *Nematoda*. De Gruyter, pp 109–152
- 861 Ndaro SGM, Olafsson E (1999) Soft-bottom fauna with emphasis on nematode assemblage structure in a tropical  
862 intertidal lagoon in Zanzibar, eastern Africa: I. spatial variability. *Hydrobiologia* 405:133–148. doi: 10.1023/A
- 863 Oksanen J, Blanchet FG, Friendly M, Kindt R, Legendre P, Mcglinn D, Minchin PR, O’hara RB, Simpson GL,  
864 Solymos P, Henry M, Stevens H, Szoecs E, Wagner H, Oksanen MJ (2017) *vegan*: community ecology  
865 package.
- 866 Pacella SR, Lebreton B, Richard P, Phillips D, DeWitt TH, Niquil N (2013) Incorporation of diet information  
867 derived from Bayesian stable isotope mixing models into mass-balanced marine ecosystem models: A case  
868 study from the Marennes-Oléron Estuary, France. *Ecol Modell* 267:127–137. doi:  
869 10.1016/j.ecolmodel.2013.07.018
- 870 Pascal P-Y (2008) *Devenir des bacteries dans les reseaux trophiques des vasieres intertidales: le cas de Brouage*  
871 (Baie de Marennes-Oleron). *Sciences du Vivant [q-bio]*. Universite de La Rochelle, 2008. Francais.
- 872 Pascal P-Y, Dupuy C, Richard P, Mallet C, du Châtelet EA, Niquil N (2009) Seasonal variation in consumption of  
873 benthic bacteria by meio- and macrofauna in an intertidal mudflat. *Limnol Oceanogr* 54:1048–1059. doi:  
874 10.4319/lo.2009.54.4.1048
- 875 Pedersen O, Borum J, Duarte CM, Fortes MD (1998) Oxygen dynamics in the rhizosphere of *Cymodocea rotundata*.  
876 *Mar Ecol Prog Ser* 169:283–288. doi: 10.3354/meps169283
- 877 Pinckney JL, Zingmark RG (1993) Modeling the annual production of intertidal benthic microalgae in estuarine  
878 ecosystems. *J Phycol* 29:396–407. doi: 10.1111/j.1529-8817.1993.tb00140.x
- 879 Platt HM, Warwick RM (1983) Free-living marine nematodes. Part 1: British enoplids. *Synopsis of the British*  
880 *Fauna (new) No. 28*. Cambridge University Press, for the Linnean Society of London and the Estuarine and  
881 *Brackish-water Sciences Association*
- 882 Platt HM, Warwick RM (1988) Free-living marine nematodes. Part II: British chromadorids. *Synopsis of the British*  
883 *Fauna (new) No. 38*.

- 884 Ravail B, Héral M, Maestrini S, Robert J-M (1988) Incidence du débit de la Charente sur la capacité biotique du  
885 bassin ostréicole du bassin de Marennes-Oléron. *J Rech Ocean* 13:48–52.
- 886 Reise K (1985) *Tidal flat ecology: An experimental approach to species interactions*, Ecological. Berlin Springer-  
887 Verlag
- 888 Reise K, Kohlus J (2008) Seagrass recovery in the Northern Wadden Sea? *Helgol Mar Res* 62:77–84. doi:  
889 10.1007/s10152-007-0088-1
- 890 Rzeznik-Orignac J, Fichet D (2012) Experimental estimation of assimilation rates of meiofauna feeding on <sup>14</sup>C-  
891 labelled benthic diatoms. *J Exp Mar Bio Ecol* 432–433:179–185. doi: 10.1016/j.jembe.2012.06.019
- 892 Rzeznik-Orignac J, Fichet D, Boucher G (2003) Spatio-temporal structure of the nematode assemblages of the  
893 Brouage mudflat (Marennes Oléron, France). *Estuar Coast Shelf Sci* 58:77–88. doi: 10.1016/S0272-  
894 7714(03)00061-1
- 895 Rzeznik-Orignac J, Boucher G, Fichet D, Richard P (2008) Stable isotope analysis of food source and trophic  
896 position of intertidal nematodes and copepods. *Mar Ecol Prog Ser* 359:145–150. doi: 10.3354/meps07328
- 897 Saint-Béat B, Dupuy C, Bocher P, Chalumeau J, De Crignis M, Fontaine C, Guizien K, Lavaud J, Lefebvre S,  
898 Montanié H, Mouget J-L, Orvain F, Pascal P-Y, Quaintenne G, Radenac G, Richard P, Robin F, Vézina AF,  
899 Niquil N (2013) Key features of intertidal food webs that support migratory shorebirds. *PLoS One* 8:1–17.  
900 doi: 10.1371/journal.pone.0076739
- 901 Seinhorst JW (1959) A rapid method for the transfer of nematodes from fixative to anhydrous glycerin.  
902 *Nematologica* 4:67–69. doi: 10.1163/187529259X00381
- 903 Semprucci F, Balsamo M (2012) Free-living marine nematodes as bioindicators: past, present and future  
904 perspectives. *Environ Res J* 6:17–36.
- 905 Semprucci F, Boi P, Manti A, Harriague AC, Rocchi M, Colantoni P, Papa S, Balsamo M (2010) Benthic  
906 communities along a littoral of the Central Adriatic Sea (Italy). *Helgol Mar Res* 64:101–115. doi:  
907 10.1007/s10152-009-0171-x
- 908 Sikora WB, Sikora JP (1982) Ecological implications of the vertical distribution of meiofauna in salt marsh  
909 sediments. *Estuar Comp* 269–282. doi: 10.1016/B978-0-12-404070-0.50022-3
- 910 Smidt ELB (1951) Animal production in the Danish Waddensea. *Meddr Kommm Danm Fisk-og Havunders, Ser Fisk*  
911 11:1–151.
- 912 Smol N, Willems KA, Govaere JCR, Sandee AJJ (1994) Composition, distribution and biomass of meiobenthos in  
913 the Oosterschelde estuary (SW Netherlands). *Hydrobiologia* 282–283:197–217. doi: 10.1007/BF00024631
- 914 Soetaert K, Vincx M, Wittoeck J, Tulkens M (1995) Meiobenthic distribution and nematode community structure in

- 915 five European estuaries. *Hydrobiologia* 311:185–206. doi: 10.1007/BF00008580
- 916 Steyaert M, Herman PMJ, Moens T, Widdows J, Vincx M (2001) Tidal migration of nematodes on an estuarine tidal  
917 flat (the Molenplaat, Schelde Estuary, SW Netherlands). *Mar Ecol Prog Ser* 224:299–304. doi:  
918 10.3354/meps224299
- 919 Steyaert M, Vanaverbeke J, Vanreusel A, Barranguet C (2003) The importance of fine-scale, vertical profiles in  
920 characterising nematode community structure. *Estuar Coast Shelf Sci* 58:353–366. doi: 10.1016/S0272-  
921 7714(03)00086-6
- 922 Steyaert M, Moodley L, Nadong T, Moens T, Soetaert K, Vincx M (2007) Responses of intertidal nematodes to  
923 short-term anoxic events. *J Exp Mar Bio Ecol* 345:175–184. doi: 10.1016/J.JEMBE.2007.03.001
- 924 Suzuki R, Shimodaira H (2015) Hierarchical clustering with p-values via multiscale bootstrap resampling.
- 925 Thornton DCO, Dong LF, Underwood GJC, Nedwell DB (2002) Factors affecting microphytobenthic biomass,  
926 species composition and production in the Colne Estuary (UK). *Aquat Microb Ecol* 27:285–300. doi:  
927 10.3354/ame027285
- 928 Tietjen J, Lee J (1977) Feeding behaviour of marine nematodes. In: Coul B (ed) *Ecology of Marine Benthos*.  
929 University of South Carolina Press, Columbia, USA, pp 21–35
- 930 Underwood GJC, Kromkamp J (1999) Primary production by phytoplankton and microphytobenthos in estuaries.  
931 *Adv Ecol Res* 29:93–139.
- 932 Underwood GJC, Paterson DM (1993) Recovery of intertidal benthic diatoms after biocide treatment and associated  
933 sediment dynamics. *J Mar Biol Assoc United Kingdom* 73:25–45. doi: 10.1017/S002531540003263X
- 934 Vafeiadou AM, Materatski P, Adão H, De Troch M, Moens T (2014) Resource utilization and trophic position of  
935 nematodes and harpacticoid copepods in and adjacent to *Zostera noltii* beds. *Biogeosciences* 11:4001–4014.  
936 doi: 10.5194/bg-11-4001-2014
- 937 Vanaverbeke J, Soetaert K, Vincx M (2004) Changes in morphometric characteristics of nematode communities  
938 during a spring phytoplankton bloom deposition. *Mar Ecol Prog Ser* 273:139–146. doi: 10.3354/meps273139
- 939 Vizzini S, Sarà G, Michener RH, Mazzola A (2002) The role and contribution of the seagrass *Posidonia oceanica*  
940 (L.) Delile organic matter for secondary consumers as revealed by carbon and nitrogen stable isotope analysis.  
941 *Acta Oecologica* 23:277–285. doi: 10.1016/S1146-609X(02)01156-6
- 942 Warwick RM (1971) Nematode associations in the Exe Estuary. *J Mar Biol Assoc United Kingdom* 51:439–454.  
943 doi: 10.1017/S0025315400031908
- 944 Warwick RM, Clarke KR (1998) Taxonomic distinctness and environmental assessment. *J Appl Ecol* 35:532–543.
- 945 Warwick RM, Price R (1979) Ecological and metabolic studies on free-living nematodes from an estuarine mud-flat.

- 946 Estuar Coast Mar Sci 9:257–271. doi: 10.1016/0302-3524(79)90039-2
- 947 Wentworth CK (1922) A scale of grade and class terms for clastic sediments. J Geol 30:377–392. doi:  
948 10.1086/622910
- 949 Wetzel MA, Fleeger JW, Powers SP (2001) Effects of hypoxia and anoxia on meiofauna: A review with new data  
950 from the Gulf of Mexico. In: Nancy N. Rabalais RET (ed) Coastal Hypoxia: Consequences for Living  
951 Resources and Ecosystems. American Geophysical Union, pp 165–184
- 952 Wieser W (1953) Die Beziehungen zwischen Mundhohlengestalt, Ernährungsweise und Vorkommen bei  
953 freilebenden marinen Nematoden. Ark Zool (Ser 2) 4:439–484.
- 954 Wieser W, Kanwisher J (1960) Growth and metabolism in a marine nematode, *Enoplus communis* Bastian. Z Vgl  
955 Physiol 43:29–36. doi: 10.1007/BF00351200
- 956