



HAL
open science

Staphylococcus aureus Nasal Colonization: An Update on Mechanisms, Epidemiology, Risk Factors, and Subsequent Infections

Adele Sakr, Fabienne Bregeon, Jean-Louis Mege, Jean-Marc Rolain, Olivier Blin

► **To cite this version:**

Adele Sakr, Fabienne Bregeon, Jean-Louis Mege, Jean-Marc Rolain, Olivier Blin. Staphylococcus aureus Nasal Colonization: An Update on Mechanisms, Epidemiology, Risk Factors, and Subsequent Infections. *Frontiers in Microbiology*, 2018, 9, 10.3389/fmicb.2018.02419 . hal-02000574

HAL Id: hal-02000574

<https://hal.science/hal-02000574v1>

Submitted on 25 Oct 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



Staphylococcus aureus Nasal Colonization: An Update on Mechanisms, Epidemiology, Risk Factors, and Subsequent Infections

Adèle Sakr^{1,2}, Fabienne Brégeon¹, Jean-Louis Mège¹, Jean-Marc Rolain¹ and Olivier Blin^{2*}

¹ Faculté de Médecine et de Pharmacie, IRD, APHM, MEPHI, IHU Méditerranée Infection, Aix-Marseille Université, Marseille, France, ² Service de Pharmacologie Clinique et Pharmacovigilance, AP-HM, Pharmacologie Intégrée et Interface Clinique et Industriel, Institut des Neurosciences Timone – UMR AMU-INSERM 1106, Aix-Marseille Université, Marseille, France

OPEN ACCESS

Edited by:

Alain Pierre Gobert,
Vanderbilt University Medical Center,
United States

Reviewed by:

Raphaël E. Duval,
Université de Lorraine, France
Michal Letek,
University of Roehampton,
United Kingdom

*Correspondence:

Olivier Blin
Olivier.BLIN@ap-hm.fr

Specialty section:

This article was submitted to
Infectious Diseases,
a section of the journal
Frontiers in Microbiology

Received: 26 July 2018

Accepted: 21 September 2018

Published: 08 October 2018

Citation:

Sakr A, Brégeon F, Mège J-L,
Rolain J-M and Blin O (2018)
Staphylococcus aureus Nasal
Colonization: An Update on
Mechanisms, Epidemiology, Risk
Factors, and Subsequent Infections.
Front. Microbiol. 9:2419.
doi: 10.3389/fmicb.2018.02419

Up to 30% of the human population are asymptotically and permanently colonized with nasal *Staphylococcus aureus*. To successfully colonize human nares, *S. aureus* needs to establish solid interactions with human nasal epithelial cells and overcome host defense mechanisms. However, some factors like bacterial interactions in the human nose can influence *S. aureus* colonization and sometimes prevent colonization. On the other hand, certain host characteristics and environmental factors can predispose to colonization. Nasal colonization can cause opportunistic and sometimes life-threatening infections such as surgical site infections or other infections in non-surgical patients that increase morbidity, mortality as well as healthcare costs.

Keywords: *Staphylococcus aureus*, nasal colonization, epidemiology, surgical site infections (SSI), nasal microbiota, predisposing factors, nasal carriage

INTRODUCTION

Staphylococcus aureus is both a human skin and mucosae commensal but also a frequent cause of serious infections with high morbidity, mortality, and healthcare-associated costs (Schmidt et al., 2015). The most frequent carriage site is the *vestibulum nasi* (or anterior nares), which serves as reservoir for the spread of the pathogen (Williams, 1963; Sivaraman et al., 2009). This bacteria can establish solid interactions with nasal epithelial cells via various proteins and many cell surface components (Wertheim et al., 2005a; Mulcahy and McLoughlin, 2016), thus transforming into persistent carriage. *S. aureus* colonizes the anterior nares of 20% to 80% of the human population (Brown et al., 2014). Nasal carriage has been shown to play a key role in the pathogenesis of *S. aureus* infections (Kluytmans et al., 1997) in patients undergoing surgery (Perl et al., 2002; Bode et al., 2010), dialysis (Kluytmans et al., 1996; Nouwen et al., 2006), and in intensive care unit (ICU) patients (Garrouste-Orgeas et al., 2001), with higher infection risks in persistent carriers (Nouwen et al., 2006).

Previously published reviews on *S. aureus* carriage have usually focused independently on colonization or infections, or have issued a specific underlying condition or surgery.

Here, we will full review recent advances in nasal microbiota composition and interspecies interactions, epidemiology, and risk factors for *S. aureus* colonization as well as the link between nasal carriage and infections both in community and nosocomial context.

NASAL MICROBIOTA AND INTERACTIONS BETWEEN BACTERIA

The adult nasal microbiota differs between individuals, but species belonging to *Corynebacterium*, *Propionibacterium*, and *Staphylococcus* genera are the most abundant bacteria (Frank et al., 2010; Human Microbiome Project Consortium, 2012; Yan et al., 2013; Kaspar et al., 2016). In a study conducted on the nasal microbiota of 178 adults, 88.2% were *Corynebacterium* carriers, 83.7% *Propionibacterium acnes* carriers, and 90.4% *Staphylococcus epidermidis* carriers. Proportional abundance varied considerably between individuals (Liu C.M. et al., 2015).

The health status may influence the nasal microbiota and vice versa. In a study involving healthy and hospitalized individuals, healthy adults harbored nares microbiota dominated by *Actinobacteria* (mainly *Propionibacterium* and *Corynebacterium* spp.) whereas patients microbiota were dominated by *S. aureus* and *S. epidermidis*. *S. aureus* colonization was negatively associated with the presence of other bacteria including *S. epidermidis* (Frank et al., 2010). Such counterweight effect between bacteria could be the result of interdependent activation-inhibition mechanisms as reviewed by Krismer et al. (2017). In fact, some bacterial species are capable of secreting anti-staphylococcal molecules modulating *S. aureus* abundance (Figure 1). For instance, *in vitro* production of H₂O₂ by *Streptococcus pneumoniae* can be bactericidal on *S. aureus* (Regev-Yochay et al., 2006; Selva et al., 2009). Recently, an *in vitro* and human study demonstrated that lugdunin, a non-ribosomal synthesized bioactive compound produced by *Staphylococcus lugdunensis*, can prevent *S. aureus* nasal colonization via a bactericidal effect (Zipperer et al., 2016).

In some cases, the bacteria-secreted molecules can modify *S. aureus* adhesion properties. Some types of *S. epidermidis* seem to be capable of synthesizing the serine protease Esp that eliminates nasal *S. aureus* in healthy humans (Iwase et al., 2010), probably by degrading staphylococcal surface proteins and human receptors critical for host-pathogen interaction (Sugimoto et al., 2013). As well, *Propionibacterium* species produce coproporphyrin III, a porphyrin metabolite that induces *S. aureus* aggregation which influences nasal colonization (Wollenberg et al., 2014).

Corynebacterium species are suggested to antagonize *S. aureus* by human cell binding competition mechanisms (Uehara et al., 2000; Lina et al., 2003). In 156 healthy volunteers, Uehara et al. (2000) observed a 71% total eradication rate of nasal *S. aureus* after performing up to 15 inoculations of a *Corynebacterium* sp. strain to the nares of *S. aureus* carriers.

Intra-species competition has also been described. In a cross-sectional clinical study, it was suggested that methicillin-sensitive *S. aureus* (MSSA) and methicillin-resistant *S. aureus* (MRSA)

compete for colonization, MSSA being protective with regard to MRSA carriage (Dall'Antonia et al., 2005). On the other hand, pre-existing nasal carriage with *S. aureus* could predispose adult patients to further staphylococcal colonization (Ghasemzadeh-Moghaddam et al., 2015).

SPREAD AND TRANSMISSION OF S. aureus


Staphylococcus aureus can be found in different body sites like the skin, rectum, vagina, gastrointestinal tract and axilla, the anterior nares appearing as the main reservoir. From a cutaneous commensal site, *S. aureus* can enter in contact with the nasal mucosa, then interact with epithelial cell ligands such as loricrin and cytokeratin 10 (K10) (Table 1). Once the host's defenses are overcome, *S. aureus* can propagate into the anterior nares so that the host becomes an *S. aureus* nasal carrier (Wertheim et al., 2005a). In human, nasal colonization may begin within the first days of life (Maayan-Metzger et al., 2017). This has been demonstrated in a cohort study evaluating nasal carriage of *S. aureus* in 100 pairs of infant-mother for a period of 6 months following delivery (Peacock et al., 2003). The carriage rate in the first 8 weeks of life was around 40–50%, thereafter it dropped to 21% at 6 months. In addition, this study found a nasal carriage concordance in 68% of infant-mother pairs attesting the role of environmental factors in *S. aureus* carriage (Peacock et al., 2003). Another study found identical strains in 80% of infant-mother pairs. In 90% of these newborns, the source of *S. aureus* was the maternal nasal strain (Leshem et al., 2012; Figure 2).

After birth, hands are the main vector for *S. aureus* transmission from surfaces to the nose (Wertheim et al., 2005a). The hypothesis of a link between hand and nose *S. aureus* carriage is supported by the double blind randomized placebo controlled trial from Reagan et al. (1991), who demonstrated that nasal decolonization with mupirocin applied to health-care workers resulted in a decrease of nose and hand carriage. In a cohort study including outpatients and healthy hospital employees, nasal carriage was evaluated by a single or several swabs. Participants completed a questionnaire about their nose picking behavior, a positive correlation between this habit and nasal carriage of *S. aureus* was found. However, it is unknown whether nose-picking patients were more frequently colonized at extra nasal sites (Wertheim et al., 2006).

Studies realized in individuals living in the same households have revealed that these people tend to carry genetically similar strains in their nares (Nouwen and Optima Grafische Communicatie, 2004; Muthukrishnan et al., 2013) suggesting horizontal transmission. Multisite MRSA carriage increases the risk for nasal MRSA colonization (Harbarth et al., 2000).

In spite infrequent, airborne transmission is another possible route of *S. aureus* dissemination (Wertheim et al., 2005a) and may play a role in hospital outbreaks (Sherertz et al., 1996).

During viral upper respiratory infections, the risk of disseminating endogenous *S. aureus* in the air increases and infection outbreaks may occur. In 1996, an MRSA outbreak involving 8 of 43 patients occurred in a surgical ICU of a



university hospital in the United States. Investigations of the cause concluded that a single physician was the source of this outbreak; he was a nasal carrier of MRSA and suffered an upper respiratory infection. To assess airborne dispersal of *S. aureus*, the authors completed their findings by an experimental clinical test on this physician and showed that transmission of the bacterium increased by 40-fold when he was infected by a rhinovirus infection than when he was not. The use of a mask significantly reduced dispersal ($P = 0.015$) (Sherertz et al., 1996).


Healthcare workers who are asymptomatic nasal carriers can sometimes be the source of MRSA outbreaks (Wang et al., 2001; Vonberg et al., 2006; Haill et al., 2013; Lamanna et al., 2017). On the other hand, in nonoutbreak situations and in presence of control measures, healthcare workers are infrequently sources of transmission of *S. aureus* (Price et al., 2017).

Mobile phones of healthcare workers may be a reservoir of *S. aureus* (Chang et al., 2017). A recent study evaluated incidence of bacterial contamination of mobile phones belonging to medical staff working in the operating room. Seventy two healthcare professionals took bacterial cultures from their phones, anterior nares, and hands. The results revealed that 31 staff had *S. aureus* isolated from their nares, 8 from their mobile

phones, and 4 from their hands. Genotyping confirmed that 7/8 of the mobile phones strains were identical to the ones isolated from the nares (Chang et al., 2017).

TABLE 1 | Major *S. aureus*-host ligands.

<i>S. aureus</i> ligand factor	Host ligand	Reference
ClfB	Loricrin, K10 (cytokeratin 10), K8 (cytokeratin 8), fibrinogen	Perkins et al., 2001; O'Brien et al., 2002; Schaffer et al., 2006; Wertheim et al., 2008; Haim et al., 2010; Mulcahy et al., 2012
IsdA	Fibrinogen, fibronectin	Clarke et al., 2004
SdrC	Unknown	Corrigan et al., 2009
SdrD	Desmoglein 1	Corrigan et al., 2009; Askarian et al., 2016
SasX	Unknown	Liu Q. et al., 2015
SasG	Unknown	Roche et al., 2003
WTA	Srec-1	Baur et al., 2014; Weidenmaier et al., 2004



MECHANISMS OF COLONIZATION

The *vestibulum nasi* or anterior part of the nares is lined by a stratified, keratinized nonciliated squamous epithelium, whereas the rest of the nasal cavity, that is, its inner part is lined with a ciliated columnar epithelium (Peacock et al., 2001; Weidenmaier et al., 2012).

Both epithelia have been described as habitats for *S. aureus* (Mulcahy et al., 2012; Baur et al., 2014) as it will be developed in this section. Intracellular localization in nasal tissue from healthy volunteers was also described (Hanssen et al., 2017). For a successful colonization, *S. aureus* expresses adhesive molecules (Burian et al., 2010), fundamental for the establishment of interactions with human cell surface components, as it was demonstrated *in vitro* and *in vivo* (Mulcahy et al., 2012; Baur et al., 2014; **Figure 3**). Major ligands interactions are listed in **Table 1**.

INTERACTION BETWEEN *S. aureus* AND THE SQUAMOUS EPITHELIUM OF THE ANTERIOR NARES


The *vestibulum nasi* is lined by skin (Cunningham, 1905; Vuyk and Watts, 2006). The uppermost layer of the epidermis is the *stratum corneum* or cornified layer (Candi et al., 2005; Eckhart et al., 2013). This layer contains keratinocytes that express proteins, such as loricrin, cytokeratin 10 (K10), involucrin, filaggrin, and other proteins (Eckhart et al., 2013).

Clumping factor B (ClfB) and iron-regulated surface determinant A (IsdA) are staphylococcal surface proteins that can adhere to cornified envelope proteins (Clarke et al., 2004, 2006; Schaffer et al., 2006; Wertheim et al., 2008; Mulcahy et al., 2012) and favor nasal colonization as it will be developed in this section.

Recently, Mulcahy et al. (2012) demonstrated that loricrin, the most abundant protein in the cornified envelope (Steinert and Marekov, 1995), was the major target ligand for ClfB during *S. aureus* nasal colonization. Nasal colonization by a ClfB⁺ *S. aureus* strain was reduced by 80% in loricrin-deficient mice compared to wild-type mice. ClfB has also been shown to interact with cytokeratin 10 (O'Brien et al., 2002; Schaffer et al., 2006; Wertheim et al., 2008), cytokeratin 8 (Haim et al., 2010), and fibrinogen (Perkins et al., 2001; **Table 1**). The role of ClfB in the adherence of *S. aureus* to the nasal epithelium has been studied *in vitro* (O'Brien et al., 2002; Mulcahy et al., 2012), in animal models (Schaffer et al., 2006; Mulcahy et al., 2012), and in human studies (Wertheim et al., 2008).

Schaffer et al. (2006) demonstrated that mutant strains of *S. aureus* deficient in ClfB resulted in a reduced nasal colonization in both mice and rats, as compared to the wild-type strain (ClfB⁺). In human, the mutant strain ClfB⁻ of *S. aureus* has been reported to be cleared faster than the wild-type strain (median 3 ± 1 vs. 7 ± 4 days, *p* = 0.006), whereas the wild-type strain persisted for 28 days after inoculation (Wertheim et al., 2008).

IsdA also plays a role in the adherence of *S. aureus* to nasal cells as demonstrated *in vitro* and *in vivo* (Clarke et al., 2004). Mutation in IsdA gene reduced the ability of the bacteria to bind



human nasal cells *in vitro* and to colonize the anterior nares in cotton rats (Clarke et al., 2006). This surface protein has also been shown to bind to fibrinogen and fibronectin (Clarke et al., 2004). The role of IsdA in nasal colonization has not been clearly demonstrated in human studies yet.

Other *S. aureus* surface proteins, such as surface protein G (SasG), SasX, and the serine-aspartate repeat proteins SdrC and SdrD may also serve as ligands to the epithelial cells (Corrigan et al., 2009; Liu Q. et al., 2015). Desmoglein 1 was identified as a host ligand for SdrD (Askarian et al., 2016). Their roles in humans have not been tested yet (Mulcahy et al., 2012).

INTERACTION IN THE INNER NASAL CAVITY

Apart from the *vestibulum nasi*, the inner part of the nasal cavity constitutes another ecological niche for *S. aureus*. The staphylococcal nonprotein adhesin, named cell wall teichoic acid (WTA) is considered as an important factor for the colonization process (Weidenmaier et al., 2004). In an *in vivo* study, WTA-deficient *S. aureus* mutants could not adhere to nasal cells and were unable to colonize cotton rat nares compared to wild type control strains (Weidenmaier et al., 2004).

In a study combining *in vitro* and *in vivo* assessments, Baur et al. (2014) studied the molecular details of WTA adhesion to

nasal cells. They first discovered that SREC-1 (a member of the F-type scavenger receptor) was expressed on epithelial cells in the inner nasal cavity of human and cotton rats. The authors further reported that SREC-1 interacted with WTA and confirmed these findings in infected cotton rats pretreated with antiSREC-1 antibody. A significant decrease in colonization was observed 8 h and 6 days after inoculation in the group receiving antiSREC-1 antibody, as compared to the controls (Baur et al., 2014). In agreement, Weidenmaier et al. (2008) previously demonstrated the role of WTA in the initial stages of *S. aureus* colonization.

INTRACELLULAR LOCALIZATION OF *S. aureus*

Intracellular localization of *S. aureus* in the nasal tissue has been described using techniques including immunohistochemical analysis, hematoxylin, and Eosin stains (Hayes et al., 2015; Ou et al., 2016, 2017). Epithelial cells, endothelial cells, and inflammatory cells, especially mast cells (Plouin-Gaudon et al., 2006; Tan et al., 2014; Hayes et al., 2015; Ou et al., 2016) have been described as habitats for intracellular *S. aureus* (ICSA). Using skin biopsies from the *vestibulum nasi* of randomly selected healthy participants, Hanssen et al. (2017) detected intracellular *S. aureus* in the *stratum spinosum*, one of the layers forming the epidermis. Intracellular localization of *S. aureus* has been reported both in patients with rhinosinusitis

(Clement et al., 2005; Plouin-Gaudon et al., 2006; Ou et al., 2016, 2017) and healthy volunteers (Hanssen et al., 2017). In the control population from the study of Ou et al. (2016), 38% had ICSA. However, in the absence of large scale population study, the true prevalence of ICSA in healthy individuals remains to determine.

Intracellular localization of *S. aureus* seems to protect the bacteria from host defense mechanisms (Clement et al., 2005) and favor antimicrobial agents' failure, as it was demonstrated in a recent study evaluating systemic and topical antibiotics. Using a cell model of *S. aureus* nasal epithelium evasion, Rigai et al. (2018) found that most of the decolonizing agents, including mupirocin, exhibited weak activity against ICSA. Elimination of ICSA was also studied in a mouse model using vancomycin. In this study, intracellular bacteria were able to establish infection even in the presence of this antibiotic (Lehar et al., 2015), whereas planktonic bacteria were eliminated by vancomycin and did not cause infection. The intracellular residency of *S. aureus* could explain re-colonization and failure in decolonization observed in some healthy carriers, and the recurrence of infections observed in patients with chronic rhinosinusitis (Clement et al., 2005; Ou et al., 2017; Rigai et al., 2018).

IMMUNE SYSTEM AND *S. aureus* COLONIZATION

Our knowledge in terms of host response during *S. aureus* nasal colonization is still limited as compared to immune response in invasive *S. aureus* infection (Brown et al., 2014). The association between the host immune system and *S. aureus* nasal carriage has been recently reviewed by Mulcahy and McLoughlin (2016). The presence of *S. aureus* in the nares seems to induce both innate and adaptive immune system. *S. aureus* can otherwise overcome host defense mechanisms. The staphylococcal protein A (SpA) and the exoprotein staphylokinase could play a key role in the bacterial adaptation to the immune system and colonization process (Peschel and Sahl, 2006; Cole et al., 2016).

Innate Immune Response

Human nasal secretions have efficient natural antimicrobial activity (Cole et al., 1999). Indeed, they contain epithelial-cells secreted antimicrobial peptides (AMPs) contributing to the first line defense in the nose. Primary nasal epithelial cells yield AMPs such as hBD3 and RNase-7 after stimulation with cytokines (IFN- γ , IL-1 β , and TNF- α) (Burgey et al., 2016). Media containing *S. aureus* induce the expression of these AMPs less efficiently than inflammatory cytokines (Burgey et al., 2016). Differential expression profiles of AMPs could be primary determinants for different *S. aureus* carriage states (Burgey et al., 2016).

Anti-staphylococcal responses can be induced by the involvement of Toll-like receptors (TLR). Hence, the role of TLR2 has been documented in nasal *S. aureus* colonization (González-Zorn et al., 2005). Carrier strains of *S. aureus* have been shown to delay the host's immune response compared to non-carrier strains due to the delayed TLR2 expression they stimulate in nasal epithelial cells (Quinn and Cole, 2007). Different polymorphisms have been found in *S. aureus* nasal

carriers and non-carriers; they target soluble or membrane-binding molecules such as genes encoding mannose-binding lectin, glucocorticoid receptor, C-reactive protein, beta-defensin 1, TLR-9, and interleukin-4 (van den Akker et al., 2006; Emonts et al., 2008; Ruimy et al., 2010; Mulcahy and McLoughlin, 2016; Nurjadi et al., 2018).

A study evaluating neutrophil depletion was performed in mice to determine the importance of neutrophil influx during carriage. Mice that received an antibody for neutrophil depletion had a significant increase in *S. aureus* colonization rate as compared to controls. This *in vivo* study highlights that neutrophil influx plays a role in clearance of *S. aureus* (Archer et al., 2013).

Adaptive Immune Response

As documented from serum analysis, *S. aureus* carriage induces an adaptive humoral immune response. Serum levels of immunoglobulin IgG and IgA specific to several staphylococcal proteins have been reported to be higher in persistent carriers than in non-carriers (van Belkum et al., 2009; Colque-Navarro et al., 2010). But evidence that these antibodies are protective against colonization are lacking.

In a murine model of nasal colonization, clearance of *S. aureus* was found to be B-cell independent but T-cell mediated. The Th17 cells, a functional T-cell lineage distinct from Th1 and Th2 are known to be involved in mucosal barriers and surface pathogen clearance. They typically produce IL-17 cytokine family. Archer et al. (2013, 2016) showed that IL-17A and IL-17F deficient mice failed to clear *S. aureus* experimentally inoculated through nasal route; IL-17A was required for AMP expression induced by nasal colonization. In a murine model, the IL-10 family cytokine IL-22 was secreted by Th17 cells and facilitated local expression of AMPs. IL-22 also reduced the expression of staphylococcal ligands loricrin and cytokeratin 10 and thus controlled nasal colonization (Mulcahy et al., 2016).

By contrast, in a human study evaluating nasal fluid inflammatory factors and nasal carriage with *S. aureus*, IL-17 was not detected (Cole et al., 2016). This was surprising and the authors suggested that IL-17 could be insoluble in nasal secretions (Cole et al., 2016). Low Th1 to Th17 cytokines ratio were found to be predictive of *S. aureus* carriage in volunteers after whole blood stimulation with *S. aureus* (Nurjadi et al., 2016). Future human studies should be conducted to better understand the role of Th17 cytokines in nasal carriage.

MICROBIOLOGY-BASED CLASSIFICATION OF NASAL CARRIERS

In order to confirm nasal carriage of *S. aureus*, samples are commonly collected using commercial dry or moistened sterile swabs, with no significant difference found between the two (Warneke et al., 2016). The protocol of sampling is not really standardized but generally consists of rubbing the swab in the anterior nares of each nostril for approximately four rotary movements (Wertheim et al., 2006; Bode et al., 2010). Swabs are then analyzed to check presence of *S. aureus*. The two

commonly used laboratory tests for the identification of *S. aureus* are the culture on chromogenic solid media – which is the less expensive test – and polymerase chain reaction, which is the gold standard and the most rapid technique for MRSA detection (Pournajaf et al., 2014). It is important to note that chronic carriage of *S. aureus* may not only result from nasal colonization. Sampling extra-nasal sites like oropharynx, rectum, wounds, axilla increases MRSA detection rate in patients at risk for *S. aureus* nosocomial infection (Miller et al., 2012; McKinnell et al., 2013).

Epidemiological studies over periods varying from 12 weeks to 3 years have described three nasal carriage patterns for *S. aureus* among healthy volunteers swabbed several times. Persistent carriers which rates ranged from 10% to 30% (Williams, 1963; Eriksen et al., 1995; VandenBergh et al., 1999; Nouwen et al., 2004; Muthukrishnan et al., 2013), non-carriers which rates ranged from 10% to 47%, and the rest who were considered intermittent carriers (Williams, 1963; Eriksen et al., 1995; VandenBergh et al., 1999; Nouwen et al., 2004). The definitions used for this classification varied from one study to another. In some studies, persistent carriers were the ones who had results of all of their swabs positive for *S. aureus* (Muthukrishnan et al., 2013). Other studies defined cut-off values for carrier-index (number of positive swabs/number of total swabs for each person) (Eriksen et al., 1995; VandenBergh et al., 1999). However, there is no clear definition on the number of swabs that should be taken and what fraction should be positive before determining carriage state.

Nouwen et al. (2004) proposed a culture rule based on the combination of qualitative and quantitative results of two consecutive nasal culture swabs taken approximately a week apart, to predict *S. aureus* nasal carrier state among healthy volunteers. When both cultures are positive over 10^3 colony forming units (cfu) a person is classified as persistent carrier. When only one of the cultures is positive, or when both cultures are positive with a low cfu, the person is considered as an intermittent carrier. van Belkum et al. (2009) performed a study in 51 volunteers with a known carriage state, who were artificially colonized with a mixture of *S. aureus* strains after a decolonization treatment and followed for 22 weeks. Median nasal bacterial survival was of 4 days in noncarriers, 14 days in intermittent carriers, and more than 154 days in persistent carriers ($P = 0.017$). The culture swabs of persistent carriers contained more bacterial loads compared to the other groups. Serum levels of anti-staphylococcal antibodies differed between persistent and nonpersistent carriers as previously stated. The authors concluded that intermittent and noncarriers share similar characteristics, and thus suggested a reclassification of *S. aureus* nasal carriers into persistent carriers and “other” carriers (van Belkum et al., 2009). Persistent carriers have been shown to have higher counts of *S. aureus* (Nouwen et al., 2004; van Belkum et al., 2009) and a higher risk of infection compared to other carriers (Wertheim et al., 2004; Nouwen et al., 2005). However, bacterial loads in persistent carriers are always variable and it is hard to determine a fixed threshold for diagnosis (Burian et al., 2010; Verhoeven et al., 2012; Nilsson et al., 2015). Persistent carriage is more common in children than in adults, but many people change their carrier state between 10

and 20 years old (Armstrong-Esther, 1976). Conclusions that persistent carriers harbor the same strain for many years, and intermittent carriers appear to have changing strains (Eriksen et al., 1995; VandenBergh et al., 1999) have been recently the subject of controversy (Muthukrishnan et al., 2013).

Cross-sectional studies yield a prevalence of approximately 20–30% of carriers in the general population which is a mix of persistent and intermittent carriers (Gorwitz et al., 2008; den Heijer et al., 2013; Saadatian-Elahi et al., 2013; Chen et al., 2017).

INDIVIDUAL RISK FACTORS FOR *S. aureus* NASAL COLONIZATION

Nasal colonization depends on host factors, such as the underlying condition or diseases (Table 2). Some studies have found that nasal carriage was more frequent in human immunodeficiency virus (HIV)-infected (Raviglione et al., 1990; Weinke et al., 1992; Kotpal et al., 2016) or obese patients (Olsen et al., 2013), compared to healthy individuals. This higher prevalence was also found among diabetic patients undergoing dialysis compared to non-diabetic patients in the same population (Luzar et al., 1990). Other diseases such as granulomatosis with polyangiitis (formerly known as Wegener’s granulomatosis), rheumatoid arthritis (Laudien et al., 2010), skin and soft tissue infections (Immergluck et al., 2017) atopic dermatitis (Breuer et al., 2002), and recurrent furunculosis (Demos et al., 2012) have been related with an increased carriage rate.

TABLE 2 | Predisposing factors for nasal carriage.

Predisposing factors for nasal carriage	Reference
HIV-infection	Raviglione et al., 1990; Weinke et al., 1992; Kotpal et al., 2016
Obesity	Olsen et al., 2013
Diabetic patients undergoing dialysis (compared to non-diabetic patients in the same population)	Luzar et al., 1990
Granulomatosis with polyangiitis	Laudien et al., 2010
Rheumatoid arthritis	Laudien et al., 2010
Skin and soft tissue infections	Immergluck et al., 2017
Recurrent furunculosis	Demos et al., 2012
Atopic dermatitis	Breuer et al., 2002
Hemoglobin in nasal secretions	Pynnonen et al., 2011
Histocompatibility antigen phenotype HLA-DR3	Kinsman et al., 1983
Polymorphisms in genes encoding for the glucocorticoid receptor, interleukin-4, C-reactive proteins, and complement inhibitor proteins	van den Akker et al., 2006; Emonts et al., 2008; Ruimy et al., 2010
Hormonal contraception use	Zanger et al., 2012
Active smokers: controversial	Olsen et al., 2012; Cole et al., 2018
Hospital workers: controversial	Elie-Turenne et al., 2010; Saadatian-Elahi et al., 2013; Chen et al., 2015; Price et al., 2017

In healthy subjects, Liu C.M. et al. (2015) found similar carriage rates in men and women, while men had higher bacterial density. Reports of a higher risk of nasal carriage of *S. aureus* among hospital workers than the rest of the population have not been confirmed (Elie-Turenne et al., 2010; Saadatian-Elahi et al., 2013; Chen et al., 2015; Price et al., 2017). The association between smoking and nasal carriage seems also controversial. In a study by Olsen et al. (2012), active smoking in healthy adults was found to be a protective factor for carriage of *S. aureus*, with a hypothesized bactericidal activity of cigarette smokes. Conversely, a recent study showed that smokers were more frequently colonized than non-smokers, and cessation from smoking improved clearance of nasal *S. aureus* in an experimental inoculation study (Cole et al., 2018). Many other host conditions have been punctually studied and reported as additional predisposing factor such as hormonal contraception (Zanger et al., 2012) and presence of hemoglobin in nasal secretions (Pynnonen et al., 2011).

At the genetic level, no correlation was found between genetic factors and *S. aureus* carriage. No significant heritability for *S. aureus* nasal colonization was detected in twins and family studies (Roghmann et al., 2011; Andersen et al., 2012). Interestingly, some polymorphisms in host inflammatory response genes have been associated with *S. aureus* nasal carriage. The presence of the histocompatibility antigen phenotype HLA-DR3 could be a predisposition (Kinsman et al., 1983).

As previously said, at the immune system level, polymorphisms in genes encoding some proteins and differential expression profiles of AMPs could be the determinants of the various carriage states.

In a study involving 93 type 1 diabetes patients, vitamin D receptor polymorphisms were determined in Deoxyribonucleic acid (DNA) extracted from peripheral blood leukocytes. Analysis showed that presence of specific alleles coding for vitamin D receptors were associated with an increased rate of *S. aureus* colonization (Panierakis et al., 2009).

NASAL CARRIAGE OF *S. aureus* AS A RISK FACTOR FOR INFECTIONS

Staphylococcus aureus nasal colonization has been identified as a major risk factor for the development of patent staphylococcal infections, whether community acquired, or nosocomial (Von Eiff et al., 2001; Wertheim et al., 2004, 2005b) which increases the risk by 2 to 10 times (Perl and Golub, 1998). The risk of infection in nasal carriers has been mainly studied in surgical patients (general, orthopedic, cardiac, and neurosurgeries) (Perl et al., 2002; Bode et al., 2010; Walsh et al., 2017), patients on hemodialysis (Kluytmans et al., 1996; Katneni and Hedayati, 2007), patients on chronic ambulatory peritoneal dialysis (CAPD) (Luzar et al., 1990), HIV-infected patients (Nguyen et al., 1999; Sissolak et al., 2002), and intensive care unit patients (Nardi et al., 2001). It has also been shown to be the primary risk factor for recurrent furunculosis, nasal colonization being present in almost 60% of individuals with furuncles and impetigo (Durupt et al., 2007) (Figure 2 and Table 3).

SURGICAL SITE INFECTIONS

Developing a post-operative infection is a multifactorial process usually combining preoperative, intraoperative, and post-operative factors (Savage and Anderson, 2013). Nasal colonization can actually be considered as a preoperative risk factor for MRSA and MSSA infections. *S. aureus* can spread from the anterior nares to other areas on the skin surface and thus contaminate the surgical wound during the operative procedure (NICE Clinical Guidelines, 2008; Savage and Anderson, 2013). It has been shown that around 80% of strains causing a staphylococcal infection at the site of surgery have molecular identity with *S. aureus* isolates in the nares of concerned patients (Perl et al., 2002).

Surgical site infections (SSI) are one of the most common post-operative complications and represent 20 to 30% of healthcare associated infections (HCAI) (Klevens et al., 2007; Magill et al., 2012; Savage and Anderson, 2013). While enterobacteria and other uro-digestive bacteria are dominant in infections after gastrointestinal, urological and gynecological surgeries (Trautman et al., 2007), *S. aureus* predominates in orthopedic and cardiac surgery settings (Lepelletier et al., 2005; Trautman et al., 2007; Muñoz et al., 2008). In orthopedic patients, biofilms can form on the implants leading to therapeutic challenges (Chen et al., 2013). According to the French Institute for Public Health Surveillance in 2014, *S. aureus* composed 19.4% of isolated germs in SSIs after coronary surgeries, 51.9% of organisms causing SSIs in orthopedic surgeries, and 29.3% of organisms causing infections in gynecologic obstetric surgery in 2014 (Le Réseau d'alerte, d'investigation et de surveillance des infections nosocomiales (Raisin), 2015). Similar implications have been also found in other countries (Saadatian-Elahi et al., 2008; Negi et al., 2015).

Several studies including case-control and multivariate analysis have identified nasal carriage of *S. aureus* an independent risk factor for SSIs (Kluytmans et al., 1995; Kalmeijer et al., 2000; Muñoz et al., 2008). In the case-control study from Kluytmans et al. (1995), cardiac surgery patients were screened for their

TABLE 3 | *S. aureus* nasal colonization, a risk factor for infections.

<i>S. aureus</i> nasal colonization, a risk factor for	Reference
Surgical site infections after orthopedic surgeries	Kalmeijer et al., 2000; Yano et al., 2000; Weiser and Moucha, 2015
Surgical site infections after cardiac surgeries	Kluytmans et al., 1995; Muñoz et al., 2008
Bacteremia in nonsurgical patients	Wertheim et al., 2004
Catheter-related infections in dialysis patients	Luzar et al., 1990; Katneni and Hedayati, 2007
<i>S. aureus</i> infections in HIV-infected patients	Nguyen et al., 1999; Sissolak et al., 2002
ICU-associated <i>S. aureus</i> infections	Honda et al., 2010
Recurrent furunculosis and impetigo	Durupt et al., 2007; Demos et al., 2012
Diabetic foot ulcer infections	Dunyach-Remy et al., 2017

nasal carriage status the day before surgery and followed for the development of an SSI. *S. aureus* wound infections occurred in 40 patients as opposed to 120 controls who did not develop infection. Nasal carriage, identified in 52% of cases as compared to 12% of controls, was found to be a significant risk factor for the development of these post-operative infections with an OR = 9.6, 95% CI (3.9–23.7). In a study involving 357 patients undergoing major heart surgery, nasal carriers had a higher incidence of SSI than non-carriers (12.5% vs. 5%, $P = 0.01$) (Muñoz et al., 2008). Similar conclusions were obtained for orthopedic patients with an incidence increasing from 3- to 11-fold (Kalmeijer et al., 2000; Yano et al., 2000; Weiser and Moucha, 2015).

Surgical site infections have an important impact on the patient and the healthcare system. These infections increase hospital stay (Leaper et al., 2004), mortality and healthcare costs and decrease health-related quality of life (Anderson et al., 2009; Lamarsalle et al., 2013; Savage and Anderson, 2013). A retrospective database analysis in France concluded that staphylococcal infections led to approximately 1.0 and 1.4 additional hospitalizations per patient, 22.1 and 22.4 additional hospital days, and an excess cost of €15,475 and €13,389 after cardiothoracic and orthopedic surgeries respectively (Schmidt et al., 2015). The rate of in-hospital mortality was 2.6 times and six times higher among infected patients than non-infected patients in cardiothoracic and orthopedic procedures (Schmidt et al., 2015).

Methicillin-resistant *S. aureus* carriage could increase the risk for the development of SSIs. In a systematic review, patients colonized with MRSA were four times more likely to develop invasive infection than patients colonized with MSSA (Safdar and Bradley, 2008). MRSA infections are reported to cause up to 40% of HCAI worldwide with particularly high incidence in the United States and many European countries (McKinnell et al., 2013) and have been shown to have increased morbidity as compared to MSSA (Anderson et al., 2009). In a study carried out by Anderson et al. (2009), patients with SSI due to MRSA had a 2.6 higher risk of dying within 3 months, the duration of hospital stay was 6 days longer, and the related cost was increased of \$23,000 compared to patients with SSI due to MSSA. On the other hand, a meta-analysis of 31 cohort studies on *S. aureus* bacteremia, showed significantly higher mortality with MRSA than with MSSA (Cosgrove et al., 2003). However, the reasons for increased fatality rate with MRSA infections are unclear. Some authors suggest that MRSA and MSSA bacteria are equally virulent but MRSA infections usually develop in patients previously treated with antibiotics. Thus, as previously suggested, the differences in patients' fatality rate may rather reflect the severity of underlying conditions than a higher bacteria-related increased virulence (Humphreys, 2012; Hraiech et al., 2013).

INFECTIONS IN NONSURGICAL PATIENTS

Nasal carriage of *S. aureus* has also been found to be a risk factor for subsequent infections in nonsurgical patients.

Wertheim et al. (2004) screened 14,008 adults who had nasal swab on admission in a nonsurgical department, 3420 (24%) were positive for *S. aureus*. The follow-up identified 81 patients who developed *S. aureus* bacteremia between 2 and 120 days after swabbing, which was three times more frequent in carriers than in non-carriers. Interestingly, the death rate from *S. aureus* bacteremia was higher in non-carriers than carriers. This could be due to the protective immunity of anti-staphylococcal antibodies (Wertheim et al., 2004; Holtfreter et al., 2006). Approximately, 80% of *S. aureus* blood isolates causing bacteremia were of endogenous origins and identical to those isolated from the anterior nares of corresponding patients, thus confirming previous report (Von Eiff et al., 2001).

A prospective cohort study evaluated the occurrence of *S. aureus* infections in 5161 patients who were screened for nasal carriage when admitted to the ICU. ICU-associated *S. aureus* infections were defined by the development of infection >48 h after their admission to the unit. These infections occurred in 113 patients and nasal colonization was associated with a 2.5 to 4.7-fold increased risk (Honda et al., 2010).

In HIV-positive patients, a prospective cohort study evaluating 231 subjects every 3 months for a minimum of two years, has reported a 6% incidence of *S. aureus* infections, nasal carriers being more at risk [$p = 0.04$, OR = 3.6 (0.9–15.4)] (Nguyen et al., 1999).

In hemodialysis and chronic peritoneal dialysis patients, most of infectious complications come from endogenous origin (Luzar et al., 1990; Ena et al., 1994). *S. aureus* is the most common isolated agent from central venous catheter-related bacteremia (Katneni and Hedayati, 2007), or exit-site infections of peritoneal dialysis catheters (Luzar et al., 1990). Nasal carriers are at increased risk of contracting these infections (Nouwen et al., 2005; Ong et al., 2017).

Staphylococcus aureus is the most frequently isolated pathogen from diabetic foot infections. A study compared the genotypic profiles of *S. aureus* strains isolated from the nares and diabetic foot ulcer infections of 276 patients. The bacterium was isolated from both sites in 36% of the population, and identical strains were found in 65% of cases. Further investigations should be performed in order to confirm the benefit of screening and treating nasal carriers in this population (Dunyach-Remy et al., 2017).

Nasal carriage of *S. aureus* constitutes a risk factor for the development of skin and soft tissue infections caused by this germ in non-hospitalized and non-diseased subjects (Chou et al., 2015). It has been shown to be the primary risk factor for recurrent furunculosis (Demos et al., 2012). Nasal carriage is also a risk factor for secondary bacterial pneumonia in patients having influenza A virus infection. The viral infection causes host physiologic changes that generates the dissemination of *S. aureus* from nasal tissue to the lungs as demonstrated in a mouse model (Reddinger et al., 2016).

PERSPECTIVES

Nasal carriage of *S. aureus* is multifactorial and can predispose carriers to subsequent infections. Nasal decolonization of carriers is therefore recommended in patients undergoing cardiothoracic and orthopedic surgeries (De Jonge et al., 2016).

A full understanding of host-pathogen interactions can help find new decolonization strategies. New fields on colonization mechanisms should be investigated. For example, the role of mycobiota starts to be described in the pathophysiology of chronic respiratory diseases (Brégeon and Rolain, 2015). Interaction of nasal *S. aureus* with the nasal fungal communities would be an interesting perspective to develop. On the other hand, some reports suggest that *S. aureus* can regulate host inflammatory gene expression (Modak et al., 2014). Epigenetics mechanisms could be interesting to investigate in order to better understand tolerance mechanisms in *S. aureus* colonization processes.

REFERENCES

- Andersen, P. S., Pedersen, J. K., Fode, P., Skov, R. L., Fowler, V. G., Stegger, M., et al. (2012). Influence of host genetics and environment on nasal carriage of *Staphylococcus aureus* in danish middle-aged and elderly twins. *J. Infect. Dis.* 206, 1178–1184. doi: 10.1093/infdis/jis491
- Anderson, D. J., Kaye, K. S., Chen, L. F., Schmader, K. E., Choi, Y., Sloane, R., et al. (2009). Clinical and financial outcomes due to methicillin resistant *Staphylococcus aureus* surgical site infection: a multi-center matched outcomes study. *PLoS One* 4:e8305. doi: 10.1371/journal.pone.0008305
- Archer, N. K., Adappa, N. D., Palmer, J. N., Cohen, N. A., Harro, J. M., Lee, S. K., et al. (2016). Interleukin-17A (IL-17A) and IL-17F are critical for antimicrobial peptide production and clearance of *Staphylococcus aureus* nasal colonization. *Infect. Immun.* 84, 3575–3583. doi: 10.1128/IAI.00596-16
- Archer, N. K., Harro, J. M., and Shirliff, M. E. (2013). Clearance of *Staphylococcus aureus* nasal carriage is T cell dependent and mediated through interleukin-17A expression and neutrophil influx. *Infect. Immun.* 81, 2070–2075. doi: 10.1128/IAI.00084-13
- Armstrong-Esther, C. A. (1976). Carriage patterns of *Staphylococcus aureus* in a healthy non-hospital population of adults and children. *Ann. Hum. Biol.* 3, 221–227. doi: 10.1080/03014467600001381
- Askarian, F., Ajayi, C., Hanssen, A.-M., van Sorge, N. M., Pettersen, I., Diep, D. B., et al. (2016). The interaction between *Staphylococcus aureus* SdrD and desmoglein 1 is important for adhesion to host cells. *Sci. Rep.* 6:22134. doi: 10.1038/srep22134
- Baur, S., Rautenberg, M., Faulstich, M., Faulstich, M., Grau, T., Severin, Y., et al. (2014). A nasal epithelial receptor for *Staphylococcus aureus* WTA governs adhesion to epithelial cells and modulates nasal colonization. *PLoS Pathog.* 10:e1004089. doi: 10.1371/journal.ppat.1004089
- Bode, L. G. M., Kluytmans, J. A. J. W., Wertheim, H. F. L., Bogaers, D., Vandendriek-Grauls, C. M. J. E., Roosendaal, R., et al. (2010). Preventing surgical-site infections in nasal carriers of *Staphylococcus aureus*. *N. Engl. J. Med.* 362, 9–17. doi: 10.1056/NEJMoa0808939
- Brégeon, F., and Rolain, J.-M. (2015). Le résistome pulmonaire. *Méd. Sci.* 31, 947–950. doi: 10.1051/medsci/20153111003
- Breuer, K., HAussler, S., Kapp, A., and Werfel, T. (2002). *Staphylococcus aureus*: colonizing features and influence of an antibacterial treatment in adults with atopic dermatitis. *Br. J. Dermatol.* 147, 55–61. doi: 10.1046/j.1365-2133.2002.04872.x
- Brown, A. F., Leech, J. M., Rogers, T. R., and McLoughlin, R. M. (2014). *Staphylococcus aureus* colonization: modulation of host immune response and

AUTHOR CONTRIBUTIONS

AS wrote the review paper. FB, J-LM, OB, and J-MR corrected the manuscript. All authors approved and revised the final version of the manuscript.

FUNDING

This study was supported by IHU Méditerranée Infection, Marseille, France and by the French Government under the “Investissements d’Avenir” (Investments for the Future) program managed by the “Agence Nationale de la Recherche” (ANR, fr: National Agency for Research) (reference: Méditerranée Infection 10-IAHU-03). This work was supported by Région Provence Alpes Côte d’Azur and European funding FEDER PRIMI. AS was supported by a CIFRE PhD Grant from the “Association Nationale de la Recherche et de la Technologie”.

- impact on human vaccine design. *Front. Immunol.* 4:507. doi: 10.3389/fimmu.2013.00507
- Burgey, C., Kern, W. V., Römer, W., and Rieg, S. (2016). Differential induction of innate defense antimicrobial peptides in primary nasal epithelial cells upon stimulation with inflammatory cytokines, Th17 cytokines or bacterial conditioned medium from *Staphylococcus aureus* isolates. *Microb. Pathog.* 90, 69–77. doi: 10.1016/j.micpath.2015.11.023
- Burian, M., Wolz, C., and Goerke, C. (2010). Regulatory adaptation of *Staphylococcus aureus* during nasal colonization of humans. *PLoS One* 5:e10040. doi: 10.1371/journal.pone.0010040
- Candi, E., Schmidt, R., and Melino, G. (2005). The cornified envelope: a model of cell death in the skin. *Nat. Rev. Mol. Cell Biol.* 6, 328–340. doi: 10.1038/nrm1619
- Chang, C.-H., Chen, S.-Y., Lu, J.-J., Chang, C.-J., Chang, Y., and Hsieh, P.-H. (2017). Nasal colonization and bacterial contamination of mobile phones carried by medical staff in the operating room. *PLoS One* 12:e0175811. doi: 10.1371/journal.pone.0175811
- Chen, A. F., Heyl, A. E., Xu, P. Z., Rao, N., and Klatt, B. A. (2013). Preoperative decolonization effective at reducing staphylococcal colonization in total joint arthroplasty patients. *J. Arthroplasty* 28, 18–20. doi: 10.1016/j.arth.2013.03.036
- Chen, B., Dai, X., He, B., Pan, K., Li, H., Liu, X., et al. (2015). Differences in *Staphylococcus aureus* nasal carriage and molecular characteristics among community residents and healthcare workers at Sun Yat-sen University, Guangzhou, Southern China. *BMC Infect. Dis.* 15:303. doi: 10.1186/s12879-015-1032-7
- Chen, B. J., Xie, X. Y., Ni, L. J., Dai, X. L., Lu, Y., Wu, X. Q., et al. (2017). Factors associated with *Staphylococcus aureus* nasal carriage and molecular characteristics among the general population at a Medical College Campus in Guangzhou, South China. *Ann. Clin. Microbiol. Antimicrob.* 16:28. doi: 10.1186/s12941-017-0206-0
- Chou, Y.-H., Lee, M.-S., Lin, R.-Y., and Wu, C.-Y. (2015). Risk factors for methicillin-resistant *Staphylococcus aureus* skin and soft-tissue infections in outpatients in Taiwan. *Epidemiol. Infect.* 143, 749–753. doi: 10.1017/S0950268814001642
- Clarke, S. R., Brummell, K. J., Horsburgh, M. J., McDowell, P. W., Mohamad, S. A. S., Stapleton, M. R., et al. (2006). Identification of *in vivo*-expressed antigens of *Staphylococcus aureus* and their use in vaccinations for protection against nasal carriage. *J. Infect. Dis.* 193, 1098–1108. doi: 10.1086/501471
- Clarke, S. R., Wiltshire, M. D., and Foster, S. J. (2004). IsdA of *Staphylococcus aureus* is a broad spectrum, iron-regulated adhesin. *Mol. Microbiol.* 51, 1509–1519. doi: 10.1111/j.1365-2958.2003.03938.x

- Clement, S., Vaudaux, P., Francois, P., Schrenzel, J., Huggler, E., Kampf, S., et al. (2005). Evidence of an intracellular reservoir in the nasal mucosa of patients with recurrent *Staphylococcus aureus* rhinosinusitis. *J. Infect. Dis.* 192, 1023–1028. doi: 10.1086/432735
- Cole, A. L., Muthukrishnan, G., Chong, C., Beavis, A., Eade, C. R., Wood, M. P., et al. (2016). Host innate inflammatory factors and staphylococcal protein A influence the duration of human *Staphylococcus aureus* nasal carriage. *Mucosal Immunol.* 9, 1537–1548. doi: 10.1038/mi.2016.2
- Cole, A. L., Schmidt-Owens, M., Beavis, A. C., Chong, C. F., Tarwater, P. M., Schaus, J., et al. (2018). Cessation from smoking improves innate host defense and clearance of experimentally inoculated nasal *S. aureus*. *Infect. Immun.* 86, e912–e917. doi: 10.1128/IAI.00912-17
- Cole, A. M., Dewan, P., and Ganz, T. (1999). Innate antimicrobial activity of nasal secretions. *Infect. Immun.* 67, 3267–3275.
- Colque-Navarro, P., Jacobsson, G., Andersson, R., Flock, J.-I., and Möllby, R. (2010). Levels of antibody against 11 *Staphylococcus aureus* antigens in a healthy population. *Clin. Vaccine Immunol.* 17, 1117–1123. doi: 10.1128/CI.00506-09
- Corrigan, R. M., Miajlovic, H., Foster, T. J., Kluytmans, J., van Belkum, A., Verbrugh, H., et al. (2009). Surface proteins that promote adherence of *Staphylococcus aureus* to human desquamated nasal epithelial cells. *BMC Microbiol.* 9:22. doi: 10.1186/1471-2180-9-22
- Cosgrove, S. E., Sakoulas, G., Perencevich, E. N., Schwaber, M. J., Karchmer, A. W., and Carmeli, Y. (2003). Comparison of mortality associated with methicillin-resistant and methicillin-susceptible *Staphylococcus aureus* bacteremia: a meta-analysis. *Clin. Infect. Dis.* 36, 53–59. doi: 10.1086/345476
- Cunningham, D. J. (1905). *Cunningham's Text-Book of Anatomy*. New York, NY?: W. Wood and company.
- Dall'Antonia, M., Coen, P. G., Wilks, M., Whiley, A., and Millar, M. (2005). Competition between methicillin-sensitive and -resistant *Staphylococcus aureus* in the anterior nares. *J. Hosp. Infect.* 61, 62–67. doi: 10.1016/j.jhin.2005.01.008
- De Jonge, S., Atema, J. J., Gans, S., Boermeester, M. A., Gomes, S. M., Solomkin, J. S., et al. (2016). Surgical site infections 1 New WHO recommendations on preoperative measures for surgical site infection prevention: an evidence-based global perspective. *Lancet Infect. Dis.* 16, e276–e287. doi: 10.1016/S1473-3099(16)30398-X
- Demos, M., McLeod, M. P., and Nouri, K. (2012). Recurrent furunculosis: a review of the literature. *Br. J. Dermatol.* 167, 725–732. doi: 10.1111/j.1365-2133.2012.11151.x
- den Heijer, C. D., van Bijnen, E. M., Paget, W. J., Pringle, M., Goossens, H., Bruggeman, C. A., et al. (2013). Prevalence and resistance of commensal *Staphylococcus aureus*, including methicillin-resistant *S. aureus*, in nine European countries: a cross-sectional study. *Lancet Infect. Dis.* 13, 409–415. doi: 10.1016/S1473-3099(13)70036-7
- Dunyach-Remy, C., Courtais-Coulon, C., DeMattei, C., Jourdan, N., Schuldiner, S., Sultan, A., et al. (2017). Link between nasal carriage of *Staphylococcus aureus* and infected diabetic foot ulcers. *Diabetes Metab.* 43, 167–171. doi: 10.1016/j.diabet.2016.09.003
- Durupt, F., Mayor, L., Bes, M., Reverdy, M.-E., Vandenesch, F., Thomas, L., et al. (2007). Prevalence of *Staphylococcus aureus* toxins and nasal carriage in furuncles and impetigo. *Br. J. Dermatol.* 157, 1161–1167. doi: 10.1111/j.1365-2133.2007.08197.x
- Eckhart, L., Lippens, S., Tschachler, E., and Declercq, W. (2013). Cell death by cornification. *Biochim. Biophys. Acta Mol. Cell Res.* 1833, 3471–3480. doi: 10.1016/j.bbamcr.2013.06.010
- Elie-Turenne, M.-C., Fernandes, H., Mediavilla, J. R., Rosenthal, M., Mathema, B., Singh, A., et al. (2010). Prevalence and characteristics of *Staphylococcus aureus* colonization among healthcare professionals in an urban teaching hospital. *Infect. Control Hosp. Epidemiol.* 31, 574–580. doi: 10.1086/652525
- Emonts, M., Uitterlinden, A. G., Nouwen, J. L., Kardys, I., Maat, M. P. M., Melles, D. C., et al. (2008). Host polymorphisms in interleukin 4, complement factor H, and C-reactive protein associated with nasal carriage of *Staphylococcus aureus* and occurrence of boils. *J. Infect. Dis.* 197, 1244–1253. doi: 10.1086/533501
- Ena, J., Boelaert, J. R., Boyken, L. D., Van Landuyt, H. W., Godard, C. A., and Herwaldt, L. A. (1994). Epidemiology of *Staphylococcus aureus* infections in patients on hemodialysis. *Infect. Control Hosp. Epidemiol.* 15, 78–81. doi: 10.2307/30145535
- Eriksen, N. H., Espersen, F., Rosdahl, V. T., and Jensen, K. (1995). Carriage of *Staphylococcus aureus* among 104 healthy persons during a 19-month period. *Epidemiol. Infect.* 115, 51–60. doi: 10.1017/S0950268800058118
- Frank, D. N., Feazel, L. M., Besesen, M. T., Price, C. S., Janoff, E. N., and Pace, N. R. (2010). The human nasal microbiota and *Staphylococcus aureus* carriage. *PLoS One* 5:e10598. doi: 10.1371/journal.pone.0010598
- Garrouste-Orgeas, M., Timsit, J. F., Kallel, H., Ben Ali, A., Dumay, M. F., Paoli, B., et al. (2001). Colonization with methicillin-resistant *Staphylococcus aureus* in ICU patients: morbidity, mortality, and glycopeptide use. *Infect. Control Hosp. Epidemiol.* 22, 687–692. doi: 10.1086/501846
- Ghasemzadeh-Moghaddam, H., Neela, V., van Wamel, W., Hamat, R. A., Shamsudin, M. N., Hussin, N. S. C., et al. (2015). Nasal carriers are more likely to acquire exogenous *Staphylococcus aureus* strains than non-carriers. *Clin. Microbiol. Infect.* 21, 998.e1–998.e7. doi: 10.1016/j.cmi.2015.07.006
- González-Zorn, B., Senna, J. P. M., Fiette, L., Shorte, S., Testard, A., Chignard, M., et al. (2005). Bacterial and host factors implicated in nasal carriage of methicillin-resistant *Staphylococcus aureus* in mice. *Infect. Immun.* 73, 1847–1851. doi: 10.1128/IAI.73.3.1847-1851.2005
- Gorwitz, R. J., Kruszon-Moran, D., McAllister, S. K., McQuillan, G., McDougal, L. K., Fosheim, G. E., et al. (2008). Changes in the prevalence of nasal colonization with *Staphylococcus aureus* in the united states, 2001–2004. *J. Infect. Dis.* 197, 1226–1234. doi: 10.1086/533494
- Hail, C., Fletcher, S., Archer, R., Jones, G., Jayarajah, M., Frame, J., et al. (2013). Prolonged outbreak of methicillin-resistant *Staphylococcus aureus* in a cardiac surgery unit linked to a single colonized healthcare worker. *J. Hosp. Infect.* 83, 219–225. doi: 10.1016/j.jhin.2012.11.019
- Haim, M., Trost, A., Maier, C. J., Achatz, G., Feichtner, S., Hintner, H., et al. (2010). Cytokeratin 8 interacts with clumping factor B: a new possible virulence factor target. *Microbiology* 156, 3710–3721. doi: 10.1099/mic.0.034413-0
- Hanssen, A.-M., Kindlund, B., Stenklev, N. C., Furberg, A.-S., Fismen, S., Olsen, R. S., et al. (2017). Localization of *Staphylococcus aureus* in tissue from the nasal vestibule in healthy carriers. *BMC Microbiol.* 17:89. doi: 10.1186/s12866-017-0997-3
- Harbarth, S., Liassine, N., Dharan, S., Herrault, P., Auckenthaler, R., and Pittet, D. (2000). Risk factors for persistent carriage of methicillin-resistant *Staphylococcus aureus*. *Clin. Infect. Dis.* 31, 1380–1385. doi: 10.1086/317484
- Hayes, S. M., Howlin, R., Johnston, D. A., Webb, J. S., Clarke, S. C., Stoodley, P., et al. (2015). Intracellular residency of *Staphylococcus aureus* within mast cells in nasal polyps: a novel observation. *J. Allergy Clin. Immunol.* 135, 1648–1651.e5. doi: 10.1016/j.jaci.2014.12.1929
- Holtfreter, S., Roschack, K., Eichler, P., Eske, K., Holtfreter, B., Kohler, C., et al. (2006). *Staphylococcus aureus* carriers neutralize superantigens by antibodies specific for their colonizing strain: a potential explanation for their improved prognosis in severe sepsis. *J. Infect. Dis.* 193, 1275–1278. doi: 10.1086/503048
- Honda, H., Krauss, M. J., Coopersmith, C. M., Kollef, M. H., Richmond, A. M., Fraser, V. J., et al. (2010). *Staphylococcus aureus* nasal colonization and subsequent infection in intensive care unit patients: does methicillin resistance matter? *Infect. Control Hosp. Epidemiol.* 31, 584–591. doi: 10.1086/652530
- Hraiech, S., Roch, A., Lepidi, H., Atieh, T., Audoly, G., Rolain, J.-M., et al. (2013). Impaired virulence and fitness of a colistin-resistant clinical isolate of *Acinetobacter baumannii* in a rat model of pneumonia. *Antimicrob. Agents Chemother.* 57, 5120–5121. doi: 10.1128/AAC.00700-13
- Human Microbiome Project Consortium (2012). Structure, function and diversity of the healthy human microbiome. *Nature* 486, 207–214. doi: 10.1038/nature11234
- Humphreys, H. (2012). *Staphylococcus aureus*: the enduring pathogen in surgery. *Surgeon* 10, 357–360. doi: 10.1016/j.surge.2012.05.003
- Immergluck, L., Jain, S., Ray, S., Mayberry, R., Satola, S., Parker, T., et al. (2017). Risk of skin and soft tissue infections among children found to be *Staphylococcus aureus* MRSA USA300 carriers. *West. J. Emerg. Med.* 18, 201–212. doi: 10.5811/westjem.2016.10.30483
- Iwase, T., Uehara, Y., Shinji, H., Tajima, A., Seo, H., Takada, K., et al. (2010). *Staphylococcus epidermidis* Esp inhibits *Staphylococcus aureus* biofilm formation and nasal colonization. *Nature* 465, 346–349. doi: 10.1038/nature09074
- Kalmeijer, M. D., van Nieuwland-Bollen, E., Bogaers-Hofman, D., and de Baere, G. A. (2000). Nasal carriage of *Staphylococcus aureus* is a major risk factor for

- surgical-site infections in orthopedic surgery. *Infect. Control Hosp. Epidemiol.* 21, 319–323. doi: 10.1086/501763
- Kaspar, U., Kriegeskorte, A., Schubert, T., Peters, G., Rudack, C., Pieper, D. H., et al. (2016). The culturome of the human nose habitats reveals individual bacterial fingerprint patterns. *Environ. Microbiol.* 18, 2130–2142. doi: 10.1111/1462-2920.12891
- Katneni, R., and Hedayati, S. S. (2007). Central venous catheter-related bacteremia in chronic hemodialysis patients: epidemiology and evidence-based management. *Nat. Clin. Pract. Nephrol.* 3, 256–266. doi: 10.1038/ncpneph0447
- Kinsman, O. S., Mckenna, R., and Noble, W. C. (1983). Association between histocompatibility antigens (HLA) and nasal carriage of *Staphylococcus aureus*. *J. Med. Microbiol.* 16, 215–220. doi: 10.1099/00222615-16-2-215
- Klevens, R. M., Edwards, J. R., Richards, C. L., Horan, T. C., Gaynes, R. P., Pollock, D. A., et al. (2007). Estimating health care-associated infections and deaths in U.S. hospitals, 2002. *Public Health Rep.* 122, 160–166.
- Kluytmans, J., van Belkum, A., and Verbrugh, H. (1997). Nasal carriage of *Staphylococcus aureus*: epidemiology, underlying mechanisms, and associated risks. *Clin. Microbiol. Rev.* 10, 505–520.
- Kluytmans, J. A., Mouton, J. W., Ijzerman, E. P., Vandenbroucke-Grauls, C. M., Maat, A. W., Wagenvoort, J. H., et al. (1995). Nasal carriage of *Staphylococcus aureus* as a major risk factor for wound infections after cardiac surgery. *J. Infect. Dis.* 171, 216–219. doi: 10.1093/infdis/171.1.216
- Kluytmans, J. A. J. W., Manders, M.-J., van Bommel, E., and Verbrugh, H. (1996). Elimination of nasal carriage of *Staphylococcus aureus* in hemodialysis patients. *Infect. Control Hosp. Epidemiol.* 17, 793–797. doi: 10.2307/30141172
- Kotpal, R. S., K. P., Bhalla, P., Dewan, R., and Kaur, R. (2016). Incidence and risk factors of nasal carriage of *Staphylococcus aureus* in HIV-infected individuals in comparison to HIV-uninfected individuals: a case-control study. *J. Int. Assoc. Provid. AIDS Care* 15, 141–147. doi: 10.1177/2325957414554005
- Krismer, B., Weidenmaier, C., Zipperer, A., and Peschel, A. (2017). The commensal lifestyle of *Staphylococcus aureus* and its interactions with the nasal microbiota. *Nat. Rev. Microbiol.* 15, 675–687. doi: 10.1038/nrmicro.2017.104
- Lamanna, O., Bongiorno, D., Bertoncillo, L., Grandesso, S., Mazzucato, S., Pozzan, G. B., et al. (2017). Rapid containment of nosocomial transmission of a rare community-acquired methicillin-resistant *Staphylococcus aureus* (CA-MRSA) clone, responsible for the Staphylococcal Scalded Skin Syndrome (SSSS). *Ital. J. Pediatr.* 43:5. doi: 10.1186/s13052-016-0323-y
- Lamarsalle, L., Hunt, B., Schauf, M., Swarczensztejn, K., and Valentine, W. J. (2013). Evaluating the clinical and economic burden of healthcare-associated infections during hospitalization for surgery in France. *Epidemiol. Infect.* 141, 2473–2482. doi: 10.1017/S0950268813000253
- Laudien, M., Gadola, S. D., Podschun, R., Hedderich, J., Paulsen, J., Reinhold-Keller, E., et al. (2010). Nasal carriage of *Staphylococcus aureus* and endonasal activity in Wegener's granulomatosis as compared to rheumatoid arthritis and chronic Rhinosinusitis with nasal polyps. *Clin. Exp. Rheumatol.* 28, 51–55.
- Le Réseau d'alerte, d'investigation et de surveillance des infections nosocomiales (Raisin) (2015). *Surveillance des Infections du Site Opératoire, France 2013*. France: Institut de veille sanitaire.
- Leaper, D. J., van Goor, H., Reilly, J., Petrosillo, N., Geiss, H. K., Torres, A. J., et al. (2004). Surgical site infection - a European perspective of incidence and economic burden. *Int. Wound J.* 1, 247–273. doi: 10.1111/j.1742-4801.2004.00067.x
- Lehar, S. M., Pillow, T., Xu, M., Staben, L., Kajihara, K. K., Vandlen, R., et al. (2015). Novel antibody-antibiotic conjugate eliminates intracellular *S. aureus*. *Nature* 527, 323–328. doi: 10.1038/nature16057
- Lepelletier, D., Perron, S., Bizouarn, P., Cailion, J., Drugeon, H., Michaud, J.-L., et al. (2005). Surgical-site infection after cardiac surgery: incidence, microbiology, and risk factors. *Infect. Control Hosp. Epidemiol.* 26, 466–472. doi: 10.1086/502569
- Leshem, E., Maayan-Metzger, A., Rahav, G., Dolitzki, M., Kuint, J., Roytman, Y., et al. (2012). Transmission of *Staphylococcus aureus* from mothers to newborns. *Pediatr. Infect. Dis. J.* 31, 360–363. doi: 10.1097/INF.0b013e318244020e
- Lina, G., Boutite, F., Tristan, A., Bes, M., Etienne, J., and Vandenesch, F. (2003). Bacterial competition for human nasal cavity colonization: role of Staphylococcal *agr* alleles. *Appl. Environ. Microbiol.* 69, 18–23. doi: 10.1128/aem.69.1.18-23.2003
- Liu, C. M., Price, L. B., Hungate, B. A., Abraham, A. G., Larsen, L. A., Christensen, K., et al. (2015). *Staphylococcus aureus* and the ecology of the nasal microbiome. *Sci. Adv.* 1:e1400216. doi: 10.1126/sciadv.1400216
- Liu, Q., Du, X., Hong, X., Li, T., Zheng, B., He, L., et al. (2015). Targeting surface protein SasX by active and passive vaccination to reduce *Staphylococcus aureus* colonization and infection. *Infect. Immun.* 83, 2168–2174. doi: 10.1128/IAI.02951-14
- Luzar, M. A., Coles, G. A., Faller, B., Slingeneyer, A., Dah, G. D., Briat, C., et al. (1990). *Staphylococcus aureus* nasal carriage and infection in patients on continuous ambulatory peritoneal dialysis. *N. Engl. J. Med.* 322, 505–509. doi: 10.1056/NEJM19900223220804
- Maayan-Metzger, A., Strauss, T., Rubin, C., Jaber, H., Dulitzky, M., Reiss-Mandel, A., et al. (2017). Clinical evaluation of early acquisition of *Staphylococcus aureus* carriage by newborns. *Int. J. Infect. Dis.* 64, 9–14. doi: 10.1016/j.ijid.2017.08.013
- Magill, S. S., Hellinger, W., Cohen, J., Kay, R., Bailey, C., Boland, B., et al. (2012). Prevalence of healthcare-associated infections in acute care hospitals in Jacksonville, Florida. *Infect. Control Hosp. Epidemiol.* 33, 283–291. doi: 10.1086/664048
- McKinnell, J. A., Huang, S. S., Eells, S. J., Cui, E., and Miller, L. G. (2013). Quantifying the impact of extranasal testing of body sites for methicillin-resistant *Staphylococcus aureus* colonization at the time of hospital or intensive care unit admission. *Infect. Control Hosp. Epidemiol.* 34, 161–170. doi: 10.1086/669095
- Miller, L. G., Eells, S. J., Taylor, A. R., David, M. Z., Ortiz, N., Zychowski, D., et al. (2012). *Staphylococcus aureus* colonization among household contacts of patients with skin infections: risk factors, strain discordance, and complex ecology. *Clin. Infect. Dis.* 54, 1523–1535. doi: 10.1093/cid/cis213
- Modak, R., Das Mitra, S., Vasudevan, M., Krishnamoorthy, P., Kumar, M., and Bhat, A. V. et al. (2014). Epigenetic response in mice mastitis: role of histone H3 acetylation and microRNA(s) in the regulation of host inflammatory gene expression during *Staphylococcus aureus* infection. *Clin. Epigenetics* 6:12. doi: 10.1186/1868-7083-6-12
- Mulcahy, M. E., Geoghegan, J. A., Monk, I. R., O'Keefe, K. M., Walsh, E. J., Foster, T. J., et al. (2012). Nasal colonisation by *Staphylococcus aureus* depends upon clumping factor B binding to the squamous epithelial cell envelope protein loricrin. *PLoS Pathog.* 8:e1003092. doi: 10.1371/journal.ppat.1003092
- Mulcahy, M. E., Leech, J. M., Renaud, J.-C., Mills, K. H., and McLoughlin, R. M. (2016). Interleukin-22 regulates antimicrobial peptide expression and keratinocyte differentiation to control *Staphylococcus aureus* colonization of the nasal mucosa. *Mucosal Immunol.* 9, 1429–1441. doi: 10.1038/mi.2016.24
- Mulcahy, M. E., and McLoughlin, R. M. (2016). Host-bacterial crosstalk determines *Staphylococcus aureus* nasal colonization. *Trends Microbiol.* 24, 872–886. doi: 10.1016/j.tim.2016.06.012
- Muñoz, P., Hortal, J., Giannella, M., Barrio, J. M., Rodríguez-Créixems, M., Pérez, M. J., et al. (2008). Nasal carriage of *S. aureus* increases the risk of surgical site infection after major heart surgery. *J. Hosp. Infect.* 68, 25–31. doi: 10.1016/j.jhin.2007.08.010
- Muthukrishnan, G., Lamers, R. P., Ellis, A., Paramanandam, V., Persaud, A. B., Tafur, S., et al. (2013). Longitudinal genetic analyses of *Staphylococcus aureus* nasal carriage dynamics in a diverse population. *BMC Infect. Dis.* 13:221. doi: 10.1186/1471-2334-13-221
- Nardi, G., Di Silvestre, A. D., De Monte, A., Massarutti, D., Proietti, A., Grazia Troncon, M., et al. (2001). Reduction in gram-positive pneumonia and antibiotic consumption following the use of a SDD protocol including nasal and oral mupirocin. *Eur. J. Emerg. Med.* 8, 203–214. doi: 10.1097/00063110-200109000-00008
- Negi, V., Pal, S., Juyal, D., Sharma, M. K., and Sharma, N. (2015). Bacteriological profile of surgical site infections and their antibiogram: a study from resource constrained rural setting of uttarakhand state, India. *J. Clin. Diagn. Res.* 9, DC17–DC20. doi: 10.7860/JCDR/2015/15342.6698
- Nguyen, M. H., Kauffman, C. A., Goodman, R. P., Squier, C., Arbeit, R. D., Singh, N., et al. (1999). Nasal carriage of and infection with *Staphylococcus aureus* in HIV-infected patients. *Ann. Intern. Med.* 130, 221–225. doi: 10.7326/0003-4819-130-3-199902020-00026
- NICE Clinical Guidelines (2008). *National Collaborating Centre for Women's and Children's Health (UK). Preoperative phase, Surgical Site Infection: Prevention*

- and Treatment of Surgical Site Infection. London: National Institute for Health and Clinical Excellence.
- Nilsson, A. C., Janson, H., Wold, H., Fugelli, A., Andersson, K., Håkangård, C., et al. (2015). LTX-109 is a novel agent for nasal decolonization of methicillin-resistant and -sensitive *Staphylococcus aureus*. *Antimicrob. Agents Chemother.* 59, 145–151. doi: 10.1128/AAC.03513-14
- Nouwen, J., Schouten, J., Schneebergen, P., Snijders, S., Maaskant, J., Koolen, M., et al. (2006). *Staphylococcus aureus* carriage patterns and the risk of infections associated with continuous peritoneal dialysis. *J. Clin. Microbiol.* 44, 2233–2236. doi: 10.1128/JCM.02083-05
- Nouwen, J. L., Fieren, M. W. J. A., Snijders, S., Verbrugh, H. A., and Van Belkum, A. (2005). Persistent (not intermittent) nasal carriage of *Staphylococcus aureus* is the determinant of CPD-related infections. *Kidney Int.* 67, 1084–1092. doi: 10.1111/j.1523-1755.2005.00174.x
- Nouwen, J. L., and Optima Grafische Communicatie (2004). *Determinants, Risks & Dynamics of Staphylococcus aureus Nasal Carriage*. Doctoral thesis, Erasmus University Rotterdam, Rotterdam.
- Nouwen, J. L., Ott, A., Kluytmans-Vandenbergh, M. F. Q., Boelens, H. A. M., Hofman, A., van Belkum, A., et al. (2004). Predicting the *Staphylococcus aureus* nasal carrier state: derivation and validation of a “culture rule”. *Clin. Infect. Dis.* 39, 806–811. doi: 10.1086/423376
- Nurjadi, D., Heeg, K., Weber, A. N. R., and Zanger, P. (2018). Toll-like receptor (TLR) -9 promotor polymorphisms and gene expression are associated with persistent *Staphylococcus aureus* nasal carriage. *Clin. Microbiol. Infect.* doi: 10.1016/j.cmi.2018.02.014 [Epub ahead of print].
- Nurjadi, D., Kain, M., Marcinek, P., Gaile, M., Heeg, K., and Zanger, P. (2016). Ratio of T-helper type 1 (Th1) to Th17 cytokines in whole blood is associated with human β -defensin 3 expression in skin and persistent *Staphylococcus aureus* nasal carriage. *J. Infect. Dis.* 214, 1744–1751. doi: 10.1093/infdis/jiw440
- O'Brien, L. M., Walsh, E. J., Massey, R. C., Peacock, S. J., and Foster, T. J. (2002). *Staphylococcus aureus* clumping factor B (ClfB) promotes adherence to human type I cyokeratin 10: implications for nasal colonization. *Cell. Microbiol.* 4, 759–770. doi: 10.1046/j.1462-5822.2002.00231.x
- Olsen, K., Danielsen, K., Wilsgaard, T., Sangvik, M., Sollid, J. U. E., Thune, I., et al. (2013). Obesity and *Staphylococcus aureus* nasal colonization among women and men in a general population. *PLoS One* 8:e63716. doi: 10.1371/journal.pone.0063716
- Olsen, K., Falch, B. M., Danielsen, K., Johannessen, M., Ericson Sollid, J. U., Thune, I., et al. (2012). *Staphylococcus aureus* nasal carriage is associated with serum 25-hydroxyvitamin D levels, gender and smoking status. The Tromsø Staph and Skin Study. *Eur. J. Clin. Microbiol. Infect. Dis.* 31, 465–473. doi: 10.1007/s10096-011-1331-x
- Ong, L. M., Ch'ng, C. C., Wee, H. C., Supramaniam, P., Zainal, H., Goh, B. L., et al. (2017). Risk of peritoneal dialysis-related peritonitis in a multi-racial asian population. *Perit. Dial. Int.* 37, 35–43. doi: 10.3747/pdi.2015.00141
- Ou, J., Bassiouni, A., Drilling, A., Psaltis, A. J., Vreugde, S., and Wormald, P. J. (2017). The persistence of intracellular *Staphylococcus aureus* in the sinuses: a longitudinal study. *Rhinol. J.* 55, 305–311. doi: 10.4193/Rhin16.218
- Ou, J., Drilling, A., Singhal, D., Tan, N. C.-W., Wallis-Hill, D., Vreugde, S., et al. (2016). Association of intracellular *Staphylococcus aureus* with prognosis in chronic rhinosinusitis. *Int. Forum Allergy Rhinol.* 6, 792–799. doi: 10.1002/alar.21758
- Panierakis, C., Goulielmos, G., Mamoulakis, D., Maraki, S., Papavasiliou, E., and Galanakis, E. (2009). *Staphylococcus aureus* nasal carriage might be associated with vitamin D receptor polymorphisms in type 1 diabetes. *Int. J. Infect. Dis.* 13, e437–e443. doi: 10.1016/j.ijid.2009.02.012
- Peacock, S. J., de Silva, I., and Lowy, F. D. (2001). What determines nasal carriage of *Staphylococcus aureus*? *Trends Microbiol.* 9, 605–610.
- Peacock, S. J., Justice, A., Griffiths, D., de Silva, G. D. I., Kantzanou, M. N., Crook, D., et al. (2003). Determinants of acquisition and carriage of *Staphylococcus aureus* in infancy. *J. Clin. Microbiol.* 41, 5718–5725. doi: 10.1128/JCM.41.12.5718-5725.2003
- Perkins, S., Walsh, E. J., Deivanayagam, C. C., Narayana, S. V., Foster, T. J., and Höök, M. (2001). Structural organization of the fibrinogen-binding region of the clumping factor B MSCRAMM of *Staphylococcus aureus*. *J. Biol. Chem.* 276, 44721–44728. doi: 10.1074/jbc.M106741200
- Perl, T. M., Cullen, J. J., Wenzel, R. P., Zimmerman, M. B., Pfäller, M. A., Sheppard, D., et al. (2002). Intranasal mupirocin to prevent postoperative *Staphylococcus aureus* infections. *N. Engl. J. Med.* 346, 1871–1877. doi: 10.1056/NEJMoa003069
- Perl, T. M., and Golub, J. E. (1998). New approaches to reduce *Staphylococcus aureus* nosocomial infection rates: treating *S. aureus* nasal carriage. *Ann. Pharmacother.* 32, S7–S16.
- Peschel, A., and Sahl, H.-G. (2006). The co-evolution of host cationic antimicrobial peptides and microbial resistance. *Nat. Rev. Microbiol.* 4, 529–536. doi: 10.1038/nrmicro1441
- Plouin-Gaudon, I., Clement, S., Huggler, E., Chaponnier, C., François, P., Lew, D., et al. (2006). Intracellular residency is frequently associated with recurrent *Staphylococcus aureus* rhinosinusitis. *Rhinology* 44, 249–254.
- Pournajaf, A., Ardebili, A., Goudarzi, L., Khodabandeh, M., Narimani, T., and Abbaszadeh, H. (2014). PCR-based identification of methicillin-resistant *Staphylococcus aureus* strains and their antibiotic resistance profiles. *Asian Pac. J. Trop. Biomed.* 4, S293–S297. doi: 10.12980/APJT.4.2014C423
- Price, J. R., Cole, K., Bexley, A., Kostiou, V., Eyre, D. W., Golubchik, T., et al. (2017). Transmission of *Staphylococcus aureus* between health-care workers, the environment, and patients in an intensive care unit: a longitudinal cohort study based on whole-genome sequencing. *Lancet Infect. Dis.* 17, 207–214. doi: 10.1016/S1473-3099(16)30413-3
- Pynnonen, M., Stephenson, R. E., Schwartz, K., Hernandez, M., and Boles, B. R. (2011). Hemoglobin promotes *Staphylococcus aureus* nasal colonization. *PLoS Pathog.* 7:e1002104. doi: 10.1371/journal.ppat.1002104
- Quinn, G. A., and Cole, A. M. (2007). Suppression of innate immunity by a nasal carriage strain of *Staphylococcus aureus* increases its colonization on nasal epithelium. *Immunology* 122, 80–89. doi: 10.1111/j.1365-2567.2007.02615.x
- Raviglione, M. C., Mariuz, P., Pablos-Mendez, A., Battan, R., Ottuso, P., and Taranta, A. (1990). High *Staphylococcus aureus* nasal carriage rate in patients with acquired immunodeficiency syndrome or AIDS-related complex. *Am. J. Infect. Control* 18, 64–69. doi: 10.1016/0196-6553(90)90083-5
- Reagan, D. R., Doebbeling, B. N., Pfäller, M. A., Sheetz, C. T., Houston, A. K., Hollis, R. J., et al. (1991). Elimination of coincident *Staphylococcus aureus* nasal and hand carriage with intranasal application of mupirocin calcium ointment. *Ann. Intern. Med.* 114, 101–106. doi: 10.7326/0003-4819-114-2-101
- Redding, R. M., Luke-Marshall, N. R., Hakansson, A. P., and Campagnari, A. A. (2016). Host physiologic changes induced by influenza a virus lead to *Staphylococcus aureus* biofilm dispersion and transition from asymptomatic colonization to invasive disease. *mBio* 7:e01235-16. doi: 10.1128/mBio.01235-16
- Regev-Yochay, G., Trzcinski, K., Thompson, C. M., Malley, R., and Lipsitch, M. (2006). Interference between *Streptococcus pneumoniae* and *Staphylococcus aureus*: *in vitro* hydrogen peroxide-mediated killing by *Streptococcus pneumoniae*. *J. Bacteriol.* 188, 4996–5001. doi: 10.1128/JB.00317-06
- Rigaill, J., Morgene, M. F., Gavid, M., Lelonge, Y., He, Z., Carricajo, A., et al. (2018). Intracellular activity of antimicrobial compounds used for *Staphylococcus aureus* nasal decolonization. *J. Antimicrob. Chemother.* doi: 10.1093/jac/dky318 [Epub ahead of print].
- Roche, F. M., Meehan, M., and Foster, T. J. (2003). The *Staphylococcus aureus* surface protein SasG and its homologues promote bacterial adherence to human desquamated nasal epithelial cells. *Microbiology* 149, 2759–2767. doi: 10.1099/mic.0.26412-0
- Roghmam, M.-C., Johnson, J. K., Stine, O. C., Lydecker, A. D., Ryan, K. A., Mitchell, B. D., et al. (2011). Persistent *Staphylococcus aureus* colonization is not a strongly heritable trait in Amish families. *PLoS One* 6:e17368. doi: 10.1371/journal.pone.0017368
- Ruimy, R., Angebault, C., Djossou, F., Dupont, C., Epelboin, L., Jarraud, S., et al. (2010). Are host genetics the predominant determinant of persistent nasal *Staphylococcus aureus* carriage in humans? *J. Infect. Dis.* 202, 924–934. doi: 10.1086/655901
- Saadatian-Elahi, M., Teyssou, R., and Vanhems, P. (2008). *Staphylococcus aureus*, the major pathogen in orthopaedic and cardiac surgical site infections: a literature review. *Int. J. Surg.* 6, 238–245. doi: 10.1016/j.ijsu.2007.05.001
- Saadatian-Elahi, M., Tristan, A., Laurent, F., Rasigade, J.-P., Bouchiat, C., Ranc, A.-G., et al. (2013). Basic rules of hygiene protect health care and lab workers from nasal colonization by *Staphylococcus aureus*: an international cross-sectional study. *PLoS One* 8:e82851. doi: 10.1371/journal.pone.0082851

- Safdar, N., and Bradley, E. A. (2008). The risk of infection after nasal colonization with *Staphylococcus aureus*. *Am. J. Med.* 121, 310–315. doi: 10.1016/j.amjmed.2007.07.034
- Savage, J. W., and Anderson, P. A. (2013). An update on modifiable factors to reduce the risk of surgical site infections. *Spine J.* 13, 1017–1029. doi: 10.1016/j.spinee.2013.03.051
- Schaffer, A. C., Solinga, R. M., Cocchiari, J., Portoles, M., Kiser, K. B., Risley, A., et al. (2006). Immunization with *Staphylococcus aureus* clumping factor B, a major determinant in nasal carriage, reduces nasal colonization in a murine model. *Infect. Immun.* 74, 2145–2153. doi: 10.1128/IAI.74.4.2145-2153.2006
- Schmidt, A., Bénard, S., and Cyr, S. (2015). Hospital cost of staphylococcal infection after cardiothoracic or orthopedic operations in France: a retrospective database analysis. *Surg. Infect.* 16, 428–435. doi: 10.1089/sur.2014.045
- Selva, L., Viana, D., Regev-Yochay, G., Trzcinski, K., Corpa, J. M., Lasa, I., et al. (2009). Killing niche competitors by remote-control bacteriophage induction. *Proc. Natl. Acad. Sci. U.S.A.* 106, 1234–1238. doi: 10.1073/pnas.0809600106
- Sherertz, R. J., Reagan, D. R., Hampton, K. D., Robertson, K. L., Streed, S. A., Hoen, H. M., et al. (1996). A cloud adult: the *Staphylococcus aureus*-virus interaction revisited. *Ann. Intern. Med.* 124, 539–547. doi: 10.7326/0003-4819-124-6-199603150-00001
- Sissolok, D., Geusau, A., Heinze, G., Witte, W., and Rotter, M. L. (2002). Risk factors for nasal carriage of *Staphylococcus aureus* in infectious disease patients, including patients infected with HIV, and molecular typing of colonizing strains. *Eur. J. Clin. Microbiol. Infect. Dis.* 21, 88–96. doi: 10.1007/s10096-001-0666-0
- Sivaraman, K., Venkataraman, N., and Cole, A. M. (2009). *Staphylococcus aureus* nasal carriage and its contributing factors. *Future Microbiol.* 4, 999–1008. doi: 10.2217/fmb.09.79
- Steinert, P. M., and Marekov, L. N. (1995). The proteins elafin, filaggrin, keratin intermediate filaments, lorricin, and small proline-rich proteins 1 and 2 are isodipeptide cross-linked components of the human epidermal cornified cell envelope. *J. Biol. Chem.* 270, 17702–17711. doi: 10.1074/jbc.270.30.17702
- Sugimoto, S., Iwamoto, T., Takada, K., Okuda, K.-I., Tajima, A., Iwase, T., et al. (2013). *Staphylococcus epidermidis* Esp degrades specific proteins associated with *Staphylococcus aureus* biofilm formation and host-pathogen interaction. *J. Bacteriol.* 195, 1645–1655. doi: 10.1128/JB.01672-12
- Tan, N. C.-W., Cooksley, C. M., Roscioli, E., Drilling, A. J., Douglas, R., Wormald, P.-J., et al. (2014). Small-colony variants and phenotype switching of intracellular *Staphylococcus aureus* in chronic rhinosinusitis. *Allergy* 69, 1364–1371. doi: 10.1111/all.12457
- Trautman, M., Stecher, J., Luz, K., Hemmer, W., Hupp, T., and Grütznert, P. A. (2007). The use of mupirocin in the eradication of *Staphylococcus aureus*. *Dtsch. Arztebl.* 104:A-3259.
- Uehara, Y., Nakama, H., Agematsu, K., Uchida, M., Kawakami, Y., Abdul Fattah, A. S. M., et al. (2000). Bacterial interference among nasal inhabitants: eradication of *Staphylococcus aureus* from nasal cavities by artificial implantation of *Corynebacterium* sp. *J. Hosp. Infect.* 44, 127–133. doi: 10.1053/jhin.1999.0680
- van Belkum, A., Verkaik, N. J., de Vogel, C. P., Boelens, H. A., Verveer, J., Nouwen, J. L., et al. (2009). Reclassification of *Staphylococcus aureus* nasal carriage types. *J. Infect. Dis.* 199, 1820–1826. doi: 10.1086/599119
- van den Akker, E. L. T., Nouwen, J. L., Melles, D. C., van Rossum, E. F. C., Koper, J. W., Uitterlinden, A. G., et al. (2006). *Staphylococcus aureus* nasal carriage is associated with glucocorticoid receptor gene polymorphisms. *J. Infect. Dis.* 194, 814–818. doi: 10.1086/506367
- VandenBergh, M. F., Yzerman, E. P., van Belkum, A., Boelens, H. A., Sijmons, M., and Verbrugh, H. A. (1999). Follow-up of *Staphylococcus aureus* nasal carriage after 8 years: redefining the persistent carrier state. *J. Clin. Microbiol.* 37, 3133–3140.
- Verhoeven, P. O., Grattard, F., Carricajo, A., Lucht, F., Cazorla, C., Garraud, O., et al. (2012). An algorithm based on one or two nasal samples is accurate to identify persistent nasal carriers of *Staphylococcus aureus*. *Clin. Microbiol. Infect.* 18, 551–557. doi: 10.1111/j.1469-0691.2011.03611.x
- Von Eiff, C., Becker, K., Machka, K., Stammer, H., and Peters, G. (2001). Nasal carriage as a source of *Staphylococcus aureus* bacteremia. Study group. *N. Engl. J. Med.* 344, 11–16. doi: 10.1056/NEJM200101043440102
- Vonberg, R.-P., Stamm-Balderjahn, S., Hansen, S., Zuschneid, I., Rüden, H., Behnke, M., et al. (2006). How often do asymptomatic healthcare workers cause methicillin-resistant *Staphylococcus aureus* outbreaks? A systematic evaluation. *Infect. Control Hosp. Epidemiol.* 27, 1123–1127. doi: 10.1086/507922
- Vuyk, H. D., Watts, S. J. (2006). “Nasal reconstruction,” in *Facial Plastic and Reconstructive Surgery*, H. D. Vuyk and P. J. F. M. Lohuis (London: Hodder Arnold), 455–480.
- Walsh, T. L., Querry, A. M., McCool, S., Galdys, A. L., Shutt, K. A., Saul, M. I., et al. (2017). Risk factors for surgical site infections following neurosurgical spinal fusion operations: a case control study. *Infect. Control Hosp. Epidemiol.* 38, 340–347. doi: 10.1017/ice.2016.307
- Wang, J.-T., Chang, S.-C., Ko, W.-J., Chang, Y.-Y., Chen, M.-L., Pan, H.-J., et al. (2001). A hospital-acquired outbreak of methicillin-resistant *Staphylococcus aureus* infection initiated by a surgeon carrier. *J. Hosp. Infect.* 47, 104–109. doi: 10.1053/jhin.2000.0878
- Warnke, P., Devidé, A., Weise, M., Frickmann, H., Schwarz, N. G., Schäffler, H., et al. (2016). Utilizing moist or dry swabs for the sampling of nasal MRSA carriers? An *in vivo* and *in vitro* study. *PLoS One* 11:e0163073. doi: 10.1371/journal.pone.0163073
- Weidenmaier, C., Goerke, C., and Wolz, C. (2012). *Staphylococcus aureus* determinants for nasal colonization. *Trends Microbiol.* 20, 243–250. doi: 10.1016/j.tim.2012.03.004
- Weidenmaier, C., Kokai-Kun, J. F., Kristian, S. A., Chanturiya, T., Kalbacher, H., Gross, M., et al. (2004). Role of teichoic acids in *Staphylococcus aureus* nasal colonization, a major risk factor in nosocomial infections. *Nat. Med.* 10, 243–245. doi: 10.1038/nm991
- Weidenmaier, C., Kokai-Kun, J. F., Kulauzovic, E., Kohler, T., Thumm, G., Stoll, H., et al. (2008). Differential roles of sortase-anchored surface proteins and wall teichoic acid in *Staphylococcus aureus* nasal colonization. *Int. J. Med. Microbiol.* 298, 505–513. doi: 10.1016/j.ijmm.2007.11.006
- Weinke, T., Schiller, R., Fehrenbach, F. J., and Pohle, H. D. (1992). Association between *Staphylococcus aureus* nasopharyngeal colonization and septicemia in patients infected with the human immunodeficiency virus. *Eur. J. Clin. Microbiol. Infect. Dis.* 11, 985–989. doi: 10.1007/BF01967787
- Weiser, M. C., and Moucha, C. S. (2015). The current state of screening and decolonization for the prevention of *Staphylococcus aureus* surgical site infection after total hip and knee arthroplasty. *J. Bone Joint Surg. Am.* 97, 1449–1458. doi: 10.2106/JBJS.N.01114
- Wertheim, H. F. L., Melles, D. C., Vos, M. C., van Leeuwen, W., van Belkum, A., Verbrugh, H. A., et al. (2005a). The role of nasal carriage in *Staphylococcus aureus* infections. *Lancet Infect. Dis.* 5, 751–762. doi: 10.1016/S1473-3099(05)70295-4
- Wertheim, H. F. L., Verveer, J., Boelens, H. A. M., van Belkum, A., Verbrugh, H. A., and Vos, M. C. (2005b). Effect of mupirocin treatment on nasal, pharyngeal, and perineal carriage of *Staphylococcus aureus* in healthy adults. *Antimicrob. Agents Chemother.* 49, 1465–1467. doi: 10.1128/AAC.49.4.1465-1467.2005
- Wertheim, H. F. L., van Kleef, M., Vos, M. C., Ott, A., Verbrugh, H. A., and Fokkens, W. (2006). Nose picking and nasal carriage of *Staphylococcus aureus*. *Infect. Control Hosp. Epidemiol.* 27, 863–867. doi: 10.1086/506401
- Wertheim, H. F. L., Vos, M. C., Ott, A., van Belkum, A., Voss, A., Kluytmans, J. A. J. W., et al. (2004). Risk and outcome of nosocomial *Staphylococcus aureus* bacteraemia in nasal carriers versus non-carriers. *Lancet* 364, 703–705. doi: 10.1016/S0140-6736(04)16897-9
- Wertheim, H. F. L., Walsh, E., Choudhury, R., Melles, D. C., Boelens, H. A. M., Miajlovic, H., et al. (2008). Key role for clumping factor B in *Staphylococcus aureus* nasal colonization of humans. *PLoS Med.* 5:e17. doi: 10.1371/journal.pmed.0050017
- Williams, R. E. (1963). Healthy carriage of *Staphylococcus aureus*: its prevalence and importance. *Bacteriol. Rev.* 27, 56–71.
- Wollenberg, M. S., Claesen, J., Escapa, I. F., Aldridge, K. L., Fischbach, M. A., and Lemon, K. P. (2014). *Propionibacterium*-produced coproporphyrin III induces *Staphylococcus aureus* aggregation and biofilm formation. *mBio* 5:e01286-14. doi: 10.1128/mBio.01286-14
- Yan, M., Pamp, S. J., Fukuyama, J., Hwang, P. H., Cho, D.-Y., Holmes, S., et al. (2013). Nasal microenvironments and interspecific interactions influence nasal microbiota complexity and *S. aureus* carriage. *Cell Host Microbe* 14, 631–640. doi: 10.1016/j.chom.2013.11.005

- Yano, M., Doki, Y., Inoue, M., Tsujinaka, T., Shiozaki, H., and Monden, M. (2000). Preoperative intranasal mupirocin ointment significantly reduces postoperative infection with *Staphylococcus aureus* in patients undergoing upper gastrointestinal surgery. *Surg. Today* 30, 16–21. doi: 10.1007/PL00010040
- Zanger, P., Nurjadi, D., Gaile, M., Gabrysch, S., and Kremsner, P. G. (2012). Hormonal contraceptive use and persistent *Staphylococcus aureus* nasal carriage. *Clin. Infect. Dis.* 55, 1625–1632. doi: 10.1093/cid/cis778
- Zipperer, A., Konnerth, M. C., Laux, C., Berscheid, A., Janek, D., Weidenmaier, C., et al. (2016). Human commensals producing a novel antibiotic impair pathogen colonization. *Nature* 535, 511–516. doi: 10.1038/nature18634

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2018 Sakr, Brégeon, Mège, Rolain and Blin. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.