



HAL
open science

How do fish and allochthonous subsidies simultaneously control food-webs?

Eric Edeline, Gérard Lacroix

► **To cite this version:**

Eric Edeline, Gérard Lacroix. How do fish and allochthonous subsidies simultaneously control food-webs?. Journées scientifiques du laboratoire Bioemco, 2009, Versailles, France. hal-02000893

HAL Id: hal-02000893

<https://hal.science/hal-02000893>



Submitted on 6 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

How do top-down and bottom-up interactions control food-webs? (Eric Edeline and Gerard Lacroix)

1. Autotrophic and heterotrophic pathways for energy transfer in aquatic food webs



GREEN: heterotrophic pathway, bacteria are the main energy mobilizers, energy transfer efficiency to higher levels is low.
BLUE: autotrophic pathway, phytoplankton is the main energy mobilizer, energy transfer efficiency to higher levels is high.

2. Predictions to be tested

The core model of aquatic food webs responds simultaneously to bottom-up (allochthonous organic matter) and top-down (fish) control.

Allochthonous organic matter and fish will both favor the heterotrophic pathway (extended model).

Exudates from the phytoplankton favors organic matter mineralization by bacteria.

The autotrophic pathway maximizes production of zooplankton and fish.

3. The experimental setup



12 experimental tanks located at the biological station of Foljuif, 4 treatments in triplicate (1: no fish, no allochthonous input; 2: no fish, allochthonous input; 3: fish, no allochthonous input; 4: fish + allochthonous input, extended model) during 1 year.

Variables to be measured:

- ▶ water physics (temperature, pH) and chemistry (N, P concentrations)
- ▶ chemical nature of the dissolved and sedimented organic matter
- ▶ concentrations in bacteria, ciliates, flagellates, phyto- and zooplankton
- ▶ fish survival, somatic growth and fecundity

4. Collaborations

