



HAL
open science

Complete Genome Sequence of the Hyperthermophilic and Piezophilic Archaeon *Thermococcus barophilus* Ch5, Capable of Growth at the Expense of Hydrogenogenesis from Carbon Monoxide and Formate

Phil M. Oger, T. G. Sokolova, D. A. Kozhevnikova, E. A. Taranov, P. Vannier, H. S. Lee, K. K. Kwon, S. G. Kang, J. H. Lee, E. A. Bonch-Osmolovskaya, et al.

► To cite this version:

Phil M. Oger, T. G. Sokolova, D. A. Kozhevnikova, E. A. Taranov, P. Vannier, et al.. Complete Genome Sequence of the Hyperthermophilic and Piezophilic Archaeon *Thermococcus barophilus* Ch5, Capable of Growth at the Expense of Hydrogenogenesis from Carbon Monoxide and Formate. *Genome Announcements*, 2016, 4 (1), 10.1128/genomeA.01534-15 . hal-02001341

HAL Id: hal-02001341

<https://hal.science/hal-02001341v1>

Submitted on 28 Nov 2024


HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Complete Genome Sequence of the Hyperthermophilic and Piezophilic Archaeon *Thermococcus barophilus* Ch5, Capable of Growth at the Expense of Hydrogenogenesis from Carbon Monoxide and Formate

Philippe Oger,^a Tatyana G. Sokolova,^b Darya A. Kozhevnikova,^b Evgeny A. Taranov,^b Pauline Vannier,^c Hyun Sook Lee,^d Kae Kyoung Kwon,^d Sung Gyun Kang,^d Jung-Hyun Lee,^d Elizaveta A. Bonch-Osmolovskaya,^b  Alexander V. Lebedinsky^b

Laboratoire de Géologie de Lyon, UMR 5276 CNRS, Ecole Normale Supérieure de Lyon, Lyon, France^a; Winogradsky Institute of Microbiology, Research Center of Biotechnology, RAS, Moscow, Russia^b; Matis ohf., Food Safety, Environment and Genetics, Reykjavik, Iceland^c; Korea Institute of Ocean Science and Technology, Ansan, South Korea^d

We report here the complete sequence and fully manually curated annotation of the genome of strain Ch5, a new member of the piezophilic hyperthermophilic species *Thermococcus barophilus*.

Received 11 November 2015 Accepted 18 November 2015 Published 14 January 2016

Citation Oger P, Sokolova TG, Kozhevnikova DA, Taranov EA, Vannier P, Lee HS, Kwon KK, Kang SG, Lee J-H, Bonch-Osmolovskaya EA, Lebedinsky AV. 2016. Complete genome sequence of the hyperthermophilic and piezophilic archaeon *Thermococcus barophilus* Ch5, capable of growth at the expense of hydrogenogenesis from carbon monoxide and formate. *Genome Announc* 4(1):e01534-15. doi:10.1128/genomeA.01534-15.

Copyright © 2016 Oger et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/3.0/).

Address correspondence to Philippe Oger, philippe.oger@ens-lyon.fr, or Alexander V. Lebedinsky, a.lebedinsky@mail.ru.

Strain Ch5, a new member of the piezophilic hydrothermal vent archaeal species *Thermococcus barophilus*, has a pressure optimum of 40 MPa. It has been isolated from a deep-sea hydrothermal field of the Mid-Atlantic Ridge (Logachev field chimney, 3,020 m depth) on medium with 0.1 g/liter yeast extract under an atmosphere of 100% CO₂, on which it grew hydrogenogenically. Strain Ch5 has been assigned to the *T. barophilus* (1) species based on 16S rRNA gene sequence identity (100%). Strain Ch5 can grow organotrophically on yeast extract (0.5 g/liter). Strain Ch5 was among the five *Thermococcus* isolates shown to be capable of formate-driven growth coupled with H₂ production (2), raising questions as to whether this capacity was determined by genetic determinants that were similar to those identified in *Thermococcus onnurineus* and *Thermococcus gammatolerans*.

The complete genome sequence was determined by a combination of 454 and Illumina sequencing of standard unpaired libraries. *De novo* assembly of the hybrid data with Newbler 2.7 yielded six contigs, which were connected by PCR amplification and Sanger sequencing. The *T. barophilus* Ch5 genome consists of a single circular chromosome of 2,388,527 bp, with an average G+C content of 41.8%. A total of 2,679 protein-coding genes were annotated on the MaGe platform (3–5).

In silico DNA-DNA hybridization of the genomes of strain Ch5 and *T. barophilus* MP^T confirmed the affiliation of strain Ch5 with the *T. barophilus* species (predicted value, 81.3% ± 2.7% using GGDC 2.0 BLAST+ and the recommended formula 2) (6). However, the genomes differ considerably in size and gene content. The Ch5 genome lacks a plasmid, is ca. 400 kb bigger, and has 310 open reading frames (ORFs) more than the type species genome (7). The *T. barophilus* species core genome is composed of 1,868 families, of which 212 families are specific to *T. barophilus* and absent from other *Thermococcales*. Eighty percent (170/212) of this *T. barophilus*-specific genome encodes conserved proteins of unknown function,

while the remainder contains complete loci coding for the degradation of maltose, mannosylglycerate, and threonine. Interestingly, the Ch5 genome contains a chemotaxis locus associated with a flagellum-encoding gene cluster dissimilar from that of strain MP. Major genome rearrangements are associated with putative integrases, transposases, or clustered regularly interspaced short palindromic repeat (CRISPR) loci. Surprisingly, the Ch5 chromosome harbors three *cdc6* homologs, one similar to that of strain MP and two probably resulting from plasmid/chromosome integration events. Strains Ch5 and MP share five highly similar hydrogenase gene clusters, one of which is adjoined by a carbon monoxide dehydrogenase gene and determines the capacity for hydrogenogenic growth on CO. Additionally, strain Ch5 harbors three more hydrogenase gene clusters, one of them encoding a hydrogenase related to F₄₂₀-reducing hydrogenases (8), and two others adjoined by formate dehydrogenase genes. One of these two hydrogenase gene clusters includes a formate transporter gene. This cluster is very similar to those described for *T. onnurineus* and *T. gammatolerans* as genetic determinants of formate-driven growth coupled with H₂ production (2).

Nucleotide sequence accession number. The GenBank accession number of the *T. barophilus* Ch5 genome sequence is CP013050. The version described here is the first version.

ACKNOWLEDGMENTS

This work was supported by an ANR Living Deep grant (ANR-2010-BLAN-1725) from the Agence Nationale de la Recherche to P.O., a grant from the Molecular and Cell Biology program of the Russian Academy of Sciences and grant no. 14-24-00165 from the Russian Scientific Foundation to E.A.B.-O., and a grant from the Korea CI Gas Refinery R&D Center program of the Ministry of Science, ICT and Future Planning in the Republic of Korea to H.S.L.

FUNDING INFORMATION

Russian Academy of Sciences (RAS) and Russian Scientific Foundation provided funding to Elizaveta A. Bonch-Osmolovskaya under grant number 14-24-00165. Agence Nationale de la Recherche (ANR) provided funding to Phil M. Oger under grant number ANR-2010-BLAN-1725. Ministry of Science, ICT and Future Planning (MSIP) provided funding to Hyun Sook Lee.

REFERENCES

- Marteinsson VT, Birrien J-, Reysenbach A-, Vernet M, Marie D, Gambacorta A, Messner P, Sleytr UB, Prieur D. 1999. *Thermococcus barophilus* sp. nov., a new barophilic and hyperthermophilic archaeon isolated under high hydrostatic pressure from a deep-sea hydrothermal vent. *Int J Syst Bacteriol* 49:351–359. <http://dx.doi.org/10.1099/00207713-49-2-351>.
- Kim YJ, Lee HS, Kim ES, Bae SS, Lim JK, Matsumi R, Lebedinsky AV, Sokolova TG, Kozhevnikova DA, Cha S, Kim S, Kwon KK, Imanaka T, Atomi H, Bonch-Osmolovskaya EA, Lee J, Kang SG. 2010. Formate-driven growth coupled with H₂ production. *Nature* 467:352–355. <http://dx.doi.org/10.1038/nature09375>.
- Vallenet D, Belda E, Calteau A, Cruveiller S, Engelen S, Lajus A, Le Fevre F, Longin C, Mornico D, Roche D, Rouy Z, Salvignol G, Scarpelli C, Thil Smith AA, Weiman M, Medigue C. 2013. MicroScope—an integrated microbial resource for the curation and comparative analysis of genomic and metabolic data. *Nucleic Acids Res* 41:D636–D647. <http://dx.doi.org/10.1093/nar/gks1194>.
- Vallenet D, Engelen S, Mornico D, Cruveiller S, Fleury L, Lajus A, Rouy Z, Roche D, Salvignol G, Scarpelli C, Medigue C. 2009. MicroScope: a platform for microbial genome annotation and comparative genomics. *Database* 2009:bap021. <http://dx.doi.org/10.1093/database/bap021>.
- Vallenet D, Labarre L, Rouy Z, Barbe V, Bocs S, Cruveiller S, Lajus A, Pascal G, Scarpelli C, Medigue C. 2006. MaGe: a microbial genome annotation system supported by synteny results. *Nucleic Acids Res* 34:53–65. <http://dx.doi.org/10.1093/nar/gkj406>.
- Meier-Kolthoff JP, Auch AF, Klenk H, Göker M. 2013. Genome sequence-based species delimitation with confidence intervals and improved distance functions. *BMC Bioinformatics* 14:60. <http://dx.doi.org/10.1186/1471-2105-14-60>.
- Vannier P, Marteinson VT, Fridjonsson OH, Oger P, Jebbar M. 2011. Complete genome sequence of the hyperthermophilic, piezophilic, heterotrophic, and carboxydophilic archaeon *Thermococcus barophilus* MP. *J Bacteriol* 193:1481–1482. <http://dx.doi.org/10.1128/JB.01490-10>.
- Jeon JH, Lim JK, Kim M, Yang T, Lee S, Bae SS, Kim YJ, Lee SH, Lee J, Kang SG, Lee HS. 2015. Characterization of the *frhAGB*-encoding hydrogenase from a non-methanogenic hyperthermophilic archaeon. *Extremophiles* 19:109–118. <http://dx.doi.org/10.1007/s00792-014-0689-y>.