



# Complete Genome Sequences of 11 Type Species from the Thermococcus Genus of Hyperthermophilic and Piezophilic Archaea

P. Oger

## ► To cite this version:

P. Oger. Complete Genome Sequences of 11 Type Species from the Thermococcus Genus of Hyperthermophilic and Piezophilic Archaea. Genome Announcements, 2018, 6 (8). hal-02001367

**HAL Id: hal-02001367**

**<https://hal.science/hal-02001367>**

Submitted on 22 Nov 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Complete Genome Sequence of 11 type-species from the *Thermococcus* genus of hyperthermophilic and piezophilic Archaea

Philippe M. Oger

Université de Lyon, INSA Lyon, CNRS UMR5240, 11 Avenue Jean Capelle, F-69621 Villeurbanne, France

1 We report the genome sequences of the type strains of the species *Thermococcus barossii*, *T. celer*, *T.*  
2 *chitonophagus*, *T. gorgonarius*, *T. pacificus*, *T. peptonophilus*, *T. profundus*, *T. radiotolerans*, *T. siculi*,  
3 *T. thioreducens* and the prototype of a possible type strain of a novel *Thermococcus* species, strain  
4 P6.

5

6 *Thermococcus celer* were isolated from surface hydrothermal vent systems in the Mediterranean (1).  
7 *Thermococcus* sp. Strain P6 was isolated in 1997 by Pr. M. Thomm (University of Kiel, Germany) in 1997  
8 from an hydrothermal site off Palaeochori Bay (Milos, Aegean Sea, Greece). *T. thioreducens* was isolated  
9 from the Rainbow hydrothermal vent system on the Mid-Atlantic ridge, 2600m below the ocean surface  
10 (2). All other species were isolated from hydrothermal vents systems in the Pacific ocean (3-9). All species  
11 exhibit very similar growth characteristics, with optimal conditions for growth ranging from 83 and 88°C  
12 (T), pH 5.8 and pH 7 and salinity between 2 and 3%.

13 DNA was extracted from cells grown in TRM medium to late exponential phase. Whole-genome  
14 shotgun sequencing was carried out by a combination of IonTorrent PGM (800 bp fragment library, 318  
15 Chip, HiQ chemistry) and Illumina HiSeq run (500bp library, 100bp paired reads), which generated at  
16 least 583k reads with a mean 389 bp length (IonTorrent) and 13 million paired 100 bp reads after QC  
17 check and trimming (10-11) for each species. The IonTorrent data were assembled de novo with Newbler  
18 2.8 (12). The paired-end Illumina data was assembled de novo with velvet (13). The pooled IonTorrent  
19 and Illumina data was assembled de novo with MIRA (14) and Newbler. The hybrid data assembly  
20 yielded between 1 and 6 contigs which were connected by PCR.

21 Chromosome sizes range from 1.52Mb in *Thermococcus* sp. strain P6 to 2.07Mb in *Thermococcus*  
22 *thioreducens*. which is similar to known genomes from this family. Only two species, *T. peptonophilus*  
23 and *T. profundus*, harbor an extrachromosomal element. The *T. profundus* plasmid belongs to the  
24 pTBMP1 family of plasmid described in the piezophilic archaeon *T. barophilus* (15). To the exception of  
25 *T. chitonophagus*, the GC% ranges between 51.7 and 56% Similar to previously sequenced  
26 *Thermococcales*, all genomes harbor a single copy of the 16S-23S operon, two copies of 5S operons and  
27 45 to 48 tRNA genes, large CRISPR loci associated with *cas* genes (*cas1*, *cas2*, *cas3*, *cas4*, *cas5*, *cas6*,  
28 *csm2*), indicating multiple contacts with mobile elements of viral or plasmidic origin. Sintony analyzes  
29 confirm the plasticity of *Thermococcales* genomes. The gene content is consistent with an anaerobic and  
30 heterotrophic lifestyle. The genome of *T. barossi* encodes the capacity for hydrogenogenic growth on  
31 carbon monoxide, as identified previously in *T. onnurienus* and *T. barophilus*.

32 The genome sequence of *T. peptonophilus* (GC%, core genome phylogeny) confirms previous  
33 reports that it may have been misplaced into the *Thermococcus* genus, and that it should be placed in the  
34 *Pyrococcus* genus instead. Strain P6 is most closely related to *Thermococcus celer* strain Vu 13 (1), as  
35 indicated by *in silico* DNA-DNA hybridization of their genomes (predicted value = 28.90% using the  
36 genome-to-genome distance calculator GGDC2.0 (16). This genomic distance suggests that strain P6  
37 defines a novel species inside the *Thermococcus* genus.

38 **Nucleotide sequence accession number.** These whole Genome Shotgun project has been deposited in  
39 GenBank under the accession numbers CP014750, CP014751, CP014854, CP014855, CP014862,  
40 CP014863, CP015101, CP015102, CP015103, CP015104, CP015105, CP015106, and CP015193.

41

## 42 **ACKNOWLEDGEMENTS**

43 This work was supported in part by the Agence Nationale de la Recherche (ANR-10-BLAN-  
44 Living deep) and a grant from the Fond Recherche of the ENS Lyon to PO. Ion Torrent

45 sequencing was performed in part at the sequencing platform of the Institut de Génomique  
46 Fonctionnelle de Lyon by Dr. Sandrine Hughes and Benjamin Gillet.

47

## 48 REFERENCES

- 49 1. Zillig W, Holz I, Janekovic D, Schäfer W, Reiter WD. 1983. The archaeobacterium *Thermococcus*  
50 *celer* represents, a novel genus within the thermophilic branch of the archaeobacteria. Syst. Appl.  
51 Microbiol. 4:88-94.
- 52 2. Pikuta EV, Marsic D, Itoh T, Bej AK, Tang J, Whitman WB, Ng JD, Garriott OK, Hoover RB.  
53 2007. *Thermococcus thio-reducens* sp. nov., a novel hyperthermophilic, obligately sulfur-reducing  
54 archaeon from a deep-sea hydrothermal vent. Int. J. Syst. Evol. Microbiol. 57:1612-8.
- 55 3. Grote R, Li L, Tamaoka J, Kato C, Horikoshi K, Antranikian G. 1999. *Thermococcus siculi* sp.  
56 nov., a novel hyperthermophilic archaeon isolated from a deep-sea hydrothermal vent at the Mid-Okinawa  
57 Trough. Extremophiles 3:55-62.
- 58 4. Huber R, Stöhr J, Hohenhaus S, Rachel R, Burggraf S, Jannasch HW, Stetter KO. 1995.  
59 *Thermococcus chitonophagus* sp. nov., a novel, chitin-degrading, hyperthermophilic archaeum from a  
60 deep-sea hydrothermal vent environment. Arch. Microbiol. 164:255-264.
- 61 5. Jolivet E, Corre E, L'Haridon S, Forterre P, Prieur D. 2004. *Thermococcus marinus* sp. nov. and  
62 *Thermococcus radiotolerans* sp. nov., two hyperthermophilic archaea from deep-sea hydrothermal vents  
63 that resist ionizing radiation. Extremophiles 8:219-27.
- 64 6. Kobayashi T, Kwak YS, Akiba T, Kudo T, Horikoshi K. 1994. *Thermococcus profundus* sp. nov.,  
65 A new hyperthermophilic archaeon isolated from a deep-sea hydrothermal vent. Syst. Appl. Microbiol.  
66 17:232-236.
- 67 7. Miroshnichenko ML, Gongadze GM, Rainey FA, Kostyukova AS, Lysenko AM, Chernych NA,  
68 Bonch-Osmolovskaya EA. 1998. *Thermococcus gorgonarius* sp. nov. and *Thermococcus pacificus* sp.  
69 nov.: heterotrophic extremely thermophilic archaea from New Zealand submarine hot vents. Int. J. Syst.  
70 Bacteriol. 48:23-29.
- 71 8. Duffaud GD, d'Hennezel OB, Peek AS, Reysenbach A-L, Kelly RM. 1998. Isolation and  
72 characterization of *Thermococcus barossii*, sp. nov., a hyperthermophilic archaeon isolated from a  
73 hydrothermal vent flange formation. Syst. Appl. Microbiol. 21:40-49.
- 74 9. Gonzalez JM, Kato C, Horikoshi K. 1995. *Thermococcus peptonophilus* sp. nov., a fast-growing,  
75 extremely thermophilic archaeobacterium isolated from deep-sea hydrothermal vents. Arch. Microbiol.  
76 164:159-164.
- 77 10. FastQC, A quality control tool for high throughput sequence data.  
78 (<http://www.bioinformatics.babraham.ac.uk/projects/>)
- 79 11. Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence  
80 data. Bioinformatics 30 (15):2114-2120.
- 81 12. Newbler, a software package for de novo DNA sequence assembly  
82 (<http://www.454.com/products/analysis-software/>)

- 83 13. D.R. Zerbino and E. Birney. 2004. Velvet: algorithms for *de novo* short read assembly using de  
84 Bruijn graphs. *Genome Res.* **18**:821-829.
- 85 14. Chevreux B., Wetter, T. and Suhai, S. Genome sequence assembly using trace signals and  
86 additional sequence information. *Computer Science and Biology: Proceedings of the German Conference*  
87 *on Bioinformatics (GCB)* 1999. 99, pp. 45-56.
- 88 15. Vannier P, Marteinsson V, Fridjonsson O, Oger P, Jebbar M. 2011. Complete genome sequence  
89 of the hyperthermophilic piezophilic, heterotrophic and carboxydophilic archaeon *Thermococcus*  
90 *barophilus* MP. *J. Bacteriol.* 193:1481–1482.
- 91 16. Meier-Kolthoff JP, Klenk H-P, Göker M. 2014. Taxonomic use of DNA G+C content and DNA–  
92 DNA hybridization in the genomic age. *Int. J. Syst. Evol. Microbiol.* **64**:352–356.