



HAL
open science

SNARE Machinery is optimized for ultra-fast fusion

Fabio Manca, Frédéric Pincet, Lev Truskinovsky, James E Rothman, Lionel Foret, Matthieu Caruel

► **To cite this version:**

Fabio Manca, Frédéric Pincet, Lev Truskinovsky, James E Rothman, Lionel Foret, et al.. SNARE Machinery is optimized for ultra-fast fusion. Proceedings of the National Academy of Sciences of the United States of America, 2019, 116 (7), pp.2435-2442. 10.1073/pnas.1820394116 . hal-02006229

HAL Id: hal-02006229

<https://hal.science/hal-02006229v1>

Submitted on 4 Feb 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

SNARE Machinery is optimized for ultra-fast fusion

Fabio Manca^a, Frederic Pincet^a, Lev Truskinovsky^b, James E. Rothman^{c,d}, Lionel Foret^a, and Matthieu Caruel^{e,1}

^a Laboratoire de Physique de l'Ecole Normale Supérieure, Ecole Normale Supérieure (ENS) - CNRS - Université PSL - Sorbonne Université - Université Paris-Diderot - Sorbonne Paris Cité, 75005 Paris, France; ^b Physique et Mécanique des Milieux Hétérogènes (PMMH), Ecole Supérieure de Physique et de Chimie Industrielles (ESPCI) - CNRS - Université PSL, 75231 Paris cedex 05, France.; ^c Department of Cell Biology, Yale University, New Haven, CT 06520, USA; ; ^d Department of Experimental Epilepsy, Institute of Neurology, University College London, London, United Kingdom; ^e Modélisation et Simulation Multi-Echelle (MSME), Université Paris-Est - CNRS, 94010 Créteil, France

SNARE proteins zipper to form SNAREpins that power vesicle fusion with target membranes in a variety of biological processes. A single SNAREpin takes about 1 second to fuse two bilayers, yet a hand-full can ensure release of neurotransmitters from synaptic vesicles much faster, in a 10th of a millisecond. We propose that, similar to the case of muscle myosins, the ultrafast fusion results from cooperative action of many SNAREpins. The coupling originates from mechanical interactions induced by confining scaffolds. Each SNAREpin is known to have enough energy to overcome the fusion barrier of 25-35 $k_B T$, however, the fusion barrier only becomes relevant when the SNAREpins are nearly completely zippered and from this state each SNAREpin can deliver only a small fraction of this energy as mechanical work. Therefore they have to act cooperatively and we show that at least 3 of them are needed to ensure fusion in less than a millisecond. However, to reach the pre-fusion state collectively, starting from the experimentally observed half-zippered metastable state, the SNAREpins have to mechanically synchronize which takes exponentially longer time as the number of SNAREpins increases. Incorporating this somewhat counter-intuitive idea in a simple coarse grained model results in the novel prediction that there should be an optimum number of SNAREpins for sub-ms fusion: 3-6 over a wide range of parameters. Interestingly, *in situ* cryo-electron microscope tomography has very recently shown that exactly six SNAREpins participate in the fusion of each synaptic vesicle. This number is in the range predicted by our theory.

SNARE | membrane fusion | protein folding | neurotransmitter release | muscle contraction

1. Introduction

Protein transport within cells relies heavily on membrane-enveloped vesicles that ferry packets of enclosed cargo (1–4). The content of the vesicles is released *via* their fusion with target membranes. This transition is impeded by repulsive forces acting when the distance between the membranes is in the range of ~ 1 nm. The encountered energy barrier is of the order of $30 k_B T$, implying that spontaneous fusion would take minutes, which is not fast enough in most biological situations (5–8). For this reason, the process is assisted by SNARE proteins (soluble N-ethylmaleimide-sensitive factor attachment protein receptors, SNAREpins) whose conformational change (zippering) exerts forces pulling the vesicle membrane towards the target membranes.

While the total free energy change associated with the zippering process is of the order of $\sim 70 k_B T$ (9), most of this energy is consumed as the SNAREpins bring the membranes into close apposition. Biologically, the initial assembly prior to fusion provides compartmental specificity (pairing the correct SNAREs together) and allows for temporal regulation (clamping). Terminal zippering is then the process which utilizes the remaining energy for bilayer fusion at the small

($\sim 1 - 2$ nm) separations where the repulsive forces become relevant. Recent studies suggest that each SNAREpin can deliver only about $5 k_B T$ of mechanical work at this stage (10, 11), which explains why it takes about 1 second for a single SNAREpin to fuse two bilayers (12, 13).

It is known however that the release of neurotransmitters from synaptic vesicle occurring at nerve endings happens considerably faster, in a 10th of a ms as is necessary to keep pace with action potentials, and ensure synchronous release (4, 14–18). A widely accepted explanation for this remarkable difference in time scales is that multiple SNAREpins would need to cooperate to accelerate fusion after being synchronously released from a clamped state. There have been indirect indications that the number of SNAREpins necessary to achieve a sub-millisecond fusion may be relatively small, ranging from 2 to 6 (19–21). Very recently, cryo-electron microscope tomography of synaptic vesicles *in situ* revealed an underlying 6-fold symmetry suggesting that exactly 6 SNAREpins are involved in such processes (22).

How so few co-operating SNAREpins manage to accelerate fusion ten thousand times (from ~ 1 s to ~ 0.1 ms) has been a mystery. Previous modeling attempts have suggested that more than 16 SNAREpins would be required (23, 24). Here we show that the key to understand how only a few SNAREpins can achieve such rapid fusion is the simple fact that they are mechanically coupled through effectively rigid common membranes. The account of such mechanical coupling leads to a striking prediction that the number of SNAREpins must be highly constrained to ensure sub-millisecond release of neurotransmitters. Quite remarkably, the predicted optimal range, 3-6, is in excellent agreement with most recent experimental

Significance Statement

We propose a new mechanistic description for the fusion of a synaptic vesicle with a target membrane executed by a team of zippering SNAREpins. In the context of neurotransmitters release, this process naturally decomposes in two steps whose rates depend on the number of SNAREpins N . The first step is the synchronized escape from the metastable half-zippered state, which get exponentially more sluggish as N increases. The second step is the fusion of the two closely tethered membranes, which is accelerated exponentially by an increase of N . The tradeoff between these two antagonist trends results in a sharply optimal number of SNAREpins $N = 3 - 6$ which ensure fusion at the physiological sub-ms timescale.

M.C. L.F. and L.T. designed research; F.M. performed research; F.M. analyzed data; F.M., F.P., L.T., J.E.R., L.F., and M.C. interpreted the results and wrote the paper

The authors declare no conflict of interest.

¹To whom correspondence should be addressed. E-mail: matthieu.caruel@u-pec.fr




Fig. 1. The fusion machinery. (A) schematic of the two membranes with two attached SNAREpins. (B) Mechanical model with $N = 4$ SNAREpins in parallel bridging the two membranes separated by the distance y ; two SNAREpins are in state c and two are in state n so $N_c = 2$. (C) Model of a single SNAREpin. (D) Fusion energy landscape. See Table 1 for the complete list of parameter values.

results (22).

We draw a fundamental analogy between the collective zippering of the SNAREpins and the *power-stroke* in a bundle of elastically coupled muscle myosin II proteins which is known to also take place at 1 ms timescale. Building upon the seminal theory of the myosin power-stroke proposed by Huxley and Simmons (25–28), we model the fusion machinery as a mechanical system where the SNAREpins are represented as snap-springs interacting through supporting membranes. The implied bi-stability is supported by recent experiments showing the presence of a metastable half-zipped state (10, 11).

The theoretical approach developed in this paper highlights the essential role of mechanical coupling among proteins undergoing conformational changes in ensuring swift, highly synchronized mechanical response. This is likely a general biological principle (27, 29).

2. Fusion Machinery

The goal of the model is to describe the dynamic coupling between the individual SNAREpins zippering and to study the associated evolution of the distance between the vesicle and a the target membrane. The assembled SNARE machinery is represented as a bundle of N parallel SNAREpins bridging the two membranes, separated by the distance y , see Fig. 1(A) and (B)]. We assume that irreversible fusion occurs when this distance reaches a critical value y_f . The characteristic length associated with the deformation of the membranes generated by a zippering SNAREpin is large compared to the typical size of the SNARE bundle, see Materials and Methods A. Hence, the membranes can be viewed as two rigid backbones cross-linked by N identically stretched SNAREpins.

Single SNAREpin as a bi-stable snap-spring The experimental work conducted in Refs. (10, 11, 30), suggests that a single SNAREpin can switch randomly between two metastable conformations: n (half-zipped) when only the N-terminal domain of the SNAREs is zippered and c (fully zippered), when both the C-terminal and the linker domain are zippered. To describe this process, we assume that the half-to fully zippered transition in a SNARE complex is similar to the pre-

Table 1. Physical parameters adopted in the model and references

Parameter	Symbol	Value	Units	Reference
Zippering distance	a	7	nm	(10)
Energy bias	e_0	28	$k_B T$	(10)
Fully zippered stiffness	κ_c	12	pN nm^{-1}	SI**
Half-zipped stiffness	κ_n	2.5	pN nm^{-1}	SI
Maximum zippering rate	k	1	MHz	(10)
Drag coefficient	η	3.8×10^{-7}	Ns m^{-1}	
FB* position	y_f	2	nm	(5, 31)
FB width	σ_f	0.3	nm	(6, 32)
FB height	e_f	26	$k_B T$	(7, 8)

*FB: Fusion Barrier; **SI: SI Appendix; $1 k_B T \approx 4 zJ$

to post-power-stroke conformational change in a myosin motor, (25, 27).

Suppose that each SNAREpin is equipped with an internal spin type degree of freedom characterizing the state of the protein, n or c . We denote by a the amount of shortening resulting from the $n \rightarrow c$ transition in the absence of external load, and by e_0 the energy difference between the two states. This parameter can be interpreted as the typical amount of mechanical work necessary to force the $c \rightarrow n$ transition (partial unzipping), see Fig. 1(C).

When the SNAREs are bound to the membranes, we assume that the rates $k_+(y)$ —associated with the $n \rightarrow c$ transition—and k_- —associated with the $c \rightarrow n$ transition—depend on the mechanical load induced by the variations of the inter-membrane distance. To specify this dependence, both states are assumed to be “elastic” in the sense that they exist, as phases, over an extended range of separations y , due to elongations of the zippered and unzipped SNARE residues, internal bonds rearrangement etc..., see Fig. 1(C). For simplicity we assume that the deformations remains in the elastic regime so that states n, c can be associated with quadratic energies $e_{n,c}(y)$, with minima located $y = \{0, a\}$, and with the lumped stiffnesses $\kappa_{n,c}$. The transitions rates are defined so that, for a given separation, they favor the state with the lowest energy and verify detailed balance. For the detailed expressions of $e_{n,c}$ and k_{\pm} , see Materials and Methods B.

Dynamics of the fusion machinery The parallel arrangement of the SNAREs implies that the conformational state of the bundle is fully characterized by N_c , the number of SNAREpins in state c . This variable evolves according to the stochastic equation $N_c(t + dt) = N_c(t) + \{1, -1, 0\}$, with the outcomes $\{1, -1, 0\}$, characterized by the probabilities $W_{+1}(y, N_c) = (N - N_c) k_+(y) dt$, $W_{-1}(y, N_c) = N_c k_-(y) dt$ and $W_0 = 1 - W_{+1} - W_{-1}$. While the SNAREpins can switch independently, the transition rates k_{\pm} are functions of the collective variable y , whose dynamics in turn depends on N_c .

To specify the coupling between the two degrees of freedom N_c and y , and thereby formulate the complete model of the fusion process, we first recall that the motion of the vesicle in the overdamped regime results from the balance between the force applied by the N SNAREpins, the membrane repulsion and the viscous drag. Taking into account the thermal fluctuations, this force balance translates into the stochastic




Fig. 2. Main results. [(A) and (B)] typical stochastic trajectories of the inter-membranes distance y (A) and the number $N_c(t)$ of SNAREpins in state c (B) obtained from the numerical simulation. (C) Average of the waiting times τ_1 (black), τ_2 (blue) and $\tau_{\text{fusion}} = \tau_1 + \tau_2$ (red) obtained from the numerical simulations (symbols) and from our effective chemical model (lines). (D) Effective free energy landscape Φ showing the three stages of fusion and the associated transition rates. Parameters are listed in Table 1.

equation,

$$\eta \dot{y} = -\frac{\partial}{\partial y} (E_{\text{snare}} + E_{\text{fusion}}) + \sqrt{2\eta k_B T} \xi(t), \quad [1]$$

where $\xi(t)$ is a standard white noise, and η is a drag coefficient representing the friction opposing the motion of the vesicle. At a given y , the force applied by the bundle derives from the sum of individual SNAREpin energies $E_{\text{snare}}(y, N_c) = N_c e_c(y) + (N - N_c) e_n(y)$. Finally, the inter-membrane repulsion, due to short-range forces between the two membranes, is schematically modeled by a Gaussian energy barrier (5) $E_{\text{fusion}}(y) = e_f \exp[-(y - y_f)^2 / (2\sigma_f^2)]$, where y_f is the critical separation, while e_f and σ_f are the height and the width of the barrier, respectively, see Fig. 1(D). The ensuing dynamics of the system unfolds in the space of two variables: the continuous one, $y(t)$, and the integer-valued one, $N_c(t)$. The associated energy landscape has a multiwell structure that accounts for the configurational states of N individuals. The response is governed by the two stochastic equations, for which initials conditions still need to be specified. We consider the initial state $N_c(t=0) = 0$, and $y(t=0) = a$, which corresponds to the configuration where the SNAREpins are at the bottom of the energy well describing state n . This configuration characterizes the system immediately after the calcium induced collapse of Synaptotagmin which triggers the full zippering of the SNAREs (33). This point is discussed in more details in Section 4.

Model parameters The model is calibrated as follows, see Table 1 and SI Appendix (A) for further details. The mechanical parameters characterizing a single SNAREpin, a , e_0 , κ_n and κ_c , are determined by using our model to reproduce the experimental results obtained from stretching tests with optical tweezers, see Refs. (9, 10). The energy bias a and e_0 are chosen to be compatible with the results obtained from these studies. The procedure used to estimate the stiffnesses $\kappa_{n,c}$ is more complex and explained in details in SI Appendix. The value of the rate k , is fixed in accordance with estimates from Refs. (10, 34). The drag coefficient is computed using the

Stokes formula $\eta = 6\pi\mu R$, where $R = 20$ nm is the vesicle radius, and $\mu = 10^{-3}$ Pa s the fluid viscosity. The corresponding characteristic timescale is $\tau_\eta = \eta a^2 / (k_B T) \approx 4.5$ μ s.

The values of the parameters y_f , σ_f and e_f are chosen to be compatible with the current literature (7, 8, 11, 12, 23, 24, 35–41). In particular, values of e_f between 26 and 34 $k_B T$ have been reported for various types of lipids. We chose 26 $k_B T$ [POPC*, see Ref. (8)] which leads to a single SNAREpin average fusion time of 1 s.

3. Results


Numerical simulations Typical stochastic trajectories, $y(t)$ and $N_c(t)$ obtained from numerical simulations, are shown in Fig. 2[(A) and (B)]. They indicate that the fusion process can be decomposed into two stages, characterized by the times τ_1 and τ_2 . During the first stage, the system remains in its initial configuration ($y \simeq a, N_c = 0$) with only isolated $n \rightarrow c \rightarrow n$ transitions. After a time τ_1 , the inter-membrane distance drops abruptly to $y \simeq 2.5$ nm, while all the SNAREpins collectively switch from state n to state c . The inserts in Fig. 2[(A) and (B)] show that this transition occurs within 10 ns after the inter-membrane distance has reached the value $y = y_*$; the irreversible collective zippering itself ($N_c = 0 \rightarrow N_c = 4$) lasting about 1 ns. After the synchronized $n \rightarrow c$ transition the inter-membrane distance remains above the threshold $y = y_f$ for a time τ_2 before fusion. The duration of the whole process is therefore $\tau_{\text{fusion}} = \tau_1 + \tau_2$.

The mean timescales $\bar{\tau}_1$ and $\bar{\tau}_2$ (obtained by averaging 10^3 stochastic trajectories) are represented as functions of the number of SNAREpins in Fig. 2(C) on a semi-logarithmic scale. Observe that $\bar{\tau}_1$ increases exponentially with N and $\bar{\tau}_2$ decreases exponentially with N . These antagonistic N -dependencies result in the average fusion time $\bar{\tau}_{\text{fusion}} = \bar{\tau}_1 + \bar{\tau}_2$ exhibiting a remarkably sharp minimum, see Fig. 2[(C), red]. With the set of parameters values reported in Table 1, this minimum is attained at $N_* = 4$ and is associated with a fusion timescale of ~ 100 μ s. In addition we obtain a fusion time

* 1-palmitoyl-2-oleoyl-sn-glycero-3-phosphocholine lipid

of the order of 1 s for a single SNAREpin. Both values are consistent with *in vitro* (12) and *in vivo* (14, 42) experimental measurements.

Fusion as a two-stage reaction To elucidate the mechanism of fusion in two stages, we present here a ‘toy’ model where the whole process is recast as two successive reactions



where IS stands for an intermediate state whose characteristics depend on the mechanical properties of the zippered SNAREpins. In this representation, the fusion is viewed as the outcome of two distinct sub-steps: the collective zippering and the topological membrane merger.

To justify such model reduction, we assume that the timescale of the $n \rightleftharpoons c$ transition is negligible compared to the timescale describing the relaxation of the vesicle position. In the corresponding limit ($kt_\eta \gg 1$) Eq. (1) can be averaged with respect to the equilibrium distribution of the variable $N_c(t)$ [see Materials and Methods D], so the original system reduces to the one-dimensional stochastic equation

$$\eta \dot{y} = - \frac{d}{dy} [N f_{\text{snare}}(y) + E_{\text{fusion}}(y)] + \sqrt{2\eta k_B T} \xi(t), \quad [3]$$

where, the energy $E_{\text{snare}}(N_c, y)$ appearing in Eq. (1)—which depends on N_c and y —is replaced by the equilibrium free energy $f_{\text{snare}}(y) = -k_B T \log \{ \exp[-e_c(y)/(k_B T)] + \exp[-e_n(y)/(k_B T)] \}$ —which depends only on y . This free energy is illustrated in Fig. 2(D) dashed line]. The overall potential $\Phi(y) = N f_{\text{snare}}(y) + E_{\text{fusion}}(y)$, driving the effective dynamics (3) is shown by the solid red line in Fig. 2(D). It exhibits two local minima representing two metastable states. The first metastable state [point A in Fig. 2(D)] is located at $y \simeq a$, where on average all the SNAREpins are in state n . The second one [point B in Fig. 2(D)] is located at $y_f < y_2 < y_*$ and represents the *intermediate* state where on average all the SNAREpins are in state c , still confronting a reduced fusion barrier.

The system evolving in this energy landscape from the initial— $y \simeq a$ —to the final— $y = y_f$ —state faces two successive energy barriers $\Delta\Phi_1$ and $\Delta\Phi_2$. With each barrier $\Delta\Phi_{1,2}$ one can associate a waiting time $\bar{\tau}_{1,2}$, that can be approximated by the Kramers formula (43–45)

$$\bar{\tau}_{1,2} = \tau_\eta \alpha_{1,2} \exp[\Delta\Phi_{1,2}/(k_B T)]. \quad [4]$$

The values of the numerical prefactors $\alpha_{1,2}$ are determined by the local curvatures of the potential Φ at its critical points and depend weakly on N , see Materials and Methods D.

The approximated timescales $\bar{\tau}_{1,2}$ are compared with the numerically computed values $\tau_{1,2}$ in Fig. 2(C) solid lines]. The excellent agreement between the two sets of results suggests that the whole fusion process can effectively be described by two successive ‘chemo-mechanical’ reactions, and that the rates in Eq. (2) can be computed from the formulas $k_{1,2} = \bar{\tau}_{1,2}^{-1}$ while the remaining rate k_{-1} is prescribed by the condition of detailed balance.

Finally, note that with the parameters reported in Table 1, $k\tau_\eta = 4.5$, which shows that our effective model is accurate even if the condition $k\tau_\eta \gg 1$ is not fully satisfied.

The peculiar dependencies of the waiting times $\tau_{1,2}$ on the number of SNAREpins N can be now understood by referring to the N -dependence of the energy barriers $\Delta\Phi_{1,2}$.

Timescale τ_2 : The cooperative action of the SNAREs reduces the time for crossing the fusion barrier. In the intermediate state [point B in Fig. 2(D)], the SNAREpins are all in state c and the pulling force they apply on the membranes is exactly balanced by the short range repulsive forces. The system remains trapped in this state until a thermal fluctuation delivers the energy $\Delta\Phi_2$ allowing the system to reach the distance $y = y_f$, where the fusion occurs.

In the absence of SNAREs, this energy difference is simply the bare fusion barrier e_f , see Fig. 1(D). When the SNAREpins are present, the total force they apply brings the two membranes in close contact, which reduces the energy barrier. This effect is amplified by an increase in the number of SNAREpins: the larger the number of SNAREpins, the larger the overall force, so the closer the membrane can be brought together, see Fig. 2(D).

Since the inter-membrane potential $E_{\text{fusion}}(y)$ decays rapidly as y increases, we can approximate the second energy barrier by $\Delta\Phi_2 \simeq e_f - Nw$, where w represents the amount of mechanical work that a single SNAREpin can deliver, see Materials and Methods E for the derivation of this result and the mathematical expression of w . According to Eq. (4), we then have

$$\bar{\tau}_2(N) \propto \exp[-Nw/(k_B T)], \quad [5]$$

hence the exponential decay of the time τ_2 with the number of SNAREpins.

With the parameters of Table 1, each SNAREpin provides a mechanical work $w \simeq 4.5 k_B T$ when it encounters the fusion barrier, which reduces the average time for fusion $\bar{\tau}_2$ by a factor of ~ 100 , see Fig. 2(D). This multiplicative effect allows fast fusion at the sub-millisecond timescale with as few as three SNAREpins. For large enough number of SNAREpins (here $N > 7$), the overall applied force surpasses the membrane repulsion and the remaining fusion barrier disappears. The obtained exponential decay of the timescale τ_2 with the number of SNAREpins suggest that the fusion could in principle proceed much faster than $\sim 100 \mu\text{s}$, being only limited by viscous forces. Considering that each vesicle can accommodate up to ~ 100 SNAREpins, one cannot rule out the possibility of neurotransmitter release occurring much faster than $100 \mu\text{s}$. Next we argue that such scenario is unlikely by showing that the fusion process gets slowed down if the number of SNAREpins becomes too large.

Timescale τ_1 : Increasing the number of SNAREpins slows down the synchronous zippering. The average time $\bar{\tau}_1$, taken for all the SNAREpins to switch from the n to the c conformation and then pull the membranes toward the bottom of the fusion barrier, exponentially increases with the number of SNAREpins N , see Fig. 2C. This dependence can be explained as follows.

As long as $y_* < y < a$, the individual transition rates are such that $k_+ < k_-$ which implies that, on average, all the SNAREpins are in state n and therefore under compression, see Fig. 1(C). This idea is in agreement with the experimental results from Ref. (9), that revealed the presence of the half-zipped metastable state. Consequently in the interval $y_* < y < a$ the average force $-df_{\text{snare}}/dy$, that the SNAREpins collectively exert on the membranes, is repulsive. Beyond the point $y \simeq y_*$, the state c is stabilized ($k_+ > k_-$) and the average force becomes attractive. Since this force is proportional to the number of SNAREpins, the waiting time before a fluctuation can provide enough energy to surpass the




Fig. 3. Effect of the intrinsic energy barrier Δe on the optimal number of SNAREpins (A) and on the associated fusion time (B). The parameters values are taken from Table 1 with $\kappa_n = 0.11 - 24 \text{ pN nm}^{-1}$.

repulsion—and overcome the barrier $\Delta\Phi_1$ —increases with N . This constraint results from the mechanical feedback induced by the membranes. The latter play the role of a rigid backbone that forces the SNAREpins to bridge approximately the same intermembrane distance, see Refs. (28, 29, 46).

To specify the N -dependence of τ_1 , we use the fact that for $y > y_*$ we can consider that $E_{\text{fusion}} = 0$, so $\Delta\Phi_1$ can be approximated by $\Delta\Phi_1 \simeq N[f_{\text{snare}}(y_*) - f_{\text{snare}}(a)] \simeq \Delta e - k_B T \log(2)$, where $\Delta e = e_n(y_*) - e_0$. According to Eq. (4), we can then write

$$\bar{\tau}_1(N) \propto \exp[N\Delta e/(k_B T)], \quad [6]$$

which shows that the timescale $\bar{\tau}_1$ increases exponentially with the number of SNAREpins.

From Eq. (6) we obtained that the first energy barrier is fully controlled by a single parameter Δe which therefore has a strong influence on both the existence and value of the optimal number of SNAREpins. To study the effect of Δe on the fusion time, we varied the parameter κ_n describing the curvature of the energy e_n . The results of our parametric study are summarized in Fig. 3. These data were obtained by using Eq. (4) to compute the intersection of the curves $\bar{\tau}_{1,2}(N)$ for each value of Δe . We checked that the results are in good agreement with direct numerical simulations. Despite the broadness of the interval of parameter values tested, the optimal number of SNAREpins remains below 10. If we consider only the cases corresponding to sub-millisecond fusion times, we obtain $N \geq 3$ with $\Delta e < 4 \text{ k}_B T$. The latter value is compatible with recent estimate of $\Delta e \approx 5 \text{ k}_B T$ for the $n \rightarrow c$ transition energy barrier, see Refs. (9, 17). Note also that the predicted optimal number of SNAREpins is robust because it corresponds to a *plateau* on the $N_*(\Delta e)$ curve, see Fig. 3(A).

Robustness of the predictions. The results presented above, were obtained for the parameters values listed in Table 1. For some of these parameters only a rough estimate is available at this stage, see SI Appendix. To test the robustness of our theoretical predictions, we computed the average waiting times $\bar{\tau}_{1,2}(N)$ from Eq. (4) for different values of four key parameters of the model, e_0 , κ_c , e_f , σ_f , see Fig. 4. For each of these parameters, the lower and the upper bounds delimit broad intervals covering the values obtained from different experimental studies.

A comparison between Fig. 4 and Fig. 2(B) shows that our results are only marginally affected by changes in the parameter values. In particular, the existence of a sharp minimum of the fusion time, associated with an optimal number




Fig. 4. Robustness of the prediction. Influence of the parameters e_0 (A), κ_c (B), e_f (C) and σ_f (D), on the timescales $\bar{\tau}_{1,2}$ and $\bar{\tau}_{\text{fusion}}$. The results were obtained using Eq. (4).

of SNAREpins, is a robust prediction. In addition, the value of the optimal number of SNAREpins, is weakly sensitive to the parameters: it always remains in between 3 and 6. Remarkably, despite the large difference between the upper and lower bounds for each of the parameters, the average fusion time remains in the sub-millisecond scale.

The energy landscape associated with the zippering of the SNARE complexes is the object of intense current research (9, 17, 47). In our model this landscape is fully characterized by only four parameters: the distance a , the energy e_0 , and the stiffnesses $\kappa_{n,c}$. While the distance a has been measured with precision in recent works (9, 10), the values of the other three parameters are still not known with certainty. Several estimates of the energy bias e_0 lying between 20 and $40 \text{ k}_B T$ can be found in the literature (9, 48). We show in Fig. 4(A) that variations within this interval affect mostly τ_1 , change the fusion time by one order of magnitude, but have almost no effect on the optimal number of SNAREpins. Currently, only indirect evaluation of the stiffnesses $\kappa_{n,c}$ can be obtained from the available data, see SI Appendix. Within the broad range of values tested in our numerical simulations we again observed only small variations of the optimal number of SNAREpins, see Fig. 3, and Fig. 4(B).

One of the most documented physical phenomena involved in the fusion process is the merging of the two membranes. The amplitude of the associated repulsion force depends, in our model, on the parameters e_f and σ_f , whose influence on the fusion time is illustrated in Fig. 4(C) and (D), respectively. As expected from the analysis presented in Section 3, changing the values of these two parameters affect only the height of energy barrier $\Delta\Phi_2$ and therefore the timescale τ_2 . Increasing e_f raises the height of the maximum of E_{fusion} [see Fig. 1(D)], while decreasing σ_f deepens the second energy well [point B in Fig. 2(D)], which results in both cases in the increase of τ_2 . This leads *in fine* to the increase of the optimal

number of SNAREs. Notice that e_f depends on the type of lipids, on the membrane curvature and is also strongly sensitive to the membrane tension (37–41). Therefore, its value can be different in different cells or experimental set-ups. In particular, we expect the *in vivo* value to be smaller than the value measured in artificial systems ($35 \text{ k}_B\text{T}$), which in general use low tension and low curvature membranes, see Ref. (8).

In conclusion, while additional experimental studies are needed to refine the calibration of the model, the above parametric study shows the robustness of the effects of the mechanical crosstalk between the SNAREpins.

4. Discussion

In this paper, we elucidated the central role played by mechanical coupling in synchronizing the activity of SNAREpins, which is necessary to enable sub-millisecond release of neurotransmitters. Our approach to the problem complements previous studies focussed predominantly on the molecular details of the single SNARE zipper transition (9, 10, 30, 37, 47, 49–53).

As a starting point we used a previously unnoticed analogy between the activity of SNARE complexes and the functioning of myosin II molecular motors. Viewed broadly, both systems ensure ultra-fast mechanical contraction. In the case of muscle, destabilization of the pre-power stroke state is the result of a mechanical bias created by an abrupt shortening of the myofibril (25, 54). In the case of SNAREs, similarly abrupt destabilization is a result of the calcium-induced removal of the synaptotagmin-based clamp, most likely when Ca^{2+} triggers disassembly of the synaptotagmin ring (55).

To pursue this analogy, we developed a variant of the power-stroke model of Huxley and Simmons (25, 27), in which the zipping is viewed as a transition between two discrete states endowed with different elastic properties. This representation is supported by recent experiments (9, 10), which provided essential data for the calibration of the model.

Our analysis of the collective behavior of N “switchers” of this type suggests that the main function of the SNARE machinery is to bring the two membranes to a distance beyond which the fusion process can proceed spontaneously. The emerging intermediate configuration, where the two membranes are sufficiently closely tethered, can be then viewed as an intermediate state in the reaction process linking the fused and unfused states. The result is a representation of the SNARE mediated fusion as a two stage reaction.

We linked the first stage of the process with the collective zipping of the SNAREpins and showed that this step gets exponentially more sluggish as the number of SNAREpins increases. This phenomenon was studied previously in the context of muscles, see Refs. (26, 29). It originates (i) from the experimentally suggested presence of a metastable half-zipped state along the zipping free energy landscape (9), and (ii) from the long-range mechanical interactions mediated by the scaffolding membranes, which create a negative feedback preventing a fast collective escape from the metastable half-zipped state.

The second stage of the process is the transition from the intermediate state to the fused state. The associated timescale τ_2 decreases exponentially with the number of SNAREpins because the larger the number of acting SNARE-

pins, the closer the membranes can be brought together in the intermediate state and therefore the higher is the energy of this state. This results in an exponential decay of the timescale τ_2 with the number of SNAREpins. Behind this phenomenon is the presence of a residual force in the configuration where the SNAREpins have reached the intermediate state. This perspective is supported by the results of Ref. (10, 11).

The antagonistic N -dependence of the rates characterizing the two stages reveals the existence of an optimal number of SNAREs N_* allowing the system to perform fusion at the physiologically appropriate timescales. Our prediction $N_* = 4 - 6$ is supported by recent *in situ* cryoelectron microscope tomography observation, see Ref. (22).

We remark that our result strongly depend on the initial configuration of the system, which we link with the structure of the fusion machinery immediately after Synaptotagmin removal by calcium. Notice that the position y_* of the barrier separating the half-zipped and the fully zippered states is such that $y_* < a$. Therefore the timescale τ_1 exists only if the initial membrane separation $y_0 > y_*$. This assumption seems to be supported by experiments, see Ref. (56). It has previously been reported that, upon approach of two membranes devoid of SNAREs, Synaptotagmin exerts repulsive force from 10 nm down to 4 nm where it becomes a repulsive wall (56). According to this result, y_0 should range between 4 and 10 nm. However it is probably slightly larger under physiological conditions because of the presence of the SNAREs. With the parameters adopted in our simulations (see Table 1), the position of the barrier is $y_f \simeq 4.5 \text{ nm}$ in accordance with Refs. (9, 10), see Fig. 2. Therefore, in all likelihood, y_0 is larger than y_* and our predictions should be valid.

Finally we mention the fact that the timescale τ_2 exponentially decrease with N seems to be supported by experimental studies reporting sub-millisecond fusion time with $N = 3 - 6$ (19–21) However, in a recent theoretical study the decay was also found to be exponential but with a much slower decay: the cooperation of at least sixteen SNAREs was predicted to be necessary to reach the physiological fusion time $\sim 100 \mu\text{s}$ (23, 24). The difference is explained by the fact that the residual work in this study is $w = 0.48 \text{ k}_B\text{T}$ instead of $4.5 \text{ k}_B\text{T}$ in our model, see Eq. (5). This difference originates from the assumption made by the authors that the zipping energy of a SNARE complex is entirely dissipated before the membranes encounter the fusion barrier. In other words, the authors have implicitly assumed that after the calcium entry, the zipping of the SNAREpins does not generate any pulling force to assist fusion and concluded that the remaining residual force is of entropic nature. Recent direct microscopic observations implying that synaptic fusion involves only six SNAREpins (22) would appear to invalidate this assumption.

In conclusion, our model describes membrane fusion by a team of mechanically interacting SNAREpins as a two stage process. We show that conventional biochemical and biophysical measurements cannot be used directly to predict the associated rates and that mechanical modeling is crucial for linking these rates with independently measured parameters. Our work emphasizes the importance of identifying mechanistic pathways and specifying mechanistic feedbacks. The main conceptual outcome of our study is the realization that in the case of synaptic fusion, SNARE proteins can perform opti-

mally only if they act collectively. The remarkable fact is that when the team is of the optimal size, such synchronization is not deterred by thermal fluctuations, which guarantees that the collective strike is simultaneously fast, strong and robust.

Finally, we mention that the synaptic fusion is only one of many biophysical processes involving mechanically-induced collective conformational changes. Other examples include ion gating in hair cells (57, 58), collective decohesion of adhesive clusters (59, 60), folding-unfolding of macromolecular hairpins (61–63) and folding of ParB-ParS complexes in DNA condensation (64, 65). In each of these situations one can identify a dominating long-range mechanical interaction making the theoretical framework developed in this paper potentially useful.

Materials and Methods

A. Rigid membrane assumption. We assume for simplicity that the vesicle and the target membranes are rigid, which implies that all the SNAREpins share the same inter-membrane distance y . This approximation is valid if the characteristic length ℓ associated with the deformation generated by a single SNAREpin is large compare to the size of the SNARE bundle. We can use the following estimate $\ell = \sqrt{\kappa/\sigma}$, where κ is the membrane rigidity and σ is the membrane tension. We have typically $\kappa \sim 20 - 50 \text{ k}_B T$ and $\sigma \sim 10^{-4} - 10^{-6} \text{ N m}^{-1}$ so $\ell \sim 30 - 120 \text{ nm}$. Since the size of the SNARE bundle is less than 10 nm, our assumption should be valid.

B. Model of a single SNAREpin. We set, for simplicity, that the energies e_n and e_c of the SNAREpins in the states n and c , respectively, depend on y quadratically, so that

$$\begin{aligned} e_n(y) &= (\kappa_n/2)(y - a)^2 + e_0, \\ e_c(y) &= (\kappa_c/2)y^2, \end{aligned} \quad [7]$$

where $\kappa_{c,n}$ represent lumped stiffnesses parameters. We denote y_* the distance where $e_n(y_*) = e_c(y_*)$, see Fig. 1(C). In the absence of external load (zero force), the stable states are located at $y = a$ and $y = 0$. In this situation the entire energy associated with the zippering process is consumed when the SNAREpin reaches state c at $y = 0$, which can then be considered as a ground state with zero energy.

The rates k_{\pm} of the $n \rightleftharpoons c$ transitions obey the detailed balance relation $k_+/k_- = \exp[(e_c - e_n)/(k_B T)]$, with the bias towards the direct transition $n \rightarrow c$, i.e. $k_+ > k_-$, at $y < y_*$ and conversely, in the direction of the reverse transition $c \rightarrow n$ at $y > y_*$, see Fig. 1(C)

For simplicity, and following (25), we consider that the transition from the high to the low energy state occurs at a constant rate k , which fixes the characteristic timescale of the conformational change. This assumption could be easily replaced by a more adequate one at the expense of introducing two additional parameters but with only a minimal impact on the results, see Ref. (28). With this assumption and using the detailed balance, we write the transition rates as

$$\begin{aligned} k_-(y) &= k, \quad k_+(y) = k \exp\{[e_n(y) - e_c(y)]/(k_B T)\}, \quad \text{if } y > y_* \\ k_+(y) &= k, \quad k_-(y) = k \exp\{[e_c(y) - e_n(y)]/(k_B T)\}, \quad \text{if } y < y_*. \end{aligned}$$

C. Numerical implementation of the model. The discrete stochastic process associated with the variable N_c was simulated as a two-state Markov chain with a fixed timestep $\Delta t = 10^{-6} t_\eta$. At each timestep the transition probabilities $W_{+1,-1,0} \Delta t$ are computed and the next event is chosen based on an acceptance-rejection condition using a random number uniformly distributed between 0 and 1. The Langevin equation was simulated using a first order explicit Euler scheme. More details about the computer algorithms can be found in SI Appendix.

D. Adiabatic elimination of the variable N_c . We consider the situation where $t_\eta \gg k^{-1}$: the characteristic time of the conformational changes is negligible compared to the timescale associated with the relaxation of the vesicle's position. In this limit, the conformational state of each SNAREpin can be considered at equilibrium. Therefore for a given position of the vesicle y , the probability of a configuration with N_c SNAREpins in state c follows the Boltzmann distribution

$$\rho(N_c; y) = \frac{1}{Z(y)} \binom{N}{N_c} \exp\{-[N_c e_c(y) + (N - N_c) e_n(y)]/(k_B T)\}, \quad [8]$$

where $\binom{N}{N_c} = \frac{N!}{N_c!(N-N_c)!}$. We then integrate Eq. (1) with respect to the distribution (8) and obtain Eq. (3). Since the energy E_{snare} is linear in N_c , our approximation results in replacing N_c by its average $\langle n_c \rangle(y) = \sum_{N_c} N_c \rho(N_c; y)$ in Eq. (1). In Eq. (4), the prefactors are given by (45) $\alpha_{1,2} = \frac{2\pi k_B T}{a^2 \sqrt{\Phi'_{1,2}(y_{\text{max}})|\Phi'_{1,2}(y_{\text{min}})|}}$, where y_{max} and y_{min} denote the positions of the considered barrier and minimum, respectively.

E. Estimation of the mechanical work w . In the intermediate state, inter membrane distance y_2 is sufficiently lower than the threshold y_* , so that the free energy can be well approximated by the energy of the state c . We then write $\Phi(y) \simeq E_{\text{fusion}}(y) + N \frac{\kappa_c}{2} y^2$ which leads to the following expression for the energy barrier separating the intermediate state and the fused state,

$$\Delta\Phi_2 = e_f \left\{ 1 - \exp\left[-(y_2 - y_f)^2/(2\sigma_f^2)\right] \right\} + N \frac{\kappa_c}{2} (y_f^2 - y_2^2).$$

By noting that y_2 verifies $\left. \frac{d\Phi(y)}{dy} \right|_{y=y_2} = 0$, we obtain $\Delta\Phi_2 = e_f - Nw$, with

$$w \simeq \kappa_c \left(y_2^2 - y_f^2 + \frac{\sigma_f^2 y_2}{y_2 - y_f} \right) \geq 0.$$

Notice that since the energy E_{fusion} decays rapidly for $y > y_f$, the parameter y_2 depends weakly on N .

ACKNOWLEDGMENTS. We would like to thank Yongli Zhang for providing us with the data used for the calibration of our model and Ben OShaughnessy for stimulating discussions. This work was supported by a European Research Council (ERC) funded grant under the European Union Qs Horizon 2020 research and innovation programme (grant agreement no. 669612) to J.E.R.

- Ivanov AI (2008) *Exocytosis and Endocytosis*. (Humana, Totowa, NJ, Totowa, NJ) Vol. 440.
- Vassilieva EV, Nusrat A (2008) Vesicular Trafficking: Molecular Tools and Targets in *Exocytosis and Endocytosis, Methods in Molecular Biology*, ed Ivanov AI. (Humana, New York) Vol. 440, pp. 3–14.
- Jahn R, Fasshauer D (2012) Molecular machines governing exocytosis of synaptic vesicles. *Nature* 490(7419):201–207.
- Südhof TC, Rothman JE (2009) Membrane fusion: grappling with SNARE and SM proteins. *Science* 323(5913):474–477.
- Rand RP, Parsegian VA (1989) Hydration forces between phospholipid bilayers. *Biochim. Biophys. Acta Rev. Biomembr.* 988(3):351–376.
- Leckband D, Israelachvili J (2001) Intermolecular forces in biology. *Q. Rev. Biophys.* 34(02):105–267.
- Ryham RJ, Klotz TS, Yao L, Cohen FS (2016) Calculating Transition Energy Barriers and Characterizing Activation States for Steps of Fusion. *Biophys. J.* 110(5):1110–1124.
- François-Martin C, Rothman JE, Pincet F (2017) Low energy cost for optimal speed and control of membrane fusion. *Proc. Natl. Acad. Sci. U.S.A.* 114(6):1238–1241.
- Zhang Y (2017) Energetics, kinetics, and pathway of SNARE folding and assembly revealed by optical tweezers. *Protein Sci.* 26(7):1252–1265.
- Gao Y, et al. (2012) Single reconstituted neuronal SNARE complexes zipper in three distinct stages. *Science* 337(6100):1340–1343.
- Zhang Z, Jackson MB (2008) Temperature Dependence of Fusion Kinetics and Fusion Pores in Ca²⁺-triggered Exocytosis from PC12 Cells. *J. Gen. Physiol.* 131(2):117–124.
- Xu W, et al. (2016) A Programmable DNA Origami Platform to Organize SNAREs for Membrane Fusion. *J. Am. Chem. Soc.* 138(13):4439–4447.
- Domanska MK, Kiessling V, Stein A, Fasshauer D, Tamm LK (2009) Single Vesicle Millisecond Fusion Kinetics Reveals Number of SNARE Complexes Optimal for Fast SNARE-mediated Membrane Fusion. *J. Biol. Chem.* 284(46):32158–32166.
- Südhof TC (2004) The synaptic vesicle cycle. *Annu. Rev. Neurosci.* 27(1):509–547.
- Camacho M, et al. (2017) Heterodimerization of Munc13 C2A domain with RIM regulates synaptic vesicle docking and priming. *Nat. Commun.* 8:15293.

16. Liu X, et al. (2016) Functional synergy between the Munc13 C-terminal C1 and C2 domains. *Elife* 5:965.
17. Li F, Tiwari N, Rothman JE (2016) Kinetic barriers to SNAREpin assembly in the regulation of membrane docking/priming and fusion. *Proc. Natl. Acad. Sci. U.S.A.* pp. 10536–10541.
18. Krishnakumar SS, et al. (2015) Re-visiting the trans insertion model for complexin clamping. *Elife* 4:1021.
19. Hua Y, Scheller RH (2001) Three SNARE complexes cooperate to mediate membrane fusion. *Proc. Natl. Acad. Sci. U.S.A.* 98(14):8065–8070.
20. Sinha R, Ahmed S, Jahn R, Klingauf J (2011) Two synaptobrevin molecules are sufficient for vesicle fusion in central nervous system synapses. *Proc. Natl. Acad. Sci. U.S.A.* 108(34):14318–14323.
21. Mohrmann R, de Wit H, Verhage M, Neher E, Sørensen JB (2010) Fast Vesicle Fusion in Living Cells Requires at Least Three SNARE Complexes. *Science* 330(6003):502–505.
22. Li X, et al. (December 18, 2018) Symmetrical organization of proteins under docked synaptic-vesicles. *FEBS Lett.*, 10.1002/1873-3468.13316.
23. Mostafavi H, et al. (2017) Entropic forces drive self-organization and membrane fusion by SNARE proteins. *Proc. Natl. Acad. Sci. U.S.A.* 84(21):201611506–5460.
24. McDargh ZA, Polley A, O'Shaughnessy B (2018) SNARE-mediated membrane fusion is a two-stage process driven by entropic forces. *FEBS Lett.* 592(21):3504–3515.
25. Huxley AF, Simmons RM (1971) Proposed mechanism of force generation in striated muscle. *Nature* 233(5321):533–538.
26. Caruel M, Allain JM, Truskinovsky L, Truskinovsky L (2013) Muscle as a metamaterial operating near a critical point. *Phys. Rev. Lett.* 110(24):248103.
27. Caruel M, Truskinovsky L (2016) Statistical mechanics of the Huxley-Simmons model. *Phys. Rev. E* 93(6):062407.
28. Caruel M, Truskinovsky L (2017) Bi-stability resistant to fluctuations. *J. Mech. Phys. Solids* 109:117–141.
29. Caruel M, Truskinovsky L (2018) Physics of muscle contraction. *Rep. Prog. Phys.* 81(3):036602.
30. Sutton RB, Fasshauer D, Jahn R, Brunger AT (1998) Crystal structure of a SNARE complex involved in synaptic exocytosis at 2.4Å resolution. *Nature* 395(6700):347–353.
31. Evans E (1991) Entropy-driven tension in vesicle membranes and unbinding of adherent vesicles. *Langmuir* 7(9):1900–1908.
32. Donaldson SH, Lee CT, Chmelka BF, Israelachvili JN (2011) General hydrophobic interaction potential for surfactant/lipid bilayers from direct force measurements between light-modulated bilayers. *Proc. Natl. Acad. Sci. U.S.A.* 108(38):15699–15704.
33. Südhof TC (2013) Neurotransmitter Release: The Last Millisecond in the Life of a Synaptic Vesicle. *Neuron* 80(3):675–690.
34. Yang WY, Gruebele M (2003) Folding at the speed limit. *Nature* 423(6936):193–197.
35. Lentz BR, Lee J (2009) Poly(ethylene glycol) (PEG)-mediated fusion between pure lipid bilayers: a mechanism in common with viral fusion and secretory vesicle release? (Review). *Mol. Membr. Biol.* 16(4):279–296.
36. Cohen FS, Melikyan GB (2004) The Energetics of Membrane Fusion from Binding, through Hemifusion, Pore Formation, and Pore Enlargement. *J. Membr. Biol.* 199(1):1–14.
37. Markvoort AJ, Marrink SJ (2011) Lipid acrobatics in the membrane fusion arena. *Curr Top Membr* 68:259–294.
38. Finkelstein A, Zimmerberg J, Cohen FS (1986) Osmotic swelling of vesicles: its role in the fusion of vesicles with planar phospholipid bilayer membranes and its possible role in exocytosis. *Annual Review of Physiology* 48(1):163–174.
39. Grafmüller A, Shillcock J, Lipowsky R (2007) Pathway of membrane fusion with two tension-dependent energy barriers. *Phys. Rev. Lett.* 98(21):218101.
40. Grafmüller A, Shillcock J, Lipowsky R (2009) The fusion of membranes and vesicles: pathway and energy barriers from dissipative particle dynamics. *Biophysical journal* 96(7):2658–2675.
41. Lee JY, Schick M (2009) Calculation of free energy barriers to the fusion of small vesicles. *Biophysical journal* 94(5):1699–1706.
42. Meinrenken CJ, Borst JGG, Sakmann B (2002) Calcium Secretion Coupling at Calyx of Held Governed by Nonuniform Channel–Vesicle Topography. *J. Neurosci.* 22(5):1648–1667.
43. Kramers H (1940) Brownian motion in a field of force and the diffusion model of chemical reactions. *Physica* 7:284–304.
44. Risken H (1988) *The Fokker-Planck equation Methods of solution and application.* (Springer, Berlin).
45. Schuss Z (2010) *Theory and application of stochastic processes: an analytical approach.* (Springer, Berlin).
46. Caruel M, Allain JM, Truskinovsky L (2015) Mechanics of collective unfolding. *J. Mech. Phys. Solids* 76:237–259.
47. Walter AM, Wiederhold K, Bruns D, Fasshauer D, Sørensen JB (2010) Synaptobrevin N-terminally bound to syntaxin–SNAP-25 defines the primed vesicle state in regulated exocytosis. *J. Cell Biol.* 188(3):401–413.
48. Li F, et al. (2007) Energetics and dynamics of SNAREpin folding across lipid bilayers. *Nat Struct Mol Biol* 14(10):890–896.
49. Risselada HJ, Grubmüller H (2012) How snare molecules mediate membrane fusion: recent insights from molecular simulations. *Current opinion in structural biology* 22(2):187–196.
50. Acuna C, et al. (2014) Microsecond Dissection of Neurotransmitter Release: SNARE-Complex Assembly Dictates Speed and Ca²⁺ Sensitivity. *Neuron* 82(5):1088–1100.
51. Hernandez JM, et al. (2012) Membrane Fusion Intermediates via Directional and Full Assembly of the SNARE Complex. *Science* 336(6088):1581–1584.
52. Li F, et al. (2014) A Half-Zippered SNARE Complex Represents a Functional Intermediate in Membrane Fusion. *J Am Chem Soc* 136(9):3456–3464.
53. Stein A, Weber G, Wahl MC, Jahn R (2009) Helical extension of the neuronal SNARE complex into the membrane. *Nature* 460(7254):525–528.
54. Ford LE, Ford LE, Huxley AF, Simmons RM (1977) Tension responses to sudden length change in stimulated frog muscle fibres near slack length. *J. Physiol.* 269(2):441–515.
55. Ramakrishnan S, et al. (2019) Synaptotagmin oligomers are necessary and can be sufficient to form a stable fusion clamp. *FEBS Lett.* in press.
56. Gruget C, et al. (2018) Rearrangements under confinement lead to increased binding energy of Synaptotagmin-1 with anionic membranes in Mg²⁺ and Ca²⁺. *FEBS Lett.* 592(9):1497–1506.
57. Martin P, Martin P, Mehta A, Hudspeth A (2000) Negative hair-bundle stiffness betrays a mechanism for mechanical amplification by the hair cell. *Proc. Natl. Acad. Sci. U.S.A.* 97(22):12026–12031.
58. Bormuth V, Barral J, Joanny JF, Jülicher F, Martin P (2014) Transduction channels' gating can control friction on vibrating hair-cell bundles in the ear. *Proc. Natl. Acad. Sci. U.S.A.* 111(20):7185–7190.
59. Yao H, Gao H (2006) Mechanics of robust and releasable adhesion in biology: Bottom-up designed hierarchical structures of gecko. *J. Mech. Phys. Solids* 54(6):1120–1146.
60. Erdmann T, Schwarz US (2007) Impact of receptor-ligand distance on adhesion cluster stability. *Eur. Phys. J.* 22(2):123–137.
61. Woodside MT, Garcia-Garcia C, Block SM (2008) Folding and unfolding single RNA molecules under tension. *Curr. Opin. Chem. Biol.* 12(6):640–646.
62. Bosaeus N, et al. (2012) Tension induces a base-paired overstretched dna conformation. *Proc. Natl. Acad. Sci. U.S.A.* 109(38):15179–15184.
63. Liphardt J, Onoa B, Smith SE, Tinoco I, Bustamante C (2001) Reversible unfolding of single rna molecules by mechanical force. *Science* 292(5517):733–737.
64. Chen BW, Lin MH, Chu CH, Hsu CE, Sun YJ (2015) Insights into ParB spreading from the complex structure of Spo0J and parS. *Proc. Natl. Acad. Sci. U.S.A.* 112(21):6613–6618.
65. Funnell BE (2016) ParB Partition Proteins: Complex Formation and Spreading at Bacterial and Plasmid Centromeres. *Front. Mol. Biosci.* 3:30067–6.