



HAL
open science

CO₂ Capture and Storage by Hydroquinone Clathrate Formation: Thermodynamic and Kinetic Studies

Romuald Coupan, Christophe Dicharry, Frédéric Plantier, Christelle Miqueu, Pierre Kaplan, Eve Péré, Mehrdji Hemati, Jean-Philippe Torrè

► **To cite this version:**

Romuald Coupan, Christophe Dicharry, Frédéric Plantier, Christelle Miqueu, Pierre Kaplan, et al.. CO₂ Capture and Storage by Hydroquinone Clathrate Formation: Thermodynamic and Kinetic Studies. 9th International Conference on Gas Hydrates (ICGH9), Jun 2017, Denver, United States. pp.0. hal-02006335

HAL Id: hal-02006335

<https://hal.science/hal-02006335>

Submitted on 4 Feb 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Open Archive Toulouse Archive Ouverte

OATAO is an open access repository that collects the work of Toulouse researchers and makes it freely available over the web where possible

This is an author's version published in: <http://oatao.univ-toulouse.fr/21871>

To cite this version:

Coupan, Romuald and Dicharry, Christophe and Plantier, Frédéric and Miqueu, Christelle and kaplan, Pierre and Péré, Eve and Hemati, Mehrdji and Torr , Jean-Philippe CO2 Capture and Storage by Hydroquinone Clathrate Formation: Thermodynamic and Kinetic Studies. (2017) In: 9th International Conference on Gas Hydrates (ICGH9), 25 June 2017 - 30 June 2017 (Denver, United States).

Any correspondence concerning this service should be sent to the repository administrator: tech-oatao@listes-diff.inp-toulouse.fr

CO₂ Capture and Storage by Hydroquinone Clathrate Formation: Thermodynamic and Kinetic Studies.

Romuald Coupan^{1*}, Christophe Dicharry¹, Frédéric Plantier¹, Christelle Miqueu¹, Pierre Kaplan², Eve Péré³, Mehrdji Hemati⁴, Jean-Philippe Torr ¹.

¹Laboratoire des Fluides Complexes et leurs R servoirs (UMR 5150), France; ²TOTAL E&P, Centre Scientifique et Technique Jean F ger, France ; ³Institut des Sciences Analytiques et de Physico-Chimie pour l'Environnement et les Mat riaux (UMR 5254), France; ⁴Laboratoire de G nie Chimique (UMR 5503), France.

*Corresponding author: romuald.coupan@univ-pau.fr

Abstract

Hydroquinone (HQ) can form a gas clathrate in specific pressure and temperature conditions in the presence of CO₂ molecules. This study presents experimental data of clathrate phase equilibrium and storage capacity for the CO₂-HQ system in the range of temperature from about 288 to 354 K. Intercalation enthalpy and entropy are determined using the obtained equilibrium data and the Langmuir adsorption model. On a kinetic point of view, CO₂-HQ clathrate formation by solid/gas reaction revealed a non-negligible effect of textural parameters on enclathration rate.

Introduction

Gas clathrates are inclusion compounds formed with host molecules self-associating and forming cages, able to retain low molecular-weight guest molecules of gas, which have recently been highlighted as an alternative way for gas storage, transportation and separation [1-2]. Gas hydrates show high gas storage capacity. However, for potential industrial applications, the hydrate-based processes require a non-negligible energetic cost as gas hydrates are usually formed at low temperature, typically a few degrees above 273 K, and high pressure of several MPa [3]. To overcome this problem, hydroquinone (HQ) has been pointed out because of its capability to form organic clathrates over a wide range of temperature and at moderate pressure (a few MPa) [4-6]. The stable form of HQ at ambient conditions of pressure and temperature is the α -HQ. Whereas, the clathrate form is the β -HQ. The maximum storage capacity of β -HQ is defined by its ideal stoichiometry of 1 molecule of gas per 3 molecules of HQ, if all the cavities are filled by one guest molecule [7]. Thus, for CO₂-HQ clathrates the maximum theoretical quantity of gas which can be stored is 67.8 L (STP) / kg^{HQ}.

This work presents experimental data on phase equilibria and kinetics of CO₂-HQ clathrate formation. Such results could be of interest for potential gas storage, transportation, or separation applications that could be developed with this compound. The equilibrium curve, the clathrate occupancy and the intercalation enthalpy and entropy are determined in the range of temperature from about 288 to 354 K. Ways for kinetic improvements are proposed and discussed. Interestingly, the use of HQ-silica composite materials as reactive medium has shown that the kinetic of HQ clathrate formation can be significantly improved.

Experimental section

Materials

HQ with purity of 99.5 mol% is provided by Acros Organics. The solvents used for the titration and impregnation experiments are butyl acetate and absolute ethanol (purities higher than 99 mol%). CO₂ gas used for the experiments (mole fraction purity of 99.995%) is purchased from Linde Gas SA. The porous supports are analytical grade spherical silica particles (SiliaSphere®) sizing 200-500 μ m with pore size of 100 nm provided by SiliCycle.

Apparatus and Procedures

The experimental apparatus used for determining equilibrium conditions of clathrates is composed by a jacketed and stirred high pressure crystallizer connected to a gas storage tank, a solvent container, and a vacuum pump [8]. This experimental set-up allows achieving isothermal titration [8], which is the method used to determine both the HQ $_{\beta}$ -HQ $_{\alpha}$ -CO $_2$ equilibrium curve and the clathrate occupancy.

The capture kinetic and the gas storage capacity measurements are performed by a gravimetric method [9] at 3.0 MPa and 323 K, using a Rubotherm magnetic suspension balance. For these measurements, the HQ is conditioned in two ways in an attempt to increase the clathrate formation kinetics. Pure HQ is ground or deposited on porous support.

The HQ deposition on porous silica support is performed in fluidized bed in a dry impregnation process [10]. This method consists in intermittently spraying a warm fluidized bed of porous silica particles with a HQ solution, allowing the coupling of the HQ solution penetration in the particle porosity and the solvent evaporation.

Results and discussion

Phase Equilibria and occupancies

The experimental equilibrium data for the CO $_2$ -HQ systems are obtained in this work from isothermal titration experiments [8]. The equilibrium curve and the clathrate occupancies at these equilibrium conditions are shown in Figure 1. Compared to the CO $_2$ hydrates equilibrium curve [11-12], it can be noted that the CO $_2$ -HQ clathrates can be formed over a wide range of temperature and at pressures not exceeding 1 MPa for temperatures close to 360 K. Regarding the obtained clathrate occupancies, it is obvious that these ones are temperature dependent. The CO $_2$ -HQ clathrate occupancy decreases with temperature, as already pointed out in literature for other HQ clathrates [13].





Figure 1: Equilibrium data for CO $_2$ -HQ system: \blacklozenge , clathrate equilibrium curve; \blacktriangle , clathrate occupancies as a function of temperature.

As there is an analogy between enclathration and adsorption, the gas-clathrate equilibrium obey a Langmuir type isotherm [13-15], where the involved reaction, given by Equation 1, corresponds to the liberation of the CO $_2$ guest molecules from the β -HQ host lattice.



The Langmuir constant C_{lang} related to this reaction (i.e. the thermodynamic equilibrium constant) depends on clathrate occupancy θ and equilibrium pressure P_{eq} , as described by Equation 2.

$$C_{\text{lang}} = P_{\text{eq}} \cdot (1 - \theta) / \theta \quad (2)$$

Thus, the intercalation enthalpy and entropy can be determined from the slope and the intercept of the linear fit of the logarithm of the Langmuir constant as a function of the inverse of equilibrium temperature (Figure 2) [13-14]. The

obtained enthalpy is $42.3 \pm 4.0 \text{ kJ/mol}^{\text{CO}_2}$ ($14.1 \pm 1.3 \text{ kJ/mol}^{\text{HQ}}$) and the associated entropy is $235 \pm 11 \text{ J/mol}^{\text{CO}_2/\text{K}}$ ($78 \pm 4 \text{ J/mol}^{\text{HQ}/\text{K}}$). To determine the total dissociation enthalpy of the CO_2 -HQ clathrate, it is necessary to consider the enthalpy of the reversion of the residual β -HQ to the α -HQ structure of about $-0.67 \text{ kJ/mol}^{\text{HQ}}$ [16]. As a result, these data indicate that the CO_2 can be recovered more easily from CO_2 -HQ clathrates than from CO_2 hydrates as the dissociation enthalpy is about $60.6 \pm 1.8 \text{ kJ/mol}^{\text{CO}_2}$ ($10.0 \pm 0.3 \text{ kJ/mol}^{\text{H}_2\text{O}}$) between the quadruples points Q1 and Q2 of this system [17].




Figure 2: Linear fit of the logarithm of the Langmuir constants of CO_2 -HQ clathrates as a function of the inverse of temperature.

Kinetic aspects

When the gas clathrate is formed by direct gas/solid reaction between the CO_2 and the HQ, the clathrate formation kinetics seems to be enhanced by increasing the gas/solid contact area [4-6]. Thus, working on the HQ conditioning (i.e. the increase of the specific area of the media and the improvement of the contact area between the CO_2 and the HQ) is very important to achieve rapid enclathration kinetics.

In this work, we have tested and compared the kinetic performances of HQ powder and of a HQ-silica composite material. The grinding of HQ gives a powder with a particle size of about $100 \mu\text{m}$. The HQ deposition on porous silica support allows developing a composite material having the HQ content of $0.44 \text{ g}^{\text{HQ}}/\text{g}^{\text{Silica}}$.

As shown in Figure 3 by SEM images of the obtained composite, it appears clearly that HQ coats uniformly the external surface of the silica particle. Moreover, HQ crystals could be present in the 100 nm pores. Indeed, as suggested by Hemati *et al.* [10] the growth phenomenon does not arise before the full filling of internal porosity.




Figure 3: SEM images of HQ-silica composite material: (a) full particle and (b) external surface.

The native, ground, and impregnated HQ are evaluated as media in CO_2 capture experiments. The amount of CO_2 captured by enclathration as a function of time is shown in Figure 4. For HQ-silica composite materials, the amount of CO_2 likely to be adsorbed on native silica (i.e. $0.34 \text{ mol}^{\text{CO}_2}/\text{kg}^{\text{Silica}}$) has been deduced from the total CO_2 gas

captured. It is worth noting that the two conditionings improve the kinetic of clathrate formation, which highlights the importance of CO₂/HQ contact area.




Figure 4: Molar quantity of CO₂ captured by enclathration as a function of time normalized by mass of HQ: (full line) native HQ, (dashed line) ground HQ, and (dotted line) HQ impregnated on silica particles.

Table 1 presents the kinetic criteria deduced from the gas capture measurements: (i) the reaction rate for the CO₂ solubilization in the α -HQ, (ii) the induction time, (iii) the enclathration reaction rate, and (iv) the characteristic time at which the clathrate occupancy is 50%.

Compared to native HQ, although grinding of HQ seems kinetically interesting, it is obvious that the HQ-silica composite material is the most efficient medium for enclathration. Indeed, for this reactive medium there is no induction period, the enclathration reaction rate is increased by a factor 52, and the time to fill 50% of the clathrate cavities is decreased by a factor 55 compared to native HQ.

Moreover, for HQ-silica composite material, the system reaches an equilibrium value of clathrate occupancy of 0.89 after approximately 3.5 days, whereas the clathrate occupancies are 0.18 and 0.34 for native and ground HQ, respectively at the same time. After about 1 month of reaction, it is found occupancies of 0.63 and 0.71 for native and ground HQ, respectively. These values are in agreement with the ones previously found for CO₂-HQ clathrates mono-crystals synthesized by crystallization from solvent [18].

Table 1: Kinetic data on the CO₂ capture by HQ clathrate formation.

HQ conditioning	r_0 (mmol ^{CO2} /kg ^{HQ} /min)	t^{ind} (h)	r_c (mmol ^{CO2} /kg ^{HQ} /min)	t_c^{50} (h)
Native	0.02	23.4	0.32	356.5
Grinding	0.16	4.2	0.52	218.9
Impregnation	-	-	16.5	6.5

Conclusion

This study brings some information on the potentiality of HQ clathrates for CO₂ capture and storage process. The measured equilibrium conditions suggest the possibility to overcome specific process limitation, as CO₂-HQ clathrates can be formed in a wide range of temperature (reaching at least 354 K) and at moderate pressure (few MPa). Furthermore, on a kinetic point of view, it is shown that the enclathration kinetic can be improved by a specific conditioning of HQ allowing to increase the gas/solid contact area. The induction period can be avoided with a non-negligible increase of enclathration rate using HQ-silica composite materials. Now, some works are necessary to evaluate the viability and feasibility of such reactive media in clathrate-based processes.

Acknowledgment

The working group involved in the ORCHIDS project, the Carnot Institute ISIFoR (Institute for the Sustainable Engineering of Fossil Resources), and Total E&P, “Gas Solutions” department are gratefully acknowledged.

References

- [1] Eslamimanesh, A. H. Mohammadi, D. Richon, P. Naidoo, D. Ramjugernath, « Application of gas hydrate formation in separation processes: A review of experimental studies”, *Journal of Chemical Thermodynamics* **2012**, 46, 62–71.
- [2] P. Babu, P. Linga, R. Kumar, P. Englezos, “A review of the hydrate based gas separation (HBGS) process for carbon dioxide pre-combustion capture”, *Energy* **2015**, 85, 261-279.
- [3] M. K. Mondal, H. K. Balsora, P. Varshney, “Progress and trends in CO₂ capture/separation technologies: A review”, *Energy* **2012**, 46, 431-441.
- [4] Y.-J. Lee, K.W. Han, J.S. Jang, T.-I. Jeon, J.Park, T. Kawamura, Y. Yamamoto, T. Sugahara, T. Vogt, J.-W. Lee, Y. Lee, J.-H. Yoon, “Selective CO₂ trapping in guest-free hydroquinone clathrate prepared by gas phase synthesis”, *ChemPhysChem* **2011**, 12, 1056 – 1059.
- [5] J.-W. Lee, P. Dotel, J. Park, J.-H. Yoon, “Separation of CO₂ from flue gases using hydroquinone clathrate compounds”, *Korean Journal of Chemical Engineering* **2015**, 12, 2507–2511.
- [6] J.-W. Lee, P. Dotel, J. Park, J.-H. Yoon, “Separation of CO₂ from flue gases using hydroquinone clathrate compounds”, *Korean Journal of Chemical Engineering* **2015**, 12, 2507–2511.
- [7] J. W. Steed, D. R. Turner, K. J. Wallace, “Core concepts in supramolecular chemistry and nanochemistry”, John Wiley & Sons: Chichester, England, **2007**, 179-194.
- [8] R. Coupan, M. Chabod, C. Dicharry, J. Diaz, C. Miqueu, J.-P. Torr , “Experimental determination of phase equilibria and occupancies for CO₂, CH₄, and N₂ hydroquinone clathrates”, *Journal of Chemical Engineering Data* **2016**, 61, 2565–2572.
- [9] F. Khaddour, A. Knorst-Fouran, F. Plantier, M. M. Pi eiro, B. Mendiboure, C. Miqueu, “A fully consistent experimental and molecular simulation study of methane adsorption on activated carbon”, *Adsorption* **2014**, 20, 649–656.
- [10] M. Hemati, R. Cherif, K. Saleh, V. Pont, “Fluidized bed coating and granulation: influence of process-related variables and physicochemical properties on the growth kinetics”, *Powder Technology* **2003**, 130, 18–34.
- [11] S. Adisasmito, R. J. Frank, E. D. Sloan, “Hydrates of carbon dioxide and methane mixtures”, *Journal of Chemical Engineering Data* **1991**, 36, 68-71.
- [12] A. Delahaye, L. Fournaison, S. Marinhas, I. Chatti, “Effect of THF on Equilibrium Pressure and Dissociation Enthalpy of CO₂ Hydrates Applied to Secondary Refrigeration”, *Industrial & Engineering Chemistry Research* **2006**, 45, 391-397.
- [13] R. L. Deming, T. L. Carlisle, B. J. Lauerman, J. T. Mucker-man, A. R. Muirhead, W. C. Child Jr., “Dissociation pressures and compositions of quinol clathrates of methane and methyl fluoride”, *Journal of Physical Chemistry* **1969**, 73, 1762–1766.
- [14] M. M. Conde, J.-P. Torr , C. Miqueu, “Revisiting the thermodynamic modelling of type I gas-hydroquinone clathrates”, *Physical Chemistry Chemical Physics* **2016**, 18, 10018–10027.

- [15] J. H. Van der Waals, J. C. Platteeuw, "Clathrate Solutions", *Advances in Chemical Physics* **1959**, 2, 1–57.
- [16] D. F. Evans, R. E. Richards, "Energies of interaction and others properties of clathrate compounds", *Proceedings of the Royal Society A* **1954**, 223, 238–250.
- [17] G. K. Anderson, "Enthalpy of dissociation and hydration number of carbon dioxide hydrate from the Clapeyron equation", *The Journal of Chemical Thermodynamics* **2003**, 35, 1171–1183.
- [18] J.-P. Torr , R. Coupan, M. Chabod, E. P r , S. Labat, A. Khoukh, R. Brown, J.-M. Sotiropoulos, H. Gornitzka, "CO₂ – Hydroquinone clathrate: Synthesis, Purification, Characterization and Crystal Structure", *Crystal Growth & Design* **2016**, 16, 5330–5338.