



HAL
open science

Estimating the robust domain of attraction for non-smooth systems using an interval Lyapunov equation

Alexandre Goldsztejn, Gilles Chabert

► **To cite this version:**

Alexandre Goldsztejn, Gilles Chabert. Estimating the robust domain of attraction for non-smooth systems using an interval Lyapunov equation. The 18th International Symposium on Scientific Computing, Computer Arithmetic, and Verified Numerical Computations, Sep 2018, Tokyo, Japan. 10.1016/j.automatica.2018.03.036 . hal-02008057

HAL Id: hal-02008057

<https://hal.science/hal-02008057v1>

Submitted on 5 Feb 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Estimating the robust domain of attraction for non-smooth systems using an interval Lyapunov equation

Alexandre Goldsztejn^{1,3} and Gilles Chabert²

¹CNRS, LS2N, Ecole Centrale de Nantes, France

²LS2N, Institut des Mines Télécom Atlantique, France

³alexandre.goldsztejn@ls2n.fr

Keywords: Nonlinear non-smooth systems; exponentially stable fixed point; domain of attraction; Lyapunov function; interval analysis.

Contribution

A symmetric positive definite (SPD) matrix P solution to the Lyapunov equation $PA + A^T P = Q$, with Q an arbitrary symmetric negative definite (SND) matrix, gives rise to a quadratic Lyapunov function $v(x) = x^T P x$, whose existence proves that the origin is an exponentially globally stable fixed point of the linear system $\dot{x} = Ax$. Solving the Lyapunov equation for the linearization of a nonlinear system can also prove the exponential stability of its fixed points. However, in this case the stability is local and the Lyapunov equation provides no information about the region of attraction.

We present an interval version of the Lyapunov equation, which allows investigating a given Lyapunov candidate function for nonlinear systems inside an explicitly given neighborhood, leading to rigorous estimates of the domain of attraction (EDA) of exponentially stable fixed points. These results are developed in the context of uncertain systems that can be non-smooth, e.g., systems with saturations. In a second step, this EDA is used to define a nonlinear optimization problem whose solution gives rise to the largest EDA (LEDA) that can be inferred using the given Lyapunov function.

A comparison with the state of the art for computing EDA based on Lyapunov functions will be presented. In particular, the proposed process is shown to outperform the approach based on sum-of-squares [1] implemented in SMRSOFT [2].

Example

We consider the following linear system with saturated input, investigated in [3]: $\dot{x}_1 = x_2$ and $\dot{x}_2 = x_1 + 5 \max\{-1, \min\{1, -2x_1 - x_2\}\}$. Linearizing the system at the stable fixed point 0 and solving the corresponding




Figure 1: The vector field of the saturated linear system. In green, the non-saturated area. The following EDA are displayed: In blue the one from [3], which is the convex hull of two ellipsoids; the gray ellipsoid (resp. black curve) is the boundary of the LEDA computed with the quadratic (resp. degree 4) Lyapunov functions.

Lyapunov equation, we select the Lyapunov candidate function $v_2(u) = 1.28x_1^2 + 0.11x_1x_2 + 0.11x_2^2$. We also consider the degree 4 Lyapunov candidate function $v_4(x) = v_2(x) + \frac{1}{15}(x_1 + x_2)^4$, aiming at a larger EDA.

The results obtained are summarized in Figure 1: The LEDA obtained using the quadratic Lyapunov function is computed in $0.05s$ (the gray ellipsoid). It is smaller than the one obtained with the degree 4 Lyapunov function, which is computed in $0.2s$ (the black curve). The latter is smaller, but still competitive with the one computed in [3] (the blue area) using a technic dedicated to linear systems with saturations.

References

- [1] Graziano Chesi. *Domain of Attraction: Analysis and Control via SOS Programming*, volume 415 of *Lecture Notes in Control and Information Sciences*. Springer, 2011.
- [2] Graziano Chesi. SMRSOFT: a matlab toolbox for optimization over polynomials and dynamical systems study via sos programming. <http://www.eee.hku.hk/~chesi>, 2011.

- [3] Tingshu Hu, Rafal Goebel, Andrew R. Teel, and Zongli Lin. Conjugate Lyapunov functions for saturated linear systems. *Automatica*, 41(11):1949 – 1956, 2005.