



HAL
open science

MEAN FIELD LIMIT FOR CUCKER-SMALE MODELS

R. Natalini, Thierry Paul

► **To cite this version:**

R. Natalini, Thierry Paul. MEAN FIELD LIMIT FOR CUCKER-SMALE MODELS. 2020. hal-02008699v2

HAL Id: hal-02008699

<https://hal.science/hal-02008699v2>

Preprint submitted on 9 Aug 2020 (v2), last revised 5 Feb 2021 (v5)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ON THE MEAN FIELD LIMIT FOR CUCKER-SMALE MODELS

R. NATALINI AND THIERRY PAUL

ABSTRACT. In this very short note, we consider the Cucker-Smale dynamical system and we derive rigorously the Vlasov-type equation introduced in [8] in the mean-field limit without using empirical measures. The vector field we consider is sublinear Lipschitz continuous and the initial data are compactly supported.

CONTENTS

1. Introduction	1
2. The results	3
3. Proofs	6
3.1. Preliminaries	6
3.2. Proof of Theorem 2.4	7
3.3. Proof of Corollary 2.6	10
3.4. Proof of Theorem 2.2	11
Appendix A. Dynamical estimates for general Cucker-Smale particle systems	12
Appendix B. Dynamical estimates for general Cucker-Smale kinetic systems	13
References	14

1. INTRODUCTION

The Cucker-Smale model [2, 3] is a very simple particles system which exhibit the so-called flocking phenomenon, namely the dynamical property of alignment of all the velocities and gathering of all the positions asymptotically when the time of evolution diverges. Its study has develop an intense activity these last years, see for example the intensive bibliography in the recent paper [6].

It consists in the following vector field on \mathbf{R}^{2dN}

$$(1) \quad \begin{cases} \dot{x}_i &= v_i \\ \dot{v}_i &= \frac{1}{N} \sum_{j=1}^N \psi(x_i - x_j)(v_i - v_j). \end{cases} \quad x_i, v_i \in \mathbf{R}^d, \quad i = 1, \dots, N$$

where $\psi : \mathbf{R}^d \rightarrow \mathbf{R}$ is a bounded Lipschitz continuous function.

Associated to (1) was introduced in [8] the following Vlasov type kinetic equation on \mathbf{R}^{2d} .

$$(2) \quad \partial_t f^t(x, v) + v \cdot \nabla_x f^t(x, v) = \nabla_v \left(f^t(x, v) \int_{\mathbf{R}^{2d}} \psi(x - y)(v - w) f^t(y, w) dy dw \right).$$

So far the kinetic non-linear equation (2) has been derived in a Lagrangian point of view, i.e. by following the trajectories solving the vector field (1) in the so-called empirical measure

$$(3) \quad \Pi_{(x_1, \dots, x_n; v_1, \dots, v_N)}(x, v) := \frac{1}{N} \sum_{i=1}^N \delta(x - x_i) \delta(v - v_i).$$

Indeed, defining

$$(4) \quad \Pi_{(x_1, \dots, x_n; v_1, \dots, v_N)}^t := \Pi_{\Phi^{-t}(x_1, \dots, x_n; v_1, \dots, v_N)},$$

one shows easily that $\Pi^t(x_1, \dots, x_n; v_1, \dots, v_N)$ solves (2) for each N .

This point of view follows directly the Dobrushin way of deriving the Vlasov equation for the large N limit of Hamiltonian vector fields. CS model (1) and its Vlasov type associated equation (2) have been extensively studied in [6] and we quote the following of their results: the function ψ is supposed to be positive, bounded, uniformly Lipschitz continuous and satisfying $(\psi(r_1) - \psi(r_2))(r_1 - r_2) \leq 0, r_1, r_2 > 0$.

(1) the solutions of (1) satisfy for all t

$$\sup_{i,j \leq N} \|x_i(t) - x_j(t)\| \leq C \quad \text{and} \quad \sup_{i,j \leq N} \|v_i(t) - v_j(t)\| \leq e^{-Dt}$$

where C and D are two positive constants depending only on

$\sup_{i,j \leq N} \|x_i(0) - x_j(0)\|$ and $\sup_{i,j \leq N} \|v_i(0) - v_j(0)\|$.

(2) let $\Pi_{(x_1, \dots, x_n; v_1, \dots, v_N)} \rightarrow \mu(0)$ as $N \rightarrow \infty$, then

$$\overline{\lim}_{N \rightarrow \infty} \sup_{t \in \mathbf{R}} W_p(\Pi_{(x_1, \dots, x_n; v_1, \dots, v_N)}^t - \mu(t)) = 0$$

where W_p is the Wasserstein distance of exponent $p \in \mathbf{N}$ (see definition in Remark 3.5 below).

(3) Let $\int_{\mathbf{R}^{2d}} \mu(0) = 1$, $\int_{\mathbf{R}^{2d}} v \mu(0) =: \bar{v}$, $\int_{\mathbf{R}^{2d}} v^2 \mu(0) < \infty$. Then, for some $E, F > 0$ and any p ,

$$\left(\int \|v - \bar{v}\|^p d\mu(t) \right)^{\frac{1}{p}} \leq E e^{-Ft}.$$

Recently was introduced a more Eulerian way of deriving the Vlasov equation associated to Hamiltonian vector fields ([5], after [4]). It consists in considering the movement of a generic particle solution of the Liouville equation associated to an Hamiltonian system, rather than considering the empirical measure associated to the Hamiltonian flow. Transposed in the (non Hamiltonian) present situation it reads as follows.

We will consider the pushforward by the flow Φ^t generated by the system (1) of a compactly supported, symmetric by permutations of the variables N -body probability density ρ_N^{in} on phase space \mathbf{R}^{2dN} .

In order to describe the movement of a ‘‘generic’’ particle in the limit of diverging number of particles, we want to perform an average on the N particles but one. This means that we consider the first marginal of

$$(5) \quad \Phi^t \# \rho_N^{in} := \rho_N^t$$

that is

$$(6) \quad \rho_{N;1}^t(x, \xi) := \int_{\mathbf{R}^{2(d-1)N}} \rho_N^t(x, \xi, x_2, \xi_2, \dots, x_N, \xi_N) dx_2 d\xi_2 \dots dx_N d\xi_N.$$

Then we will prove that, when ρ_N^{in} is factorized as $\rho_N^{in} = (\rho^{in})^{\otimes N}$, where ρ^{in} is a probability measure on \mathbf{R}^{2d} , the marginal of ρ_n^t tends, as $N \rightarrow \infty$, to the solution of the Vlasov type effective non-linear equation (2).

$$(7) \quad \partial_t \rho^t(x, v) + v \cdot \nabla_x \rho^t = \nabla_v \cdot (G_{\rho^t} \rho^t), \quad \rho^{t=0} = \rho^{in} \in L^1(\mathbf{R}^{2d}, dx dv),$$

with

$$(8) \quad G_{\rho}(x, v) = \int_{\mathbf{R}^{2d}} \psi(x - y)(v - w) \rho(y, w) dy dw.$$

In fact, we can prove (see Remark 3.5 below) that the marginals at any order n of ρ_N^t as defined by

$$(9) \quad \rho_{N;n}^t(x_1, \xi_1, \dots, x_n, \xi_n) := \int_{\mathbf{R}^{2dN(-n)}} \rho_N^t(x_i, \xi_i, x_2, \xi_2, \dots, x_N, \xi_N) dx_{n+1} d\xi_{n+1} \dots dx_N d\xi_N,$$

tends, in Wasserstein topology of any exponent $p \geq 1$, to the n th tensorial power of the solution ρ^t of the Vlasov equation, i.e. $\rho_{N;n}^t \rightarrow (\rho^t)^{\otimes n}$. Nevertheless, for sake of clarity of this short note, we will present in detail only the case $n = 1$.

This result (for $n = 1$), i.e. Theorem 2.2 below, has to be put in correspondence with the item 2 of the results of [6] just described, but there several differences. First of all our results hold true for more general initial data than the empirical measures. On the other side, our result will not be uniform in time as in item 1, but this inconvenient will be compensated by the fact that we will get an explicit rate of convergence in the asymptotism $N \rightarrow \infty$.

Our methods will also apply to systems of the general form

$$(10) \quad \begin{cases} \dot{x}_i = v_i \\ \dot{v}_i = \frac{1}{N} \sum_{j=1}^N \gamma(x_i - x_j, v_i - v_j). \end{cases} \quad x_i, v_i \in \mathbf{R}^d, \quad i = 1, \dots, N$$

where $\gamma(x, v) : \mathbf{R}^{2d} \rightarrow \mathbf{R}^d$ is a Lipschitz continuous function, bounded in (x, v) , Corollary 2.5 or bounded in x and sublinear in v , Theorem 2.4 and Corollary 2.6.

In the sequel of the paper, we will denote by $\text{Lip}(\gamma)_{(x,v)}$ the (local) Lipschitz constant of γ at the point $(x, v) \in \mathbf{R}^{2d}$ and we will suppose, without loss of generality, that there exists $\gamma_0 > 0$ such that, for all $(x, v) \in \mathbf{R}^{2d}$,

$$\begin{aligned} \gamma(x, v) &\leq \gamma_0 |v| \\ \text{Lip}(\gamma)_{(x,v)} &\leq \gamma_0 |v| \end{aligned}$$

where $|\cdot|$ denotes the Euclidean norm on \mathbf{R}^d .

We show in Apeendix A the gobal existence of the solutions of (10), with an exponential growth in time with respect to the initial conditions. Uniqueness is of course a consequence of (the iteration in time of) the Cauchy-Lipschitz Theorem.

The Cucker-Smale vector fields satisfies this hypothesis with $\gamma_0 = \|\psi\|_{L^\infty}$.

Note that the following Vlasov type kinetic equation on \mathbf{R}^d is obviously associated to (10), in a natural way:

$$(11) \quad \partial_t f^t(x, v) + v \cdot \nabla_x f^t(x, v) = \nabla_v \cdot \left(f^t(x, v) \int_{\mathbf{R}^{2d}} \gamma(x - y, v - w) f^t(y, w) dy dw \right).$$

Let us immediatly notice that our assumptions on γ contain the two assumptions in Theorem 2.3 in [9]. The first one because of the Lipschitz and sublinearity properties of γ , and the second one thanks to the fact, see [10, 11], that

$$\sup_{|\text{lip}(\varphi)| \leq 1} \left| \int \varphi(\mu - \nu) \right| \leq W_1(\mu, \nu)$$

where W_1 is the Wasserstien distance of exponent 1, as defined in Remark 3.5 in the present paper. Therefore, there exist a unique continuous solution ρ^t of (11) $\rho^t \in C_0(\mathbf{R}, \mathcal{P}_c(\mathbf{R}^{2d}))$ where $\mathcal{P}_c(\mathbf{R}^{2d})$ is the space of compactly supported probability measures on \mathbf{R}^{2d} .

2. THE RESULTS

In this section we will state our main results.

Let us start this section by recalling the definition of the second order Wasserstein distance $\text{dist}_{\text{MK},2}$ (see [1, 10, 11]).

Definition 2.1 (quadratic Wasserstein distance). *The Wasserstein distance of order two between two probability measures μ, ν on \mathbf{R}^m with finite second moments is defined as*

$$W_2(\mu, \nu)^2 = \inf_{\gamma \in \Gamma(\mu, \nu)} \int_{\mathbf{R}^m \times \mathbf{R}^m} |x - y|^2 \gamma(dx, dy)$$

where $\Gamma(\mu, \nu)$ is the set of probability measures on $\mathbf{R}^m \times \mathbf{R}^m$ whose marginals on the two factors are μ and ν . E

That is to say that elements γ of $\Gamma(\mu, \nu)$, called couplings or transportation maps or transport maps or just maps, depending of the authors, satisfy, for any test functions a and b in $C_0(\mathbf{R}^d)$,

$$\int_{\mathbf{R}^{2m}} a(x)\gamma(dx, dy) = \int_{\mathbf{R}^m} a(x)\mu(dx), \quad \int_{\mathbf{R}^{2m}} b(y)\gamma(dx, dy) = \int_{\mathbf{R}^m} b(y)\nu(dy).$$

The symmetry property in μ, ν is obvious and the separability property is easily proven by taking the optimal coupling between μ and itself equal to

$$(12) \quad \gamma = \mu\delta(x - y).$$

Conversely, if $W_2(\mu, \nu)^2 = \int_{\mathbf{R}^m \times \mathbf{R}^m} |x - y|^2 \gamma(dx, dy)$ for some γ so that $x = y$ γ a.e. and, for any Borel function φ ,

$$\int \varphi(x)\mu(dx) = \int \varphi(x)\gamma(dx, dy) = \int \varphi(y)\gamma(dx, dy) = \int \varphi(y)\nu(dy) \Rightarrow \mu = \nu.$$

The proof of the triangular inequality is much more involved, see again [1, 10, 11]).

We can now state the main result of this little note, proven in Section 3.4.

Theorem 2.2 (Cucker-Smale model).

Let ρ^t be the solution of (7) with an initial condition $\rho^{in} \in L^1(\mathbf{R}^{2d})$ compactly supported and let $\rho_{N;1}^t$ be the first marginal of $\rho_N^t := \Phi^t \# (\rho^{in})^{\otimes N}$, as defined in (6).

Then, for all $N > 1$, $t \in \mathbf{R}$,

$$W_2(\rho_{N;1}^t, \rho^t) \leq 4 \frac{\|\psi\|_\infty \|v\|_{L^\infty(\rho^{in})}}{N^{\frac{1}{2}}} \left(\frac{e^{Lt} - 1}{L} \right)^{\frac{1}{2}}$$

with

$$L := 2(1 + 4\|\psi\|_\infty^2 (2|\bar{v}| + |\text{supp}[\rho^{in}]|^2)), \quad \bar{v} = \int v \rho^{in} dx dv$$

(by $\|v\|_{L^\infty(\text{supp}[\rho^{in}])}$ we mean $\sup_{(x,v) \in \text{supp}[\rho^{in}]} (|v|)$).

Remark 2.3. There exists an equivalent result for higher orders marginals of $\rho(t)$ and other Monge-Kantorovich distances but we prefer in this little note to concentrate on the case of first marginal and quadratic Wasserstein distance. The proof in the more general situation is very close to the one presented here. See Remark 3.5 for some details.

Theorem 2.2 is actually a corollary of the following more general result, proven in Section 3.2

Theorem 2.4 (General Cucker-Smale model with sublinear force).

Let ρ^t be the solution of (11), γ bounded on \mathbf{R}^{2d} , with an initial condition $\rho_1^{in} \in L^1(\mathbf{R}^{2d})$ compactly supported and Φ^t the flow defined by the system (10). Finally, let $\rho_{N;1}^t$ be the first marginal of $\rho_N^t := \Phi^t \# (\rho^{in})^{\otimes N}$, as defined in (6).

Then, for all $N > 1$, $t \in \mathbf{R}$,

$$W_2(\rho_{N;1}^t, \rho^t) \leq C(t)N^{-\frac{1}{2}}$$

with

$$C(t) := \left(4 \int_0^t \sup_{\substack{k,l=1,dots,N \\ (Y,\Xi) \in \text{supp}[\rho_N^t]}} |\gamma(y_k - y_l, \xi_k - \xi_l)|^2 e^{\int_s^t L(u) du} ds \right)^{\frac{1}{2}},$$

$$L(t) := 2(1 + 2 \sup_{(x,v),(y,\xi) \in \text{supp}[\rho^t]} \text{Lip}(\gamma)_{(x-y, v-\xi)}^2).$$

The following result is elementary derived from Theorem 2.4.

Corollary 2.5 (General Cucker-Smale model with bounded force).

Let ρ^t be the solution of (11), γ globally Lipschitz and bounded on \mathbf{R}^{2d} , with an initial condition $\rho^{in} \in L^1(\mathbf{R}^{2d})$ and Φ^t the flow defined by the system (10). Finally, let $\rho_{N;1}^t$ be the first marginal of $\rho_N^t := \Phi^t \# (\rho^{in})^{\otimes N}$, as defined in (6).

Then, for all $N > 1, t \in \mathbf{R}$,

$$W_2(\rho_{N;1}^t, \rho^t) \leq C(t)N^{-\frac{1}{2}}$$

with

$$C(t) := \left(4\|\gamma\|_\infty \frac{e^{\Lambda t} - 1}{\Lambda} \right)^{\frac{1}{2}} \quad \Lambda := 2(1 + 2\text{Lip}(\gamma)^2).$$

Note that no need of compactness of the support of ρ^{in} is needed any more in Corollary 2.5.

The boundedness in space and sublinearity in velocities of γ allow to control the increasing of the flow $\Phi(t)$, and leads to our next result, whose proof is given in Section 3.3 below.

Corollary 2.6. Under the same hypothesis as in Theorem 2.4 we have that, for all $N > 1, t \in \mathbf{R}$,

$$W_2(\rho_{N;1}^t, \rho^t) \leq C(t)N^{-\frac{1}{2}}$$

with

$$C(t) = \left(2\gamma_0 \|v\|_{L^\infty(\text{supp}[\rho^{in}])}^2 e^{4\gamma_0 t} 4\gamma_0 (\|v\|_{L^\infty(\text{supp}[\rho^{in}])} + \|v\|_{L^1(\text{supp}[\rho^{in}])})^2 e^{4t} (e^{4\gamma_0 t} - 1) \right)^{\frac{1}{2}},$$

Let us finish this section of results by a simple remark. Flocking properties of the solution of the Vlasov kinetic equation (2) can be derived from the corresponding properties of the particle system (1), as in [7, 6]. But they can also be derived by a direct PDE study of (2), as in [9]. This suggests a kind of inverse questioning: suppose one determines some Flocking properties for the solution of a general kinetic Vlasov equation, e.g. (11). Does this infer on Flocking properties for the corresponding particle system, e.g. (10)? Our Corollary 2.6 gives some insight on this problem, as it tells us quantitatively how close is the solution of the kinetic equation (and its tensorial powers) to the marginals of any orders of the pushforward by the corresponding particle flow of a general N particle density.

More precisely, suppose that the solution $\rho(t)$ of (11) satisfies the following property:

$$(13) \quad \text{supp}[\rho(t)] \subset B(\bar{x} + t\bar{v}, X) \times B(\bar{v}, V e^{-\alpha t})$$

for some positive constants $\bar{x}, X, \bar{v}, V, \alpha$.

This implies that $L(t)$ in Theorem 2.4 can be easily estimated by

$$L(t) \leq 2(1 + 8(\bar{v}^2 + V^2 e^{-2\alpha t})) \leq 2(1 + 8(\bar{v}^2 + V^2))$$

and, therefore, in the same Theorem,

$$(14) \quad C(t) \leq \gamma_0 V_\infty^2 \frac{e^{(4\gamma_0 + 2(1 + 8(\bar{v}^2 + V^2))t) - 1}}{4\gamma_0 + 2(1 + 8(\bar{v}^2 + V^2))}$$

(note that in (14), $C(t)$ has an exponential growth, and now a double exponential one as in Corollary 2.6).

Of course the first marginal $\rho_{N;1}^t$ of the pushforward of $(\rho^{in})^{\otimes N}$ by the flows induced by the system (10) has no reason for being compactly supported, as we didn't impose anything on the particle flow. Nevertheless, we have the following result provide a weak version of the property (13).

Proposition 2.7. For all $t, \epsilon > 0$ let us defined

$$N_{t,\epsilon} := \left(\frac{C(t)}{\epsilon} \right)^2,$$

where $C(t)$ is the function defined by (14).

Then, for any $N \geq N_{t,\epsilon}$ and any Lipschitz function φ on \mathbf{R}^{2d} of Lipschitz constant smaller than 1,

$$\int_{\mathbf{R}^{2d} \setminus B(\bar{x} + t\bar{v}, X) \times B(\bar{v}, V e^{-\alpha t})} \varphi(x, v) \rho_{N,1}(t) dx dv \leq \epsilon.$$

Proof. It is an immediate consequence of Theorem 2.4 or Theorem 2.6, and the fact that the the Waaserstein distance metrizes the weak topology in the sense that

$$\sup_{\text{Lip}(\varphi) \leq 1} \int (\mu - \nu) \varphi(x, v) dx dv \leq W_2(\mu, \nu)$$

for any probability measures μ and ν □

3. PROOFS

3.1. Preliminaries. Let us first recall the general situation we are dealing with, in order also to fix the notations.

We consider on \mathbf{R}^{2dN} the following Cucker-Smale type vector field

$$(15) \quad \begin{aligned} \dot{x}_i &= v_i \\ \dot{v}_i &= G_i(t, X, V), \quad i = 1, \dots, N \end{aligned}$$

where

$$(16) \quad G_i(t, X, V) = \frac{1}{N} \sum_{j=1}^N \gamma(x_i - x_j, v_i - v_j).$$

Here the function $\gamma(x, v) : \mathbf{R}^{2d} \rightarrow \mathbf{R}^d$ is a Lipschitz, bounded in x and sublinear in v , continuous function., such that $\gamma(x, v), \text{Lip}(\gamma)_{(x,v)} \leq \gamma_0$.

We used the notation $X = (x_1, \dots, x_N), V = (v_1, \dots, v_N)$.

In fact, we are rather interested in the Liouville equation associated to (15) [8], namely

$$(17) \quad \partial_t \rho_N^t + v \cdot \nabla_x \rho_N^t = \sum_{i=1}^N \nabla_{v_i} (G_i \rho_N^t), \quad \rho_N^{t=0} = \rho_N^{in}$$

with $\rho_N^{in} \in \mathcal{P}(\mathbf{R}^{2dN})$.

Although the argument is standard, let us recall why the solution of (17) is equal to the pushforward of ρ_N^{in} by the flow Φ_t generated by (15): integrating ρ_N^{in} against a test φ function composed by Φ_t gives

$$(18) \quad \partial_t \int \varphi(\Phi_t(X, V)) \rho_N^{in}(X, V) dX dV = \int \varphi(X, V) \partial_t (\Phi_t \# \rho_N^{in}(X, V)) dX dV$$

Moreover

$$(19) \quad \begin{aligned} \partial_t \int \varphi(\Phi_t(X, V)) \rho_N^{in}(X, V) dX dV &= \int ((-V, G) \cdot \nabla_{(X,V)} \varphi(X, V)) \phi_t \# \rho_N^{in}(X, V) dX dV \\ &= - \int \varphi(X, V) \nabla_{(X,V)} \cdot (V, G) \Phi_t \# \rho_N^{in}(X, V) dX dV \end{aligned}$$

So that $\Phi_t \# \rho_N^{in}$ solves (17).

We want to prove that the marginals of ρ_N^t tend, as $N \rightarrow \infty$, to the solution of a Vlasov type equation.

Such Vlasov-type equation associated to (17) has (introduced in [8] for the Cucker-Smale model) reads

$$(20) \quad \partial_t \rho^t(x, v) + v \cdot \nabla_x \rho^t(x, v) = \nabla_v \cdot (G_{\rho^t} \rho^t(x, v)), \quad \rho^{t=0} = \rho_1^{in} \in L^1(\mathbf{R}^{2d}, dx dv),$$

with

$$(21) \quad G_{\rho}(x, v) = \int_{\mathbf{R}^{2d}} \gamma(x - y, v - w) \rho(y, w) dy dw.$$

We can now state the proof of the main results of this little note.

3.2. Proof of Theorem 2.4. The proof will be articulated around the following four lemmas.

Let $\pi_N^{in} \in \Pi((\rho^{in})^{\otimes N}, (\rho^{in})^{\otimes N})$ satisfy

$$(22) \quad T_\sigma \# \pi_N^{in} = \pi_N^{in}, \quad \text{for each } \sigma \in \mathfrak{S}_N,$$

where \mathfrak{S}_N is the group of permutations of N elements and

$$\begin{aligned} & T_\sigma(x_1, v_1, \dots, x_N, v_N, y_1, \xi_1, \dots, y_N, \xi_N) \\ &= (x_{\sigma(1)}, v_{\sigma(1)}, \dots, x_{\sigma(N)}, v_{\sigma(N)}, y_{\sigma(1)}, \xi_{\sigma(1)}, \dots, y_{\sigma(N)}, \xi_{\sigma(N)}). \end{aligned}$$

We will denote $X = (x_1, \dots, x_N), V = (v_1, \dots, v_N), Y = (y_1, \dots, y_N), \Xi = (\xi_1, \dots, \xi_N)$.

The following first Lemma will be one of the keys of the proof of Theorem 2.2. It consists in considering in evolving a coupling π_N^{in} of two initial conditions of the Liouville (17) and Vlasov (20) equations by the two dynamics of each factor of π_N^{in} .

Lemma 3.1. *Let $\pi_N(t)$ be the solution of*

$$(23) \quad \partial_t \pi_N + V \cdot \nabla_X \pi_N + \Xi \cdot \nabla_Y \pi_N = \sum_{i=1}^N (\nabla_{\xi_i} \cdot (G_i(Y, \Xi) \pi_N) + \nabla_{v_i} \cdot (G_{\rho^t}(x_i, v_i) \pi_N))$$

with $\pi_N(0) = \pi_N^{in}$.

Then, for all $t \in \mathbf{R}$, $\pi_N(t)$ is a coupling between ρ_N^t and $(\rho^t)^{\otimes N}$. Moreover $\pi_N(t)$ is invariant by permutations T_σ .

Proof. By taking the two marginals of the two sides of the equality, one get that they satisfy the two Liouville and Vlasov equations. The result is then obtained by uniqueness of the solutions of both equations. \square

Lemma 3.2. *Let*

$$\begin{aligned} D_N(t) &:= \int \frac{1}{N} \sum_{j=1}^N (|x_j - y_j|^2 + |v_j - \xi_j|^2) \pi_N(t) \\ &= \frac{1}{N} \int ((X - Y)^2 + (V - \Xi)^2) \pi_N(t). \end{aligned}$$

Then

$$\frac{dD_N}{dt} \leq L(t) D_N + \frac{1}{N} \sum_{j=1}^N \int |G_{\rho^t}(y_i, \xi_i) - G_i(Y, \Xi)|^2 \rho_N^t dY d\Xi,$$

with

$$L(t) := 2(1 + 2 \sup_{\substack{i=1, \dots, N \\ (Y, \Xi, X, V) \in \text{supp}[\pi_N(t)] \\ ((y, \xi) \in \text{supp}[\rho^t]}} \text{Lip}(\gamma)_{(x_i - y, \xi_i - \xi)}^2).$$

Proof. We first notice that

$$\frac{dD_N}{dt} = \frac{2}{N} \int \left((\Xi - V) \cdot (X - Y) + \sum_{i=1}^N (v_i - \xi_i) \cdot (G_i(Y, \Xi) - G_{\rho^t}(x_i, v_i)) \right) d\pi_N$$

Using $2uv \leq u^2 + v^2$ we get

$$\begin{aligned} \frac{dD_N}{dt} &\leq \frac{1}{2N} \int \left((X - Y)^2 + 2(V - \Xi)^2 + \sum_{i=1}^N |G_i(Y, \Xi) - G_{\rho^t}(x_i, v_i)|^2 \right) d\pi_N \\ (24) \quad &\leq D_N(t) + \frac{1}{2N} \int \left(\sum_{i=1}^N |G_i(Y, \Xi) - G_{\rho^t}(x_i, v_i)|^2 \right) d\pi_N. \end{aligned}$$

Let us add to $G_i(Y, \Xi) - G_{\rho^t}(x_i, v_i)$ the null term $(G_{\rho^t}(y_i, \xi_i) - G_{\rho^t}(y_i, \xi_i))$ so that

$$|G_i(Y, \Xi) - G_{\rho^t}(x_i, v_i)|^2 \leq 2(|G_i(Y, \Xi) - G_{\rho^t}(y_i, \xi_i)|^2 + |G_{\rho^t}(y_i, \xi_i) - G_{\rho^t}(x_i, v_i)|^2)$$

Let us first estimate

$$\begin{aligned} |G_{\rho^t}(y_i, \xi_i) - G_{\rho^t}(x_i, v_i)| &= \left| \int (\gamma(y_i - y, \xi_i - \xi) - \gamma(x_i - y, v_i - \xi)) \rho^t(y, \xi) dy d\xi \right| \\ &\leq \int |(x_i - y_i, v_i - \xi_i)| \text{Lip}(\gamma)_{(x_i - y, \xi_i - \xi)} \rho^t(y, \xi) dy d\xi \\ &\leq \sup_{(y, \xi) \in \text{supp}[\rho^t]} \text{Lip}(\gamma)_{(x_i - y, \xi_i - \xi)} \int |(x_i - y_i, v_i - \xi_i)| \rho^t(y, \xi) dy d\xi \\ &= \sup_{(y, \xi) \in \text{supp}[\rho^t]} \text{Lip}(\gamma)_{(x_i - y, \xi_i - \xi)} |(x_i - y_i, v_i - \xi_i)| \end{aligned}$$

Therefore, by convexity,

$$|G_{\rho^t}(y_i, \xi_i) - G_{\rho^t}(x_i, v_i)|^2 \leq 2 \sup_{(y, \xi) \in \text{supp}[\rho^t]} \text{Lip}(\gamma)_{(x_i - y, \xi_i - \xi)}^2 (|x_i - y_i|^2 + |v_i - \xi_i|^2)$$

and

$$\begin{aligned} &\frac{1}{N} \int \sum_{i=1}^N |G_{\rho^t}(y_i, \xi_i) - G_{\rho^t}(x_i, v_i)|^2 d\pi_N \\ &\leq \sup_{\substack{i=1, \dots, N \\ (Y, \Xi: X, V) \in \text{supp}[\pi_N(t)] \\ ((y, \xi) \in \text{supp}[\rho^t])}} \text{Lip}(\gamma)_{(x_i - y, \xi_i - \xi)}^2 \frac{1}{N} \int \sum_{i=1}^N (|x_i - y_i|^2 + |v_i - \xi_i|^2) d\pi_N \\ &= \sup_{\substack{i=1, \dots, N \\ (Y, \Xi: X, V) \in \text{supp}[\pi_N(t)] \\ ((y, \xi) \in \text{supp}[\rho^t])}} \text{Lip}(\gamma)_{(x_i - y, \xi_i - \xi)}^2 D_N(t) \end{aligned}$$

And the lemma follows by (24) and Lemma 3.1. □

It remains to estimate in (24) the term

$$\int \frac{1}{N} \sum_{j=1}^N |G_{\rho^t}(y_j, \xi_j) - G_i(Y, \Xi)|^2 \rho_N^t dY d\Xi.$$

The following result is a straightforward extension of Lemma 3.3. in [5] with the special value $p = 2$ and d replaced by $2d$.

Lemma 3.3. *Let π be a compactly supported probability density on \mathbf{R}^{2dN} and F be a vector field on \mathbf{R}^{2d} .*

For each $j = 1, \dots, N$, one has

$$(25) \quad \int \left| F \star \rho(y_j, \xi_j) - \frac{1}{N} \sum_{k=1}^N F(y_j - y_k, \xi_j - \xi_k) \right|^2 \rho_N^t dY d\Xi \leq \frac{16}{N} \sup_{\substack{k, l=1, \dots, N \\ (Y, \Xi) \in \text{supp}[\rho_N^t]}} |F(y_k - y_l, \xi_k - \xi_l)|^2.$$

Proof. There are two differences between the statements in Lemma 3.3 and Lemma 3.3. in [5]:

- the term $\sup_{\substack{k, l=1, \dots, N \\ (Y, \Xi) \in \text{supp}[\rho_N^t]}} |F(y_k - y_l, \xi_k - \xi_l)|^2$ used in the right hand side of (25) instead of the norm

$L^\infty(\mathbf{R}^{2d})$ used in [5]. This difference is unessential since the integration in the right hand side of (25) is effectively over $\text{supp}[\rho_N^t]$.

- the measure ρ_n^t in the left hand-side of (25) instead of the measure $\rho^{\otimes N}$ used in [5]. It is clear from the (beginning) of the proof of Theorem 3.3. in [5] that it works exactly the same way with ρ_N^t . \square

Lemma 3.3 with $F(x, v) = \gamma(x, v)$ together with Lemma 3.2 gives immediatly that

$$\frac{dD_N}{dt} \leq L(t)D_N + \frac{16}{N} \sup_{\substack{k,l=1,\dots,N \\ (Y,\Xi) \in \text{supp}[\rho_N^t]}} |\gamma(y_k - y_l, \xi_k - \xi_l)|^2$$

and, by Gronwall's inequality,

$$(26) \quad D_N(t) \leq D_N(0)e^{\int_0^t L(s)ds} + \frac{16}{N} \int_0^t \sup_{\substack{k,l=1,\dots,N \\ (Y,\Xi) \in \text{supp}[\rho_N^t]}} |\gamma(y_k - y_l, \xi_k - \xi_l)|^2 e^{\int_s^t L(u)du} ds.$$

Let us denote by $(\pi_N(t))_1$ the measure on $\mathbf{R}^{2d} \times \mathbf{R}^{2d}$ defined, for every test function $\varphi(x_1, v_1; y_1, \xi_1)$, by

$$\int_{\mathbf{R}^{2dN} \times \mathbf{R}^{2dN}} \varphi \pi_N(dX, dV; dY d\Xi) = \int_{\mathbf{R}^{2d} \times \mathbf{R}^{2d}} \varphi(x_1, v_1; y_1, \xi_1) (\pi_N(t))_1(dx_1, dv_1; dy_1, d\xi_1)$$

We now notice the following obvious fact.

Lemma 3.4. $(\pi_N(t))_1$ is a coupling between $\rho_{N;1}^t$ and ρ^t .

Since $\pi_N(t)$ is symmetric by permutations, one has easily that

$$(27) \quad D_N(t) = \int (|x_1 - y_1|^2 + |v_1 - \xi_1|^2) (\pi_N(t))_1$$

and Lemma 3.4 immediatly implies that

$$(28) \quad D_N(t) \geq W_2(\rho_1(t), (\rho(t))_1)^2,$$

so that, by Lemma 3.3,

$$(29) \quad W_2(\rho_1(t), (\rho(t))_1)^2 \leq D_N(0)e^{\int_0^t L(s)ds} + \frac{16}{N} \int_0^t \sup_{\substack{k,l=1,\dots,N \\ (Y,\Xi) \in \text{supp}[\rho_N^t]}} |\gamma(y_k - y_l, \xi_k - \xi_l)|^2 e^{\int_s^t L(u)du} ds.$$

Remember that (29) is valid for $D(0)$ as defined in Lemma 3.2 for *any* π_N coupling $(\rho_1^{in})^{\otimes N}$ with itself. Since W_2 is a distance, one has, for any of these π^{op} ,

$$W_2((\rho_1^{in})^{\otimes N}, (\rho_1^{in})^{\otimes N})^2 = \int ((X - Y)^2 + (V - \Xi)^2) \pi_N^{op} = ND_N(0) = 0$$

so that

$$W_2(\rho_1(t), (\rho(t))_1)^2 \leq \frac{16}{N} \int_0^t \sup_{\substack{k,l=1,\dots,N \\ (Y,\Xi) \in \text{supp}[\rho_N^t]}} |\gamma(y_k - y_l, \xi_k - \xi_l)|^2 e^{\int_s^t L(u)du} ds.$$

Finally, choosing for π^{op} the optimal coupling

$$\pi^{op}(Y, \Xi, X, V) := (\rho_i^{in})^{\otimes N}(X, V) \delta(X - Y) \delta(V - \Xi),$$

one get immediatly that

$$(Y, \Xi : X, V) \in \text{supp}[\pi_N(t)] \Rightarrow (X, V) \in \text{supp}[\rho^t]^{\otimes N} \Rightarrow (x_i, v_i) \in \text{supp}[\rho^t], \quad i = 1, \dots, N.$$

and therefore

$$L(t) = \mathbf{L}(t) := 2(1 + 2 \sup_{(x,v),(y,\xi) \in \text{supp}[\rho^t]} \text{Lip}(\gamma)_{(x-y, v-\xi)}^2).$$

Theorem 2.4 is proven.

Remark 3.5 (Higher order Wasserstein and marginals). *As we wanted to leap this note as short as possible, we expressed our results only for the first marginal $\rho_{N;1}$ in the 2-Wasserstein topology, but the method developed in [5] allows as well, with the same kind of modification than the ones used before in this section, to the higher cases. Let us remind, for $p \geq 1$, the definition of $W_p(\mu, \nu)$ for two positive measures μ, ν (cf. Definition 2.1*

$$W_p(\mu, \nu)^p = \inf_{\gamma \in \Gamma(\mu, \nu)} \int_{\mathbf{R}^m \times \mathbf{R}^m} |x - y|^p \gamma(dx, dy),$$

and, for any probability measure ρ on \mathbf{R}^{2dN} and $n = 1, \dots, N$, the definition of the n th marginal $\rho_{N;n}$ defined on \mathbf{R}^{2dn}

$$\begin{aligned} \rho_{N;n}(x_1, \dots, x_n, v_1, \dots, v_n) := \\ \int_{\mathbf{R}^{2d(N-n)}} \rho((x_1, \dots, x_n, x_{n+1}, \dots, x_N, v_1, \dots, v_n, v_{n+1}, \dots, v_N) dx_{n+1} \dots dx_N dv_{n+1} \dots dv_N. \end{aligned}$$

One gets, for each $p \geq 1$, $N \geq 1$ and $n = 1, \dots, N$,

$$\frac{1}{n} W_p(\rho_{N;n}^t, (\rho^t)^{\otimes n})^p \leq D_{p,n}(t) N^{-\min(p/2, 1)}$$

with

$$D_{p,n}(t) = 2^{2p} \max(1, p-1) ([p/2] + 1) \int_0^t \sup_{\substack{k,l=1, \text{dots}, N \\ (Y, \Xi) \in \text{supp}[\rho_{N;n}^t]}} |\gamma(y_k - y_l, \xi_k - \xi_l)|^p e^{2 \max(1, p-1) \int_s^t L(u) du} ds$$

where

$$L(u) = 1 + 2^{p-1} \sup_{(x,v), (y,\xi) \in \text{supp}[\rho^t]} \text{Lip}(\gamma)_{(x-y, v-\xi)}^p.$$

The main changes in the proof are the use of the Young inequality

$$puv^{p-1} \leq u^p + (p-1)v^p \leq \max(1, p-1)(u^p + v^p)$$

instead of $2uv \leq u^2 + v^2$ before (24), the convexity of $|\cdot|^p$ for $p \geq 1$ and a variant of the contents of Lemma 3.3 which reads

$$\begin{aligned} & \int \left| F \star \rho(x_j, v_j) - \frac{1}{N} \sum_{k=1}^N F(x_j - x_k, v_i - v_j) \right|^p \rho^{\otimes N} dX dV \\ & \leq \frac{2[p/2] + 2}{N^{\min(p/2, 1)}} \sup_{\substack{k,l=1, \text{dots}, N \\ (Y, \Xi) \in \text{supp}[\rho_{N;n}^t]}} |F(y_k - y_l, \xi_k - \xi_l)|^p. \end{aligned}$$

Finally, the statement of Lemma 3.4 becomes now easily that $\pi_N(t)_n$ is a coupling between $\rho_{N;n}^t$ and $(\rho^t)^{\otimes n}$, where $\pi_N(t)_n$ is defined through by

$$\int_{\mathbf{R}^{2dN} \times \mathbf{R}^{2dN}} \varphi \pi_N(dX, dV; dY d\Xi) = \int_{\mathbf{R}^{2d} \times \mathbf{R}^{2d}} \varphi \pi_N(t)_n(d(x, v; y, \xi)_1 \dots d(x, v; y, \xi)_n)$$

for every test function $\varphi((x, v; y, \xi)_1, \dots, (x, v; y, \xi)_n)$,

3.3. Proof of Corollary 2.6. We get to estimates $L(t)$ and $C(t)$ as given in Theorem 2.4 out of the estimates established in Appendices A and B.

Let us recall that

$$L(t) := 2(1 + 2 \sup_{(x,v), (y,\xi) \in \text{supp}[\rho^t]} \text{Lip}(\gamma)_{(x-y, v-\xi)}^2) = 2(1 + 2\|\psi\|_\infty^2 \sup_{(x,v), (y,\xi) \in \text{supp}[\rho^t]} |v - \xi|^2)$$

Thanks to (36), we get

$$\begin{aligned}
\mathbf{L}(t) &:= 2(1 + 2 \sup_{(x,v),(y,\xi) \in \text{supp}[\rho^t]} \text{Lip}(\gamma)_{(x-y,v-\xi)}^2) \\
&\leq 2(1 + 2 \sup_{(x,v),(y,\xi) \in \text{supp}[\rho^t]} \gamma_0^2 |v - \xi|^2) \leq 4(1 + 2 \sup_{(x,v) \in \text{supp}[\rho^t]} \gamma_0^2 |v|^2) \\
&= 4(1 + 2 \sup_{(x,v) \in \text{supp}[\rho^{in}]} \gamma_0^2 |\Phi(t)(x,v)|^2) \\
&\leq 4(1 + 4\gamma_0^2 e^{4\gamma_0 t} (\|v\|_{L^\infty(\text{supp}[\rho^{in}])} + \|v\|_{L^1(\text{supp}[\rho^{in}])})^2)
\end{aligned}$$

Moreover, (34) gives

$$\begin{aligned}
\sup_{\substack{k,l=1,\dots,N \\ (Y,\Xi) \in \text{supp}[\rho_N^t]}} |\gamma(y_k - y_l, \xi_k - \xi_l)|^2 &\leq 2\gamma_0^2 \sup_{\substack{k=1,\dots,N \\ (Y,\Xi) \in \text{supp}[\rho_N^t]}} |\xi_k|^2 \\
&= 2\gamma_0^2 \sup_{\substack{k=1,\dots,N \\ (X,V) \in \text{supp}[(\rho^{in})^{\otimes N}]} |v_k(t)|^2 \\
&\leq 2\gamma_0^2 \sup_{\substack{k=1,\dots,N \\ (X,V) \in \text{supp}[(\rho^{in})^{\otimes N}]} |v_k|^2 e^{4\gamma_0 t} \\
&= 2\gamma_0^2 \|v\|_{L^\infty(\text{supp}[\rho^{in}])}^2 e^{4\gamma_0 t}
\end{aligned}$$

(here $(X(t), V(t))$ solves (10).

Therefore

$$\begin{aligned}
C(t)^2 &:= 4 \int_0^t \sup_{\substack{k,l=1,\dots,N \\ (Y,\Xi) \in \text{supp}[\rho_N^t]}} |\gamma(y_k - y_l, \xi_k - \xi_l)|^2 e^{\int_s^t \mathbf{L}(u) du} ds, \\
&\leq 8\gamma_0^2 \|v\|_{L^\infty(\text{supp}[\rho^{in}])}^2 \int_0^t e^{4(\gamma_0 s + t - s)} e^{16\gamma_0^2 (\|v\|_{L^\infty(\text{supp}[\rho^{in}])} + \|v\|_{L^1(\text{supp}[\rho^{in}])})^2 (e^{4\gamma_0 t} - e^{4\gamma_0 s}) / 4\gamma_0} ds \\
&\leq 2\gamma_0 \|v\|_{L^\infty(\text{supp}[\rho^{in}])}^2 e^{e^{4\gamma_0 t} 4\gamma_0 (\|v\|_{L^\infty(\text{supp}[\rho^{in}])} + \|v\|_{L^1(\text{supp}[\rho^{in}])})^2} e^{4t} (e^{4\gamma_0 t} - 1).
\end{aligned}$$

Corollary 2.6 is proven.

3.4. Proof of Theorem 2.2. In order to prove Theorem 2.2, we need to estimate, in the Cucker-Smale particular case, that is when

$$\gamma(x, v) = \psi(x)v,$$

the two quantities

$$\begin{aligned}
C(t) &:= \left(4 \int_0^t \sup_{\substack{k,l=1,\dots,N \\ (Y,\Xi) \in \text{supp}[\rho_N^t]}} |\gamma(y_k - y_l, \xi_k - \xi_l)|^2 e^{\int_s^t \mathbf{L}(u) du} ds \right)^{\frac{1}{2}}, \\
\mathbf{L}(t) &:= 2(1 + 2 \sup_{(x,v),(y,\xi) \in \text{supp}[\rho^t]} \text{Lip}(\gamma)_{(x-y,v-\xi)}^2)
\end{aligned}$$

$$(30) \quad \mathbf{L}(t) := 2(1 + 2 \sup_{(x,v),(y,\xi) \in \text{supp}[\rho^t]} \text{Lip}(\gamma)_{(x-y,v-\xi)}^2) = 2(1 + 2\|\psi\|_\infty^2 \sup_{(x,v),(y,\xi) \in \text{supp}[\rho^t]} |v - \xi|^2)$$

and

$$C(t) := \left(4 \int_0^t \sup_{\substack{k,l=1,\dots,N \\ (Y,\Xi) \in \text{supp}[\rho_N^t]}} |\gamma(y_k - y_l, \xi_k - \xi_l)|^2 e^{\int_s^t \mathbf{L}(u) du} ds \right)^{\frac{1}{2}}$$

that is

$$(31) \quad \sup_{\substack{k,l=1,\dots,N \\ (Y,\Xi) \in \text{supp}[\rho_N^t]}} |\gamma(y_k - y_l, \xi_k - \xi_l)|^2 = \|\psi\|_\infty^2 \sup_{\substack{k,l=1,\dots,N \\ (Y,\Xi) \in \text{supp}[\rho_N^t]}} |\xi_k - \xi_l|^2$$

To estimate $\sup_{(x,v),(y,\xi) \in \text{supp}[\rho^t]} |v - \xi|^2 \leq 2 \sup_{(x,v),(y,\xi) \in \text{supp}[\rho^t]} (|v|^2 + |\xi|^2)$ we will use Lemma 3.2 in [9] which stipulates that

$$\text{supp}_v[\rho^t] \subset B(\bar{v}, V(t))$$

with $\bar{v} = \int v \rho^{in} dx dv$ and $\frac{d}{dt} V(t) \leq 0$.

Therefore $\|v\|_{L^\infty(\text{supp}[\rho^t])} \leq |\bar{v}| + V(0)$ so that, since one can take $V(0) = |\bar{v}| + |\text{supp}[\rho^{in}]|$,

$$\sup_{(x,v),(y,\xi) \in \text{supp}[\rho^t]} |v - \xi|^2 \leq 2\|\psi\|_\infty^2 (2|\bar{v}| + |\text{supp}[\rho^{in}]|^2).$$

and

$$L(t) \leq 2(1 + 4\|\psi\|_\infty^2 (2|\bar{v}| + |\text{supp}[\rho^{in}]|^2)) := L.$$

Moreover,

$$\sup_{\substack{k,l=1,\dots,N \\ (Y,\Xi) \in \text{supp}[\rho_N^t]}} |\xi_k - \xi_l|^2 = \sup_{\substack{k,l=1,\dots,N \\ (X(0),V(0)) \in \text{supp}[(\rho^{in})^{\otimes N}]}} |v_k(t) - v_l(t)|^2$$

where $x_i(t0, v_i(t))$, $i = 1, \dots, N$ satisfies (1).

We will need just a very little part of the stability results expressed by Ha, Kim and Zhuang, namely the following lemma, taken from formula (8) in Lemma 2.1 in [6]:

$$\frac{d}{dt} \sup_{k,l=1,\dots,N} |v_k(t) - v_l(t)| \leq 0, \quad \forall t,$$

which leads naturally to

$$\sup_{\substack{k,l=1,\dots,N \\ (Y,\Xi) \in \text{supp}[\rho_N^t]}} |\xi_k - \xi_l|^2 \leq 4\|v\|_{L^\infty(\rho^{in})}^2$$

which implies

$$C(t) \leq 4 \left(\|\psi\|_\infty^2 \|v\|_{L^\infty(\rho^{in})}^2 \frac{e^{Lt} - 1}{L} \right)^{\frac{1}{2}}.$$

Theorem 2.2 is proved.

APPENDIX A. DYNAMICAL ESTIMATES FOR GENERAL CUCKER-SMALE PARTICLE SYSTEMS

In this section, we give global estimates on the flow $\Phi^N(t)$ generated by (10).

We have, for each $i = 1, \dots, N$,

$$(32) \quad \begin{aligned} \frac{d}{dt} |v_i| &\leq |\dot{v}_i| \\ &\leq \frac{1}{N} \sum_{j=1}^N |\gamma(x_i - v_j, v_i - v_j)| \leq \frac{1}{N} \sum_{j=1}^N \gamma_0 |v_i - v_j| \\ &\leq \frac{1}{N} \sum_{j=1}^N \gamma_0 (|v_j| + |v_i|), \end{aligned}$$

so that

$$\frac{d}{dt} \sum_{i=1}^N |v_i| \leq 2\gamma_0 \sum_{i=1}^N |v_i|.$$

and, by Gronwall inequality,

$$(33) \quad \sum_{i=1}^N |v_i(t)| \leq \sum_{i=1}^N |v_i(0)| e^{2\gamma_0 t}.$$

Turning back to (32), we get by (33), for each $i = 1, \dots, N$,

$$\begin{aligned} \frac{d}{dt}(|v_i|) &\leq \gamma_0 |v_i| + \frac{\gamma_0}{N} \sum_{j=1}^N |v_j| \leq \gamma_0 |v_i| + \gamma_0 e^{2\gamma_0 t} \frac{1}{N} \sum_{j=1}^N |v_j(0)| \\ &\leq \gamma_0 |v_i| + \gamma_0 e^{2\gamma_0 t} \max_{j=1, \dots, N} |v_j(0)|. \end{aligned}$$

Therefore, again by Gronwall Lemma and uniformly in N ,

$$(34) \quad \begin{aligned} |v_i(t)| &\leq e^{\gamma_0 t} \left(|v_i(0)| + \gamma_0 \max_{j=1, \dots, N} |v_j(0)| \frac{e^{\gamma_0 t} - 1}{\gamma_0} \right) \\ &\leq \max_{j=1, \dots, N} |v_j(0)| e^{2\gamma_0 t}, \quad i = 1, \dots, N. \end{aligned}$$

Finally, (10) gives immediatly that

$$\| |x_1(t)| - |x_i(0)| \| \leq \max_{j=1, \dots, N} |v_j(0)| \frac{e^{2\gamma_0 t} - 1}{2\gamma_0}, \quad i = 1, \dots, N.$$

APPENDIX B. DYNAMICAL ESTIMATES FOR GENERAL CUCKER-SMALE KINETIC SYSTEMS

Let us recall that, according to Theorem 2.3 in [9], there exist a diffeomorphism $\Phi(t)$ on \mathbf{R}^{2d} such that the solution ρ^t of (11) is given by

$$\rho(t) = \Phi(t) \# \rho^{in}.$$

Moreover, $\Phi(t)$ solves the system

$$\dot{\Phi}(t) := \begin{pmatrix} \dot{\Phi}_x(t) \\ \dot{\Phi}_v(t) \end{pmatrix} = \begin{pmatrix} \Phi_v(t) \\ (\gamma * \rho^t)(\Phi(t)) := \int \gamma(\Phi_x(t) - y, \Phi_v(t) - \xi) \rho^t(y, \xi) dx d\xi \end{pmatrix}$$

Therefore,

$$(35) \quad \frac{d}{dt} |\Phi_v(t)| \leq |\dot{\Phi}_v(t)| \leq \gamma_0 (\|\rho^t\|_1 |\Phi_v(t)|) + \int |\xi| \rho^t(y, \xi) dy d\xi.$$

Obviously $\|\rho^t\|_{L^1} = \|\rho^{in}\|_{L^1} = 1$.

Lemma B.1.

$$\int |v| \rho^t dx dv \leq e^{2\gamma_0 t} \int |v| \rho^{in} dx dv.$$

Proof.

$$\begin{aligned} \frac{d}{dt} \int |v| \rho^t dx dv &\leq \left| \int |v| \dot{\rho}^t dx dv \right| \\ &\leq \left| \int |v| \nabla_v \left(\int \gamma(x - y \cdot v - \xi) \rho^t(y, \xi) \rho^t(x, v) dx dv dy d\xi \right) \right| \\ &\leq \int \left| \frac{v}{|v|} \gamma_0 (|v| + |\xi|) \rho^t(y, \xi) \rho^t(x, v) dx dv dy d\xi \right| \\ &\leq 2\gamma_0 \int |v| \rho^t dx dv \end{aligned}$$

The result follows by Gronwall inequality. \square

Thanks to Lemma B.1, (35) becomes

$$\frac{d}{dt}|\Phi_v(t)| \leq \gamma_0|\Phi_v(t)| + \gamma_0 e^{2\gamma_0 t} \int |v|\rho^{in} dx dv,$$

and, by Gronwall Lemma, we have

$$\begin{aligned} |\Phi_v(t)(x, v)| &\leq e^{\gamma_0 t}(|v| + (e^{\gamma_0 t} - 1) \int |v|\rho^{in} dx dv) \\ (36) \qquad \qquad \qquad &\leq e^{2\gamma_0 t}(\|v\|_{L^\infty(\text{supp}[\rho^{in}])} + \|v\|_{L^1(\text{supp}[\rho^{in}])}). \end{aligned}$$

Acknowledgements. The work of Thierry Paul was partly supported by LIA LYSM (co-funded by AMU, CNRS, ECM and INdAM).

REFERENCES

- [1] L. Ambrosio, N. Gigli, G. Savaré, “Gradient Flows in Metric Spaces and in the Space of Probability Measures”, Second Edition, Birkhäuser Verlag, Berlin, 2008.
- [2] F. Cucker and S. Smale, On the mathematics of emergence. *Japan. J. Math.* **2**, 197227, (2007).
- [3] F. Cucker and S. Smale, Emergent behavior in flocks. *IEEE Trans. Automat. Control* **52**, 852–862, 2007.
- [4] F. Golse, C. Mouhot, V. Ricci: *Empirical measures and Vlasov hierarchies*, *Kinetic and Related Models* **6** (2013), 919–943.
- [5] F. Golse, C. Mouhot, T. Paul, On the Mean-Field and Classical Limits of Quantum Mechanics, , *Communication in Mathematical Physics* **343**, 165-205 (2016).
- [6] S.-Y. Ha, J. Kim, X. Zhang, Uniform stability of the Cucker-Smale model and applications to the mean-field limit, *Kinetic and Related Models* **11** (5), 1157–1181 (2018)
- [7] S.-Y. Ha and J.-G. Liu, A simple proof of Cucker-Smale flocking dynamics and mean-field limit, *Commun. Math. Sci.*, **7** (2009), 297–325.
- [8] S.-Y. Ha, E. Tadmor, From particle to kinetic and hydrodynamic descriptions of flocking, *Kinet. Relat. Models* **1** (2008) 415-435.
- [9] B. Piccoli, F. Rossi, E. Trélat, *Control to flocking of the kinetic Cucker-Smale model*, *SIAM J. MATH. ANAL.* . Vol. 47, No. 6, pp. 4685-4719.
- [10] C. Villani: “Topics in Optimal Transportation”, Amer. Math. Soc., Providence (RI), 2003.
- [11] C. Villani: “Optimal Transport. Old and New”, Springer-Verlag, Berlin, Heidelberg, 2009.

(R.N) IAC, VIA DEI TAURINI, 19, 00185 ROMA RM ITALY
E-mail address: roberto.natalini@cnr.it

(T.P.) CNRS & LJLL SORBONNE UNIVERSIT 4 PLACE JUSSIEU 75005 PARIS, FRANCE FRANCE
E-mail address: thierry.paul@upmc.fr