



**HAL**  
open science

## On The Mean Field limit for Cucker-Smale models

R. Natalini, Thierry Paul

► **To cite this version:**

| R. Natalini, Thierry Paul. On The Mean Field limit for Cucker-Smale models. 2020. hal-02008699v3

**HAL Id: hal-02008699**

**<https://hal.science/hal-02008699v3>**

Preprint submitted on 26 Aug 2020 (v3), last revised 5 Feb 2021 (v5)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# ON THE MEAN FIELD LIMIT FOR CUCKER-SMALE MODELS

R. NATALINI AND THIERRY PAUL

ABSTRACT. In this short note, we consider the Cucker-Smale dynamical system and we derive rigorously the Vlasov-type equation introduced in [8] in the mean-field limit without using empirical measures. The vector field we consider is sublinear Lipschitz continuous and the initial data are compactly supported.

## CONTENTS

1.	Introduction	1
2.	The results	4
3.	Proofs	7
3.1.	Preliminaries	7
3.2.	Proof of Theorem 2.4	9
3.3.	Proof of Corollary 2.6	13
3.4.	Proof of Theorem 2.2	14
Appendix A.	Dynamical estimates for general Cucker-Smale particle systems	16
Appendix B.	Dynamical estimates for general Cucker-Smale kinetic systems	16
References		17

## 1. INTRODUCTION

The Cucker-Smale model [2, 3] is a particles system which exhibit the so-called flocking phenomenon, namely the dynamical property of alignment of all the velocities and gathering of all the positions asymptotically when the time of evolution diverges. Its study has developed an intense activity these last years, see for example the intensive bibliography in the recent paper [6].

It consists in the following vector field on  $\mathbf{R}^{2dN}$

$$(1) \quad \begin{cases} \dot{x}_i = v_i \\ \dot{v}_i = \frac{1}{N} \sum_{j=1}^N \psi(x_i - x_j)(v_i - v_j). \end{cases} \quad x_i, v_i \in \mathbf{R}^d, \quad i = 1, \dots, N$$

where  $\psi : \mathbf{R}^d \rightarrow \mathbf{R}$  is a bounded Lipschitz continuous function.

Associated to (1), the following Vlasov type kinetic equation on  $\mathbf{R}^{2d}$  was introduced in [8]:

$$(2) \quad \partial_t f^t(x, v) + v \cdot \nabla_x f^t(x, v) + \nabla_v \cdot \left( f^t(x, v) \int_{\mathbf{R}^{2d}} \psi(x - y)(v - w) f^t(y, w) dy dw \right) = 0.$$

So far the kinetic non-linear equation (2) has been derived in a Lagrangian point of view, i.e. by following the trajectories solving the vector field (1) in the so-called empirical measure

$$(3) \quad \Pi_{(x_1, \dots, x_n; v_1, \dots, v_N)}(x, v) := \frac{1}{N} \sum_{i=1}^N \delta(x - x_i) \delta(v - v_i).$$

Indeed, defining

$$(4) \quad \Pi_{(x_1, \dots, x_n; v_1, \dots, v_N)}^t := \Pi_{\Phi^{-t}(x_1, \dots, x_n; v_1, \dots, v_N)},$$

one shows easily that  $\Pi^t(x_1, \dots, x_n; v_1, \dots, v_N)$  solves (2) for each  $N$ .

This point of view follows directly the Dobrushin way of deriving the Vlasov equation for the large  $N$  limit of Hamiltonian vector fields. The Cucker-Smale model (1) and its Vlasov type associated equation (2) have been extensively studied in [6] and we quote the following of their results: the function  $\psi$  is supposed to be positive, bounded, uniformly Lipschitz continuous and satisfying  $(\psi(r_1) - \psi(r_2))(r_1 - r_2) \leq 0, r_1, r_2 > 0$ .

(1) the solutions of (1) satisfy for all  $t$

$$\sup_{i, j \leq N} \|x_i(t) - x_j(t)\| \leq C \quad \text{and} \quad \sup_{i, j \leq N} \|v_i(t) - v_j(t)\| \leq e^{-Dt}$$

where  $C$  and  $D$  are two positive constants depending only on  $\sup_{i, j \leq N} \|x_i(0) - x_j(0)\|$  and  $\sup_{i, j \leq N} \|v_i(0) - v_j(0)\|$ .

(2) let  $\Pi_{(x_1, \dots, x_n; v_1, \dots, v_N)} \rightarrow \mu(0)$  as  $N \rightarrow \infty$ , then

$$\overline{\lim}_{N \rightarrow \infty} \sup_{t \in \mathbf{R}} W_p(\Pi_{(x_1, \dots, x_n; v_1, \dots, v_N)}^t - \mu(t)) = 0$$

where  $W_p$  is the Wasserstein distance of exponent  $p \in \mathbf{N}$  (see definition in Remark 3.5 below).

(3) Let  $\int_{\mathbf{R}^{2d}} \mu(0) = 1$ ,  $\int_{\mathbf{R}^{2d}} v \mu(0) =: \bar{v}$ ,  $\int_{\mathbf{R}^{2d}} v^2 \mu(0) < \infty$ . Then, for some  $E, F > 0$  and any  $p$ ,

$$\left( \int \|v - \bar{v}\|^p d\mu(t) \right)^{\frac{1}{p}} \leq E e^{-Ft}.$$

Recently, a more Eulerian way of deriving the Vlasov equation associated to Hamiltonian vector fields was introduced ([5], after [4]). It consists in considering the movement of a generic particle solution to the Liouville equation associated to an Hamiltonian system, rather than considering the empirical measure associated to the Hamiltonian flow. Transposed in the (non Hamiltonian) present situation it reads as follows.

We will consider the pushforward<sup>1</sup> by the flow  $\Phi^t$  generated by the system (1) of a compactly supported, symmetric by permutations of the variables  $N$ -body probability density  $\rho_N^{in}$  on phase space  $\mathbf{R}^{2dN}$ .

<sup>1</sup>We recall that the pushforward of a measure  $\mu$  by a measurable function  $\Phi$  is  $\Phi\#\mu$  defined by  $\int \varphi d(\Phi\#\mu) := \int (\varphi \circ \Phi) d\mu$  for any measurable function  $\varphi$ .

In order to describe the movement of a “generic” particle in the limit of diverging number of particles, we want to perform an average on the  $N$  particles but one. This means that we consider the first marginal of

$$(5) \quad \Phi^t \# \rho_N^{in} := \rho_N^t$$

that is

$$(6) \quad \rho_{N;1}^t(x, \xi) := \int_{\mathbf{R}^{2(d-1)N}} \rho_N^t(x, \xi, x_2, \xi_2, \dots, x_N, \xi_N) dx_2 d\xi_2 \dots dx_N d\xi_N.$$

Then we will prove that, when  $\rho_N^{in}$  is factorized as  $\rho_N^{in} = (\rho^{in})^{\otimes N}$ , where  $\rho^{in}$  is a probability measure on  $\mathbf{R}^{2d}$ , the marginal of  $\rho_N^t$  tends, as  $N \rightarrow \infty$ , to the solution of the effective non-linear Vlasov type equation (2):

$$(7) \quad \partial_t \rho^t(x, v) + v \cdot \nabla_x \rho^t + \nabla_v (G_{\rho^t} \rho^t) = 0, \quad \rho^{t=0} = \rho^{in} \in L^1(\mathbf{R}^{2d}, dx dv),$$

with, for any  $L^1$  function  $\rho$ ,

$$(8) \quad G_\rho(x, v) = \int_{\mathbf{R}^{2d}} \psi(x - y)(v - w) \rho(y, w) dy dw.$$

In fact, we can prove (see Remark 3.5 below) that the marginals at any order  $n$  of  $\rho_N^t$  as defined by

$$(9) \quad \rho_{N;n}^t(x_1, \xi_1, \dots, x_n, \xi_n) := \int_{\mathbf{R}^{2dN(-n)}} \rho_N^t(x_i, \xi_i, x_2, \xi_2, \dots, x_N, \xi_N) dx_{n+1} d\xi_{n+1} \dots dx_N d\xi_N,$$

tends, in Wasserstein topology of any exponent  $p \geq 1$ , to the  $n$ th tensorial power of the solution  $\rho^t$  of the Vlasov equation, i.e.  $\rho_{N;n}^t \rightarrow (\rho^t)^{\otimes n}$ . Nevertheless, for sake of clarity of this short note, we will present in detail only the case  $n = 1$ .

This result (for  $n = 1$ ), i.e. Theorem 2.2 below, has to be put in correspondence with the item 2 of the results of [6] just described, but there several differences. First of all our results hold true for more general initial data than the empirical measures. On the other side, our result will not be uniform in time as in item 1, but this inconvenient will be compensated by the fact that we will get an explicit rate of convergence in the asymptotism  $N \rightarrow \infty$ .

Our methods will also apply to systems of the general form

$$(10) \quad \begin{cases} \dot{x}_i = v_i \\ \dot{v}_i = \frac{1}{N} \sum_{j=1}^N \gamma(x_i - x_j, v_i - v_j). \end{cases} \quad x_i, v_i \in \mathbf{R}^d, \quad i = 1, \dots, N$$

where  $\gamma(x, v) : \mathbf{R}^{2d} \rightarrow \mathbf{R}^d$  is a Lipschitz continuous function, bounded in  $(x, v)$ , Corollary 2.5 or bounded in  $x$  and sublinear in  $v$ , Theorem 2.4 and Corollary 2.6.

In the following, we will denote by  $\text{Lip}(\gamma)_{(x,v)}$  the (local) Lipschitz constant of  $\gamma$  at the point  $(x, v) \in \mathbf{R}^{2d}$  and we will suppose, without loss of generality, that there exists  $\gamma_0 > 0$  such that, for all  $(x, v) \in \mathbf{R}^{2d}$ ,

$$\begin{aligned} \gamma(x, v) &\leq \gamma_0 |v| \\ \text{Lip}(\gamma)_{(x,v)} &\leq \gamma_0 |v| \end{aligned}$$

where  $|\cdot|$  denotes the Euclidean norm on  $\mathbf{R}^d$ .

We will show in Appendix A the global existence of the solutions to (10), with an exponential growth in time with respect to the initial conditions. Of course, uniqueness is a consequence of (the iteration in time of) the Cauchy-Lipschitz Theorem.

The Cucker-Smale vector fields satisfies this assumptions with  $\gamma_0 = \|\psi\|_{L^\infty}$ .

Note that the following Vlasov type kinetic equation on  $\mathbf{R}^d$  is obviously associated to (10), in a natural way:

$$(11) \quad \partial_t f^t(x, v) + v \cdot \nabla_x f^t(x, v) + \nabla_v \left( f^t(x, v) \int_{\mathbf{R}^{2d}} \gamma(x - y, v - w) f^t(y, w) dy dw \right) = 0.$$

Let us immediatly notice that our assumptions on  $\gamma$  contain the two assumptions in Theorem 2.3 in [9]. The first one because of the Lipschitz and sublinearity properties of  $\gamma$ , and the second one thanks to the fact, see [10, 11], that

$$\sup_{|\text{Lip}(\varphi)| \leq 1} \left| \int \varphi(\mu - \nu) \right| \leq W_1(\mu, \nu)$$

where  $W_1$  is the Wasserstien distance of exponent 1, as defined in Remark 3.5 in the present paper. Therefore, thanks to the results of [9], there exist a unique continuous solution to (11) in  $C_0(\mathbf{R}, \mathcal{P}_c(\mathbf{R}^{2d}))$ , where  $\mathcal{P}_c(\mathbf{R}^{2d})$  is the space of compactly supported probability measures on  $\mathbf{R}^{2d}$ .

## 2. THE RESULTS

In this section we will state our main results, but let us start by recalling the definition of the second order Wasserstein distance  $\text{dist}_{\text{MK},2}$  (see [1, 10, 11]).

**Definition 2.1** (quadratic Wasserstein distance). *The Wasserstein distance of order two between two probability measures  $\mu, \nu$  on  $\mathbf{R}^m$  with finite second moments is defined as*

$$W_2(\mu, \nu)^2 = \inf_{\gamma \in \Gamma(\mu, \nu)} \int_{\mathbf{R}^m \times \mathbf{R}^m} |x - y|^2 \gamma(dx, dy)$$

where  $\Gamma(\mu, \nu)$  is the set of probability measures on  $\mathbf{R}^m \times \mathbf{R}^m$  whose marginals on the two factors are  $\mu$  and  $\nu$ . *E*

*That is to say that elements  $\gamma$  of  $\Gamma(\mu, \nu)$ , called couplings or transportation maps or transport maps or just maps, depending of the authors, satisfy, for any test functions  $a$  and  $b$  in  $C_0(\mathbf{R}^d)$ ,*

$$\int_{\mathbf{R}^{2m}} a(x) \gamma(dx, dy) = \int_{\mathbf{R}^m} a(x) \mu(dx), \quad \int_{\mathbf{R}^{2m}} b(y) \gamma(dx, dy) = \int_{\mathbf{R}^m} b(y) \nu(dy).$$

The symmetry property in  $\mu, \nu$  is obvious and the separability property is easily proven by taking the optimal coupling between  $\mu$  and itself equal to

$$(12) \quad \gamma = \mu \delta(x - y).$$

Conversely, if  $W_2(\mu, \nu)^2 = 0$ , then  $\int_{\mathbf{R}^m \times \mathbf{R}^m} |x - y|^2 \gamma(dx, dy) = 0$  for some  $\gamma$  so that  $x = y$   $\gamma$  a.e. and, for any Borel function  $\varphi$ ,

$$\int \varphi(x) \mu(dx) = \int \varphi(x) \gamma(dx, dy) = \int \varphi(y) \gamma(dx, dy) = \int \varphi(y) \nu(dy) \Rightarrow \mu = \nu.$$

The proof of the triangular inequality is much more involved, see again [1, 10, 11]).

We can now state the main result of this note, proven in Section 3.4.

**Theorem 2.2** (Cucker-Smale model).

Let  $\rho^t$  be the solution to (7) with an initial condition  $\rho^{in} \in L^1(\mathbf{R}^{2d})$  compactly supported and let  $\rho_{N;1}^t$  be the first marginal of  $\rho_N^t := \Phi^t \# (\rho^{in})^{\otimes N}$ , as defined in (6).

Then, for all  $N > 1$ ,  $t \in \mathbf{R}$ ,

$$W_2(\rho_{N;1}^t, \rho^t) \leq 4 \frac{\|\psi\|_\infty^2 (2|\bar{v}| + |\text{supp}[\rho^{in}]|)}{N^{\frac{1}{2}}} \left( \frac{e^{Lt} - 1}{L} \right)^{\frac{1}{2}}$$

with

$$L := 2(1 + 8\|\psi\|_\infty^2 \|v\|_{L^\infty(\text{supp}[\rho^{in}])}^2), \quad \bar{v} = \int v \rho^{in} dx dv$$

(by  $\|v\|_{L^\infty(\text{supp}[\rho^{in}])}$  we mean  $\sup_{(x,v) \in \text{supp}[\rho^{in}]} (|v|)$ ).

**Remark 2.3.** There exists an equivalent result for higher orders marginals of  $\rho(t)$  and other Monge-Kantorovich distances but we prefer in this note to concentrate on the case of first marginal and quadratic Wasserstein distance. The proof in the more general situation is very close to the one presented here. See Remark 3.5 for some details.

Theorem 2.2 is actually a corollary of the following more general result, proven in Section 3.2

**Theorem 2.4** (General Cucker-Smale model with sublinear force).

Let  $\rho^t$  be the solution to (11),  $\gamma$  bounded on  $\mathbf{R}^{2d}$ , with an initial condition  $\rho_1^{in} \in L^1(\mathbf{R}^{2d})$  compactly supported and  $\Phi^t$  the flow defined by the system (10). Finally, let  $\rho_{N;1}^t$  be the first marginal of  $\rho_N^t := \Phi^t \# (\rho^{in})^{\otimes N}$ , as defined in (6).

Then, for all  $N > 1$ ,  $t \in \mathbf{R}$ ,

$$W_2(\rho_{N;1}^t, \rho^t) \leq C(t) N^{-\frac{1}{2}}$$

with

$$C(t) := \left( 4 \int_0^t \sup_{(x,v), (x',v') \in \text{supp}[\rho^s]} |\gamma(x - x', v - v')|^2 e^{\int_s^t L(u) du} ds \right)^{\frac{1}{2}},$$

$$\underline{L}(t) := 2(1 + 2 \min \left( \sup_{\substack{i,l=1,\dots,N \\ (Y,\Xi) \in \text{supp}[\rho_N^t]}} \text{Lip}(\gamma)_{(y_i - y_l, \xi_i - \xi_l)}^2, \sup_{(x,v), (x',v') \in \text{supp}[\rho^t]} \text{Lip}(\gamma)_{(x - x', v - v')}^2 \right)).$$

The following result is elementarily derived from Theorem 2.4.

**Corollary 2.5** (General Cucker-Smale model with bounded globally Lipschitz force).

Let  $\rho^t$  be the solution to (11),  $\gamma$  globally Lipschitz and bounded on  $\mathbf{R}^{2d}$ , with an initial condition  $\rho^{in} \in L^1(\mathbf{R}^{2d})$  and  $\Phi^t$  the flow defined by the system (10). Finally, let  $\rho_{N;1}^t$  be the first marginal of  $\rho_N^t := \Phi^t \# (\rho^{in})^{\otimes N}$ , as defined in (6).

Then, for all  $N > 1, t \in \mathbf{R}$ ,

$$W_2(\rho_{N;1}^t, \rho^t) \leq C(t)N^{-\frac{1}{2}}$$

with

$$C(t) := \left( 4\|\gamma\|_\infty \frac{e^{\Lambda t} - 1}{\Lambda} \right)^{\frac{1}{2}} \quad \Lambda := 2(1 + 2\text{Lip}(\gamma)^2).$$

Note that no need of compacity of the support of  $\rho^{in}$  is needed any more in Corollary 2.5.

The boundness in space and sublinearity in velocities of  $\gamma$  allow to control the increasing of the flow  $\Phi(t)$ , and leads to our next result, whose proof is given in Section 3.3 below.

**Corollary 2.6.** Under the same hypothesis as in Theorem 2.4 we have that, for all  $N > 1, t \in \mathbf{R}$ ,

$$W_2(\rho_{N;1}^t, \rho^t) \leq C(t)N^{-\frac{1}{2}}$$

with

$$C(t) = 2\gamma_0(\|v\|_{L^\infty(\text{supp}[\rho^{in}])} + \|v\|_{L^1(\text{supp}[\rho^{in}])}) \left( 2e^{e^{4\gamma_0 t} 4\gamma_0(\|v\|_{L^\infty(\text{supp}[\rho^{in}])} + \|v\|_{L^1(\text{supp}[\rho^{in}])})^2} e^{4t}(e^{4\gamma_0 t} - 1) \right)^{\frac{1}{2}},$$

Let us finish this section of results by a simple remark. Flocking properties of the solution of the Vlasov kinetic equation (2) can be derived from the corresponding properties of the particle system (1), as in [7, 6]. But they can also be derived by a direct PDE study of (2), as in [9]. This suggest a kind of inverse questioning: suppose one determines some flocking properties for the solution of a general kinetic Vlasov equation, e.g. (11). Does this infer on flocking properties for the corresponding particle system, e.g. (10)? Our Corollary 2.6 gives some insight on this problem, as it tells us quantitatively how close is the solution of the kinetic equation (and its tensorial powers) to the marginals of any orders of the pushforward obey the corresponding particle flow of a general  $N$  particle density.

More precisely, suppose that the solution  $\rho^t$  to (11) satisfies the following property:

$$(13) \quad \text{supp}[\rho^t] \subset B(\bar{x} + t\bar{v}, X) \times B(\bar{v}, Ve^{-\alpha t})$$

for some positive constants  $\bar{x}, X, \bar{v}, V, \alpha$ .

$$C(t) := \left( 4 \int_0^t \sup_{(x,v),(x',v') \in \text{supp}[\rho^t]} |\gamma(x-x', v-v')|^2 e^{\int_s^t L(u) du} ds \right)^{\frac{1}{2}},$$

$$\underline{L}(t) := 2(1 + 2 \min \left( \sup_{\substack{i,l=1,\dots,N \\ (Y,\Xi) \in \text{supp}[\rho_N^t]}} \text{Lip}(\gamma)_{(y_i-y_l, \xi_i-\xi_l)}, \sup_{(x,v),(x',v') \in \text{supp}[\rho^t]} \text{Lip}(\gamma)_{(x-x', v-v')} \right)).$$

This implies that  $\underline{L}(t)$  in Theorem 2.4 can be easily estimated by

$$\underline{L}(t) \leq 2(1 + 8\gamma_0^2(\bar{v}^2 + V^2 e^{-2\alpha t})) \leq 2(1 + 8\gamma_0^2(\bar{v}^2 + V^2))$$

and, therefore, in the same Theorem,

$$(14) \quad C(t) \leq 4\gamma_0(\bar{v}^2 + V^2) \frac{e^{2(1+8(\bar{v}^2+V^2))t} - 1}{2(1 + 8(\bar{v}^2 + V^2))}.$$

Note that in (14),  $C(t)$  has an exponential growth, and not a double exponential one as in Corollary 2.6.

Of course the first marginal  $\rho_{N,1}^t$  of the pushforward of  $(\rho^{in})^{\otimes N}$  by the flows induced by the system (10) has no reason for being compactly supported, as we did not impose anything on the particle flow. Nevertheless, the following result provides a weak version of the property (13).

**Proposition 2.7.** *For all  $t, \epsilon > 0$  let us define*

$$N_{t,\epsilon} := \left( \frac{C(t)}{\epsilon} \right)^2,$$

where  $C(t)$  is the function defined by (14).

Then, for any  $N \geq N_{t,\epsilon}$  and any Lipschitz function  $\varphi$  on  $\mathbf{R}^{2d}$  of Lipschitz constant smaller than 1,

$$\int_{\mathbf{R}^{2d} \setminus B(\bar{x}+t\bar{v}, X) \times B(\bar{v}, V e^{-\alpha t})} \varphi(x, v) \rho_{N,1}(t) dx dv \leq \epsilon.$$

*Proof.* It is an immediate consequence of Theorem 2.4 or Theorem 2.6, and the fact that the Wasserstein distance induces a metric for the weak topology in the sense that

$$\sup_{\text{Lip}(\varphi) \leq 1} \int (\mu - \nu) \varphi(x, v) dx dv \leq W_2(\mu, \nu)$$

for any probability measures  $\mu$  and  $\nu$  □

### 3. PROOFS

**3.1. Preliminaries.** Let us first recall the general situation we are dealing with, in order also to fix the notations.

We consider on  $\mathbf{R}^{2dN}$  the following Cucker-Smale type vector field

$$(15) \quad \begin{aligned} \dot{x}_i &= v_i \\ \dot{v}_i &= G_i(X, V), \quad i = 1, \dots, N \end{aligned}$$



where

$$(16) \quad G_i(X, V) = \frac{1}{N} \sum_{j=1}^N \gamma(x_i - x_j, v_i - v_j).$$

Here the function  $\gamma(x, v) : \mathbf{R}^{2d} \rightarrow \mathbf{R}^d$  is a Lipschitz, bounded in  $x$  and sublinear in  $v$ , continuous function., such that  $\gamma(x, v), \text{Lip}(\gamma)_{(x,v)} \leq \gamma_0$ .

We used the notation  $X = (x_1, \dots, x_N), V = (v_1, \dots, v_N)$ .

In fact, we are rather interested in the Liouville equation associated to (15) [8], namely

$$(17) \quad \partial_t \rho_N^t + v \cdot \nabla_x \rho_N^t + \sum_{i=1}^N \nabla_{v_i} (G_i \rho_N^t) = 0, \quad \rho_N^{t=0} = \rho_N^{in}$$

with  $\rho_N^{in} \in \mathcal{P}(\mathbf{R}^{2dN})$ .

Although the argument is standard, let us recall why the solution of (17) is equal to the pushforward of  $\rho_N^{in}$  by the flow  $\Phi_t$  generated by (15): integrating  $\rho_N^{in}$  against a test  $\varphi$  function composed by  $\Phi_t$  gives

$$(18) \quad \partial_t \int \varphi(\Phi_t(X, V)) \rho_N^{in}(X, V) dX dV = \int \varphi(X, V) \partial_t (\Phi_t \# \rho_N^{in}(X, V)) dX dV$$

On the other side

$$(19) \quad \begin{aligned} \partial_t \int \varphi(\Phi_t(X, V)) \rho_N^{in}(X, V) dX dV &= \int (\dot{\Phi}_t \cdot \nabla_{(X,V)} \varphi)(\Phi_t(X, V)) \rho_N^{in}(X, V) dX dV \\ &= \int ((V, G) \cdot \nabla_{(X,V)} \varphi)(\Phi_t(X, V)) \rho_N^{in}(X, V) dX dV \\ &= \int ((V, G) \cdot \nabla_{(X,V)} \varphi)(X, V) (\Phi_t \# \rho_N^{in}(X, V)) dX dV \\ &= - \int \varphi(X, V) (\nabla_{(X,V)} \cdot (V, G)) \Phi_t \# \rho_N^{in}(X, V) dX dV \end{aligned}$$

So that (18) and (19) implies that  $\Phi_t \# \rho_N^{in}$  solves (17).

We want to prove that the marginals of  $\rho_N^t$  tend, as  $N \rightarrow \infty$ , to the solution of a Vlasov type equation.

Such Vlasov-type equation associated to (17) has (introduced in [8] for the Cucker-Smale model) reads

$$(20) \quad \partial_t \rho^t(x, v) + v \cdot \nabla_x \rho^t(x, v) + \nabla_v (G_\rho^t \rho^t(x, v)) = 0, \quad \rho^{t=0} = \rho_1^{in} \in L^1(\mathbf{R}^{2d}, dx dv),$$

with

$$(21) \quad G_\rho(x, v) = \int_{\mathbf{R}^{2d}} \gamma(x - y, v - w) \rho(y, w) dy dw.$$

We can now prove the main results of this paper.

**3.2. Proof of Theorem 2.4.** The proof will be articulated around the following four lemmas.

Let  $\pi_N^{in} \in \Pi((\rho^{in})^{\otimes N}, (\rho^{in})^{\otimes N})$  satisfy

$$(22) \quad T_\sigma \# \pi_N^{in} = \pi_N^{in}, \quad \text{for each } \sigma \in \mathfrak{S}_N,$$

where  $\mathfrak{S}_N$  is the group of permutations of  $N$  elements and

$$\begin{aligned} T_\sigma(x_1, v_1, \dots, x_N, v_N, y_1, \xi_1, \dots, y_N, \xi_N) \\ = (x_{\sigma(1)}, v_{\sigma(1)}, \dots, x_{\sigma(N)}, v_{\sigma(N)}, y_{\sigma(1)}, \xi_{\sigma(1)}, \dots, y_{\sigma(N)}, \xi_{\sigma(N)}). \end{aligned}$$

We will denote  $X = (x_1, \dots, x_N)$ ,  $V = (v_1, \dots, v_N)$ ,  $Y = (y_1, \dots, y_N)$ ,  $\Xi = (\xi_1, \dots, \xi_N)$ .

The following first Lemma will be one of the keys of the proof of Theorem 2.2. It consists in considering in evolving a coupling  $\pi_N^{in}$  of two initial conditions of the Liouville (17) and Vlasov (20) equations by the two dynamics of each factor of  $\pi_N^{in}$ .

**Lemma 3.1.** *Let  $\pi_N(t)$  be the (measure) solution to*

$$(23) \quad \partial_t \pi_N + V \cdot \nabla_X \pi_N + \Xi \cdot \nabla_Y \pi_N + \sum_{i=1}^N (\nabla_{\xi_i} \cdot (G_i(Y, \Xi) \pi_N) + \nabla_{v_i} \cdot (G_{\rho^t}(x_i, v_i) \pi_N)) = 0$$

with  $\pi_N(0) = \pi_N^{in}$ .

Then, for all  $t \in \mathbf{R}$ ,  $\pi_N(t)$  is a coupling between  $\rho_N^t$  and  $(\rho^t)^{\otimes N}$ . Moreover  $\pi_N(t)$  is invariant by permutations  $T_\sigma$ .

*Proof.* By taking the two marginals of the two sides of the equality, one get that they satisfy the two Liouville and Vlasov equations. The result is then obtained by uniqueness of the solutions of both equations.  $\square$

**Lemma 3.2.** *Let*

$$\begin{aligned} D_N(t) &:= \int \frac{1}{N} \sum_{j=1}^N (|x_j - y_j|^2 + |v_j - \xi_j|^2) d\pi_N(t) \\ &= \frac{1}{N} \int ((X - Y)^2 + (V - \Xi)^2) d\pi_N(t). \end{aligned}$$

Then

$$\frac{dD_N}{dt} \leq L(t) D_N + \frac{1}{N} \sum_{j=1}^N \int |G_{\rho^t}(x_i, v_i) - G_i(X, V)|^2 (\rho^t)^{\otimes N} dX dV,$$

with

$$(24) \quad L(t) := 2(1 + 2 \min \left( \sup_{\substack{i,l=1,\dots,N \\ X,V,Y,\Xi \in \text{supp}[\pi_N(t)]}} \text{Lip}(\gamma)_{(x_i-x_l, v_i-v_l)}^2, \sup_{\substack{i,l=1,\dots,N \\ X,V,Y,\Xi \in \text{supp}[\pi_N(t)]}} \text{Lip}(\gamma)_{(y_i-y_l, \xi_i-\xi_l)}^2 \right)).$$

*Proof.* We first notice that

$$\frac{dD_N}{dt} = \frac{2}{N} \int \left( (V - \Xi) \cdot (X - Y) + \sum_{i=1}^N (v_i - \xi_i) \cdot (G_{\rho^t}(x_i, v_i) - G_i(Y, \Xi)) \right) d\pi_N$$

Using  $2uv \leq u^2 + v^2$  we get

$$\begin{aligned}
\frac{dD_N}{dt} &\leq \frac{1}{N} \int \left( (X - Y)^2 + 2(V - \Xi)^2 + \sum_{i=1}^N |G_i(Y, \Xi) - G_{\rho^t}(x_i, v_i)|^2 \right) d\pi_N \\
(25) \quad &\leq D_N(t) + \frac{1}{N} \int \left( \sum_{i=1}^N |G_i(Y, \Xi) - G_{\rho^t}(x_i, v_i)|^2 \right) d\pi_N.
\end{aligned}$$

Let us add to  $G_i(Y, \Xi) - G_{\rho^t}(x_i, v_i)$  the null term  $(G_i(X, V) - G_i(X, V))$  so that

$$|G_i(Y, \Xi) - G_{\rho^t}(x_i, v_i)|^2 \leq 2(|G_i(Y, \Xi) - G_i(X, V)|^2 + |G_i(X, V) - G_{\rho^t}(x_i, v_i)|^2)$$

Let us first estimate

$$|G_i(Y, \Xi) - G_i(X, V)|^2 \leq \min(\text{Lip}(G_i)_{(X,V)}^2, \text{Lip}(G_i)_{(Y,\Xi)}^2)(|X - Y|^2 + |V - \Xi|^2).$$

By (16), we have

$$\begin{aligned}
&\min(\text{Lip}(G_i)_{(X,V)}^2, \text{Lip}(G_i)_{(Y,\Xi)}^2) \\
&\leq \min\left(\sup_{l=1,\dots,N} \text{Lip}(\gamma(x_i - x_l, v_i - v_l))^2, \sup_{l=1,\dots,N} \text{Lip}(\gamma(y_i - y_l, \xi_i - \xi_l))^2\right)
\end{aligned}$$

Therefore, by convexity,

$$\begin{aligned}
&\frac{2}{N} \int \sum_{i=1}^N |G_i(Y, \Xi) - G_i(X, V)|^2 \pi_N(dX, dV, dY, d\Xi) \\
&\leq 4 \min\left(\sup_{\substack{i,l=1,\dots,N \\ X,V,Y,\Xi \in \text{supp}[\pi_N(t)]}} \text{Lip}(\gamma(x_i - x_l, v_i - v_l))^2, \sup_{\substack{i,l=1,\dots,N \\ X,V,Y,\Xi \in \text{supp}[\pi_N(t)]}} \text{Lip}(\gamma(y_i - y_l, \xi_i - \xi_l))^2\right) \\
&\quad \times D_N(t)
\end{aligned}$$

And the lemma follows by (25).  $\square$

It remains to estimate in (25) the term

$$\begin{aligned}
&\frac{1}{N} \int \sum_{j=1}^N |G_j(X, V) - G_{\rho^t}(x_j, v_j)|^2 \pi_N^t(dX, dV, dY, d\Xi) \\
&= \frac{1}{N} \int \sum_{j=1}^N |G_j(X, V) - G_{\rho^t}(x_j, v_j)|^2 (\rho^t)^{\otimes N} dX dV
\end{aligned}$$

The following result is a straightforward extension of Lemma 3.3. in [5] with the special value  $p = 2$  and  $d$  replaced by  $2d$ .

**Lemma 3.3.** *Let  $\rho$  be a compactly supported probability density on  $\mathbf{R}^{2d}$  and let  $F$  be a vector field on  $\mathbf{R}^{2d}$ .*

For each  $j = 1, \dots, N$ , one has

$$(26) \quad \int \left| F \star \rho(x_j, v_j) - \frac{1}{N} \sum_{k=1}^N F(x_j - x_k, v_j - v_k) \right|^2 \rho^{\otimes N} dX dV \leq \frac{4}{N} \sup_{(x,v) \in \text{supp}[\rho]} |F(x_j - x, v_j - v)|^2.$$

*Proof.* Let us denote

$$(27) \quad \nu(x, v) := F \star \rho(x_j, v_j) - F(x_j - x, v_j - v)$$

One has

$$\begin{aligned} & \int \left| F \star \rho(x_j, v_j) - \frac{1}{N} \sum_{k=1}^N F(x_j - x_k, v_j - v_k) \right|^2 \rho^{\otimes N} dX dV \\ &= \int \left| \frac{1}{N} \sum_{k=1}^N \nu(x_k, v_k) \right|^2 \rho^{\otimes N} dX dV \\ &= \frac{1}{N^2} \sum_{k,l=1,\dots,N} \int \nu(x_k, v_k) \nu(x_l, v_l) \rho^{\otimes N} dX dV \\ &= \frac{1}{N^2} \sum_{k \neq l=1,\dots,N} \int \nu(x_k, v_k) \nu(x_l, v_l) \rho(x_k, v_k) dx_k dv_k \rho(x_l, v_l) dx_l dv_l \\ &+ \frac{1}{N^2} \sum_{k=1,\dots,N} \int \nu(x_k, v_k)^2 \rho(x_k, v_k) dx_k dv_k \\ &= \frac{1}{N^2} \sum_{k=1,\dots,N} \int \nu(x_k, v_k)^2 \rho(x_k, v_k) dx_k dv_k, \leq \frac{1}{N} \sup_{(x,v) \in \text{supp}[\rho]} \nu(x, v)^2 \end{aligned}$$

since  $\int \rho(x, v) dx dv = 1$  and, by (27),

$$\int \nu(x, v) \rho(x, v) dx dv = 0.$$

By (27) again,  $|\nu(x, v)| \leq 2 \sup_{(x,v) \in \text{supp}[\rho]} |F(x_j - x, v_j - v)|$  and the lemma is proved.  $\square$

Lemma 3.3 with  $F(x, v) = \gamma(x, v)$  together with Lemma 3.2 gives immediatly that

$$\frac{dD_N}{dt} \leq L(t) D_N + \frac{4}{N} \sup_{(x,v),(x',v') \in \text{supp}[\rho^t]} |\gamma(x - x', v - v')|^2$$

and, by Gronwall's inequality,

$$(28) \quad D_N(t) \leq D_N(0) e^{\int_0^t L(s) ds} + \frac{4}{N} \int_0^t \sup_{(x,v),(x',v') \in \text{supp}[\rho^t]} |\gamma(x - x', v - v')|^2 e^{\int_s^t L(u) du} ds.$$

Let us denote by  $(\pi_N(t))_1$  the measure on  $\mathbf{R}^{2d} \times \mathbf{R}^{2d}$  defined, for every test function  $\varphi(x_1, v_1; y_1, \xi_1)$ , by

$$\int_{\mathbf{R}^{2dN} \times \mathbf{R}^{2dN}} \varphi \pi_N(dX, dV; dY d\xi) = \int_{\mathbf{R}^{2d} \times \mathbf{R}^{2d}} \varphi(x_1, v_1; y_1, \xi_1) (\pi_N(t))_1(dx_1, dv_1; dy_1, d\xi_1)$$

We now notice the following obvious fact.

**Lemma 3.4.**  $(\pi_N(t))_1$  is a coupling between  $\rho_{N;1}^t$  and  $\rho^t$ .

Since  $\pi_N(t)$  is symmetric by permutations, one has easily that

$$(29) \quad D_N(t) = \int (|x_1 - y_1|^2 + |v_1 - \xi_1|^2) d(\pi_N(t))_1$$

and Lemma 3.4 immediately implies that

$$(30) \quad D_N(t) \geq W_2(\rho_1(t), (\rho(t))_1)^2,$$

so that, by Lemma 3.3,

$$(31) \quad W_2(\rho_1(t), (\rho(t))_1)^2 \leq D_N(0) e^{\int_0^t L(s) ds} + \frac{16}{N} \int_0^t \sup_{\substack{k,l,1,dots,N \\ (Y,\Xi) \in \text{supp}[\rho_N^t]}} |\gamma(y_k - y_l, \xi_k - \xi_l)|^2 e^{\int_s^t L(u) du} ds.$$

Remember that (31) is valid for  $D(0)$  as defined in Lemma 3.2 for *any*  $\pi_N$  coupling  $(\rho_1^{in})^{\otimes N}$  with itself. Since  $W_2$  is a distance, one has, for any of these  $\pi^{op}$ ,

$$W_2((\rho_1^{in})^{\otimes N}, (\rho_1^{in})^{\otimes N})^2 = \int ((X - Y)^2 + (V - \Xi)^2) d\pi_N^{op} = ND_N(0) = 0$$

so that

$$W_2(\rho_1(t), (\rho(t))_1)^2 \leq \frac{4}{N} \int_0^t \sup_{(x,v),(x',v') \in \text{supp}[\rho^t]} |\gamma(x - x', v - v')|^2 e^{\int_s^t L(u) du} ds.$$

Finally, choosing for  $\pi^{op}$  the optimal coupling (see (12))

$$\pi^{op}(Y, \Xi, X, V) := (\rho_i^{in})^{\otimes N}(X, V) \delta(X - Y) \delta(V - \Xi),$$

one get immediatly that

$$\begin{aligned} (Y, \Xi; X, V) \in \text{supp}[\pi_N(t)] &\Rightarrow (Y, \Xi) \in \text{supp}[\rho_N^t] \text{ and} \\ &\Rightarrow (X, V) \in \text{supp}[(\rho^t)^{\otimes N}] \Leftrightarrow (x_i, v_i) \in \text{supp}[\rho^t], \quad i = 1, \dots, N. \end{aligned}$$

Therefore, after (24)

$$(32) \quad \mathbf{L}(t) := 2(1 + 2 \min \left( \sup_{\substack{i,l=1,\dots,N \\ (Y,\Xi) \in \text{supp}[\rho_N^t]}} \text{Lip}(\gamma)_{(y_i - y_l, \xi_i - \xi_l)}^2, \sup_{(x,v),(x',v') \in \text{supp}[\rho^t]} \text{Lip}(\gamma)_{(x - x', v - v')}^2 \right)).$$

Theorem 2.4 is proven.

**Remark 3.5** (Higher order Wasserstein and marginals). *As we wanted to leap this note as short as possible, we expressed our results only for the first marginal  $\rho_{N;1}$  in the 2-Wasserstein topology, but the method developed in [5] allows as well, with the same kind of modification than the ones used before in this section, to the higher cases. Let us remind, for  $p \geq 1$ , the definition of  $W_p(\mu, \nu)$  for two positive measures  $\mu, \nu$  (cf. Definition 2.1*

$$W_p(\mu, \nu)^p = \inf_{\gamma \in \Gamma(\mu, \nu)} \int_{\mathbf{R}^m \times \mathbf{R}^m} |x - y|^p \gamma(dx, dy),$$

and, for any probability measure  $\rho$  on  $\mathbf{R}^{2dN}$  and  $n = 1, \dots, N$ , the definition of the  $n$ th marginal  $\rho_{N;n}$  defined on  $\mathbf{R}^{2dn}$

$$\rho_{N;n}(x_1, \dots, x_n, v_1, \dots, v_n) := \int_{\mathbf{R}^{2d(N-n)}} \rho((x_1, \dots, x_n, x_{n+1}, \dots, x_N, v_1, \dots, v_n, v_{n+1}, \dots, v_N)) dx_{n+1} \dots dx_N dv_{n+1} \dots dv_N.$$

One gets, for each  $p \geq 1$ ,  $N \geq 1$  and  $n = 1, \dots, N$ ,

$$\frac{1}{n} W_p(\rho_{N;n}^t, (\rho^t)^{\otimes n})^p \leq D_{p,n}(t) N^{-\min(p/2, 1)}$$

with

$$D_{p,n}(t) = 2^{2p} \max(1, p-1) ([p/2]+1) \int_0^t \sup_{\substack{k,l=1,\dots,N \\ (Y,\Xi) \in \text{supp}[\rho_N^t]}} |\gamma(y_k - y_l, \xi_k - \xi_l)|^p e^{2 \max(1, p-1) \int_s^t L(u) du} ds$$

where

$$L(u) = 1 + 2^{p-1} \sup_{(x,v),(y,\xi) \in \text{supp}[\rho^t]} \text{Lip}(\gamma)_{(x-y, v-\xi)}^p.$$

The main changes in the proof are the use of the Young inequality

$$p u v^{p-1} \leq u^p + (p-1) v^p \leq \max(1, p-1) (u^p + v^p)$$

instead of  $2uv \leq u^2 + v^2$  before (25), the convexity of  $|\cdot|^p$  for  $p \geq 1$  and a variant of the contents of Lemma 3.3 which reads

$$\begin{aligned} & \int \left| F \star \rho(x_j, v_j) - \frac{1}{N} \sum_{k=1}^N F(x_j - x_k, v_i - v_j) \right|^p \rho^{\otimes N} dX dV \\ & \leq \frac{2[p/2] + 2}{N^{\min(p/2, 1)}} \sup_{\substack{k,l=1,\dots,N \\ (Y,\Xi) \in \text{supp}[\rho_N^t]}} |F(y_k - y_l, \xi_k - \xi_l)|^p. \end{aligned}$$

Finally, the statement of Lemma 3.4 becomes now easily that  $\pi_N(t)_n$  is a coupling between  $\rho_{N;n}^t$  and  $(\rho^t)^{\otimes n}$ , where  $\pi_N(t)_n$  is defined through by

$$\int_{\mathbf{R}^{2dN} \times \mathbf{R}^{2dN}} \varphi \pi_N(dX, dV; dY d\Xi) = \int_{\mathbf{R}^{2d} \times \mathbf{R}^{2d}} \varphi \pi_N(t)_n(d(x, v; y, \xi)_1 \dots d(x, v; y, \xi)_n)$$

for every test function  $\varphi((x, v; y, \xi)_1, \dots, (x, v; y, \xi)_n)$ ,

**3.3. Proof of Corollary 2.6.** We get to estimates  $L(t)$  and  $C(t)$  as given in Theorem 2.4 out of the estimates established in Appendices A and B.

Let us recall that

$$\begin{aligned}
\underline{L}(t) &:= 2(1 + 2 \min ( \sup_{\substack{i,l=1,\dots,N \\ (Y,\Xi) \in \text{supp}[\rho_N^t]}} \text{Lip}(\gamma)_{(y_i-y_l, \xi_i-\xi_l)}^2, \sup_{(x,v),(x',v') \in \text{supp}[\rho^t]} \text{Lip}(\gamma)_{(x-x',v-v')}^2 )). \\
&\leq 2(1 + 2 \sup_{(x,v),(x',v') \in \text{supp}[\rho^t]} \text{Lip}(\gamma)_{(x-x',v-v')}^2). \\
&\leq 2(1 + 2\gamma_0^2 \sup_{(x,v),(x',v') \in \text{supp}[\rho^t]} |v - v'|^2) \\
&\leq 2(1 + 8\gamma_0^2 \sup_{(x,v) \in \text{supp}[\rho^t]} |v|^2) \\
&\leq 2(1 + 8\gamma_0^2 \sup_{(x,v) \in \text{supp}[\rho^{in}]} |\Phi_v(t)(x, v)|^2)
\end{aligned}$$

Thanks to (37), we get

$$\underline{L}(t) \leq 2(1 + 16\gamma_0^2 e^{4\gamma_0 t} (\|v\|_{L^\infty(\text{supp}[\rho^{in}])} + \|v\|_{L^1(\text{supp}[\rho^{in}])})^2)$$

By the same argument, we get

$$\begin{aligned}
C(t)^2 &:= 4 \int_0^t \sup_{(x,v),(x',v') \in \text{supp}[\rho^t]} |\gamma(x - x', v - v')|^2 e^{\int_s^t L(u) du} ds, \\
&\leq 32\gamma_0^2 (\|v\|_{L^\infty(\text{supp}[\rho^{in}])} + \|v\|_{L^1(\text{supp}[\rho^{in}])})^2 \\
&\quad \times \int_0^t e^{4(\gamma_0 s + t - s)} e^{16\gamma_0^2 (\|v\|_{L^\infty(\text{supp}[\rho^{in}])} + \|v\|_{L^1(\text{supp}[\rho^{in}])})^2 (e^{4\gamma_0 t} - e^{4\gamma_0 s}) / 4\gamma_0} ds \\
&\leq 8\gamma_0^2 (\|v\|_{L^\infty(\text{supp}[\rho^{in}])} + \|v\|_{L^1(\text{supp}[\rho^{in}])})^2 e^{4\gamma_0 t} 4\gamma_0 (\|v\|_{L^\infty(\text{supp}[\rho^{in}])} + \|v\|_{L^1(\text{supp}[\rho^{in}])})^2 e^{4t} (e^{4\gamma_0 t} - 1).
\end{aligned}$$

Corollary 2.6 is proven.

**3.4. Proof of Theorem 2.2.** In order to prove Theorem 2.2, we need to estimate, in the Cucker-Smale particular case, that is when

$$\gamma(x, v) = \psi(x)v,$$

the two quantities

$$\begin{aligned}
C(t) &:= \left( 4 \int_0^t \sup_{(x,v),(x',v') \in \text{supp}[\rho^t]} |\gamma(x-x', v-v')|^2 e^{\int_s^t L(u) du} ds \right)^{\frac{1}{2}}, \\
&\leq 2 \|\psi\|_\infty \left( \int_0^t \sup_{(x,v),(x',v') \in \text{supp}[\rho^t]} |v-v'|^2 e^{\int_s^t L(u) du} ds \right)^{\frac{1}{2}}, \\
\underline{L}(t) &:= 2(1 + 2 \min \left( \sup_{\substack{i,l=1,\dots,N \\ (Y,\Xi) \in \text{supp}[\rho_N^t]}} \text{Lip}(\gamma)_{(y_i-y_l, \xi_i-\xi_l)}^2, \sup_{(x,v),(x',v') \in \text{supp}[\rho^t]} \text{Lip}(\gamma)_{(x-x', v-v')}^2 \right)). \\
&\leq 2(1 + 2 \sup_{\substack{i,l=1,\dots,N \\ (Y,\Xi) \in \text{supp}[\rho_N^t]}} \text{Lip}(\gamma)_{(y_i-y_l, \xi_i-\xi_l)}^2) \\
&\leq 2(1 + 2 \|\psi\|_\infty^2 \sup_{\substack{i,l=1,\dots,N \\ (Y,\Xi) \in \text{supp}[\rho_N^t]}} |\xi_i - \xi_l|^2)
\end{aligned}$$

We will need just a very little part of the stability results expressed by Ha, Kim and Zhuang, namely the following lemma, taken from formula (8) in Lemma 2.1 in [6]:

$$\frac{d}{dt} \sup_{k,l=1,\dots,N} |v_k(t) - v_l(t)| \leq 0, \quad \forall t,$$

which leads naturally to

$$\sup_{\substack{i,l=1,\dots,N \\ (Y,\Xi) \in \text{supp}[\rho_N^t]}} |\xi_i - \xi_l|^2 \leq 4 \|v\|_{L^\infty(\text{supp}[\rho^{in}])}^2.$$

This implies

$$\underline{L}(t) \leq 2(1 + 8 \|\psi\|_\infty^2 \|v\|_{L^\infty(\text{supp}[\rho^{in}])}^2) := L$$

We will estimate  $\sup_{(x,v),(y,\xi) \in \text{supp}[\rho^t]} |v - \xi|^2 \leq 2 \sup_{(x,v),(y,\xi) \in \text{supp}[\rho^t]} (|v|^2 + |\xi|^2)$  wthanks to Lemma 3.2 in [9] which stipulates that

$$\text{supp}_v[\rho^t] \subset B(\bar{v}, V(t))$$

with  $\bar{v} = \int v \rho^{in} dx dv$  and  $\frac{d}{dt} V(t) \leq 0$ .

Therefore  $\|v\|_{L^\infty(\text{supp}[\rho^t])} \leq |\bar{v}| + V(0)$  so that, since one can take  $V(0) = |\bar{v}| + |\text{supp}[\rho^{in}]|$ ,

$$\sup_{(x,v),(x',v') \in \text{supp}[\rho^t]} |v - v'|^2 \leq 2 \|\psi\|_\infty^2 (2|\bar{v}| + |\text{supp}[\rho^{in}]|)^2.$$

and

$$\begin{aligned}
C(t) &\leq 2 \|\psi\|_\infty^2 \left( 2 \int_0^t (2|\bar{v}| + |\text{supp}[\rho^{in}]|)^2 e^{L(t-s)} ds \right)^{\frac{1}{2}} \\
&= 2 \|\psi\|_\infty^2 \left( 2(2|\bar{v}| + |\text{supp}[\rho^{in}]|)^2 \frac{e^t - 1}{L} \right)^{\frac{1}{2}}
\end{aligned}$$

Theorem 2.2 is proved.



## APPENDIX A. DYNAMICAL ESTIMATES FOR GENERAL CUCKER-SMALE PARTICLE SYSTEMS

In this section, we give global estimates on the flow  $\Phi^N(t)$  generated by (10). We have, for each  $i = 1, \dots, N$ ,

$$\begin{aligned}
 \frac{d}{dt}|v_i| &\leq |\dot{v}_i| \\
 &\leq \frac{1}{N} \sum_{j=1}^N |\gamma(x_i - v_j, v_i - v_j)| \leq \frac{1}{N} \sum_{j=1}^N \gamma_0 |v_i - v_j| \\
 (33) \quad &\leq \frac{1}{N} \sum_{j=1}^N \gamma_0 (|v_j| + |v_i|),
 \end{aligned}$$

so that

$$\frac{d}{dt} \sum_{i=1}^N |v_i| \leq 2\gamma_0 \sum_{i=1}^N |v_i|.$$

and, by Gronwall inequality,

$$(34) \quad \sum_{i=1}^N |v_i(t)| \leq \sum_{i=1}^N |v_i(0)| e^{2\gamma_0 t}.$$

Turning back to (33), we get by (34), for each  $i = 1, \dots, N$ ,

$$\begin{aligned}
 \frac{d}{dt}(|v_i|) &\leq \gamma_0 |v_i| + \frac{\gamma_0}{N} \sum_{j=1}^N |v_j| \leq \gamma_0 |v_i| + \gamma_0 e^{2\gamma_0 t} \frac{1}{N} \sum_{j=1}^N |v_j(0)| \\
 &\leq \gamma_0 |v_i| + \gamma_0 e^{2\gamma_0 t} \max_{j=1, \dots, N} |v_j(0)|.
 \end{aligned}$$

Therefore, again by Gronwall Lemma and uniformly in  $N$ ,

$$\begin{aligned}
 |v_i(t)| &\leq e^{\gamma_0 t} \left( |v_i(0)| + \gamma_0 \max_{j=1, \dots, N} |v_j(0)| \frac{e^{\gamma_0 t} - 1}{\gamma_0} \right) \\
 (35) \quad &\leq \max_{j=1, \dots, N} |v_j(0)| e^{2\gamma_0 t}, \quad i = 1, \dots, N.
 \end{aligned}$$

Finally, (10) gives immediatly that

$$\| |x_1(t)| - |x_i(0)| \| \leq \max_{j=1, \dots, N} |v_j(0)| \frac{e^{2\gamma_0 t} - 1}{2\gamma_0}, \quad i = 1, \dots, N.$$

## APPENDIX B. DYNAMICAL ESTIMATES FOR GENERAL CUCKER-SMALE KINETIC SYSTEMS

Let us recall that, according to Theorem 2.3 in [9], there exist a diffeomorphism  $\Phi(t)$  on  $\mathbf{R}^{2d}$  such that the solution  $\rho^t$  of (11) is given by

$$\rho(t) = \Phi(t) \# \rho^{in}.$$

Moreover,  $\Phi(t)$  solves the system

$$\dot{\Phi}(t) := \begin{pmatrix} \dot{\Phi}_x(t) \\ \dot{\Phi}_v(t) \end{pmatrix} = \begin{pmatrix} \Phi_v(t) \\ (\gamma * \rho^t)(\Phi(t)) := \int \gamma(\Phi_x(t) - y, \Phi_v(t) - \xi) \rho^t(y, \xi) dx d\xi \end{pmatrix}$$

Therefore,

$$(36) \quad \frac{d}{dt} |\Phi_v(t)| \leq |\dot{\Phi}_v(t)| \leq \gamma_0 (\|\rho^t\|_1 |\Phi_v(t)| + \int |\xi| \rho^t(y, \xi) dy d\xi).$$

Obviously  $\|\rho^t\|_{L^1} = \|\rho^{in}\|_{L^1} = 1$ .

**Lemma B.1.**

$$\int |v| \rho^t dx dv \leq e^{2\gamma_0 t} \int |v| \rho^{in} dx dv.$$

*Proof.*

$$\begin{aligned} \frac{d}{dt} \int |v| \rho^t dx dv &\leq \left| \int |v| \dot{\rho}^t dx dv \right| \\ &\leq \left| \int |v| \nabla_v \left( \int \gamma(x - y, v - \xi) \rho^t(y, \xi) \rho^t(x, v) dx dv dy d\xi \right) \right| \\ &\leq \int \left| \frac{v}{|v|} \gamma_0 (|v| + |\xi|) \rho^t(y, \xi) \rho^t(x, v) dx dv dy d\xi \right| \\ &\leq 2\gamma_0 \int |v| \rho^t dx dv \end{aligned}$$

The result follows by Gronwall inequality. □

Thanks to Lemma B.1, (36) becomes

$$\frac{d}{dt} |\Phi_v(t)| \leq \gamma_0 |\Phi_v(t)| + \gamma_0 e^{2\gamma_0 t} \int |v| \rho^{in} dx dv,$$

and, by Gronwall Lemma, we have

$$(37) \quad \begin{aligned} |\Phi_v(t)(x, v)| &\leq e^{\gamma_0 t} (|v| + (e^{\gamma_0 t} - 1) \int |v| \rho^{in} dx dv) \\ &\leq e^{2\gamma_0 t} (\|v\|_{L^\infty(\text{supp}[\rho^{in}])} + \|v\|_{L^1(\text{supp}[\rho^{in}])}). \end{aligned}$$

**Acknowledgements.** The work of Thierry Paul was partly supported by LIA LYSM (co-funded by AMU, CNRS, ECM and INdAM).

## REFERENCES

- [1] L. Ambrosio, N. Gigli, G. Savaré, “Gradient Flows in Metric Spaces and in the Space of Probability Measures”, Second Edition, Birkhäuser Verlag, Berlin, 2008.
- [2] F. Cucker and S. Smale, On the mathematics of emergence. *Japan. J. Math.* **2**, 197227, (2007).
- [3] F. Cucker and S. Smale, Emergent behavior in flocks. *IEEE Trans. Automat. Control* **52**, 852–862, 2007.
- [4] F. Golse, C. Mouhot, V. Ricci: *Empirical measures and Vlasov hierarchies*, *Kinetic and Related Models* **6** (2013), 919–943.
- [5] F. Golse, C. Mouhot, T. Paul, On the Mean-Field and Classical Limits of Quantum Mechanics, *Communication in Mathematical Physics* **343**, 165–205 (2016).

- [6] S.-Y. Ha, J. Kim, X. Zhang, Uniform stability of the Cucker-Smale model and applications to the mean-field limit, *Kinetic and Related Models* **11** (5), 1157–1181 (2018)
- [7] S.-Y. Ha and J.-G. Liu, A simple proof of Cucker-Smale flocking dynamics and mean-field limit, *Commun. Math. Sci.*, **7** (2009), 297–325.
- [8] S.-Y. Ha, E. Tadmor, From particle to kinetic and hydrodynamic descriptions of flocking, *Kinet. Relat. Models* **1** (2008) 415-435.
- [9] B. Piccoli, F. Rossi, E. Trélat, *Control to flocking of the kinetic Cucker-Smale model*, *SIAM J. MATH. ANAL.* . Vol. 47, No. 6, pp. 46854719.
- [10] C. Villani: “Topics in Optimal Transportation”, Amer. Math. Soc., Providence (RI), 2003.
- [11] C. Villani: “Optimal Transport. Old and New”, Springer-Verlag, Berlin, Heidelberg, 2009.

(R.N) IAC, VIA DEI TAURINI, 19, 00185 ROMA RM ITALY  
*Email address:* `roberto.natalini@cnr.it`

(T.P.) CNRS & LJLL SORBONNE UNIVERSIT 4 PLACE JUSSIEU 75005 PARIS, FRANCE FRANCE  
*Email address:* `thierry.paul@upmc.fr`