



**HAL**  
open science

## The SOPHIE search for northern extrasolar planets XIII. Two planets around M-dwarfs Gl617A and Gl96

M. J. Hobson, R. F. Díaz, X. Delfosse, N. Astudillo-Defru, I. Boisse, F. Bouchy, X. Bonfils, T. Forveille, N. Hara, L. Arnold, et al.

► **To cite this version:**

M. J. Hobson, R. F. Díaz, X. Delfosse, N. Astudillo-Defru, I. Boisse, et al.. The SOPHIE search for northern extrasolar planets XIII. Two planets around M-dwarfs Gl617A and Gl96. *Astronomy and Astrophysics - A&A*, 2018, 618, pp.A103. 10.1051/0004-6361/201832732 . hal-02009759

**HAL Id: hal-02009759**

**<https://hal.science/hal-02009759v1>**

Submitted on 6 Feb 2019

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# The SOPHIE search for northern extrasolar planets<sup>★</sup>

## XIII. Two planets around M-dwarfs Gl617A and Gl96

M. J. Hobson<sup>1</sup>, R. F. Díaz<sup>2,3</sup>, X. Delfosse<sup>4</sup>, N. Astudillo-Defru<sup>5</sup>, I. Boisse<sup>1</sup>, F. Bouchy<sup>5</sup>, X. Bonfils<sup>4</sup>, T. Forveille<sup>4</sup>, N. Hara<sup>5,6</sup>, L. Arnold<sup>7</sup>, S. Borgniet<sup>4</sup>, V. Bourrier<sup>5</sup>, B. Brugger<sup>1</sup>, N. Cabrera<sup>4</sup>, B. Courcol<sup>1</sup>, S. Dalal<sup>8</sup>, M. Deleuil<sup>1</sup>, O. Demangeon<sup>9</sup>, X. Dumusque<sup>5</sup>, D. Ehrenreich<sup>5</sup>, G. Hébrard<sup>8,7</sup>, F. Kiefer<sup>8</sup>, T. Lopez<sup>1</sup>, L. Mignon<sup>4</sup>, G. Montagnier<sup>8,7</sup>, O. Mousis<sup>1</sup>, C. Moutou<sup>1,11</sup>, F. Pepe<sup>5</sup>, J. Rey<sup>5</sup>, A. Santerne<sup>1</sup>, N. Santos<sup>9,10</sup>, M. Stalport<sup>5</sup>, D. Ségransan<sup>5</sup>, S. Udry<sup>5</sup>, and P.A. Wilson<sup>12,8</sup>

<sup>1</sup> Aix Marseille Université, CNRS, Laboratoire d'Astrophysique de Marseille UMR 7326, 13388 Marseille cedex 13, France  
e-mail: melissa.hobson@lam.fr

<sup>2</sup> Universidad de Buenos Aires, Facultad de Ciencias Exactas y Naturales. Buenos Aires, Argentina

<sup>3</sup> CONICET - Universidad de Buenos Aires. Instituto de Astronomía y Física del Espacio (IAFE). Buenos Aires, Argentina

<sup>4</sup> Univ. Grenoble Alpes, CNRS, IPAG, 38000 Grenoble, France

<sup>5</sup> Observatoire Astronomique de l'Université de Genève, 51 Chemin des Maillettes, 1290 Versoix, Switzerland

<sup>6</sup> ASD/IMCCE, CNRS-UMR8028, Observatoire de Paris, PSL, UPMC, 77 Avenue Denfert-Rochereau, 75014 Paris, France

<sup>7</sup> Observatoire de Haute Provence, CNRS, Aix Marseille Université, Institut Pythéas UMS 3470, 04870 Saint-Michel-l'Observatoire, France

<sup>8</sup> Institut d'Astrophysique de Paris, UMR7095 CNRS, Université Pierre & Marie Curie, 98bis boulevard Arago, 75014 Paris, France

<sup>9</sup> Instituto de Astrofísica e Ciências do Espaço, Universidade do Porto, CAUP, Rua das Estrelas, 4150-762 Porto, Portugal

<sup>10</sup> Departamento de Física e Astronomia, Faculdade de Ciências, Universidade do Porto, Rua do Campo Alegre, 4169-007 Porto, Portugal

<sup>11</sup> Canada-France-Hawaii Telescope Corporation, 65-1238 Mamalahoa Hwy, Kamuela, HI 96743, USA

<sup>12</sup> Leiden Observatory, Leiden University, Postbus 9513, 2300 RA Leiden, The Netherlands

Received January 30, 2018, accepted June 27, 2018

### ABSTRACT

We report the detection of two exoplanets and a further tentative candidate around the M-dwarf stars Gl96 and Gl617A, based on radial velocity measurements obtained with the SOPHIE spectrograph at the Observatoire de Haute Provence. Both stars were observed in the context of the SOPHIE exoplanet consortium's dedicated M-dwarf subprogramme, which aims to detect exoplanets around nearby M-dwarf stars through a systematic survey. For Gl96, we present the discovery of a new exoplanet at 73.9 d with a minimum mass of 19.66 earth masses. Gl96 b has an eccentricity of 0.44, placing it among the most eccentric planets orbiting M stars. For Gl617A we independently confirm a recently reported exoplanet at 86.7 d with a minimum mass of 31.29 earth masses. Both Gl96 b and Gl617A b are potentially within the habitable zone, though Gl96 b's high eccentricity may take it too close to the star at periapsis.

**Key words.** Techniques: radial velocities – planetary systems – stars: late-type – stars: individual: Gl617A, Gl96

## 1. Introduction

M-dwarf stars are both interesting and promising targets for exoplanet hunts. They are the most common stars in the Galaxy, and studies suggest their planet occurrence rates are high (e.g. Bonfils et al. 2013, Dressing & Charbonneau 2015). Moreover, they are interesting candidates for habitable planet searches. Their relatively small masses (0.07 – 0.6 $M_{\odot}$ , Reid & Hawley 2005) mean that small planets will still induce detectable signals, while their faintness compared to G-type stars means the habitable zone is located closer to the stars. Hence, low-mass short-period habitable planets are easier to detect for M-dwarfs than for sun-like stars - e.g. Gl 667C c, the first habitable-zone Earth-size planet around an M-dwarf (Delfosse et al. 2013); LHS 1140 b, one of the most recently detected ones, orbiting the brightest M-

dwarf with a transiting planet in the habitable zone (Dittmann et al. 2017); TRAPPIST-1 e, f, g, three potentially habitable telluric planets in a seven-planet system (Gillon et al. 2017); GJ 273, with two super-Earths one of which is in the habitable zone (Astudillo-Defru et al. 2017c); K2-18b, a transiting habitable-zone planet whose mass was characterised by RV follow-up, revealing a density that may correspond to a rocky planet with extended atmosphere or to a water world (Foreman-Mackey et al. 2015; Cloutier et al. 2017).

Currently, 146 exoplanets around main sequence M-dwarf stars are known, compared to 997 planets around FGK stars<sup>1</sup>. However, this number is expected to grow, as several current or near-future projects have M-dwarf stars as part of their (or their sole) primary targets, e.g. SPIRou (Artigau et al. 2014), TESS (NASA mission, launch 2018, Ricker 2016), TRAPPIST (e.g.

<sup>★</sup> Based on observations collected with the SOPHIE spectrograph on the 1.93-m telescope at Observatoire de Haute-Provence (CNRS), France by the SOPHIE Consortium.

<sup>1</sup> Retrieved on Oct 11, 2017, from "The Exoplanet Encyclopaedia" 1995-2017, considering only the stars for which spectral type is reported in the catalogue, and filtering out those not on the main sequence.

Gillon et al. 2017), CARMENES (e.g. Quirrenbach & Carrenes Consortium 2016), HADES (e.g. Affer et al. 2016), NIRPS (Bouchy et al. 2017), ExTrA (Bonfils et al. 2015). Given the relatively low number of detected exoplanets around M-dwarfs, each new detection provides valuable information on the population that can be used to refine observing strategies.

Since 2006, the SOPHIE exoplanet consortium has been carrying out several planet-hunting programmes using the SOPHIE spectrograph at the Observatoire de Haute-Provence (Bouchy et al. 2009a). Sub-programme 3, or SP3, consists of a systematic survey of nearby M-dwarfs which aims at detecting habitable SuperEarths and Neptunes, constraining the statistics of planets around M-dwarfs, and finding potentially transiting companions. A complete description of the SP3 is beyond the scope of this paper; the programme will be presented in detail in a forthcoming paper.

In this work, we report the results of the SP3 study of two M-dwarfs, G196 and G1617A. Section 2 presents the observations. In Section 3 we describe the analysis of the data. Our results are presented in Section 4, together with an analysis of the Hipparcos photometry of these stars in Section 5, and discussed in Section 6.

## 2. Observations

G196 and G1617A were observed with the SOPHIE spectrograph as part of the SOPHIE consortium search for exoplanets around M-dwarfs. SOPHIE is a fibre-fed, environmentally stabilised, cross-dispersed echelle spectrograph mounted on the 193 cm telescope at the Observatoire de Haute-Provence (Perruchot et al. 2008). In 2011, SOPHIE was upgraded by inserting an octagonal-section fibre in the fibre link (Perruchot et al. 2011; Bouchy et al. 2013). The upgraded spectrograph is known as SOPHIE+.

The observations presented here were performed after the 2011 upgrade, using the high-resolution (HR) mode of the spectrograph, for a resolving power of  $\lambda/\delta\lambda \approx 75000$ . The SOPHIE spectrograph provides two modes for fibre B: *thosimult* mode, in which a simultaneous ThAr calibration is performed and is used to trace the spectrograph drift during the night, and *objAB* mode, in which fibre B is used to monitor the sky brightness. The choice of modes for the SP3 targets depends on the brightness of the stars (with a limit at  $V=9$ ). G196, which is above this limit, was observed in the *objAB* mode to control for possible moonlight contamination and avoid any potential ThAr contamination, with a calibration lamp spectrum obtained immediately before each observation to monitor potential drifts. G1617A, which is brighter, was observed in the *thosimult* mode, since it is bright enough that the contamination by moonlight can be neglected, and contamination from the ThAr spectrum on fibre B is negligible.

Observations were gathered between 2011 and 2017. In total, 79 spectra were obtained for G196 with SOPHIE+, with a median exposure time of 1800s and a median SNR at 550 nm of 82.7. 163 spectra were obtained for G1617A, with a median exposure time of 900s and a median SNR at 550 nm of 84.1.

The SOPHIE pipeline (Bouchy et al. 2009a) was used to reduce and extract the spectra. The spectra were then cross-correlated with an M3 stellar spectral mask in order to obtain the cross-correlation functions (CCFs), from which radial velocities (RV), FWHM, contrast, and bisectors were measured. The mask was built from the median of a large number of spectra of G1581 obtained with HARPS (La Silla, ESO) and degraded to the SOPHIE spectral resolution.


For G1617A, we removed 9 spectra with  $\text{SNR} < 35$ . For G196, we removed a total of 8 observations: 2 spectra following the same SNR criteria as for G1617A, and 6 spectra due to moon contamination.

## 3. Data Analysis

Radial velocities for stabilised spectrographs with ThAr-derived wavelength calibration have usually been obtained by the CCF method, in which the spectra are correlated with a weighted binary mask (see Queloz 1995 and Pepe et al. 2002 for full descriptions of the method). Although the CCF method is very effective for FGK stars, which have strong spectral lines and a well defined continuum, this is not the case for M-dwarfs, where numerous overlapping molecular bands complicate the continuum determination. For these stars, the use of these binary masks which target only clearly defined lines underutilises the Doppler information present in the spectrum. Therefore, other methods have been developed to better exploit this information, such as template-matching using a true stellar template (as done, e.g., by the HARPS-TERRA code of Anglada-Escudé & Butler 2012). Template-matching also allows a more precise removal of the telluric lines, and of any parts of the spectra that are not compliant with the template or that have no spectral information. In this work, we make use of an algorithm developed by N. Astudillo-Defru (Astudillo-Defru et al. 2015, Astudillo-Defru et al. 2017c) for this purpose; it constructs stellar and telluric templates from the observed spectra, discards the telluric-contaminated zones, and derives the radial velocity by  $\chi^2$  minimisation, using the RV determined by the CCF method as a first guess. We applied this method to all the SOPHIE+ spectra.

Before employing the template-matching procedure, we performed a correction for the charge transfer inefficiency (CTI) effect, following the characterisation of Bouchy et al. (2009b). Once the RVs were determined, we added a further correction for the instrumental drift using the simultaneous ThAr calibration (in the case of G1617A) or an interpolation between ThAr calibrations performed before and after the exposure (for G196).

The SOPHIE+ spectrograph also presents long-term variations of the zero-point, an effect described in Courcol et al. (2015). In order to correct it, the authors iteratively construct a master RV time series from RV constant stars. This master series is used to correct the measurements of a given target by subtracting the velocity in the master series, interpolated at the dates of observation. We followed the procedure described in Courcol et al. (2015) to construct an analogous master for the SP3 programme, using the ensemble of SP3 targets with at least 10 SOPHIE+ observations plus the four 'super-constants' from the SP1 programme (defined by Courcol et al. 2015) as a starting point. We chose to use primarily SP3 targets when constructing our master correction in order to mitigate any potential bias or offset due to differing spectral types. The final master employs 14 stars: the SP1 (G-type) constants HD185144, HD9407, HD22154, and HD89269A; three M-dwarfs which are systematically observed for all observation seasons and considered as our SP3 constants, G1411, G1514, and G1686; and the additional SP3 stars G239-25, G1133, G115A, G1436, G1521, G1694, and G1728 (all additional stars used for the constant correction have a corrected rms after the first iteration lower than 3 m/s, as defined by Courcol et al. 2015). We also tested a master constructed using only the SP3 targets, but found the resulting RVs had higher dispersion (around 1.5 m/s higher on average). Figure 1 shows the master used for the zero-point drift correction and the RVs from which it was derived.



**Fig. 1.** Zero-point drift correction (red line) and radial velocities of the 14 stars used to construct it (black dots). The vertical lines indicate identified jumps and their causes (see Section 3 for details). The zero-point correction spans 6 years; it has a dispersion of 3.15 m/s, and a peak-to-peak variation of 16.9 m/s.

The zero-point drift correction clearly reflects several instrument modifications: a jump at 55872 (6/11/2011) corresponding to a ThAr calibration lamp change; a jump at 56274 (12/12/2012) following the installation of octagonal fibres after the double scrambler; a second ThAr lamp-change related jump at 56690 (01/02/2014); a jump at 56730 (13/03/2014) after the installation of a new calibration unit; a jump at 56941 (10/10/2014) due to a change in the current of the ThAr calibration lamp; and a jump at 57435 corresponding to the installation of a new thermal regulation. These events are indicated in Figure 1. An additional long-term drift is also seen.

The first two events described were also noted by Courcol et al. (2015). We do not, however, find the jump at 56775 which the authors correlated at the time to the recoating of the secondary mirror (including when we regarded a test master constructed using only the template-matching derived RVs of the SP1 super-constants employed by Courcol et al. 2015). We have identified the actual cause of the jump at 56775 as an instability of the wavelength solution on blue spectral orders, whose weight changed after mirror recoating. As this effect is not present in the template-matching procedure, the jump no longer appears.

The full corrected radial velocity sets for G196 and G1617A are given in Appendix A, in Tables A.1, A.2 respectively.

### 3.1. Activity Indicators

In order to study the stellar activity of our targets, we calculated several indicators. Two of the most widely used spectral indicators of chromospheric activity are the  $H\alpha$  index, which measures the flux in the  $H\alpha$  line, and the  $\log(R'_{HK})$  index, which is based on the flux in the Ca II H and K emission lines. We computed the  $H\alpha$  index following the definitions of Boisse et al. (2009). For the Ca lines, we followed Boisse et al. (2010) to obtain the S-index scaled to Mount Wilson values. The  $\log(R'_{HK})$  index was originally defined by Noyes et al. (1984), using a photometric correction based on the B-V index. However, in that work the conversion from the S-index to  $\log(R'_{HK})$  was not calibrated for redder M-dwarfs, and the B-V index is not ideal for M-dwarfs which are too faint in the B band. Therefore, we used the calibra-

tions of Astudillo-Defru et al. (2017a) for M-dwarfs to calculate the  $\log(R'_{HK})$  index employing V-K colours.

Gomes da Silva et al. (2011) carried out a study of activity indices for M-dwarfs, finding that - in addition to the  $H\alpha$  and  $\log(R'_{HK})$  indices - the Na I D1 and D2 lines correlate well with stellar activity in these stars. Therefore, we also calculated the NaI index as defined by the authors.

The CCF bisector is also known to correlate with stellar activity for short rotational periods. We obtained the bisector for our observations from the SOPHIE pipeline.

### 3.2. Stellar Parameters

The stellar parameters are listed in Table 1. Spectral types were obtained from Gaidos et al. (2014); masses from Delfosse et al. (2000); metallicities and luminosities from Gaidos & Mann (2014); temperatures from Mann et al. (2015) when available, and if not from Gaidos & Mann (2014). Magnitudes and colour indices were taken from Zacharias et al. (2012), except for the K magnitude which is from Cutri et al. (2003). For G1617A, a GAIA DR1 parallax is available, while for G196 we take the HIPPARCOS parallax. The mean and standard deviation of  $\log(R'_{HK})$  were calculated from the SOPHIE spectra. We used the  $\log(R'_{HK}) - \log(P_{rot})$  relation from Astudillo-Defru et al. (2017a) to estimate the rotation period from the mean  $\log(R'_{HK})$ , with error bars calculated by propagation.

**Table 1.** Stellar parameters.

Parameter	G1617A	G196
Spectral Type	M1 <sup>a</sup>	M2 <sup>a</sup>
V	8.896 <sup>b</sup>	9.345 <sup>b</sup>
B-V	1.010 <sup>b</sup>	1.519 <sup>b</sup>
V-K	3.943 <sup>c</sup>	3.791 <sup>c</sup>
Mass [ $M_{\odot}$ ]	$0.60 \pm 0.07^d$	$0.60 \pm 0.07^d$
$\Pi$ [mas]	$93.15 \pm 0.23^e$	$83.75 \pm 1.14^f$
$\log(R'_{HK})$	$-4.75 \pm 0.14$	$-4.77 \pm 0.06$
$P_{rot}$ [d]	$28.8 \pm 6.1$	$29.6 \pm 2.8$
$T_{eff}$ [k]	$4156 \pm 73^g$	$3785 \pm 62^h$
$L_{\star}$ [ $L_{\odot}$ ]	$0.1069 \pm 0.0153^g$	$0.0888 \pm 0.0135^g$
Fe/H [dex]	$0.19 \pm 0.08^g$	$0.14 \pm 0.08^g$

**Notes.** <sup>(a)</sup> Source: Gaidos et al. (2014). <sup>(b)</sup> Source: Zacharias et al. (2012). <sup>(c)</sup> Source: V: Zacharias et al. (2012), K: Cutri et al. (2003). <sup>(d)</sup> Source: Delfosse et al. (2000). <sup>(e)</sup> Source: Gaia Collaboration et al. (2016). <sup>(f)</sup> Source: van Leeuwen (2007). <sup>(g)</sup> Source: Gaidos & Mann (2014). <sup>(h)</sup> Source: Mann et al. (2015).

### 3.3. Radial velocity analysis

To analyse the radial velocities, we employed the DACE (Data and Analysis Center for Exoplanets) web platform<sup>2</sup>, which is based at the University of Geneva. This platform allows the generation of generalized Lomb-Scargle periodograms and the calculation of false alarm probabilities (FAP). Signals can be fit by keplerian models (DACE employs the formalism of Delisle et al. 2016 for this purpose), and polynomial drifts and stellar jitter can be added. DACE also provides a MCMC analysis facility, described in Díaz et al. (2014) and Díaz et al. (2016b).

<sup>2</sup> Available at <https://dace.unige.ch>




Fig. 2. Time series of the radial velocities calculated with template-matching for G196 from the SOPHIE+ measurements.

## 4. Results

### 4.1. G196

The RVs calculated with template-matching from the SOPHIE+ observations of G196 were analysed using DACE. The time series and periodogram of the G196 RVs are shown in Figures 2 and 3 respectively. Figure 3 also shows the periodogram of the zero-point correction applied, and of the data prior to this correction. In Figure 4 we show periodograms of the activity indicators  $H\alpha$ ,  $\log(R'_{HK})$ , NaI, and the CCF bisector.

The RV periodogram shows a peak at 75d below 1% FAP, which bootstrap resampling places below 0.05% FAP. There is no corresponding peak for any of the activity indicators, nor does the zero-point drift correction applied show any signal at this period. Additionally, the signal also appears in the uncorrected time series periodogram. We also applied an I1 periodogram, as defined by Hara et al. (2017); this technique searches for a number of signals simultaneously, using compressed sensing techniques. As such, it is much less prone to aliases and other problems arising in the traditional periodogram. The resulting periodogram is shown in Figure 6, where the signal at 74 days can clearly be seen to dominate the data; the FAP of this signal is conservatively estimated (using an analytical formula from Baluev 2008) as  $\log_{10}(FAP) = -2.6072$ . The remaining signals are consistent with the stellar rotation period and half this period, suggesting they originate in stellar activity.

We fit this signal by a keplerian model with DACE. The highest peak in the residuals, at 29d, is only below 50% FAP, while a second peak at 14d at similar FAP corresponds to half this period (Figure 5). Additionally, the highest peak is close to the peaks seen at 28-29d well below 1% FAP in the  $H\alpha$  and  $\log(R'_{HK})$  indices' periodograms. Furthermore, these periods are consistent with the estimated stellar rotation period of  $29.6 \pm 2.8$ d (see Section 3.2). Therefore, we cannot justify treating it as a potential second planet.

We used DACE to carry out an MCMC analysis of the single-planet model for G196, in order to better constrain the parameters. The results are summarized in Table 2. Figure 8 shows the phase-folded data points. The best-fit solution results in a rather highly eccentric orbit. In order to analyse whether contamination from stellar-activity driven RV variations is influencing the results, we tested two approaches: the addition of a keplerian fit to the 29d peak, and a red-noise model. The two-keplerian model did not modify the planetary parameters greatly; in particular, the resulting eccentricity is of  $0.46^{+0.22}_{-0.14}$ , which is indis-






Fig. 3. Periodogram of (top) the radial velocities calculated with template-matching for G196 from the SOPHIE+ measurements, corrected from the zero-point drift; (middle) the time series of the master correction for the zero-point drift applied; (bottom) the uncorrected radial velocities for G196 (prior to the application of the master correction). The horizontal lines correspond to 50%, 10%, and 1% FAP respectively.

tinguishable from the one-keplerian result within error bars. The red noise was modelled using a Gaussian process with a quasi-periodic kernel (see details of the model in, e.g., Astudillo-Defru et al. 2017b). We informed the model of the rotational period of the star by including an informative prior on the corresponding hyper-parameter. The posterior distribution of this parameter is narrower, and we find  $P_{rot} = 28.4 \pm 1.4$  days, where the uncertainties correspond to the  $1-\sigma$  credible interval. Concerning the planet eccentricity, the maximum-a-posteriori estimate is 0.50, in agreement with the model without red noise. The inclusion of correlated noise, on the other hand, seems to allow for lower values of the eccentricity: the 95-% highest-density interval extends between 0.0 and 0.68 (Fig. 7), showing that in the presence of red noise, the eccentricity is effectively unconstrained.

To quantify the significance of the detection, we estimated the posterior odds ratio (POR) between several competing models. If we assume equal prior probability for all models, the POR reduces to the ratio of marginal likelihoods, i.e. the Bayes factor ( $BF$ ). We used the estimation introduced by Perrakis et al. (2014), based on importance sampling, to compute the marginal likelihoods for a model without any Keplerian (k0), a model with a quadratic long-term trend (k0d2), and a model with a single Keplerian (k1). All three models included an additional white noise component, whose amplitude was an additional nuisance parameter. The results are shown in Figure 9 as a function of the sample size used for the estimation. As expected, the Perrakis estimator is biased, but for sample sizes larger than around




**Fig. 4.** Periodograms of activity indicators for G196: from top to bottom, H $\alpha$  index, log( $R'_{HK}$ ) index, NaI index, and CCF bisector. The horizontal lines correspond to 50%, 10%, and 1% FAP respectively. For the NaI index and CCF bisector, no horizontal lines are visible because the entire periodogram is beneath the 50% FAP line.




**Fig. 5.** Periodogram of the residuals of a Keplerian fit with  $P=75$ d to the radial velocities calculated with template-matching for G196 from the SOPHIE+ measurements. The horizontal line corresponds to 50% FAP level.

3000, the bias is negligible. We find logarithmic Bayes factors  $\log(BF_{k1;k0}) = 5.21 \pm 0.06$ , and  $\log(BF_{k1;k0d2}) = 7.91 \pm 0.06$ , for the comparison between  $k1$  and  $k0$ , and  $k1$  and  $k0d2$ , respectively, where the reported values are the empirical means and standard deviations obtained by repeating the calculation for each model 5000 times, and drawing 1000 random pairs. The resulting distributions are very approximately normal. The analysis shows, therefore, that the posterior probability of a model with a single Keplerian is much larger than any of the competing models. This is so in spite of the strong penalisation the Bayes factor gives to more complex models.



**Fig. 6.** 11 periodogram of the SOPHIE RVs for G196. The signal at 74 days is clearly predominant, while the other two peaks are probably related to activity.




**Fig. 7.** Posterior probability of the orbital eccentricity of G196 b under the model including correlated noise.


**Table 2.** Best-fitted solution for the planetary system orbiting G196.

Param.	Units	G196 b
$P$	[d]	$73.94^{+0.33}_{-0.38}$
$K$	[ $\text{m s}^{-1}$ ]	$4.69^{+0.72}_{-0.62}$
$e$		$0.44^{+0.09}_{-0.11}$
$\omega$	[deg]	$339.58^{+12.45}_{-14.52}$
$T_P$	[d]	$55556.39^{+10.57}_{-8.98}$
$T_C$	[d]	$55568.90^{+11.56}_{-9.82}$
$A_r$	[AU]	$0.291^{+0.005}_{-0.005}$
$M \cdot \sin i$	[ $M_{\text{Earth}}$ ]	$19.66^{+2.42}_{-2.30}$
$\gamma_{\text{SOPHIE}}$	[ $\text{m s}^{-1}$ ]	$-37874.84^{+0.31}_{-0.32}$
$\sigma_{\text{JIT}}$	[ $\text{m s}^{-1}$ ]	$3.45^{+0.93}_{-0.91}$
$\sigma_{(O-C)}$	[ $\text{m s}^{-1}$ ]	3.37
log(Post)		$-196.77^{+1.68}_{-2.46}$

**Notes.** For each parameter, the median of the posterior is reported, with error bars computed from the MCMC chains using a 68.3% confidence interval.  $\sigma_{O-C}$  corresponds to the weighted standard deviation of the residuals around this best solutions. log(Post) is the posterior likelihood. All the parameters probed by the MCMC can be found in Appendix B, Table B.1



**Fig. 8.** Phase-folded radial velocities of Gl96 for a one-planet model with  $P=74.7$ d.



**Fig. 9.** *Left:* Marginal likelihood, estimated by the Perrakis method, vs. sample size for three models: a single Keplerian plus white noise (k1; blue), a quadratic drift plus white noise (k0d2; green), and pure white noise (k0; orange). The one-Keplerian model is clearly favoured by the data. The error bars correspond to the 95% confidence interval. *Right:* histogram of 1000 MonteCarlo realisations of the Bayes Factor between model k1 and k0 (orange) and k0d2 (green); the solid curves are normal distributions with the mean and variance equal to those of the Monte-Carlo sample.


One of the goals of the SP3 programme is to detect habitable planets around M-dwarf stars. The habitable zone calculator<sup>3</sup> based on the work of Kopparapu et al. (2013b) places Gl96 b inward of the conservative HZ, though within the optimistic one; however, due to its high eccentricity it would probably move inward of the optimistic HZ at periaapsis (Figure 10). To better quantify the habitability of this eccentric planet, we follow the method employed by Díaz et al. (2016a), who calculated the mean incident flux over an orbit as defined by Williams & Pollard (2002), and compared it with the limits given by Kopparapu et al. (2013a). For Gl96 b, the mean incident flux is  $\langle F \rangle / F_{\oplus} = 1.168$ , placing it between the Recent Venus and Runaway Greenhouse limits.

#### 4.2. Gl617A


A planet at 86.54 d around this star was recently announced by the CARMENES team (Reiners et al. 2017). In this section, we present an independent detection of this planet from our SOPHIE data, and describe a further potential candidate at 500 d.

We analysed the RVs calculated by template-matching from the SOPHIE+ observations of Gl617A with the DACE platform. The time series and periodogram of the Gl617A RVs are shown in Figures 11 and 12 respectively. Figure 12 also shows the periodograms of the zero-point correction applied and of the uncor-

<sup>3</sup> Located at <https://depts.washington.edu/naivpl/content/hz-calculator>



**Fig. 10.** Calculated orbit for Gl96 b with respect to the optimistic and conservative habitable zones, as defined by Kopparapu et al. (2013b).




**Fig. 11.** Time series of the radial velocities calculated with template-matching for Gl617A from the SOPHIE+ measurements.

rected data (i.e. the Gl617A RVs prior to the application of this correction).

In Figure 13 we show periodograms of the activity indicators  $H\alpha$ ,  $\log(R'_{HK})$ , NaI, and the CCF bisector. The  $H\alpha$  index exhibits a periodicity at around 21.8 days, probably related to the rotational period of the star, as well as long-period peaks. The  $\log(R'_{HK})$  and NaI periodograms present only very long-period peaks, in the 500-1000 day range. The CCF bisector shows a small peak at 10.9 days, which is close to half the rotation period and therefore is probably due to stellar activity.

The RV periodogram shows a very strong signal at 86d well below 1% FAP, which bootstrap resampling places below 0.01% FAP, and which is not present in any of the activity indicators, nor in the periodogram of the zero-point correction applied. We employed DACE to fit it with a keplerian model. Figure 14 shows the periodogram of the residuals. There are two signals below 10% FAP: one at 21d and one at around 500d, and one further signal below 50 % FAP at 29d. The 21d signal is probably due to stellar activity, as it coincides with a signal below 1% FAP in the  $H\alpha$  periodogram, and is close to the estimated rotation period (Table 1), although there is no signal at 21d in the other activity indicators. The 29d signal is removed by the addition of a quadratic drift, suggesting it may be an artifact of the window function (although the period may also point to moon contam-



**Fig. 12.** Periodogram of: (top) the radial velocities calculated with template-matching for Gl617A from the SOPHIE+ measurements, corrected from the zero-point drift; (middle) the master correction for the zero-point drift applied; (bottom) the uncorrected radial velocities for Gl617A (prior to the application of the master correction). The horizontal lines correspond to 50%, 10%, and 1% FAP respectively.


ination; as this star is observed with simultaneous wavelength calibration, we do not have a measure of the sky).

The long-period signal is intriguing; the addition of a linear or quadratic drift does not affect it, whereas a test keplerian model removes it completely and results in a low-eccentricity keplerian fit. While the H $\alpha$ , log(R'<sub>HK</sub>), and NaI indices' periodograms show long-period peaks, these are not well fitted by the model obtained from the RVs. More concerning is the periodogram of the master correction, which shows a peak at 500 days; nevertheless, we note that the long-period signal is also present in the uncorrected RVs (see Fig. 12). However, we stress that this signal is of only moderate significance. We therefore present it as a tentative detection to be confirmed, not a definite planet.


As for Gl96, we also applied an 11 periodogram, which is shown in Figure 15; the signal at 86 days dominates the data, with  $\log_{10}(FAP) = -17.1869$ . The remaining signals at 21d and  $\sim 500$ d are consistent with the discussion above.

In order to explore the parameter space more thoroughly, we employed the MCMC sampler implemented in DACE. We tested two models: a single keplerian at 86d, and a two-keplerian model with initial periods at 86d and 500d, both including a quadratic drift. The resulting best-fit solutions are summarised in Table 3.

The planet at 86d is consistent in both models, with practically all parameters indistinguishable within error bars between the two best-fitted solutions. The two-keplerian model actually provides a better fit to the data, with a lower  $\sigma_{(O-C)}$  and BIC. Figures 16 and 17 show the phase-folded RVs for Gl617A b



**Fig. 13.** Periodograms of activity indicators for Gl617A: from top to bottom, H $\alpha$  index, log(R'<sub>HK</sub>) index, NaI index, and CCF bisector. The horizontal lines correspond to 50%, 10%, and 1% FAP respectively.




**Fig. 14.** Periodogram of the residuals of a keplerian fit with P=86d to the radial velocities calculated with template-matching for Gl617A from the SOPHIE+ measurements. The horizontal lines correspond to 50% and 10% FAP respectively.


and Gl617A c respectively, using the parameters derived from the two-keplerian model. In the one-keplerian model, the drift is significant; removing it slightly decreases the amplitude and increases the eccentricity of the keplerian fit. In the two-keplerian model, however, the drift is only marginally significant. This is coherent with the fact that in the one-keplerian model, the drift may attempt to absorb the long-period signal that is fitted by a keplerian in the second model.

We compared the parameters obtained by our analysis with those recently presented by Reiners et al. (2017). The orbital period and semi-major axis are compatible within error bars; however, we find a somewhat larger mass and distinctly larger eccentricity from our data. The mass reported by Reiners et al. (2017) of  $M \cdot \sin i = 24.7^{+2.4}_{-1.8} M_{\text{Earth}}$  is compatible with our cal-






**Fig. 15.** I1 periodogram of the SOPHIE RVs for Gl617A. The signal at 86 days is clearly predominant, while the one at 21 days is probably related to activity. The  $\sim 500$ d signal is discussed in the text.




**Fig. 16.** Phase-folded radial velocities of Gl617A for a  $P=86$ d planet, using the parameters derived from the 2-keplerian model.




**Fig. 17.** Phase-folded radial velocities of Gl617A for a  $P=497$ d planet, using the parameters derived from the 2-keplerian model.

culated mass at  $2\sigma$ . This is true for both the one-keplerian and the two-keplerian models, with the eccentricity higher for the one-keplerian model (see Table 3).

As for Gl96, we employed the habitable zone calculator to estimate the location of the HZ for Gl617A. The orbits calculated by the MCMC analysis place Gl617A b at 0.32AU, closer to the star than the conservative inner limit, but within the optimistic HZ (Figure 18). This differs from the results of Reiners et al. (2017). As Gl617A b is moderately eccentric, we also calculate the mean incident flux (as done for Gl96 b) in order to better quantify its habitability. The mean incident flux is  $\langle F \rangle / F_{\oplus} = 1.053$ , placing it between the Recent Venus and Runaway Greenhouse limits as defined by Kopparapu et al. (2013a).



**Fig. 18.** Calculated orbit for Gl617A b with respect to the optimistic and conservative habitable zones, as defined by Kopparapu et al. (2013b).



**Fig. 19.** Periodograms of: (top) the combined SOPHIE, CARMENES, and KECK radial velocities of Gl617A; (bottom) the residuals of a keplerian fit with  $P=86$ d to the data. The horizontal lines correspond to 50%, 10%, and (top periodogram only) 1% FAP respectively.

#### 4.2.1. Combination with CARMENES and KECK data

We combined our observations of Gl617A with the CARMENES data presented by Reiners et al. (2017) and the KECK data of Butler et al. (2017). Figure 19 shows the resulting periodogram, with a strong signal at 86d, to which we fit a keplerian model using DACE.; and the periodogram of the residuals of this fit. The only strong signal is at 21d, and presumably corresponds to the stellar activity as discussed previously. The 500d signal that is present in the SOPHIE data is not in evidence in the combined observations, suggesting it may be spurious.

The posterior distribution of the model parameters was sampled using an MCMC algorithm implemented in DACE. A single-keplerian model with a quadratic drift was used. The resulting best-fit solution is summarised in Table 4, and the phase-folded combined data is shown in Figure 20.

We compared the parameters resulting from the analysis of the combined data with those we obtained from SOPHIE data

**Table 3.** Best-fitted solutions for the planetary system orbiting Gl617A - 1-keplerian and 2-keplerian models plus drift.

Param.	Units	Gl617A b <sup>1</sup>	Gl617A b <sup>2</sup>	Gl617A c
$P$	[d]	86.93 <sup>+0.18</sup> <sub>-0.19</sub>	86.72 <sup>+0.20</sup> <sub>-0.18</sub>	496.90 <sup>+35.45</sup> <sub>-21.82</sub>
$K$	[m s <sup>-1</sup> ]	6.56 <sup>+0.41</sup> <sub>-0.39</sub>	6.57 <sup>+0.36</sup> <sub>-0.38</sub>	3.16 <sup>+0.43</sup> <sub>-0.42</sub>
$e$		0.32 <sup>+0.08</sup> <sub>-0.09</sub>	0.23 <sup>+0.07</sup> <sub>-0.08</sub>	0.15 <sup>+0.15</sup> <sub>-0.10</sub>
$\omega$	[deg]	102.20 <sup>+10.04</sup> <sub>-11.48</sub>	97.25 <sup>+13.55</sup> <sub>-13.46</sub>	-311.97 <sup>+980.42</sup> <sub>-63.26</sub>
$T_P$	[d]	55466.60 <sup>+4.75</sup> <sub>-4.32</sub>	55470.17 <sup>+4.97</sup> <sub>-4.85</sub>	55214.75 <sup>+146.26</sup> <sub>-182.87</sub>
$T_C$	[d]	55465.26 <sup>+4.47</sup> <sub>-4.08</sub>	55469.18 <sup>+4.30</sup> <sub>-4.41</sub>	55275.54 <sup>+109.85</sup> <sub>-121.34</sub>
$Ar$	[AU]	0.324 <sup>+0.006</sup> <sub>-0.006</sub>	0.323 <sup>+0.006</sup> <sub>-0.005</sub>	1.036 <sup>+0.051</sup> <sub>-0.036</sub>
$M.\sin i$	[M <sub>Earth</sub> ]	30.56 <sup>+2.10</sup> <sub>-2.12</sub>	31.29 <sup>+2.20</sup> <sub>-2.15</sub>	27.26 <sup>+3.84</sup> <sub>-3.72</sub>
$\gamma_{SOPHIE}$	[m s <sup>-1</sup> ]	-18737.35 <sup>+5.38</sup> <sub>-5.74</sub>	-18715.26 <sup>+5.58</sup> <sub>-5.63</sub>	
$lin$	[m s <sup>-1</sup> yr <sup>-1</sup> ]	7.71 <sup>+2.42</sup> <sub>-2.3</sub>	-3.13 <sup>+2.65</sup> <sub>-2.34</sub>	
$quad$	[m s <sup>-1</sup> yr <sup>-2</sup> ]	-0.86 <sup>+0.23</sup> <sub>-0.25</sub>	0.30 <sup>+0.27</sup> <sub>-0.26</sub>	
$\sigma_{JIT}$	[m s <sup>-1</sup> ]	4.86 <sup>+0.56</sup> <sub>-0.62</sub>	3.71 <sup>+0.49</sup> <sub>-0.46</sub>	
$\sigma_{(O-C)}$	[m s <sup>-1</sup> ]	4.55	3.90	
log (Post)		-466.65 <sup>+1.82</sup> <sub>-2.56</sub>	-440.08 <sup>+2.28</sup> <sub>-3.07</sub>	
BIC		105.93	118.52	

**Notes.** <sup>(1)</sup> Parameters obtained from the 1-keplerian model for Gl617A b. <sup>(2)</sup> Parameters obtained from the 2-keplerian model for Gl617A b. For each parameter, the median of the posterior is reported, with error bars computed from the MCMC chains using a 68.3% confidence interval.  $\sigma_{O-C}$  corresponds to the weighted standard deviation of the residuals around this best solution. log (Post) is the posterior likelihood. All the parameters probed by the MCMC can be found in Appendix B, Tables B.3 (1-keplerian model) and B.5 (2-keplerian model).

alone, and with those presented by Reiners et al. (2017). In all three cases, the orbital periods are compatible at the  $1\sigma$  level (though slightly larger for the SOPHIE and combined data).  $M.\sin i$  is also compatible at the  $1\sigma$  level between SOPHIE data alone and the combined data, but Reiners et al. (2017) present a slightly smaller value of  $M.\sin i = 24.7^{+2.4}_{-1.8} M_{\text{Earth}}$ , only compatible with the others at the  $2\sigma$  level. The eccentricity is also different; the one derived from the SOPHIE data alone is larger, while those of the combined data and of Reiners et al. (2017) are compatible at  $1\sigma$ . The amplitude of the signal, correspondingly, differs, being largest for the SOPHIE data alone and smallest in Reiners et al. (2017)'s analysis; this is coherent with the fact that for fixed  $e$ ,  $K$  grows with  $M.\sin i$ , and for fixed  $M.\sin i$   $K$  grows with  $e$  (see e.g. Seager 2010).

Recently Feng et al. (2018) published a analysis of this system in the Research Note of the AAS, also employing the CARMENES, HIRES, and SOPHIE data. Their use of our SOPHIE data, however, is biased by the fact that they do not take into account the CTI effect and zero-point drift described in Section 3 that we correct for in this work.

## 5. Photometry

Stellar activity can be reflected not only in the radial velocity but in the photometric observations of a star, where we may also hope to find signals linked to the rotation period. In order to analyse whether this is the case for our targets, we obtained the Hipparcos photometry for both from ESA (1997).


For Gl96, we retrieved 121 measurements over a time span of three years. Fig. 21 shows the photometric data points and the corresponding periodogram. No signals below 50% FAP can be found in the periodogram. There is a small peak at 29d, the period found in the RV, H $\alpha$ , and log( $R'_{\text{HK}}$ ) periodograms that is presumed related to activity, but it is not particularly relevant in the

**Table 4.** Best-fitted solution for the planetary system orbiting Gl617A, from SOPHIE, CARMENES, and KECK combined data.


Param.	Units	Gl617A b
$P$	[d]	86.78 <sup>+0.16</sup> <sub>-0.15</sub>
$K$	[m s <sup>-1</sup> ]	5.83 <sup>+0.22</sup> <sub>-0.24</sub>
$e$		0.07 <sup>+0.04</sup> <sub>-0.04</sub>
$\omega$	[deg]	97.22 <sup>+30.39</sup> <sub>-41.83</sub>
$T_P$	[d]	55468.26 <sup>+8.52</sup> <sub>-10.67</sub>
$T_C$	[d]	55466.91 <sup>+4.12</sup> <sub>-4.11</sub>
$Ar$	[AU]	0.323 <sup>+0.005</sup> <sub>-0.006</sub>
$M.\sin i$	[M <sub>Earth</sub> ]	28.55 <sup>+1.49</sup> <sub>-1.45</sub>
$\gamma_{CARMENES}$	[m s <sup>-1</sup> ]	0.40 <sup>+1.64</sup> <sub>-1.60</sub>
$\gamma_{KECK-PUB_1}$	[m s <sup>-1</sup> ]	-1.76 <sup>+2.63</sup> <sub>-2.68</sub>
$\gamma_{SOPHIE}$	[m s <sup>-1</sup> ]	-18721.59 <sup>+1.54</sup> <sub>-1.50</sub>
$\sigma_{JIT}$	[m s <sup>-1</sup> ]	15.89 <sup>+4.14</sup> <sub>-1.82</sub>
$\sigma_{(O-C)}$	[m s <sup>-1</sup> ]	3.87
log (Post)		-939.95 <sup>+2.11</sup> <sub>-2.76</sub>
BIC		1668.19

**Notes.** Same notes as Table 3. All the parameters probed by the MCMC can be found in Appendix B, Table B.7

periodogram of photometric measurements which is dominated by a forest of peaks at around 2d. This set of peaks is probably linked to the sampling - as can be seen in the upper panel of Fig. 21, this star appears to have been observed in groups of several measurements within two days, with the groups set weeks or months apart.



**Fig. 20.** Phase-folded radial velocities of G1617A for a  $P=86d$  planet, using the combined SOPHIE, CARMENES, and KECK data.




**Fig. 21.** Hipparcos photometry for G196 (top) and its corresponding periodogram (bottom).

For G1617A, we retrieved 103 measurements over a time span of two and a half years. The photometric data points and the corresponding periodogram are shown in Fig. 22. No signals below 50% FAP can be found in the periodogram. Nevertheless, it is worth remarking that the two highest peaks are at 10.4d and 20d; these values are very close to the 21d period seen in the RV and  $H\alpha$  periodograms and half this period.

For both stars, we note that the HIPPARCOS photometry has a relatively high RMS (2.62% for G196, 1.97% for G1617A). Activity signals at the level attained by our RV measurements would fall well below this, so any photometric activity tracers may be absorbed in the uncertainty of the data.

## 6. Discussion and Conclusions

We have presented the detection of a new Neptune-like exoplanet orbiting the M dwarf G196 and the independent detection of a second Neptune-like exoplanet orbiting the M dwarf G1617A for which we refine the planetary parameters, using the SOPHIE+ spectrograph on a 1.93m telescope. The planets have




**Fig. 22.** Hipparcos photometry for G1617A (top) and its corresponding periodogram (bottom).

minimum masses of 29 and 31 Earth masses and orbital periods of 74 and 87 days respectively, and are located close to the inner limit of the HZ. For G196 we find no evidence for further planetary companions. G1617A shows an intriguing signal of moderate significance at  $\sim 500d$  in the periodogram that is best fit by a keplerian model. For G1617A, we also analysed the combination of our data with that from CARMENES (Reiners et al. 2017) and KECK (Butler et al. 2017). The 87d signal is clear in the combined data, though the resulting planetary parameters differ slightly from those obtained by SOPHIE data alone. The 500d signal, however, is no longer significant, which would suggest it is spurious. We may also suspect the influence of a magnetic cycle here; complementary observations in polarimetry with SPIRou should help to resolve the question.

As mentioned in Section 1, some 146 exoplanets around M-dwarf stars are presently known, of which 75 were detected by radial velocities and 33 by transits. The two planets presented here fall in the mid-to-long period and mid-to-high mass ranges of this sample, as shown in Figure 23 (median period = 13.5 d, median mass =  $14.3 M_{\text{Earth}}$ ). G196 b is among the most eccentric known planets around M-dwarf stars (mean  $e = 0.12$ ), surpassed only by Wolf 1061 d (Wright et al. 2016) and GJ 317 c (Johnson et al. 2007), with eccentricities of 0.55 and 0.81 respectively.

Both host stars are metal-rich, as noted in Table 1. This is consistent with a trend found for M-dwarfs hosting planets to be preferentially metal-rich (e.g. Courcol et al. 2016, Hobson et al. 2017). Additionally, the masses determined for the planets are compatible with the upper mass boundary determined for Neptune-like planets around M-dwarfs by Courcol et al. (2016), which corresponds to around 35 Earth masses for the metallicities of these stars.

Low-mass planets orbiting M-dwarfs are mainly found in multi-planet systems. Follow-up observations with SPIRou in spectropolarimetry will permit us to disentangle the stellar activity and planetary signals, refine the mass and identify possible additional rocky planets.



**Fig. 23.** Orbital period versus minimum mass for planets around M-dwarfs detected by radial velocities or transits. Each point represents a planet, with Gl96 b indicated by a pentagon and Gl617A b by a diamond. The colours indicate the orbital eccentricity.

**Acknowledgements.** We warmly thank the OHP staff for their support on the observations. XD, XB, IB and TF received funding from the French Programme National de Physique Stellaire (PNPS) and the Programme National de Planétologie (PNP) of CNRS (INSU). NCS and OD were supported by Fundação para a Ciência e a Tecnologia (FCT, Portugal) through national funds and by FEDER through COMPETE2020 in the context of the projects and grants reference UID/FIS/04434/2013 & POCI-01-0145-FEDER-007672, PTDC/FIS-AST/1526/2014 & POCI-01-0145-FEDER-016886, and IF/00169/2012/CP0150/CT0002. This publication makes use of The Data & Analysis Center for Exoplanets (DACE), which is a facility based at the University of Geneva (CH) dedicated to extrasolar planets data visualisation, exchange and analysis. DACE is a platform of the Swiss National Centre of Competence in Research (NCCR) PlanetS, federating the Swiss expertise in Exoplanet research. The DACE platform is available at <https://dace.unige.ch>. We thank the anonymous referees for their comments which helped to improve the manuscript.

## References

Affer, L., Micela, G., Damasso, M., et al. 2016, *A&A*, 593, A117  
 Anglada-Escudé, G. & Butler, R. P. 2012, *ApJS*, 200, 15  
 Artigau, É., Kouach, D., Donati, J.-F., et al. 2014, in *Proc. SPIE*, Vol. 9147, Ground-based and Airborne Instrumentation for Astronomy V, 914715  
 Astudillo-Defru, N., Bonfils, X., Delfosse, X., et al. 2015, *A&A*, 575, A119  
 Astudillo-Defru, N., Delfosse, X., Bonfils, X., et al. 2017a, *A&A*, 600, A13  
 Astudillo-Defru, N., Díaz, R. F., Bonfils, X., et al. 2017b, *A&A*, 605, L11  
 Astudillo-Defru, N., Forveille, T., Bonfils, X., et al. 2017c, *A&A*, 602, A88  
 Baluev, R. V. 2008, *MNRAS*, 385, 1279  
 Boisse, I., Eggenberger, A., Santos, N. C., et al. 2010, *A&A*, 523, A88  
 Boisse, I., Moutou, C., Vidal-Madjar, A., et al. 2009, *A&A*, 495, 959  
 Bonfils, X., Almenara, J. M., Jocou, L., et al. 2015, in *Proc. SPIE*, Vol. 9605, Techniques and Instrumentation for Detection of Exoplanets VII, 96051L  
 Bonfils, X., Delfosse, X., Udry, S., et al. 2013, *A&A*, 549, A109  
 Bouchy, F., Díaz, R. F., Hébrard, G., et al. 2013, *A&A*, 549, A49  
 Bouchy, F., Doyon, R., Artigau, É., et al. 2017, *The Messenger*, 169, 21  
 Bouchy, F., Hébrard, G., Udry, S., et al. 2009a, *A&A*, 505, 853  
 Bouchy, F., Isambert, J., Lovis, C., et al. 2009b, in *EAS Publications Series*, Vol. 37, *EAS Publications Series*, ed. P. Kern, 247–253  
 Butler, R. P., Vogt, S. S., Laughlin, G., et al. 2017, *AJ*, 153, 208  
 Cloutier, R., Astudillo-Defru, N., Doyon, R., et al. 2017, *A&A*, 608, A35  
 Courcol, B., Bouchy, F., & Deleuil, M. 2016, *MNRAS*, 461, 1841  
 Courcol, B., Bouchy, F., Pepe, F., et al. 2015, *A&A*, 581, A38  
 Cutri, R. M., Skrutskie, M. F., van Dyk, S., et al. 2003, *VizieR Online Data Catalog*, 2246  
 Delfosse, X., Bonfils, X., Forveille, T., et al. 2013, *A&A*, 553, A8  
 Delfosse, X., Forveille, T., Ségransan, D., et al. 2000, *A&A*, 364, 217  
 Delisle, J.-B., Ségransan, D., Buchschacher, N., & Alesina, F. 2016, *A&A*, 590, A134  
 Díaz, R. F., Almenara, J. M., Santerne, A., et al. 2014, *MNRAS*, 441, 983

Díaz, R. F., Rey, J., Demangeon, O., et al. 2016a, *A&A*, 591, A146  
 Díaz, R. F., Ségransan, D., Udry, S., et al. 2016b, *A&A*, 585, A134  
 Dittmann, J. A., Irwin, J. M., Charbonneau, D., et al. 2017, *Nature*, 544, 333  
 Dressing, C. D. & Charbonneau, D. 2015, *ApJ*, 807, 45  
 ESA, ed. 1997, *ESA Special Publication*, Vol. 1200, *The HIPPARCOS and TYCHO catalogues. Astrometric and photometric star catalogues derived from the ESA HIPPARCOS Space Astrometry Mission*  
 Feng, F., Jones, H. R. A., & Tuomi, M. 2018, *Research Notes of the American Astronomical Society*, 2, 23  
 Foreman-Mackey, D., Montet, B. T., Hogg, D. W., et al. 2015, *ApJ*, 806, 215  
 Gaia Collaboration, Brown, A. G. A., Vallenari, A., et al. 2016, *A&A*, 595, A2  
 Gaidos, E. & Mann, A. W. 2014, *ApJ*, 791, 54  
 Gaidos, E., Mann, A. W., Lépine, S., et al. 2014, *MNRAS*, 443, 2561  
 Gillon, M., Triau, A. H. M. J., Demory, B.-O., et al. 2017, *Nature*, 542, 456  
 Gomes da Silva, J., Santos, N. C., Bonfils, X., et al. 2011, *A&A*, 534, A30  
 Hara, N. C., Boué, G., Laskar, J., & Correia, A. C. M. 2017, *MNRAS*, 464, 1220  
 Hobson, M. J., Jofré, E., García, L., Petrucci, R., & Gómez, M. 2017, *ArXiv e-prints* [arXiv:1711.04878]  
 Johnson, J. A., Butler, R. P., Marcy, G. W., et al. 2007, *ApJ*, 670, 833  
 Kopparapu, R. K., Ramirez, R., Kasting, J. F., et al. 2013a, *ApJ*, 770, 82  
 Kopparapu, R. K., Ramirez, R., Kasting, J. F., et al. 2013b, *ApJ*, 765, 131  
 Mann, A. W., Feiden, G. A., Gaidos, E., Boyajian, T., & von Braun, K. 2015, *ApJ*, 804, 64  
 Noyes, R. W., Hartmann, L. W., Baliunas, S. L., Duncan, D. K., & Vaughan, A. H. 1984, *ApJ*, 279, 763  
 Pepe, F., Mayor, M., Galland, F., et al. 2002, *A&A*, 388, 632  
 Perrakis, K., Ntzoufras, I., & Tsonas, E. G. 2014, *Computational Statistics & Data Analysis*, 77, 54  
 Perruchot, S., Bouchy, F., Chazelas, B., et al. 2011, in *Proc. SPIE*, Vol. 8151, Techniques and Instrumentation for Detection of Exoplanets V, 815115  
 Perruchot, S., Kohler, D., Bouchy, F., et al. 2008, in *Proc. SPIE*, Vol. 7014, Ground-based and Airborne Instrumentation for Astronomy II, 70140J  
 Queloz, D. 1995, in *IAU Symposium*, Vol. 167, *New Developments in Array Technology and Applications*, ed. A. G. D. Philip, K. Janes, & A. R. Uppgren, 221  
 Quirrenbach, A. & Carmones Consortium. 2016, in *COSPAR Meeting*, Vol. 41, 41st COSPAR Scientific Assembly  
 Reid, I. N. & Hawley, S. L. 2005, *New light on dark stars : red dwarfs, low-mass stars, brown dwarfs*  
 Reiners, A., Ribas, I., Zechmeister, M., et al. 2017, *ArXiv e-prints* [arXiv:1712.05797]  
 Ricker, G. R. 2016, *AGU Fall Meeting Abstracts*  
 Seager, S. 2010, *Exoplanets*  
 "The Exoplanet Encyclopaedia". 1995-2017, *The Exoplanet Encyclopaedia*, <http://exoplanet.eu/>  
 van Leeuwen, F. 2007, *A&A*, 474, 653  
 Williams, D. M. & Pollard, D. 2002, *International Journal of Astrobiology*, 1, 61  
 Wright, D. J., Wittenmyer, R. A., Tinney, C. G., Bentley, J. S., & Zhao, J. 2016, *ApJ*, 817, L20  
 Zacharias, N., Finch, C. T., Girard, T. M., et al. 2012, *VizieR Online Data Catalog*, 1322

## **Appendix A: Radial Velocities**

In this Appendix, we present for each star the full set of radial velocities derived by template-matching, with the short-term instrumental drift and long-term zero-point corrections applied.

## **Appendix B: MCMC analysis - Full Probed Parameters**

### *Appendix B.1: Gl96*

We present the full set of parameters probed by the DACE MCMC analysis for Gl96, and the derived physical parameters.

### *Appendix B.2: Gl617A*

We present the full set of parameters probed by the DACE MCMC analysis for Gl617A, and the derived physical parameters.

**Table B.1.** Parameters probed by the MCMC used to fit the RV measurements of Gl96.

Param.	Units	Max(Like)	Med	Mod	Std	CI(15.85)	CI(84.15)	CI(2.275)	CI(97.725)	Prior
<b>Likelihood</b>										
log (Post)		-193.285494	-196.771970	-195.921884	1.870757	-199.232726	-195.096198	-202.542797	-194.075043	
log (Like)		-191.922894	-195.623787	-195.175969	1.973270	-198.178506	-193.846439	-201.754278	-192.754170	
log (Prior)		-1.362600	-1.078578	-1.136538	0.427678	-1.579763	-0.619581	-2.223648	-0.226630	
$M_{\star}$	[ $M_{\odot}$ ]	0.621188	0.600424	0.596743	0.026436	0.570211	0.630397	0.540296	0.660160	$\mathcal{U}$
$\sigma_{JIT}$	[ $m s^{-1}$ ]	3.00	3.45	3.21	0.93	2.54	2.30	1.78	0.77	$\mathcal{U}$
$\gamma_{SOPHIE}$	[ $m s^{-1}$ ]	-37874.88	-37874.84	-37874.90	0.28	-37875.15	-37874.53	-37875.50	-37874.22	$\mathcal{U}$
$P$	[d]	73.939773	73.937730	73.993314	0.325983	73.553342	74.264366	73.092750	74.719936	$\mathcal{U}$
$K$	[ $m s^{-1}$ ]	5.20	4.69	4.42	0.60	4.07	5.41	3.50	6.26	$\mathcal{U}$
$e$		0.495245	0.439034	0.430347	0.088190	0.332797	0.529659	0.200769	0.609855	$\mathcal{U}$
$\omega$	[deg]	341.181282	339.576769	338.551057	11.790040	325.056172	352.022293	307.050528	358.379033	$\mathcal{U}$
$\lambda_0$	[deg]	64.471037	63.383026	67.891413	45.293155	9.598258	108.428683	-55.974137	170.848142	$\mathcal{U}$

**Notes.** The maximum likelihood solution (Max(Like)), the median (Med), mode (Mod) and standard deviation (Std) of the posterior distribution for each parameter is shown, as well as the 68.3% (CI(15.85),CI(84.15)) and 95.45% (CI(2.275),CI(97.725)) confidence intervals. The prior for each parameter can be of type:  $\mathcal{U}$ : uniform,  $\mathcal{N}$ : normal,  $\mathcal{SN}$ :split normal,  $\mathcal{TN}$ : truncated normal.

**Table B.2.** Physical parameters derived from the MCMC chains used fit the RV measurements of Gl96.

Param.	Units	Max(Like)	Med	Mod	Std	CI(15.85)	CI(84.15)	CI(2.275)	CI(97.725)	Prior
<b>Likelihood</b>										
$M_{\star}$	[ $M_{\odot}$ ]	0.621188	0.600424	0.596743	0.026436	0.570211	0.630397	0.540296	0.660160	$\mathcal{U}$
$P$	[d]	73.939773	73.937730	73.993314	0.325983	73.553342	74.264366	73.092750	74.719936	$\mathcal{U}$
$K$	[ $m s^{-1}$ ]	5.20	4.69	4.42	0.60	4.07	5.41	3.50	6.26	$\mathcal{U}$
$e$		0.495245	0.439034	0.430347	0.088190	0.332797	0.529659	0.200769	0.609855	$\mathcal{U}$
$\omega$	[deg]	341.181282	339.576769	338.551057	11.790040	325.056172	352.022293	307.050528	358.379033	$\mathcal{U}$
$T_P$	[d]	55556.833035	55556.392386	55553.872826	9.006008	55547.408500	55566.959102	55534.517800	55579.539962	
$T_C$	[d]	55567.178726	55568.899883	55564.573169	9.663639	55559.078770	55580.463887	55546.269103	55593.191399	
$Ar$	[AU]	0.294178	0.290799	0.290868	0.004383	0.285732	0.295728	0.280513	0.300435	
$M.\sin i$	[ $M_{Jup}$ ]	0.067975	0.061855	0.058986	0.006479	0.054626	0.069477	0.047715	0.077017	
$M.\sin i$	[ $M_{Earth}$ ]	21.602453	19.657445	18.745664	2.058966	17.359998	22.079752	15.163668	24.475879	

**Notes.** The maximum likelihood solution (Max(Like)), the median (Med), mode (Mod) and standard deviation (Std) for the posterior distribution of each parameter is shown, as well as the 68.3% (CI(15.85),CI(84.15)) and 95.45% (CI(2.275),CI(97.725)) confidence intervals. The prior for each parameter can be of type:  $\mathcal{U}$ : uniform,  $\mathcal{N}$ : normal,  $\mathcal{SN}$ :split normal,  $\mathcal{TN}$ : truncated normal.

**Table B.3.** Parameters probed by the MCMC used to fit the RV measurements of Gl617A - 1-keplerian model plus quadratic drift.

Param.	Units	Max(Like)	Med	Mod	Std	CI(15.85)	CI(84.15)	CI(2.275)	CI(97.725)	Prior
<b>Likelihood</b>										
log (Post)		-462.619011	-466.651637	-466.520190	1.964561	-469.209353	-464.831362	-472.561763	-463.627502	
log (Like)		-461.941145	-466.049364	-465.293525	2.015104	-468.693270	-464.197613	-472.107721	-462.923059	
log (Prior)		-0.677866	-0.555253	-0.500711	0.254783	-0.874032	-0.292873	-1.220718	-0.097352	
$M_{\star}$	[ $M_{\odot}$ ]	0.611414	0.600358	0.595467	0.026063	0.570942	0.630052	0.540715	0.658600	$\mathcal{U}$
$\sigma_{JIT}$	[ $m s^{-1}$ ]	4.44	4.86	4.73	0.52	4.30	4.24	3.78	3.55	$\mathcal{U}$
$\gamma_{SOPHIE}$	[ $m s^{-1}$ ]	-18739.60	-18737.35	-18736.94	4.89	-18743.09	-18731.97	-18748.80	-18726.19	$\mathcal{U}$
$lin$	[ $m s^{-1} yr^{-1}$ ]	8.53	7.71	7.09	2.09	5.41	10.13	3.01	12.66	$\mathcal{U}$
$quad$	[ $m s^{-1} yr^{-2}$ ]	-0.93	-0.86	-0.84	0.21	-1.11	-0.63	-1.36	-0.38	$\mathcal{U}$
$P$	[d]	86.916721	86.925044	86.904924	0.164238	86.730726	87.102317	86.523585	87.278783	$\mathcal{U}$
$K$	[ $m s^{-1}$ ]	6.70	6.56	6.47	0.35	6.16	6.97	5.78	7.37	$\mathcal{U}$
$e$		0.349308	0.316142	0.319096	0.073835	0.229602	0.396643	0.132376	0.468753	$\mathcal{U}$
$\omega$	[deg]	101.004334	102.200583	103.915448	9.807382	90.720509	112.245164	75.691327	123.345529	$\mathcal{U}$
$\lambda_0$	[deg]	238.987027	239.918335	240.420912	16.482165	220.387042	257.567720	199.451375	275.038557	$\mathcal{U}$

**Notes.** The maximum likelihood solution (Max(Like)), the median (Med), mode (Mod) and standard deviation (Std) of the posterior distribution for each parameter is shown, as well as the 68.3% (CI(15.85),CI(84.15)) and 95.45% (CI(2.275),CI(97.725)) confidence intervals. The prior for each parameter can be of type:  $\mathcal{U}$ : uniform,  $\mathcal{N}$ : normal,  $\mathcal{SN}$ :split normal,  $\mathcal{TN}$ : truncated normal.

**Table B.4.** Physical parameters derived from the MCMC chains used fit the RV measurements of Gl617A - 1-keplerian model plus quadratic drift.

Param.	Units	Max(Like)	Med	Mod	Std	CI(15.85)	CI(84.15)	CI(2.275)	CI(97.725)	Prior
<b>Likelihood</b>										
$M_\star$	[ $M_\odot$ ]	0.611414	0.600358	0.595467	0.026063	0.570942	0.630052	0.540715	0.658600	$\mathcal{U}$
$P$	[d]	86.916721	86.925044	86.904924	0.164238	86.730726	87.102317	86.523585	87.278783	$\mathcal{U}$
$K$	[ $\text{m s}^{-1}$ ]	6.70	6.56	6.47	0.35	6.16	6.97	5.78	7.37	$\mathcal{U}$
$e$		0.349308	0.316142	0.319096	0.073835	0.229602	0.396643	0.132376	0.468753	$\mathcal{U}$
$\omega$	[deg]	101.004334	102.200583	103.915448	9.807382	90.720509	112.245164	75.691327	123.345529	$\mathcal{U}$
$T_P$	[d]	55466.686102	55466.601086	55465.672814	4.040069	55462.278510	55471.350467	55457.295504	55476.587804	
$T_C$	[d]	55465.481743	55465.258184	55464.278744	3.805920	55461.181843	55469.727111	55456.842749	55474.584025	
$A_r$	[AU]	0.325940	0.323965	0.324425	0.004713	0.318459	0.329242	0.312843	0.334318	
$M.\sin i$	[ $M_{\text{Jup}}$ ]	0.098576	0.096167	0.094936	0.005835	0.089490	0.102775	0.083150	0.110012	
$M.\sin i$	[ $M_{\text{Earth}}$ ]	31.327394	30.561882	30.170671	1.854279	28.440031	32.661895	26.425117	34.961725	

**Notes.** The maximum likelihood solution (Max(Like)), the median (Med), mode (Mod) and standard deviation (Std) for the posterior distribution of each parameter is shown, as well as the 68.3% (CI(15.85),CI(84.15)) and 95.45% (CI(2.275),CI(97.725)) confidence intervals. The prior for each parameter can be of type:  $\mathcal{U}$ : uniform,  $\mathcal{N}$ : normal,  $\mathcal{SN}$ : split normal,  $\mathcal{TN}$ : truncated normal.

**Table B.5.** Parameters probed by the MCMC used to fit the RV measurements of Gl617A - 2-keplerian model plus quadratic drift.

Param.	Units	Max(Like)	Med	Mod	Std	CI(15.85)	CI(84.15)	CI(2.275)	CI(97.725)	Prior
<b>Likelihood</b>										
log(Post)		-434.164980	-440.075963	-440.047078	2.381702	-443.144186	-437.799233	-446.780069	-436.227907	
log(Like)		-433.765721	-439.525314	-439.289290	2.390269	-442.596771	-437.243499	-446.189031	-435.601729	
log(Prior)		-0.399259	-0.486225	-0.368662	0.298923	-0.882288	-0.243982	-1.520647	-0.083642	
$M_\star$	[ $M_\odot$ ]	0.581188	0.598699	0.589123	0.026717	0.569082	0.629830	0.538640	0.660030	$\mathcal{U}$
$\sigma_{JIT}$	[ $\text{m s}^{-1}$ ]	3.47	3.71	3.83	0.42	3.25	3.22	2.86	2.68	$\mathcal{U}$
$\gamma_{SOPHIE}$	[ $\text{m s}^{-1}$ ]	-18714.90	-18715.26	-18716.78	4.92	-18720.89	-18709.68	-18726.32	-18703.97	$\mathcal{U}$
$\dot{\ln}$	[ $\text{m s}^{-1} \text{ yr}^{-1}$ ]	-3.13	-2.98	-2.46	2.19	-5.47	-0.48	-8.10	1.92	$\mathcal{U}$
$\dot{quad}$	[ $\text{m s}^{-1} \text{ yr}^{-2}$ ]	0.32	0.30	0.24	0.23	0.04	0.57	-0.22	0.85	$\mathcal{U}$
$P$	[d]	86.685971	86.716536	86.644378	0.163173	86.533594	86.911495	86.364597	87.082150	$\mathcal{U}$
$K$	[ $\text{m s}^{-1}$ ]	6.51	6.57	6.62	0.33	6.19	6.93	5.83	7.32	$\mathcal{U}$
$e$		0.232050	0.230947	0.204544	0.066766	0.152388	0.303804	0.070509	0.372129	$\mathcal{U}$
$\omega$	[deg]	91.237907	97.248455	96.102520	12.239918	83.790120	110.796187	66.715021	127.067212	$\mathcal{U}$
$\lambda_0$	[deg]	215.291405	220.718337	216.160185	16.196583	202.266759	239.462980	185.195600	257.350058	$\mathcal{U}$
$P$	[d]	485.429776	496.902333	484.032769	26.820241	475.087228	532.349347	456.095460	579.472904	$\mathcal{U}$
$K$	[ $\text{m s}^{-1}$ ]	3.38	3.16	3.03	0.37	2.74	3.59	2.33	4.03	$\mathcal{U}$
$e$		0.134236	0.146812	0.021121	0.110116	0.043096	0.298966	0.005614	0.458703	$\mathcal{U}$
$\omega$	[deg]	-358.202334	-311.969148	-380.751216	423.482663	-375.233350	668.453999	-478.847476	714.163530	$\mathcal{U}$
$\lambda_0$	[deg]	240.692904	274.875882	256.683399	77.426560	207.978823	379.031065	146.361972	491.545801	$\mathcal{U}$

**Notes.** The maximum likelihood solution (Max(Like)), the median (Med), mode (Mod) and standard deviation (Std) of the posterior distribution for each parameter is shown, as well as the 68.3% (CI(15.85),CI(84.15)) and 95.45% (CI(2.275),CI(97.725)) confidence intervals. The prior for each parameter can be of type:  $\mathcal{U}$ : uniform,  $\mathcal{N}$ : normal,  $\mathcal{SN}$ : split normal,  $\mathcal{TN}$ : truncated normal.



**Table B.6.** Physical parameters derived from the MCMC chains used fit the RV measurements of Gl617-A - 2-keplerian model plus quadratic drift.

Param.	Units	Max(Like)	Med	Mod	Std	CI(15.85)	CI(84.15)	CI(2.275)	CI(97.725)	Prior
<b>Likelihood</b>										
$M_*$	[ $M_\odot$ ]	0.581188	0.598699	0.589123	0.026717	0.569082	0.629830	0.538640	0.660030	$\mathcal{U}$
$P$	[d]	86.685971	86.716536	86.644378	0.163173	86.533594	86.911495	86.364597	87.082150	$\mathcal{U}$
$K$	[ $\text{m s}^{-1}$ ]	6.51	6.57	6.62	0.33	6.19	6.93	5.83	7.32	$\mathcal{U}$
$e$		0.232050	0.230947	0.204544	0.066766	0.152388	0.303804	0.070509	0.372129	$\mathcal{U}$
$\omega$	[deg]	91.237907	97.248455	96.102520	12.239918	83.790120	110.796187	66.715021	127.067212	$\mathcal{U}$
$T_P$	[d]	55470.128617	55470.170612	55469.066335	4.360219	55465.320275	55475.139385	55459.955671	55480.110606	
$T_C$	[d]	55469.947886	55469.184485	55469.744721	3.803915	55464.775258	55473.489374	55460.507190	55477.410804	
$Ar$	[AU]	0.319911	0.323187	0.321331	0.004833	0.317754	0.328762	0.311972	0.333917	
$M.\sin i$	[ $M_{\text{Jup}}$ ]	0.096018	0.098450	0.097920	0.005996	0.091680	0.105364	0.085691	0.113023	
$M.\sin i$	[ $M_{\text{Earth}}$ ]	30.514385	31.287511	31.118932	1.905555	29.135990	33.484633	27.232456	35.918729	
$P$	[d]	485.429776	496.902333	484.032769	26.820241	475.087228	532.349347	456.095460	579.472904	$\mathcal{U}$
$K$	[ $\text{m s}^{-1}$ ]	3.38	3.16	3.03	0.37	2.74	3.59	2.33	4.03	$\mathcal{U}$
$e$		0.134236	0.146812	0.021121	0.110116	0.043096	0.298966	0.005614	0.458703	$\mathcal{U}$
$\omega$	[deg]	-358.202334	-311.969148	-380.751216	423.482663	-375.233350	668.453999	-478.847476	714.163530	$\mathcal{U}$
$T_P$	[d]	55177.869828	55214.752874	55283.022634	135.078647	55031.879715	55361.011891	54952.197623	55438.122406	
$T_C$	[d]	55276.199082	55275.541145	55277.491526	107.612883	55154.204758	55385.387258	55055.822719	55543.038427	
$Ar$	[AU]	1.008817	1.035866	1.026245	0.039843	0.999836	1.087041	0.965852	1.152976	
$M.\sin i$	[ $M_{\text{Jup}}$ ]	0.090084	0.085769	0.082725	0.010332	0.074076	0.097851	0.062734	0.109579	
$M.\sin i$	[ $M_{\text{Earth}}$ ]	28.628775	27.257285	26.290082	3.283456	23.541287	31.097041	19.936952	34.824225	

**Notes.** The maximum likelihood solution (Max(Like)), the median (Med), mode (Mod) and standard deviation (Std) for the posterior distribution of each parameter is shown, as well as the 68.3% (CI(15.85),CI(84.15)) and 95.45% (CI(2.275),CI(97.725)) confidence intervals. The prior for each parameter can be of type:  $\mathcal{U}$ : uniform,  $\mathcal{N}$ : normal,  $\mathcal{SN}$ : split normal,  $\mathcal{TN}$ : truncated normal.

**Table B.7.** Parameters probed by the MCMC used to fit the combined SOPHIE, CARMENES, and KECK RV measurements of Gl617A.

Param.	Units	Max(Like)	Med	Mod	Std	CI(15.85)	CI(84.15)	CI(2.275)	CI(97.725)	Prior
<b>Likelihood</b>										
log(Post)		-934.737509	-939.951019	-940.359405	2.181801	-942.707691	-937.845539	-946.160275	-936.299385	
log(Like)		-931.716450	-937.193692	-936.740517	2.633389	-940.751319	-934.848899	-944.875654	-933.276083	
log(Prior)		-3.021059	-3.015167	-3.129632	1.055714	-3.056399	-0.160778	-3.122142	-0.006795	
$M_*$	[ $M_\odot$ ]	0.617017	0.599486	0.595975	0.025964	0.569661	0.628938	0.539536	0.658046	$\mathcal{U}$
$\sigma_{JIT}$	[ $\text{m s}^{-1}$ ]	14.15	15.89	15.31	2.25	14.07	20.03	12.57	21.49	$\mathcal{U}$
$\gamma_{\text{CARMENES}}$	[ $\text{m s}^{-1}$ ]	0.51	0.40	0.14	1.44	-1.20	2.03	-2.85	3.80	$\mathcal{U}$
$\gamma_{\text{KECK-PUB}_1}$	[ $\text{m s}^{-1}$ ]	-2.55	-1.76	-1.60	2.37	-4.44	0.87	-7.25	3.72	$\mathcal{U}$
$\gamma_{\text{SOPHIE}}$	[ $\text{m s}^{-1}$ ]	-18721.42	-18721.59	-18721.85	1.34	-18723.09	-18720.05	-18724.63	-18718.48	$\mathcal{U}$
$\dot{\ln}$	[ $\text{m s}^{-1} \text{ yr}^{-1}$ ]	-0.20	-0.12	-0.21	0.40	-0.57	0.32	-1.06	0.79	$\mathcal{U}$
$\dot{quad}$	[ $\text{m s}^{-1} \text{ yr}^{-2}$ ]	0.01	-0.00	-0.02	0.04	-0.05	0.04	-0.09	0.09	$\mathcal{U}$
$P$	[d]	86.717889	86.776472	86.776265	0.135383	86.630677	86.933133	86.478695	87.110679	$\mathcal{U}$
$K$	[ $\text{m s}^{-1}$ ]	5.77	5.83	5.76	0.20	5.59	6.05	5.38	6.28	$\mathcal{U}$
$e$		0.067519	0.071525	0.072428	0.035663	0.030243	0.112694	0.004864	0.159102	$\mathcal{U}$
$\omega$	[deg]	74.887523	97.219922	98.412412	34.567998	55.385453	127.614449	-16.938101	176.384852	$\mathcal{U}$
$\lambda_0$	[deg]	220.983105	227.841194	223.343831	14.690130	211.859018	244.846508	195.405489	263.911937	$\mathcal{U}$

**Notes.** The maximum likelihood solution (Max(Like)), the median (Med), mode (Mod) and standard deviation (Std) for the posterior distribution of each parameter is shown, as well as the 68.3% (CI(15.85),CI(84.15)) and 95.45% (CI(2.275),CI(97.725)) confidence intervals. The prior for each parameter can be of type:  $\mathcal{U}$ : uniform,  $\mathcal{N}$ : normal,  $\mathcal{SN}$ : split normal,  $\mathcal{TN}$ : truncated normal.

**Table B.8.** Physical parameters derived from the MCMC chains used fit the combined SOPHIE, CARMENES, and KECK RV measurements of Gl617A.

Param.	Units	Max(Like)	Med	Mod	Std	CI(15.85)	CI(84.15)	CI(2.275)	CI(97.725)	Prior
<b>Likelihood</b>										
$M_*$	[ $M_\odot$ ]	0.617017	0.599486	0.595975	0.025964	0.569661	0.628938	0.539536	0.658046	$\mathcal{U}$
$P$	[d]	86.717889	86.776472	86.776265	0.135383	86.630677	86.933133	86.478695	87.110679	$\mathcal{U}$
$K$	[ $\text{m s}^{-1}$ ]	5.77	5.83	5.76	0.20	5.59	6.05	5.38	6.28	$\mathcal{U}$
$e$		0.067519	0.071525	0.072428	0.035663	0.030243	0.112694	0.004864	0.159102	$\mathcal{U}$
$\omega$	[deg]	74.887523	97.219922	98.412412	34.567998	55.385453	127.614449	-16.938101	176.384852	$\mathcal{U}$
$T_P$	[d]	55464.808054	55468.264643	55469.553885	9.043231	55457.599658	55476.783117	55440.927623	55489.460498	
$T_C$	[d]	55467.985292	55466.914469	55465.796807	3.657568	55462.799889	55471.033793	55458.170838	55475.281484	
$A_r$	[AU]	0.326432	0.323420	0.322034	0.004684	0.317906	0.328658	0.312283	0.333674	
$M.\sin i$	[ $M_{\text{Jup}}$ ]	0.090837	0.089847	0.090072	0.004054	0.085275	0.094534	0.080783	0.099046	
$M.\sin i$	[ $M_{\text{Earth}}$ ]	28.867969	28.553526	28.624925	1.288239	27.100269	30.042914	25.672842	31.476762	

**Notes.** The maximum likelihood solution (Max(Like)), the median (Med), mode (Mod) and standard deviation (Std) for the posterior distribution of each parameter is shown, as well as the 68.3% (CI(15.85),CI(84.15)) and 95.45% (CI(2.275),CI(97.725)) confidence intervals. The prior for each parameter can be of type:  $\mathcal{U}$ : uniform,  $\mathcal{N}$ : normal,  $\mathcal{SN}$ : split normal,  $\mathcal{TN}$ : truncated normal.

**Table A.1.** Radial velocities for Gl96 derived by template-matching, with the short-term instrumental drift and long-term zero-point corrections applied.

BJD [-2400000d]	RV [km/s]	sigma RV [km/s]
55813.6543	-37.8703	0.0023
55916.385	-37.8778	0.003
56177.6458	-37.8799	0.002
57260.6283	-37.8687	0.0018
57261.6026	-37.8647	0.0017
57262.591	-37.8692	0.0017
57263.5689	-37.8714	0.0018
57264.6339	-37.8793	0.0019
57269.613	-37.8702	0.0021
57272.5499	-37.8704	0.003
57285.486	-37.8727	0.0023
57287.5502	-37.8739	0.0018
57289.513	-37.8721	0.0025
57291.5016	-37.8723	0.0023
57292.6058	-37.88	0.0017
57318.4756	-37.8718	0.002
57319.5631	-37.8764	0.0021
57324.5429	-37.8759	0.0023
57325.525	-37.8747	0.0017
57326.4449	-37.8705	0.0017
57327.4813	-37.8652	0.0018
57341.5201	-37.8722	0.0023
57342.4705	-37.8749	0.0018
57349.3667	-37.875	0.002
57352.4482	-37.8775	0.0026
57353.411	-37.8796	0.0021
57373.4177	-37.8781	0.0018
57375.413	-37.8766	0.002
57631.6103	-37.867	0.002
57632.6444	-37.8706	0.0023
57634.637	-37.8721	0.0021
57635.6266	-37.8683	0.0018
57639.5888	-37.8729	0.0021
57640.5853	-37.8747	0.002
57642.596	-37.8722	0.0024
57652.5243	-37.8814	0.0024
57656.5486	-37.8768	0.0025
57657.5581	-37.8723	0.0025
57664.4734	-37.8716	0.0024
57665.4864	-37.875	0.0022
57666.5107	-37.8758	0.0017
57667.5688	-37.8812	0.0027
57669.5344	-37.8772	0.0021
57670.4846	-37.8803	0.0024
57671.4838	-37.8748	0.0025
57677.513	-37.8789	0.0016
57680.5849	-37.8735	0.0022
57700.6179	-37.8657	0.0022
57701.4932	-37.8712	0.0024
57709.5002	-37.874	0.0017
57712.2714	-37.8743	0.0026
57729.3458	-37.8747	0.0017
57730.4299	-37.8755	0.0017
57731.4116	-37.8782	0.0017
57732.4296	-37.8821	0.0018
57733.3579	-37.8835	0.0017
57734.3638	-37.8815	0.0024
57740.4115	-37.8775	0.0019
57741.3933	-37.8712	0.0019
57758.299	-37.875	0.0025

Table A.1. continued.

BJD [ $-2400000d$ ]	RV [ $km/s$ ]	sigma RV [ $km/s$ ]
57761.2904	-37.8822	0.0023
57762.3098	-37.8812	0.0017
57765.2701	-37.8794	0.0019
57770.3366	-37.8726	0.0027
57772.3558	-37.8708	0.0017
57801.3467	-37.8733	0.0031
57815.3111	-37.8793	0.0019
57822.3095	-37.885	0.0029
57823.3328	-37.8796	0.0021
57825.2954	-37.8803	0.0024
57828.2853	-37.877	0.0022
57832.2892	-37.8687	0.0019

**Table A.2.** Radial velocities for Gl617A derived by template-matching, with the short-term instrumental drift and long-term zero-point corrections applied.

BJD [-2400000d]	RV [km/s]	sigma RV [km/s]
55827.3208	-18.7247	0.0019
55982.7022	-18.714	0.0018
56328.6907	-18.7186	0.0019
56370.663	-18.7214	0.002
56372.632	-18.7201	0.0022
56377.596	-18.7262	0.0023
56405.4505	-18.7182	0.002
56419.4838	-18.7232	0.002
56425.4627	-18.7237	0.0028
56469.5016	-18.4846	0.0465
56471.4651	-18.7236	0.0026
56487.4569	-18.7161	0.0022
56488.4484	-18.7198	0.0056
56490.3756	-18.7147	0.0022
56496.4612	-18.7101	0.0023
56497.4503	-18.7158	0.002
56500.3679	-18.7166	0.0019
56501.48	-18.7186	0.0036
56502.3675	-18.713	0.0022
56519.3387	-18.7259	0.0021
56521.3385	-18.7281	0.0018
56523.3515	-18.7262	0.0019
56556.3177	-18.7195	0.0021
56557.3098	-18.7184	0.0019
56558.2799	-18.7188	0.0018
56583.2842	-18.7161	0.0021
56613.2241	-18.7205	0.002
56624.2446	-18.7294	0.0033
56625.2369	-18.7243	0.0019
56626.2231	-18.7218	0.0023
56629.2163	-18.7274	0.0024
56694.6636	-18.7255	0.0024
56705.6963	-18.7306	0.0024
56711.7062	-18.726	0.0023
56712.6984	-18.725	0.002
56730.5614	-18.7189	0.002
56732.6738	-18.7207	0.0029
56733.6147	-18.7156	0.0018
56761.5278	-18.7143	0.002
56777.4867	-18.7315	0.0024
56779.5828	-18.7281	0.0021
56782.5313	-18.7304	0.0019
56793.4986	-18.7205	0.0019
56794.5888	-18.7242	0.0019
56800.5492	-18.7373	0.0033
56815.4608	-18.7189	0.0018
56818.4207	-18.7249	0.0019
56835.3756	-18.7293	0.0058
56841.3896	-18.7216	0.0021
56844.4113	-18.7214	0.002
56850.4049	-18.7169	0.0024
56854.42	-18.715	0.0019
56859.4538	-18.7286	0.0031
56862.5723	-18.7348	0.0048
56876.3988	-18.7228	0.0027
56885.4011	-18.7334	0.003
56904.3149	-18.7264	0.0023
56914.3578	-18.7178	0.0024
56926.3241	-18.7209	0.0017
56929.2964	-18.7251	0.0017

Table A.2. continued.

BJD [-2400000d]	RV [km/s]	sigma RV [km/s]
56939.3032	-18.713	0.0021
56941.2947	-18.7143	0.0021
56942.2937	-18.7233	0.0017
56962.2331	-18.7236	0.0017
56968.2461	-18.7293	0.0021
56999.2152	-18.7108	0.0032
57002.2308	-18.7135	0.0057
57072.6911	-18.7282	0.002
57079.64	-18.7204	0.0022
57084.6695	-18.7188	0.0026
57087.6397	-18.7157	0.0022
57089.6958	-18.7118	0.0018
57090.615	-18.7094	0.0024
57091.5817	-18.7152	0.0019
57115.5282	-18.73	0.003
57119.5769	-18.7259	0.0023
57438.6677	-18.722	0.0023
57465.6133	-18.7231	0.0019
57466.5738	-18.7289	0.0018
57469.5975	-18.7307	0.0019
57471.637	-18.7295	0.0023
57474.5271	-18.7266	0.0023
57476.5225	-18.7293	0.0018
57496.5362	-18.7298	0.0021
57498.4581	-18.7274	0.002
57502.5692	-18.7208	0.0029
57507.5044	-18.7187	0.0019
57510.5087	-18.7192	0.0031
57512.5605	-18.7145	0.0031
57521.4872	-18.4686	0.0648
57523.4672	-18.7198	0.0045
57524.4615	-18.7157	0.0036
57526.6088	-18.7141	0.0019
57528.5508	-18.714	0.0021
57529.4749	-18.716	0.0018
57530.5157	-18.7174	0.0021
57544.5369	-18.7181	0.0023
57547.4552	-18.7158	0.0019
57548.4152	-18.7183	0.0023
57549.4331	-18.4442	0.0485
57550.5206	-18.721	0.002
57553.3989	-18.7221	0.0025
57557.4901	-18.7244	0.0017
57559.4385	-18.7276	0.0024
57562.4873	-18.4644	0.0478
57562.5082	-18.7308	0.0018
57564.5346	-18.7248	0.0026
57566.4621	-18.7266	0.003
57568.4546	-18.7204	0.002
57571.3651	-18.7236	0.0025
57574.4738	-18.7253	0.0025
57575.4626	-18.7195	0.0028
57582.4238	-18.7254	0.0027
57583.3905	-18.726	0.0025
57586.4968	-18.7188	0.0018
57593.393	-18.7072	0.0029
57594.408	-18.713	0.0024
57597.3821	-18.7161	0.0023
57614.3752	-18.71	0.0019
57618.4374	-18.7158	0.0026
57621.3838	-18.7191	0.003

Table A.2. continued.

BJD [-2400000d]	RV [km/s]	sigma RV [km/s]
57622.4412	-18.7201	0.0043
57633.3421	-18.7199	0.0019
57635.32	-18.7217	0.002
57637.3095	-18.7181	0.0024
57638.307	-18.7154	0.0023
57639.2986	-18.4182	0.0437
57639.3109	-18.717	0.0023
57641.3044	-18.7198	0.0021
57652.2945	-18.727	0.0021
57656.3939	-18.7278	0.0024
57795.702	-18.7188	0.0018
57800.6382	-18.7199	0.002
57814.6062	-18.7299	0.0022
57820.6203	-18.733	0.0025
57823.6354	-18.7319	0.0017
57825.6004	-18.731	0.002
57827.5652	-18.7296	0.0018
57831.6704	-18.727	0.0023
57849.5691	-18.7246	0.0018
57851.6214	-18.7276	0.0016
57852.6711	-18.7255	0.0019
57853.6085	-18.7269	0.0016
57860.5126	-18.7242	0.0032
57861.5489	-18.72	0.0028
57883.518	-18.7135	0.0017
57886.5506	-18.7163	0.0017
57888.4005	-18.7187	0.0019
57914.4292	-18.7264	0.0019
57916.4513	-18.7279	0.002
57917.4221	-18.7333	0.0016
57918.4669	-18.7285	0.002
57918.5051	-18.7413	0.0019
57922.4145	-18.7238	0.0017