



**HAL**  
open science

## About biomass overyielding of mixed cultures in batch processes

Alain Rapaport, Thibault Nidelet, Sahar El Aida, Jérôme Harmand

### ► To cite this version:

Alain Rapaport, Thibault Nidelet, Sahar El Aida, Jérôme Harmand. About biomass overyielding of mixed cultures in batch processes. 2019. hal-02010420v1

**HAL Id: hal-02010420**

**<https://hal.science/hal-02010420v1>**

Preprint submitted on 7 Feb 2019 (v1), last revised 7 Oct 2019 (v2)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# About biomass overyielding of mixed cultures in batch processes

A. Rapaport <sup>a,\*</sup> T. Nidelet <sup>b</sup> S. El Aidar <sup>a,b</sup> J. Harmand <sup>c</sup>

<sup>a</sup>*MISTEA, Univ. Montpellier, INRA, Montpellier SupAgro, France.*

<sup>b</sup>*SPO, Univ. Montpellier, INRA, Montpellier SupAgro, France.*

<sup>c</sup>*LBE, Univ. Montpellier, INRA, Narbonne, France.*

---

## Abstract

We study mechanisms that can produce an increase of biomass production in batch processes when considering mixed cultures, compared to pure cultures. We show that growth thresholds or variable yields can produce ‘overyielding’, while this is not possible in the classical batch model with multiple species. We give sufficient conditions on the characteristics of the species to obtain overyielding, and illustrate these theoretical results with numerical simulations. This work provides new insights on species complementary in models of mixed cultures, without having to consider direct interactions terms between species as, for instance in the well known Generalized Lotka-Volterra model.

*Key words:* batch culture, complex ecosystems, diversity, overyielding, transient, optimization.

---

## 1 Introduction

Microorganisms play an important role in many ecosystems on earth, such as carbon or nitrogen biochemical cycles, or food chains [26]. Today, many industrial bio-processes rely on bacteria or yeasts to transform matter into high-value products, such as in the food industry (winery, bakery, cheese dairy...). Cultivation is operated either in continuous (chemostat) [15], batch [6] or fed-batch modes [19]. Although questions related to biodiversity (exclusion,

---

\* Corresponding author.

*Email addresses:* [alain.rapaport@inra.fr](mailto:alain.rapaport@inra.fr) (A. Rapaport),  
[thibault.nidelet@inra.fr](mailto:thibault.nidelet@inra.fr) (T. Nidelet), [saharelaida@gmail.com](mailto:saharelaida@gmail.com) (S. El Aidar),  
[jerome.harmand@inra.fr](mailto:jerome.harmand@inra.fr) (J. Harmand).

coexistence, persistence of species) have been already well investigated for continuous mode (see e.g. [28]), batch cultures have received less fundamental attention with respect to the study of diversity. One of the main reasons is much probably that most of the industrial batch processes rely on a single species that have been carefully selected for its characteristics (such as yield conversion, growth kinetics, products composition). Another possible explanation is the relative recent availability of genetic tools to discriminate with accuracy the ecosystems composition in these application fields.

Yet, in many cases, a better performance has been observed with mixed cultures when compared to the performances obtained with a single species. When the criterion is the conversion into biomass or into a product of the bioreaction, we shall say that such situations exhibit an 'overyielding'. Several natural batch ecosystems rely on a certain diversity for their functioning, such as the ones one can find in cheese dairy [22] or animal rumen [5] activities. In such ecosystems, a number of species interact together and are complementary one of each others in terms of functioning. Finally, in many cases, a complex ecosystem exhibits better stability and robustness properties with respect to unexpected situations. It is why species interactions have been studied in order to explain how the main functions of the ecosystems can be ensured by concomitant species. Nevertheless, it appears that few works have investigated the role of ecosystems diversity - from a theoretical framework - in terms of performances of a given function, such as the biomass production. In industrial applications, the paradigm that the best performances are necessarily obtained by the selection of a single species - *the best one* - seems to remain quite strong. Let us underline that differently to the continuous mode, for which competitive exclusion predicts that under perfectly well-controlled environment the best species excludes the other ones at steady state, the dynamics of batch culture is fundamentally transitory. Therefore, dominance alternation among species can occur.

The objective of the present work is to study with existing mathematical models [20, 25] mechanisms that can explain overyielding in batch cultures. Here we will say that there is biomass overyielding in a batch process if the total production of biomass using an inoculum composed of multiple species is greater than the biomass production obtained when the inoculum is made up of a single species. In other words, we study the role of the diversity of the inoculum with respect to process performances in terms of biomass production. For simplicity, we restrict our attention to single step reactions, which means that we investigate possible mechanisms of overyielding in any biological system where the biomass growth is limited by a single limiting substrate. In other terms, we study biomass overyielding in the framework of the competition of  $n$  species on a single resource.

The paper is organized as follows. In the next Section 2, we first introduce assumptions and modeling framework to give a precise mathematical definition of overyielding. In Section 3 we consider the most classical batch model and show, under the simplest hypotheses, that overyielding is not possible (the greatest biomass production is obtained when the ‘best’ species is cultivated alone). For this simplest case however, it is shown that results hold for a very important class of growth functions. Then, we consider more sophisticated models from the literature and highlight key mechanisms for biomass overyielding to occur: constant yield with a threshold in the growth functions of species in Section 4 and variable yields in Section 5. For these two classes of models, we give precise conditions for which overyielding is possible. Summary and comparison of possibilities to obtain overyielding are given in Section 6. Finally, numerical examples which illustrate the possibilities of overyielding for these models are then provided and discussed in Section 7.

## 2 Model, assumptions and definitions

Under appropriate environmental conditions (temperature, pH,...), we consider a vessel which contains all the biotic resources necessary for microorganisms to grow, except a single limiting resource. At initial time, one or several species and the resource are introduced (in fermentation processes, these are typically one or several yeasts and sugar). We shall denote by  $X_i(t)$ ,  $i \in \{1, \dots, n\}$  the concentrations (or densities) of microorganisms of species  $i$ , and by  $S(t)$  the concentration of the resource. The general expression of classical consumers-resources model which describes the time evolution of the reactions that take place between  $X_i$  and  $S$  is given by the following system:

$$\begin{aligned} \dot{X}_i &= \mu_i(S)X_i \quad (i = 1 \dots n) \\ \dot{S} &= - \sum_{i=1}^n \frac{1}{y_i(S)} \mu_i(S)X_i \end{aligned} \tag{1}$$

where  $\mu_i(\cdot)$  is the specific growth rate function and  $y_i(\cdot)$  the yield conversion factor of species  $i$ . In microbiology, four phases are usually described in batch processes: 1. the lag phase, 2. the growth phase, 3. the stationary phase and 4. the decline phase. As our objective here is to focus on the production of the growth phase, we neglect with this model the lag phase, assuming that at initial time all the species are already in their ‘active’ phase. We also do not consider the decline phase, and assume that the mortality of micro-organisms is negligible during the growth and stationary phase. However, we shall allow  $y(\cdot)$  to be possibly resource dependent, as this is commonly met in literature (see e.g. [2, 25]). We make the following assumptions on the functions  $\mu_i$  and  $y_i$ .

## Hypothesis 1

(1) The functions  $\mu_i(\cdot)$  are Lipschitz continuous from  $\mathbb{R}_+$  to  $\mathbb{R}_+$  and there exist growth thresholds  $\underline{S}_i \geq 0$  ( $i = 1 \cdots n$ ) such that

$$\mu_i(S) = 0, \quad S \in [0, \underline{S}_i], \quad \mu_i(S) > 0, \quad S > \underline{S}_i$$

(2) The functions  $y_i(\cdot)$  are Lipschitz continuous from  $\mathbb{R}_+$  to  $\mathbb{R}_+$ , with  $y_i(S) > 0$  for any  $S \in \mathbb{R}_+$  ( $i = 1 \cdots n$ ).

The threshold  $\underline{S}_i$  represents the minimal value of the substrate concentration under which a growth cannot occur [32]. This value is often imposed to be equal to 0, as in the classical Monod model. However, in practice it is rare to have observations for low values of the substrate concentration to identify such threshold. Nevertheless, it has been observed that for some strains, this threshold is not negligible [29, 31].

Under this hypothesis, one can easily check that solutions of the system of differential equations (1) are well defined and bounded at any positive time, whatever the initial condition is in  $\mathbb{R}_+^{n+1}$ .

Recall that we shall speak of overyielding when the production of biomass of an inoculum composed of multiple species is greater than the production obtained when the inoculum is made up of a single species. Let us formalize this definition from a mathematical viewpoint: let us denote by  $\mathcal{S}$  the simplex in the positive orthant of  $\mathbb{R}^n$

$$\mathcal{S} := \left\{ p \in \mathbb{R}_+^n; \sum_i p_i = 1 \right\}$$

and the vertices  $\sigma_i$  ( $i = 1 \cdots n$ ) such that

$$\sigma_{i,i} = 1, \quad \sigma_{i,j} = 0, \quad j \neq i$$

We shall also denote by  $X$  the vector in  $\mathbb{R}_+^n$  of components  $X_i$ .

**Definition 2** Let  $B_0, S_0$  be two positive numbers. For any  $p_0 \in \mathcal{S}$ , consider the solution of (1) for the initial condition

$$X(0) = p_0 B_0, \quad S(0) = S_0$$

and define the number

$$R(p_0) := B_\infty - B_0 \quad \text{where} \quad B_\infty := \lim_{t \rightarrow +\infty} B(t) \quad \text{with} \quad B(t) = \sum_i^n X_i(t)$$

System (1) is said to present overyielding for the pair  $(B_0, S_0)$  when the inequality

$$\max_{p_0 \in \mathcal{S}} R(p_0) > \max_{i \in \{1 \dots n\}} R(\sigma_i) \quad (2)$$

is fulfilled.

In other words, overyielding in this precise framework means that there exists a composition of the initial biomass with different species that gives a greater production of biomass than the one we obtain when the initial biomass is made up of a single species.

In the definition of the function  $R$ , we consider asymptotic values  $B_\infty$  of the biomass, whereas we have previously mentioned that the model do not take into consideration the decline phase of the biomass which occurs for large times. Therefore, those asymptotic values may not be reached in practice. However, if one considers that the stationary phase corresponds to biomass levels close to these asymptotic values, one may reasonably assume that the comparison order between the biomass productions determined with the asymptotic values of the model is also verified at the stationary phase.

**Remark 3** *In some works [9, 27], the weaker condition  $R(p_0) > \sum_{i=1}^n p_{0,i} R(\sigma_i)$  is considered as the definition for overyielding, and condition (2) refers then as transgressive overyielding. We believe that this difference makes sense when the criterion is static or deals with steady-states. Here the proportions of species in the consortium are changing with time, which makes us choose a stricter definition of overyielding, as we shall see later on.*

To study the possibilities of biomass overyielding, it is useful to first characterize the residual concentration of substrate, with the following lemma.

**Lemma 4** *For any initial condition in  $\mathbb{R}_+^n$ , the solution of (1) verifies*

$$S_\infty := \lim_{t \rightarrow +\infty} S(t) = \min \left( S(0), \min_{i=1 \dots n} \{ \underline{S}_i \text{ s.t. } X_i(0) > 0 \} \right)$$

**PROOF.** Posit  $l = \min(S(0), \min_{i=1 \dots n} \{ \underline{S}_i \text{ s.t. } X_i(0) > 0 \})$ .

Notice first, from equations (1), that  $S(\cdot)$  is a decreasing function bounded from below by 0. Therefore, it admits a limit  $S_\infty \leq S(0)$ . Moreover, the functions  $X_i(\cdot)$  are non decreasing. One can then write

$$\sum_{i=1}^n \frac{\mu_i(S(t))}{y_i(S(t))} X_i(t) \geq \sum_{i=1}^n \min_{S \in [S_\infty, S(0)]} \left\{ \frac{\mu_i(S)}{y_i(S)} \right\} X_i(0), \quad t \geq 0$$

If  $S_\infty > l$ , there exists  $j \in \{1, \dots, n\}$  such that  $X_j(0) > 0$  with  $l = \underline{S}_j < S(0)$ . Then, let

$$\eta = \min_{S \in [S_\infty, S(0)]} \frac{\mu_j(S)}{y_j(S)} X_j(0) > 0$$

and one has  $\dot{S}(t) \leq -\eta$  for any  $t > 0$ , which contradicts the positivity of the variable  $S$  for any time.

If  $S_\infty < l$ , there exists  $\tilde{S} \in (0, l)$  and  $T > 0$  such that  $S(T) = \tilde{S}$ . Then,  $(X(T), \tilde{S})$  is a steady-state of (1), which contradicts the uniqueness of solutions of (1).

We conclude that the equality  $S_\infty = l$  is satisfied.

**Remark 5** *When there is no threshold i.e.  $\underline{S}_i = 0$ , we find the well-known fact that there is no residual substrate at the end of the reaction.*

### 3 Overyielding in the classical ‘batch competition model’

We consider here the model (1) with constant yields  $y_i(\cdot) = Y_i$  and growth functions with no threshold i.e.  $\underline{S}_i = 0$ .

**Proposition 6** *Let  $B_0, S_0$  be two positive numbers.*

1. *One has*

$$R(\sigma_i) = Y_i S_0, \quad i = 1 \dots n$$

2. *The system (1) does not present any overyielding, and best species  $i^*$  are the ones with the highest  $Y_{i^*}$ .*

**PROOF.** From Lemma 4, one has  $S_\infty = 0$  whatever is the initial composition  $p_0$  of the biomass. From (1), one can write

$$\frac{d}{dt} \left( \sum_{i=1}^n \frac{X_i(t)}{Y_i} + S(t) \right) = 0$$

and then, integrating between  $t = 0$  and  $t = +\infty$ , one obtains

$$\sum_{i=1}^n \frac{X_i(\infty) - X_i(0)}{Y_i} = S_0$$

Posit

$$q_i = \frac{X_i(\infty) - X_i(0)}{Y_i S_0}, \quad i = 1 \dots n$$

Then, for any initial proportion  $p_0 \in \mathcal{S}$  of the species, one has

$$R(p_0) = \sum_{i=1}^n X_i(\infty) - X_i(0) = \sum_{i=1}^n q_i Y_i S_0$$

For the proportions  $\sigma_i$  ( $i = 1 \cdots n$ ), one has

$$R(\sigma_i) = Y_i S_0, \quad i = 1 \cdots n$$

which proves the point 1. One can then write

$$R(p_0) = \sum_{i=1}^n q_i R(\sigma_i)$$

As the numbers  $q_i$  belongs to  $[0, 1]$  and their sum is equal to 1, we conclude that any  $R(p_0)$  cannot be above the largest  $R(\sigma_i)$  with  $i \in \{1, \cdots, n\}$ , which proves point 2.

**Remark 7** Notice that Proposition 6 remains valid whatever the expressions of  $\mu_i(\cdot)$  are. The only condition is that these functions are positive when  $S$  is positive, so that one has  $\underline{S}_i = 0$ . Indeed, the only condition for the result to hold is that the terms  $\mu_i(\cdot)$  appearing in the dynamics of  $X_i$  be the same than the ones appearing negatively in the dynamics of  $S_i$ , whatever are the growth rates under the condition that  $S_\infty = 0$ : they can involve any state of the system or any external ‘inputs’ or environmental variables like the pH or the oxygen concentration, or even include direct interaction terms. From a chemical engineering viewpoint, this result is easily understandable: since our performance index - the biomass production - is essentially related to the yield of the biomass, the final state does not depend on the transitory but only on the initial resource available.

In the next sections, we study two different mechanisms for biomass overyielding to appear.

#### 4 Overyielding with constant yields and growth thresholds

We denote by capital letters functions  $y_i(\cdot)$  when constant, that is  $y_i(S) = Y_i$ .

**Proposition 8** Fix  $B_0, S_0$  two positive numbers.

1. One has

$$R(\sigma_i) = Y_i \max(0, S_0 - \underline{S}_i), \quad i = 1 \cdots n \quad (3)$$



2. Let  $j \in \{1, \dots, n\}$  be such that  $R(\sigma_j) = \max_{i \in \{1, \dots, n\}} R(\sigma_i) > 0$ . If there exists  $k \in \{1, \dots, n\} \setminus \{j\}$  such that  $Y_k > Y_j$ , then for  $S_0 > \underline{S}_k$ , the model (1) presents an overyielding.

**PROOF.**

1. For pure cultures i.e. proportions  $p_0$  equal to one of the  $\sigma_i$ ,  $i \in \{1, \dots, n\}$ , one has from equations (1)

$$\frac{d}{dt}(X_i(t) + Y_i S(t)) = 0$$

Integrating this equality between  $t = 0$  and  $t = +\infty$  gives

$$X_i(\infty) - X_i(0) = Y_i(S_0 - S_\infty)$$

where  $S_\infty = \min(S_0, \underline{S}_i)$  by Lemma 4. Therefore, one obtains the equality  $R(\sigma_i) = Y_i \max(0, S_0 - \underline{S}_i)$ .

2. Consider an initial proportion  $p_0$  with the presence of the species  $j$  and  $k$  only. Note that  $Y_k > Y_j$  implies necessarily to have  $\underline{S}_k > \underline{S}_j$  (as  $R(\sigma_j) > R(\sigma_k)$ ). When  $S_0 > \underline{S}_k$ , the two species grow and one has  $S_\infty = \underline{S}_j$  (by Lemma 4). From equations (1), we write

$$\frac{d}{dt} \left( \frac{X_j(t)}{Y_j} + \frac{X_k(t)}{Y_k} + S(t) \right) = 0$$

and by integration, one has

$$\frac{X_j(\infty) - X_j(0)}{Y_j} + \frac{X_k(\infty) - X_k(0)}{Y_k} = S_0 - \underline{S}_j$$

From  $Y_k > Y_j$ , one gets

$$R(p_0) = X_j(\infty) + X_k(\infty) - X_j(0) - X_k(0) > Y_j(S_0 - \underline{S}_j) = R(\sigma_j)$$

which shows that a composition of species  $j$  and  $k$  gives a better production than the best species  $j$  alone.

This result shows that having different growth thresholds could be a way to obtain overyielding, playing with a *complementary effect* among species. Typically, this happens when a species  $j$  has the best biomass production among single species, and another species  $k$  has an even better conversion factor  $Y_k$  but suffers from a relatively large growth threshold  $\underline{S}_k$  (so that it can no longer grow when the resource level  $S$  is below  $\underline{S}_k$ ). Then, the presence

of species  $k$  in addition to species  $j$  boosts the performances of the ecosystem when  $S$  is above  $\underline{S}_k$  due to the better conversion factor of species  $k$ . Later on, species  $j$  carries alone on the conversion of the resource, ensuring a better total production at the end when compared to situations where  $k$  or  $j$  are alone.

**Remark 9** *When the numbers  $\underline{S}_i$ ,  $i \in \{1, \dots, n\}$  are all identical, one can straightforwardly extend the result of Proposition 6 to show that no overyielding is possible.*

The mechanism we have identified for overyielding is based on growth thresholds, assuming that the species do not consume any substrate when there is no growth. We investigate now another mechanism that could produce overyielding.

## 5 Overyielding with variable yields

In this section, we explore another mechanism based on variable yields, that can produce overyielding. To properly separate this mechanism from the one induced by growth thresholds that has been studied in Section 4, we come back to the hypothesis used in Section 3 and consider growth thresholds  $\underline{S}_i$  all equal to 0.

Evidence of variable yield cultures could be deduced from observed oscillations in both continuous [7, 10] or batch cultures [17, 33], based on the fact that models without variable yields cannot exhibit oscillations [1, 18]. Moreover, models with several species and variable yields have been also investigated in the literature but for continuous cultures [14, 23] and not - at the best of author's knowledge - to characterizing overyielding in batch, as we do in the present work.

Several biological mechanisms could justify a variable yield, such as extracellular material or transporters [8, 12]. However, the most frequently encountered justification is related to maintenance terms [16, 24, 25] Typically, in continuous culture, a maintenance term  $mX$  (where  $m$  is a positive constant) is subtracted from the substrate kinetics, as follows:

$$\begin{aligned}\dot{X} &= \mu(S)X - DX \\ \dot{S} &= -\frac{1}{Y}\mu(S)X + D(S_{in} - S) - mX\end{aligned}\tag{4}$$

where  $D$  represents the dilution rate. Nevertheless, although this formalism has been successfully validated in chemostats [16], this writing is not satisfactory in batch mode (i.e. when  $D = 0$ ) because the solutions of the differential

equations (4) can take negative values of  $S$ . This issue has been addressed by several authors that have proposed more complex models, incorporating microbial death [3, 30, 34]. However, non-constant maintenance terms have been observed experimentally (see e.g. [4, 13]). This is why we consider here the maintenance term  $m$  as a function of  $S$  with  $m(0) = 0$  (which gives no substrate consumption at  $S = 0$ ) instead of a constant coefficient. One can take, for instance, an expression of the form  $m(S) = \min(kS, m_c)$  (where  $k$  and  $m_c$  are constant parameters) to recover the constant case  $m = m_c$  when  $S > m_c/k$ . Therefore, the model (1) with constant yield  $Y_i$  becomes

$$\begin{aligned} \dot{X}_i &= \mu(S_i)X_i \quad (i = 1 \cdots n) \\ \dot{S} &= - \sum_{i=1}^n \frac{1}{Y_i} \mu_i(S)X_i - m_i(S)X_i \end{aligned} \quad (5)$$

We shall consider the following assumption

**Assumption 10** *The function  $m_i(\cdot)$  are Lipschitz continuous from  $\mathbb{R}_+$  to  $\mathbb{R}_+$  with  $m_i(0) = 0$  and  $m_i(S) > 0$  for  $S > 0$ . Moreover, for each  $i \in \{1, \dots, n\}$ , one has*

$$\exists \lim_{S \rightarrow 0, S > 0} \frac{m_i(S)}{\mu_i(S)} < +\infty$$

Under this assumption, one can define for each  $i \in \{1 \cdots n\}$ , the function

$$\alpha_i(S) := \frac{m_i(S)}{\mu_i(S)}, \quad S \geq 0$$

and consider functions  $y_i(\cdot)$  (denoted in lowercase letters) defined as follows

$$y_i(S) := \frac{Y_i}{1 + \alpha_i(S)Y_i}, \quad S \geq 0$$

Then, formally, one can check that system (5) is equivalent to (1).

**Proposition 11** *Fix  $B_0, S_0$  two positive numbers.*

1. *One has*

$$R(\sigma_i) = \int_0^{S_0} y_i(s) ds, \quad i = 1 \cdots n \quad (6)$$

2. *Assume there exist  $j, k$  in  $\{1, \dots, n\}$  such that*

(i) *One has  $\lim_{S_0 \rightarrow +\infty} R(\sigma_j) = +\infty$ , and for  $S_0$  large enough, one has  $R(\sigma_j) = \max_{i \in \{1 \cdots n\}} R(\sigma_i)$ .*

(ii) *There exist positive numbers  $S^*, \tilde{S}$  with  $\tilde{S} \geq S^*$  such that*

$$(a) (y_j(S) - y_k(S))(S^* - S) > 0, \forall S \neq S^*$$

$$(b) \exists \varepsilon > 0 \text{ s.t. } y_k(S) \geq y_j(S) + \varepsilon, \mu_k(S) \geq \mu_j(S), \forall S \geq \tilde{S}$$

Then, for any  $B_0 > 0$ , the model (5) presents a biomass overyielding for  $S_0$  large enough.

## PROOF.

1. For pure cultures, equations (1) give

$$\frac{d}{dt}(X_i(t) + y_i(S(t))S(t)) = 0$$

Integrating this equality between  $t = 0$  and  $t = +\infty$  gives

$$X_i(\infty) - X_i(0) = \int_0^\infty y_i(S(t))S(t)dt$$

By Lemma 4 one has  $S_\infty = 0$  and from (1)  $S(\cdot)$  is a decreasing function from  $[0, \infty]$  to  $[0, S_0]$ . Therefore, we obtain

$$R(\sigma_i) = \int_0^{S_0} y_i(s)ds$$

2. Choose  $B_0 > 0$  and  $S_0 > \tilde{S}$ . We shall consider ecosystems with species  $j$  and  $k$  only. From Lemma 4, we know that  $S(t)$  tends to 0 when  $t$  tends to  $+\infty$ . As  $S(\cdot)$  is decreasing with time, there exists  $t^* > \tilde{t} > 0$  such that  $S(\tilde{t}) = \tilde{S}$  and  $S(t^*) = S^*$ . We proceed in three steps.

*Step 1:*  $S(t) \in [0, S^*]$ .

From equations (1), let us write  $\dot{S}$  as follows

$$\dot{S} = -\frac{\dot{X}_j}{y_j(S)} - \frac{\dot{X}_k}{y_k(S)} = -\frac{\dot{X}_j + \dot{X}_k}{y_j(S)} + \left( \frac{1}{y_j(S)} - \frac{1}{y_k(S)} \right) \dot{X}_k$$

and define the number

$$\eta = - \min_{s \in [0, S^*]} \left( 1 - \frac{y_j(s)}{y_k(s)} \right)$$

From hypothesis (ii)-(a),  $\eta$  is a positive number. Then, one can write

$$\dot{B}(t) = \dot{X}_j(t) + \dot{X}_k(t) \geq -y_j(S(t))\dot{S}(t) - \eta\dot{X}_k(t), \quad t \geq t^*$$

and integrating this last inequality between  $t = t^*$  and  $t = +\infty$ , one obtains

$$B_\infty - B(t^*) \geq \int_0^{S^*} y_j(s) ds - \eta(X_k(\infty) - X_k(t^*))$$

As  $B_\infty - B(t^*) > X_k(\infty) - X_k(0)$  is fulfilled, one obtains

$$B_\infty - B(t^*) > \underbrace{\frac{1}{1+\eta}}_{\delta_j} \int_0^{S^*} y_j(s) ds \quad (7)$$

*Step 2:*  $S(t) \in [\tilde{S}, S_0]$

In a similar way than previously, we write

$$\dot{B} = -y_j(S)\dot{S} + \left(1 - \frac{y_j(S)}{y_k(S)}\right) \dot{X}_k$$

and define the number

$$\gamma = \inf_{s \geq \tilde{S}} \left(1 - \frac{y_j(s)}{y_k(s)}\right)$$

which is positive by hypothesis (ii)-(b). One has then

$$B(\tilde{t}) - B_0 \geq \int_{\tilde{S}}^{S_0} y_j(s) ds + \gamma(X_k(\tilde{t}) - X_k(0))$$

Consider now the proportion variable  $p_k = X_k/B$  with  $p_k(0) = p_{0,k} > 0$ . A straightforward computation, from equations (1), gives

$$\dot{p}_k = (\mu_k(S(t)) - \mu_j(S(t)))p_k(1 - p_k)$$

From hypothesis (ii)-(b), we conclude that  $p_k(\cdot)$  is a non decreasing function on  $[0, \tilde{t}]$ . Therefore, one has

$$X_k(\tilde{t}) - X_k(0) \geq p_{0,k}(B(\tilde{t}) - B_0)$$

which yields

$$B(\tilde{t}) - B_0 \geq \underbrace{\frac{1}{1 - \gamma p_{0,k}}}_{\delta_k} \int_{\tilde{S}}^{S_0} y_j(s) ds \quad (8)$$

*Step 3:*  $S(t) \in [S^*, \tilde{S}]$ .

From hypothesis (ii)-(a), one has  $y_k(S(t)) > y_j(S(t))$  for any  $t \in [\tilde{t}, t^*)$ , which allows to write

$$\dot{S}(t) = -\frac{\dot{X}_k(t)}{y_k(S(t))} - \frac{\dot{X}_j(t)}{y_j(S(t))} > -\frac{\dot{B}(t)}{y_j(S(t))}, \quad t \in [\tilde{t}, t^*)$$

Integrating between  $t = \tilde{t}$  and  $t = t^*$ , one obtains

$$B(t^*) - B(\tilde{t}) > \int_{S^*}^{\tilde{S}} y_j(s) ds \quad (9)$$

Finally, gathering inequalities (7)-(8)-(9), we get the inequality

$$B_\infty - B_0 > \delta_k \int_{\tilde{S}}^{S_0} y_j(s) ds + \int_{S^*}^{\tilde{S}} y_j(s) ds + \delta_j \int_0^{S^*} y_j(s) ds$$

that we write as follows

$$B_\infty - B_0 > R(\sigma_j) + \underbrace{(\delta_k - 1) \int_{\tilde{S}}^{S_0} y_j(s) ds - (1 - \delta_j) \int_0^{S^*} y_j(s) ds}_{\Delta}$$

Notice that  $\delta_j$  is smaller than one while  $\delta_k$  is larger than 1. Under hypothesis (i), the term  $\Delta$  is then positive when  $S_0$  is large enough, which implies that the biomass production is larger than  $R(\sigma_j)$ , and thus an overyielding.

The condition (ii) of Proposition 11 requires a complementarity of species  $j$  and  $k$  in terms of yield, depending on the level of resource. Notice that differently to the constant yield case (Proposition 8), a condition on the difference between the growth functions is also required here.

## 6 Summary and comparison

We summarize On Fig. 1 the results obtained in Sections 3, 4 and 5.

	<i>classical model</i>	<i>threshold model</i>	<i>variable yield model</i>
growth	any without threshold on $S$	threshold on $S$	function of $S$ without threshold
yield	constant	constant	variable
	no overyielding	overyielding when the best species does not have the highest yield	overyielding when the best species does not have for large $S$ the highest yield and best growth rate

Fig. 1. Overyielding conditions

One can see, as already underlined in Remark 7, that under constant yields, it is not possible to have overyielding whatever are the growth functions and their dependency, but on the single condition that growth functions are non null when  $S > 0$ . Conversely, threshold is not sufficient to have overyielding: a complementary between species with both high yield and high threshold versus lower yield and lower threshold is required. Notice that here also no particular condition on the kinetics, apart the existence of thresholds, is required. On the contrary, the consideration of variable yield requires more sophisticated complementarity conditions involving kinetics conditions for large values of  $S$ . Although different thresholds or different variable yields both reflect complementary between species, the underlying mechanisms in the transient are quite different.

## 7 Numerical simulations and discussions

In this section, we illustrate the results of Propositions 8 and 11 on numerical simulations with growth functions of the literature. For simplicity and ease of results interpretation, we show examples of overyielding with two species only. Of course, more sophisticated examples with more species can be provided.

### 7.1 Case of constant yields

According to Proposition 8, we consider non-null growth thresholds. Take for instance two species with growth function of Moser type [21] with threshold:

$$\mu_i(S) = \begin{cases} 0, & S < \underline{S}_i \\ \frac{\mu_{max,i} S^{\alpha_i}}{K^{\alpha_i} + S^{\alpha_i}}, & S \geq \underline{S}_i \quad (\alpha_i > 1) \end{cases}$$

Other choices of growth functions are possible. The Moser expression possesses the advantage to be smooth at  $S = \underline{S}_i$ . Parameter values of the two species are given in Table 7.1, while the graphs of their growth functions are plotted in Fig. 2.

	$Y_i$	$\underline{S}_i$	$\mu_{max,i}$	$K_i$	$\alpha_i$
$i = 1$	1	0.5	0.3	0.7	2
$i = 2$	1.4	1	0.6	0.5	2

Table 1

Parameter values of the two species.




Fig. 2. Graphs of the two growth rate functions.

Consider now initial conditions with  $B_0 = 0.01$  and  $S_0 = 2$ . One can check that  $R(\sigma_1) = Y_1 \max(0, S_0 - \underline{S}_1) = 1.5 > R(\sigma_2) = Y_2 \max(0, S_0 - \underline{S}_2) = 1.4$ . Therefore, species 1 is the best one when alone. However, one has  $Y_2 > Y_1$  and according to Proposition 8, any mixture of both species brings a better final biomass production. This is verified numerically, as illustrated in Table 7.1 and Fig. 3, with for instance the initial proportion  $p_0 = 0.4$ .

$p_0$	$X_1(\infty)$	$X_2(\infty)$	$S(\infty)$	$B(\infty)$
1	1.5	0	0.5	1.5
0	0	1.4	1.5	1.4
0.4	$\simeq 0.664$	$\simeq 1.169$	0.5	$\simeq 1.832$

Table 2

Comparison of final states between pure and mixed cultures.

In Table 7.1, one can observe that one has  $S(\infty) = \underline{S}_i$ ,  $B(\infty) = R(\sigma_i)$  ( $i = 1, 2$ ) as expected from Lemma 4 and formula (3). For the mixed culture, one has  $S(\infty) = \min(\underline{S}_1, \underline{S}_2)$ , consistently with Lemma 4. As predicted by Proposition 8, the mixed culture presents an overyielding with an increase of the production about 21.7% compared to the best species alone. This example shows that even with species of similar performances in pure culture, the mixed culture could bring a significantly better biomass production. In Fig. 3, one can see that for the particular choice of the growth rate functions  $\mu_i$ , the second species grows the fastest in mixed culture, as long as the level of substrate is above its growth threshold  $\underline{S}_2$ , in a similar way of what it does in pure culture. On the contrary, the first species is somewhat penalized in the competition and increases less quickly than if it were alone. This is due to the fact that the growth rate of the second species is the largest for large values of substrate (see Fig. 2). But, differently to the second species, the first species keeps growing when  $S$  goes below  $\underline{S}_2$ , and as a consequence, its stationary phase is reached later. As noted in the last remark of section 3, it is worth to be underlined that the final biomass in pure culture does not depend on the growth rate






Fig. 3. From left to right: pure culture of species 1, pure culture of species 2 and mixed culture with both species.

function (the growth rate impacts the kinetics but not the yield). In mixed culture, although overyielding is systematic, the interplay between the growth rates impacts the final gain. Its mathematical analysis appears to be quite complex, because of richness of the nonlinear nature of the growth functions.

## 7.2 Case of variable yields

We have considered here a first species with a Monod law and a decreasing yield against a second species with a Moser law and a constant yield. Decreasing variable yields are typically met in alcohol fermentation [11, 13]. The expressions of the growth and yield functions are given in Table 7.2 and their graphs are depicted in Fig. 4.

	$\mu_i(S)$	$y_i(S)$
$i = 1$	$\frac{0.5 S}{0.1 + S}$	$1.8 e^{-0.2 S}$
$i = 2$	$\frac{0.8 S^4}{625 + S^4}$	0.8

Table 3

The growth and yield functions considered for the simulations.




Fig. 4. Graphs of the growth rate functions (on the left) and yield functions (right).

One can see that hypothesis (ii)-(a) of Proposition 11 is verified with  $S^* = (\log(1.8) - \log(0.8))/0.2 \simeq 4.055$ . Moreover the graphs of the growth functions cross at  $\tilde{S}$ , which is the largest root of  $0.6S^4 + 0.16S^3 - 625 = 0$ , found numerically to be  $\tilde{S} \simeq 5.616$ , which is larger than  $S^*$ . Hypothesis (ii)-(b) of Proposition 11 is then fulfilled.

We have considered initial conditions with  $B_0 = 1$  and  $S_0 = 10$  (larger than  $\tilde{S}$ ). Table 7.2 gives the final values for pure and mixed cultures. One can observe that species 1 has the best production when alone, and that hypothesis (i) of Proposition 11 is fulfilled. These figures show that although the species

$p_0$	$X_1(\infty)$	$X_2(\infty)$	$S(\infty)$	$B(\infty)$
1	$\simeq 8.782$	0	0	$\simeq 8.782$
0	0	$\simeq 8.599$	0	$\simeq 8.599$
0.3	$\simeq 5.443$	$\simeq 4.490$	0	$\simeq 9.933$

Table 4

Comparison of final states between pure and mixed cultures.

exhibit similar performances in pure culture (the advantage of the production of species 1 over species 2 is quite slight), an overyielding occurs in mixed culture, with a gain approximately equal to 13.1%. The trajectories depicted in Fig. 5 provide some additional information about the transients. In accordance with Lemma 4, there is no residual substrate over infinite horizon. However, one can see that species 2 alone reaches much slower the stationary phase than species 1. This is due to its low kinetics for small value of  $S$ . In mixed culture, species 1 contributes to a faster consumption of the substrate. The complementarity between species is illustrated here with the species 2 having a better conversion rate for large value of  $S$  (that is at the beginning of the batch process), while species 1 takes over species 2 when the resource  $S$  takes smaller values. This feature seems comparable to the simulations with constant yields. However, the interplay between the species dynamics is here much more complex, as the initial value  $S_0$  has to be large enough (see Proposition 11).




Fig. 5. From left to right: pure culture of species 1, pure culture of species 2 and mixed culture with both species.

## 8 Conclusion

In this work, we have investigated mechanisms that could explain a gain in biomass production with mixed cultures of micro-organisms, compared to pure cultures, that we defined as “overyielding”. Our main message is that the nature of the non-linearities of the growth characteristics (specific growth rate functions and yields) is enough to produce overyielding. In particular, there is no need to have direct interaction between species to explain overyielding. It was first shown that overyielding is not possible with the classical batch models, such as the Monod model with multi-species. To play with complementarity between species, we have to face some additional features. We have exhibited two kinds of situations, with features reported in the literature :

- (1) different growth thresholds and (constant) yields (see Proposition 8),
- (2) variable yields coupled with conditions on the growth functions (see Proposition 11),

that have been illustrated with numerical simulations. The underlying mechanism is based on the utilization of the resources during the transients, with a succession of stages where each species contributes to its best. Of course, those mechanisms can be added to other ones to explain overyielding. The analysis we performed here allowed to check a potential overyielding from the single

knowledge of growth characteristics in pure culture, which can then guide the choices of experiments among the combinatorial of possible mixtures.

## References

- [1] P. Agrawal, C. Lee, H. Lim, and D. Ramkrishna. Theoretical investigations of dynamic behavior of isothermal continuous stirred tank biological reactors. *Chemical Engineering Science*, 37(3):453–462, 1982.
- [2] J. Arino, S. Pilyugin, and G. Wolkowicz. Considerations on yield, nutrient uptake, cellular growth, and competition in chemostat models. *Canadian Applied Mathematics Quarterly*, 11(2):107–142, 2003.
- [3] H. Beeftink, R. van der Heijden, and J. Heijnen. Maintenance requirements: energy supply from simultaneous endogenous respiration and substrate consumption. *FEMS Microbiology Letters*, 73(3):203 – 209, 1990.
- [4] W. Beyeler, P. L. Rogers, and A. Fiechter. A simple technique for the direct determination of maintenance energy coefficient: An example with *Zygomonas mobilis*. *Applied Microbiology and Biotechnology*, 19(4):277–280, 1984.
- [5] Y. F. Cheng, J. E. Edwards, G. G. Allison, W.-Y. Zhu, and M. K. Theodorou. Diversity and activity of enriched ruminal cultures of anaerobic fungi and methanogens grown together on lignocellulose in consecutive batch culture. *Bioresource Technology*, 100(20):4821–4828, 2009.
- [6] A. Cinar, G. Birol, S. Parulekar, and C. Undey. *Batch Fermentation: Modeling, Monitoring, and Control*. Taylor & Francis, CRC Press, 2003.
- [7] P. Crooke, C. Wei, and R. Tanner. The effect of the specific growth rate and yield expressions on the existence of oscillatory behavior of a continuous fermentation model. *Chem. Eng. Commun.*, 6:333–347, 1980.
- [8] R. David, D. Dochain, J.-R. Mouret, A. Vande Wouwer, and J.-M. Sablayrolles. Nitrogen-backed modeling of wine-making in standard and nitrogen-added fermentations. *Bioprocess and Biosystems Engineering*, 37(1):5–16, 2014.
- [9] D. Dochain, P. De Leenheer, and A. Rapaport. About Transgressive Over-Yielding in the Chemostat. In *MATHMOD 2012 - 7th Vienna International Conference on Mathematical Modelling*, pages 653–658. IFAC Proceedings Volumes 45 (2), 2012.
- [10] A. Dorofeev, M. Glagolev, T. Bondarenko, and N. Panikov. Observation and explanation of the unusual growth kinetics of *Arthrobacter globiformis*. *Microbiology*, 24:24–31, 1992.
- [11] S. El Aida. Modélisation mathématique et exploration numérique de la fermentation en œnologie. MSc Thesis in mathematics, Université de Pau et des Pays de l’Adour, France, 2018.
- [12] C. Essajee and R. Tanner. The effect of extracellular variables on the stability of the continuous baker’s yeast-ethanol fermentation process.

- Process Biochemistry*, 14:16–25, 1979.
- [13] G. Goma, R. Moletta, and M. Novak. Comments on the “maintenance coefficient” changes during alcohol fermentation. *Biotechnology Letters*, 1(10):415–420, 1979.
  - [14] J. Grover. Constant- and variable-yield models of population growth: Responses to environmental variability and implications for competition. *Journal of Theoretical Biology*, 158(4):409–428, 1992.
  - [15] J. Harmand, C. Lobry, A. Rapaport, and T. Sari. *The Chemostat: Mathematical Theory of Microorganism Cultures*, volume 1 of *Chemical Engineering series / Chemostat and bioprocesses set*. Wiley, 2017.
  - [16] J. J. Heijnen and J. A. Roels. A macroscopic model describing yield and maintenance relationships in aerobic fermentation processes. *Biotechnology and Bioengineering*, 23(4):739–763, 1981.
  - [17] Y. Lenbury, P. Crooke, and R. Tanner. Relating damped oscillations to sustained limit cycles describing real and ideal batch fermentation processes. *BioSystems*, 19:15–22, 1986.
  - [18] Y. Lenbury, J. J. Stepan, D.-H. Park, and R. D. Tanner. Modelling oscillatory behavior in batch fermentations. *Acta Biotechnologica*, 6(1):45–53, 1986.
  - [19] H. C. Lim and H. S. Shin. *Fed-Batch Cultures: Principles and Applications of Semi-Batch Bioreactors*. Cambridge Series in Chemical Engineering. Cambridge University Press, 2013.
  - [20] R. McKellar and X. Lu. *Modeling Microbial Responses in Food*. Taylor & Francis, CRC Press, 2003.
  - [21] H. Moser. *The Dynamics of Bacterial Populations Maintained in the Chemostat*. Carnegie Institution of Washington Publication, 1958.
  - [22] J. Mounier, C. Monnet, T. Vallaey, R. Arditi, A.-S. Sarthou, A. Hélias, and F. Irlinger. Microbial interactions within a cheese microbial community. *Applied and Environmental Microbiology*, 74(1):172–181, 2008.
  - [23] S. Pilyugin and P. Waltman. Multiple limit cycles in the chemostat with variable yield. *Math. Biosciences*, 182:151–166, 2003.
  - [24] S. J. Pirt. The maintenance energy of bacteria in growing cultures. *Proceedings of the Royal Society of London. Series B. Biological Sciences*, 163(991):224–231, 1965.
  - [25] S. J. Pirt. *Principles of microbe and cell cultivation*. Blackwell Scientific Publications., Oxford, UK, 1975.
  - [26] T. Satyanarayana and G. Kunze. *Yeast Biotechnology: Diversity and Applications*. Springer Netherlands, 2009.
  - [27] B. Schmid, A. Hector, P. Saha, and M. Loreau. Biodiversity effects and transgressive overyielding. *Journal of Plant Ecology*, 1(2):95–102, 2008.
  - [28] H. L. Smith and P. Waltman. *The Theory of the Chemostat: Dynamics of Microbial Competition*. Cambridge Studies in Mathematical Biology. Cambridge University Press, 1995.
  - [29] M. Tros, T. Bosma, G. Schraa, and A. Zehnder. Measurement of minimum substrate concentration ( $s_{min}$ ) in a recycling fermentor and its prediction

- from the kinetic parameters of pseudomonas strain b13 from batch and chemostat cultures. *Appl Environ Microbiol.*, 62(10):3655–61, 1996.
- [30] P. van Bodegom. Microbial maintenance: A critical review on its quantification. *Microbial Ecology*, 53(4):513–523, 2007.
- [31] N. van Uden. Transport-limited fermentation and growth of *saccharomyces cerevisiae* and its competitive inhibition. *Archiv für Mikrobiologie*, 58(2):155–168, 1967.
- [32] N. van Uden. Transport-limited growth in the chemostat and its competitive inhibition; a theoretical treatment. *Archiv für Mikrobiologie*, 58(2):145–154, 1967.
- [33] J. VanBriesen. Evaluation of methods to predict bacterial yield using thermodynamics. *Biodegradation*, 13:171–190, 2002.
- [34] G. Wang and W. M. Post. A theoretical reassessment of microbial maintenance and implications for microbial ecology modeling. *FEMS Microbiology Ecology*, 81(3):610–617, 2012.