



HAL
open science

A few Multiresolution Schemes for the Black-Scholes equation

Philippe Poncet, Remi Abgrall

► **To cite this version:**

Philippe Poncet, Remi Abgrall. A few Multiresolution Schemes for the Black-Scholes equation. CEM-RACS 2000, Jul 2000, Marseille, France. hal-02011193

HAL Id: hal-02011193

<https://hal.science/hal-02011193>

Submitted on 7 Feb 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A few Multiresolution Schemes for the Black-Scholes equation

Philippe Poncet* Rémi Abgrall†

January 29, 2002

Abstract

In this note, we apply multiresolution techniques to a discretisation of the linear Black-Scholes equation, which is a convection diffusion equation. In particular, we describe a few numerical schemes in order to perform one step of decimation and reconstruction. This suggests a new discretisation method of this equation, which should easily extend to more than one dimension, and lead to very efficient numerical schemes.

1 Introduction

We are interested in the numerical simulation of the following equation :

$$\frac{\partial u}{\partial \tilde{t}} + \frac{1}{2}\sigma^2 S^2 \frac{\partial^2 u}{\partial S^2} = r \left(u - S \frac{\partial u}{\partial S} \right) \quad S \in \mathbb{R}^+ \text{ and } \tilde{t} \in [0, T]$$

$$u(S, T) = [S - K]^+ \quad S \in \mathbb{R}^+ \text{ and } \tilde{t} = T$$

where $S(\tilde{t})$ is a market price, $u(S, \tilde{t})$ is the option premium for a market price S at time \tilde{t} , and K is the price “at-the-money”.

By means of the changes of variable $x = \log S$ and $t = T - \tilde{t}$, the last equation is transformed into

$$\frac{\partial u}{\partial t} - v \frac{\partial u}{\partial x} - \frac{\sigma^2}{2} \frac{\partial^2 u}{\partial x^2} + ru = 0 \quad x \in \mathbb{R} \text{ and } t \in [0, T]$$

$$u(x, 0) = [e^x - K]^+ \quad x \in \mathbb{R} \text{ and } t = 0$$

which is a convection diffusion equation submitted to an initial condition with $v = r - \frac{\sigma^2}{2}$.

Since $\sigma^2 > 0$, the singularity of the initial condition is immediately smoothed, and the solution belongs to $C^\infty(\mathbb{R}) \times \mathbb{R}^+$. Thus one can reasonably expect that very few information is needed to represent the solution.

*Laboratoire de Modélisation et de Calcul (LMC-IMAG), Université Grenoble I, BP 53, Grenoble cedex 9, France

†Mathématiques Appliquées de Bordeaux, Université Bordeaux I, 351 Cours de la Libération, 33 405 Talence cedex, France

The x range must be reduced from \mathbb{R} to a window $[a, b]$, hence the problem with boundary conditions :

$$\left\{ \begin{array}{l} \frac{\partial u}{\partial t} - v \frac{\partial u}{\partial x} - \frac{\sigma^2}{2} \frac{\partial^2 u}{\partial x^2} + ru = 0 \quad (x, t) \in [a, b] \times [0, T] \\ u(a, t) = 0 \quad \text{and} \quad e^{-b} \frac{\partial u}{\partial x}(b, t) = 1 \quad \text{for } t \in [0, T] \\ u(x, 0) = [e^x - K]^+ \quad \text{for } t = 0 \end{array} \right. \quad (1)$$

Several numerical schemes are considered : explicit schemes, both in physical space (equation 3) and in coefficient space with complet multiresolution (equation 4), and implicit schemes (equation 6 in physical space and equation 7 in coefficient space with one step of multiresolution). Most of these different strategies for solving the linear Black-Sholes equation have been developed during the summer program of CEMRACS 2000 (cf. [4]).

2 Multiresolution analysis

2.1 Decimation and reconstruction

The material of this section is taken from [6, 7]. In [3], this technique was extended to the representation of function to unstructured meshes, and applied to fluid mechanics problems in [1, 2].

One considers a set of embedded meshes

$$\mathcal{U}_R \subset \dots \subset \mathcal{U}_{L-1} \subset \mathcal{U}_L$$

where \mathcal{U}_L is the finest mesh, *i.e.* the physical space, with N_L points, and \mathcal{U}_R the roughest mesh considered.

To simplify the text, we assume that $N_L = 2^L + 1$ and

$$\#\mathcal{U}_j = 2^j + 1 \quad \text{for all } R \leq j \leq L$$

but we can handle more general cases (as explained in [7] or [3]), where any value of N_L is admissible.

The loss of information from a mesh to another is completed so that the initial $2^L + 1$ datas can always be recovered : let \mathcal{D}_j be these completions. They are defined as sets of 2^j elements¹.

Hence a natural definition of global meshes \mathcal{M}_j that contain $2^L + 1$ informations :

$$\mathcal{M}_j = \mathcal{U}_j \oplus \bigoplus_{k=j}^{L-1} \mathcal{D}_k$$

One step of *decimation*, *i.e.* the transit from a mesh \mathcal{U}_j to a mesh $\mathcal{U}_{j-1} \oplus \mathcal{D}_{j-1}$, is given by

$$\left\{ \begin{array}{l} w_p^{j-1} = u_{2p}^j \quad \text{for } p = 0 \dots 2^{j-1} \\ d_p^{j-1} = u_{2p+1}^j - \frac{u_{2p}^j + u_{2p+2}^j}{2} \quad \text{for } p = 0 \dots 2^{j-1} - 1 \quad (\text{called } \textit{details}) \end{array} \right.$$

¹Indeed, 2^j datas are lost when we reduce the mesh from \mathcal{U}_{j+1} to \mathcal{U}_j .




Fig. 1 : One step of decimation and reconstruction.

while *reconstruction* is given by

$$\begin{cases} u_{2^p}^j = u_p^{j-1} & \text{for } p = 0 \dots 2^{j-1} \\ u_{2^{p+1}}^j = \frac{u_p^{j-1} + u_{p+1}^{j-1}}{2} + d_p^{j-1} & \text{for } p = 0 \dots 2^{j-1} - 1 \end{cases}$$

Let S_j be the decimation matrix sample at level j :

$$S_j = \left[\begin{array}{cccc|ccc} 1 & 0 & \dots & & 0 & \dots & \\ \frac{1}{2} & \frac{1}{2} & 0 & \dots & 1 & 0 & \dots \\ 0 & 1 & 0 & \dots & 0 & \dots & \\ 0 & \frac{1}{2} & \frac{1}{2} & 0 & 0 & 1 & 0 \\ & & \ddots & \ddots & & \ddots & \ddots \\ & & & 0 & 1 & 0 & \dots & 0 \\ & & & & \frac{1}{2} & \frac{1}{2} & & \\ & & & & 0 & 1 & & \\ & & & & & & & 0 \end{array} \right] \in \mathbb{M}_{(2^j+1) \times (2^j+1)}$$

and reconstruction sample

$$S_j^{-1} = \left[\begin{array}{cccc|ccc} 1 & 0 & \dots & & & & \\ 0 & 0 & 1 & \dots & & & \\ & & & \ddots & & & \\ & & & & 1 & 0 & 0 \\ & & & & & 0 & 1 \\ \hline -\frac{1}{2} & 1 & -\frac{1}{2} & \dots & & & \\ & & -\frac{1}{2} & 1 & -\frac{1}{2} & & \\ & & & \ddots & & & \\ & & & & -\frac{1}{2} & 1 & -\frac{1}{2} \end{array} \right] \in \mathbb{M}_{(2^j+1) \times (2^j+1)}$$

Hence it is possible to define the $(2^L + 1) \times (2^L + 1)$ decimation operators by :

$$\left\{ \begin{array}{l} M_{j-1} = \left[\begin{array}{c|c} S_j & 0 \\ \hline 0 & Id_{2^L-2^{L-j}} \end{array} \right] \times M_j \\ M_{L-1} = S_L \quad \text{and} \quad M_L = Id_{2^L+1} \end{array} \right.$$

and the full reconstruction operators :

$$\left\{ \begin{array}{l} M_{j-1}^{-1} = M_j^{-1} \times \left[\begin{array}{c|c} S_j^{-1} & 0 \\ \hline 0 & Id_{2^L-2^{L-j}} \end{array} \right] \\ M_{L-1}^{-1} = S_L^{-1} \quad \text{and} \quad M_L^{-1} = Id_{2^L+1} \end{array} \right.$$

M_{L-j} is the operator that makes j steps of decimation, from mesh \mathcal{M}_L to \mathcal{M}_{L-j} . Moreover, M_R is the operator of full decimation, from the physical space \mathcal{M}_L to coefficient space \mathcal{M}_R .

2.2 An explicit scheme

This kind of scheme has very strong CFL conditions. Nevertheless it is interesting to study because of its simplicity. The explicit scheme related to equation (1) can be written

$$\left\{ \begin{array}{l} \frac{U_i^{n+1} - U_i^n}{\Delta t} - \frac{F_{i+1/2}^n - F_{i-1/2}^n}{\Delta x} - \frac{\sigma^2 U_{i+1}^n - 2U_i^n + U_{i-1}^n}{2\Delta x^2} + rU_i = 0 \\ \text{with } F_{i+1/2}^n = \frac{1}{2}(vU_i^n + vU_{i+1}^n + |v|(U_{i+1}^n - U_i^n)) \\ U_0^{n+1} = 0 \quad \text{and} \quad U_{2^L}^{n+1} = U_{2^L-1}^{n+1} + \Delta x e^{x_{max}} \\ U_i^0 = [e^{x_{min}+i\Delta x} - K]^+ \end{array} \right. \quad (2)$$

Let be

$$\left\{ \begin{array}{l} \alpha = \frac{\Delta t}{\Delta x} |v| \\ \mu = \frac{\Delta t}{\Delta x^2} \frac{\sigma^2}{2} \\ \beta = 1 - r\Delta t \end{array} \right.$$

If $v > 0$ (most of the time) the scheme (2) can be written

$$U^{n+1} = A \times U^n + V \quad (3)$$

with

$$A = \begin{bmatrix} 0 & 0 & \dots & 0 \\ \mu & \beta - \alpha - 2\mu & \mu + \alpha & \\ & \mu & \beta - \alpha - 2\mu & \mu + \alpha \\ & & \ddots & \ddots & \ddots \\ & & & \mu & \beta - \alpha - 2\mu & \mu + \alpha \\ & & & \mu & \beta - \alpha - 2\mu & \mu + \alpha \end{bmatrix} \quad \text{and} \quad V = \Delta x e^{x_{max}} \begin{bmatrix} 0 \\ 0 \\ 0 \\ \vdots \\ 0 \\ 1 \end{bmatrix}$$

This is the explicit scheme in physical space.

It is possible to write this scheme in the coefficient space, using the decimation operator. Let \mathcal{M}_R be a minimal mesh². Then M_R is the full decimation operator :

$$\begin{array}{ccc} U^n & \xrightarrow{A U + V} & U^{n+1} \\ \uparrow M_R^{-1} & & \downarrow M_R \\ D^n & \xrightarrow{B D + W} & D^{n+1} \end{array}$$

The scheme written in the coefficient space is then

$$D^{n+1} = B \times D^n + W \quad (4)$$

with $B = M_R \times A \times M_R^{-1}$ and $W = M_R \times V$. The main drawbacks of this explicit scheme are, on the one hand, a lot of time steps, and on the other hand a lot of matrix multiplications.

2.3 Compression

Let \mathcal{M}_R be the minimal admissible mesh and ϵ a tolerance.

At any time, it is possible to truncate the solution in coefficient space : given a set of details $\{d_k^j\} = \mathcal{D}_j$ at a level j , we define

$$\tilde{d}_k^j = \begin{cases} d_k^j & \text{if } |d_k^j| > \epsilon_j \\ 0 & \text{otherwise} \end{cases}$$

with $\epsilon_j = \frac{\epsilon}{2^{j-R}}$.

Then the reconstructed solution after truncation $\tilde{U}^n = M_R \times \tilde{D}^n$ satisfies

$$\|\tilde{U}^n - U^n\|_\infty \leq \epsilon$$

Since $U^n \in \mathbb{R}^{2^L+1}$, we need $2^L + 1$ informations to represent U^n and D^n . After truncation, less information is needed within a tolerance ϵ . This amount of information is quantified by the *compression*

²The mesh \mathcal{M}_0 contains only two points, so it can be useful to consider a cap in decimation.

factor

$$\Lambda = \frac{2^L}{2^{2R} + \sum_{j=R}^{L-1} \#\left\{d_k^j \text{ such as } |d_k^j| > \epsilon_j, k = 0 \dots 2^j - 1\right\}}$$

3 Implicit Scheme

3.1 A possible implicitation

One of the possible implicitations of the scheme (2) is

$$\left\{ \begin{array}{l} \frac{U_i^{n+1} - U_i^n}{\Delta t} - \frac{F_{i+1/2}^n - F_{i-1/2}^n}{\Delta x} - \frac{\sigma^2 U_{i+1}^{n+1} - 2U_i^{n+1} + U_{i-1}^{n+1}}{2\Delta x^2} + rU_i = 0 \\ \text{with } F_{i+1/2}^n = \frac{1}{2}(vU_i^n + vU_{i+1}^n + |v|(U_{i+1}^n - U_i^n)) \\ U_0^{n+1} = 0 \quad \text{and} \quad U_{2^L}^{n+1} = U_{2^L-1}^{n+1} + \Delta x e^{x_{max}} \\ U_i^0 = [e^{x_{min} + i\Delta x} - K]^+ \end{array} \right. \quad (5)$$

This scheme can be written

$$AU^{n+1} = BU^n + V \quad (6)$$

as

$$\begin{bmatrix} 1 & 0 & \dots \\ -\mu & 1+2\mu & -\mu \\ & -\mu & 1+2\mu & -\mu \\ & & \ddots & \ddots & \ddots \\ & & & -\mu & 1+2\mu & -\mu \\ & & & & 0 & -1 & 1 \end{bmatrix} U^{n+1} = \begin{bmatrix} 0 & 0 & \dots \\ 0 & \beta - \alpha & \alpha \\ 0 & 0 & \beta - \alpha & \alpha \\ & & \ddots & \ddots \\ & & & \beta - \alpha & \alpha \\ & & & & \dots & 0 & 0 \end{bmatrix} U^n + \begin{bmatrix} 0 \\ 0 \\ 0 \\ \vdots \\ 0 \\ \Delta x e^{x_{max}} \end{bmatrix}$$

3.2 One step of multiresolution

Unlike the explicit scheme, we do not need to express the equation in the coefficient space. If we express the equation as a vector with firstly even nodes and then odds, the right member (flux and sources) can be written³

³First and last nodes are even, respectively 0 and 2^L . This vector P^n can be computed very fast.

$$P^n = \begin{bmatrix} 0 & & & & & & & & & & 0 \\ 0 & 0 & \beta - \alpha & \alpha & & & & & & & \\ & & & \ddots & \ddots & & & & & & \\ & & & & \ddots & \ddots & & & & & \\ & & & & & \beta - \alpha & \alpha & 0 & 0 & & \\ \hline 0 & & & & & & & \beta - \alpha & \alpha & & \\ \hline 0 & \beta - \alpha & \alpha & 0 & \dots & & & & & & \\ 0 & 0 & 0 & \beta - \alpha & \alpha & & & & & & \\ & & & & \ddots & \ddots & & & & & \\ & & & & & 0 & 0 & \beta - \alpha & \alpha & 0 & 0 \end{bmatrix} U^n + \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ \frac{e^{x_m \alpha x} \Delta x}{0} \\ 0 \\ \vdots \\ 0 \end{bmatrix} \begin{array}{l} \leftarrow \text{first node, left boundary} \\ \\ \leftarrow \text{even nodes} \\ \\ \leftarrow \text{last node, right boundary} \\ \\ \leftarrow \text{odd nodes} \end{array}$$

The implicit scheme can be written as follows

$$(A \times S_L^{-1}) D^{n+1} = B U^n + V$$

which means, in the base “even-then-odd” (*i.e.* after permutation)

$$C \times D^{n+1} = P^n \tag{7}$$

with

$$C = \left[\begin{array}{cccccc|cccccc} 1 & 0 & 0 & & & 0 & 0 & 0 & \dots & \dots & 0 \\ -\mu/2 & 1 + \mu & -\mu/2 & 0 & & 0 & -\mu & -\mu & 0 & \dots & 0 \\ 0 & -\mu/2 & 1 + \mu & -\mu/2 & 0 & \vdots & 0 & -\mu & -\mu & 0 & \vdots \\ \vdots & \ddots & \ddots & \ddots & \ddots & 0 & \vdots & \ddots & \ddots & \ddots & 0 \\ \vdots & & & & -\mu/2 & 1 + \mu & -\mu/2 & & & -\mu & -\mu \\ \hline 0 & \dots & \dots & 0 & -1/2 & 1/2 & 0 & \dots & 0 & \dots & -1 \\ \hline 1/2 & 1/2 & 0 & & & 0 & 1 + 2\mu & 0 & 0 & \dots & 0 \\ 0 & 1/2 & 1/2 & 0 & & 0 & 0 & 1 + 2\mu & 0 & \dots & 0 \\ \vdots & \ddots & \ddots & \ddots & \ddots & 0 & \vdots & 0 & \ddots & 0 & \vdots \\ \vdots & & & 1/2 & 1/2 & \vdots & \vdots & & & & \vdots \\ 0 & \dots & \dots & 0 & 1/2 & 1/2 & 0 & \dots & \dots & 0 & 1 + 2\mu \end{array} \right]$$

whose block sizes are

$$\left[\begin{array}{c|c} (2^{L-1} + 1) \times (2^{L-1} + 1) & (2^{L-1} + 1) \times (2^{L-1}) \\ \hline (2^{L-1}) \times (2^{L-1} + 1) & (2^{L-1}) \times (2^{L-1}) \end{array} \right]$$

3.3 Example

As an example, we compute the premium of the Call “C90 dec 98” on MATIF Notional Bonds, on october 17th 1990, 44 days before exercise. The updated market price was $K^* = 97.98$.

This corresponds to the datas :

Technical datas				Market datas				
L	N_L	S_{min}	S_{max}	σ	r	K	K^*	T
10	1025	88	108	7.5%	7.5%	98.0	97.98	0.12

The exact Black formula gives a premium of 1.00, and the implicit multiresolution scheme gives 1.00 : The numerical interpolated value is

$$u(e^{-rT}K^*, 0) = 1.004$$

with a tick of 0.01.




Fig. 2 : Premium of C90 dec 98 on MATIF Notional Bonds, using one step of multiresolution with the implicit scheme. Points are the evaluations of the exact Black formula.

References

- [1] R. Abgrall. Multiresolution analysis in unstructured meshes : application to CFD. In K. W. Morton and M.J. Baines, editors, *Numerical Methods for Fluid Dynamics*, volume V. Oxford university Press, 1996.
- [2] R. Abgrall. Some preliminary results in multiresolution analysis in unstructured meshes for compressible flows simulations. In Chetverushkin, Désideri, Kuznetsov, Muzafarov, Périaux, and Pironneau, editors, *Experimentation, Modelling and Computation in Flow, Turbulence and Combustion*, volume 2. Wiley, 1997.
- [3] R. Abgrall and A. Harten. Multiresolution Representation in Unstructured Meshes. *Siam J. Numer Anal.*, à paraitre, Dec 1998. Rapport INRIA 2342, Septembre 1994.

- [4] R. Abgrall, P. Poncet, M. Briani and C. Godard. A first step toward Multiresolution Analysis for the Black-Scholes equation. *CEMRACS*, Marseille, France, 2000.
- [5] G. Chiavassa and R. Donat. Numerical experiments with point values multiresolution for 2D compressible flows. *Siam J. Scien. Comp.*, submitted, 2000.
- [6] A. Harten. Discrete multiresolution analysis and generalized wavelets. *J. Appl. Numer. Math.*, 12:153–192, 1993.
- [7] A. Harten. Multiresolution representation of data : a general framework. *SINUM*, 33:1205–1256, 1996.
- [8] B. Sjögreen. Numerical experiments with the multiresolution scheme for compressible Euler solvers. *J. Comp. Phys.*, 117, 1995.