



HAL
open science

Uniform Asymptotic and Convergence Estimates for the Jin–Xin Model Under the Diffusion Scaling

Roberta Bianchini

► **To cite this version:**

Roberta Bianchini. Uniform Asymptotic and Convergence Estimates for the Jin–Xin Model Under the Diffusion Scaling. *SIAM Journal on Mathematical Analysis*, 2018, 50 (2), pp.1877-1899. hal-02012705

HAL Id: hal-02012705

<https://hal.science/hal-02012705>

Submitted on 9 Feb 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **UNIFORM ASYMPTOTIC AND CONVERGENCE ESTIMATES FOR**
2 **THE JIN-XIN MODEL UNDER THE DIFFUSION SCALING***

3 ROBERTA BIANCHINI [†]

4 **Abstract.** We obtain sharp decay estimates in time in the context of Sobolev spaces, for smooth
5 solutions to the one dimensional Jin-Xin model under the diffusion scaling, which are uniform with
6 respect to the singular parameter of the scaling. This provides convergence to the limit nonlinear
7 parabolic equation both for large time, and for the vanishing singular parameter. The analysis is
8 performed by means of two main ingredients. First, a crucial change of variables highlights the
9 dissipative property of the Jin-Xin system, and allows to observe a faster decay of the dissipative
10 variable with respect to the conservative one, which is essential in order to close the estimates. Next,
11 the analysis relies on a deep investigation on the Green function of the linearized Jin-Xin model,
12 depending on the singular parameter, combined with the Duhamel formula in order to handle the
13 nonlinear terms.

14 **Key words.** Relaxation, Green analysis, asymptotic behavior, dissipation, global existence,
15 decay estimates, diffusive scaling, conservative-dissipative form, BGK models.

16 **AMS subject classifications.** 35L40, 35L45, 35K55.

17 **1. Introduction.** We consider the following scaled version of the Jin-Xin ap-
18 proximation for systems of conservation laws in [10]:

19 (1)
$$\begin{cases} \partial_t u + \partial_x v = 0, \\ \varepsilon^2 \partial_t v + \lambda^2 \partial_x u = f(u) - v, \end{cases}$$

20 where $\lambda > 0$ is a positive constant, u, v depend on $(t, x) \in \mathbb{R}^+ \times \mathbb{R}$ and take values in
21 \mathbb{R} , while $f(u) : \mathbb{R} \rightarrow \mathbb{R}$ is a Lipschitz function such that $f(0) = 0$, and $f'(0) = a$, with
22 a a constant value independent of ε, λ . The diffusion limit of this system for $\varepsilon \rightarrow 0$
23 has been studied in [9, 4], where the convergence to the following equations is proved:

24 (2)
$$\begin{cases} \partial_t u + \partial_x v = 0 \\ v = f(u) - \lambda^2 \partial_x u. \end{cases}$$

25 From [17, 4], it is well-known that system (1) can be written in BGK formulation, [3],
26 by means of the linear change of variables:

27 (3)
$$u = f_1^\varepsilon + f_2^\varepsilon, \quad v = \frac{\lambda}{\varepsilon} (f_1^\varepsilon - f_2^\varepsilon).$$

28 Precisely, the BGK form of (1) reads:

29 (4)
$$\begin{cases} \partial_t f_1^\varepsilon + \frac{\lambda}{\varepsilon} \partial_x f_1^\varepsilon = \frac{1}{\varepsilon^2} (M_1(u) - f_1^\varepsilon), \\ \partial_t f_2^\varepsilon - \frac{\lambda}{\varepsilon} \partial_x f_2^\varepsilon = \frac{1}{\varepsilon^2} (M_2(u) - f_2^\varepsilon), \end{cases}$$

*Submitted to the editors October 16th, 2017.

[†]Università degli Studi di Roma "Tor Vergata", via della Ricerca Scientifica 1, I-00133 Rome, Italy - Istituto per le Applicazioni del Calcolo "M. Picone", Consiglio Nazionale delle Ricerche, via dei Taurini 19, I-00185 Rome, Italy, (bianchin@mat.uniroma2.it). Phone: +390649937340. Current address: UMPA, ENS-Lyon 46, allée d'Italie 69364 Lyon Cedex 07, France (roberta.bianchini@ens-lyon.fr)

30 where the so-called Maxwellians are:

$$31 \quad (5) \quad M_1(u) = \frac{u}{2} + \frac{\varepsilon f(u)}{2\lambda}, \quad M_2(u) = \frac{u}{2} - \frac{\varepsilon f(u)}{2\lambda}.$$

32 According to the theory on diffusive limits of the Boltzmann equation and related
 33 BGK models, see [7, 23], we consider fluctuations of the Maxwellian functions as
 34 initial data for the Cauchy problem associated with system (1). Namely, given a
 35 function $\bar{u}_0(x)$, depending on the spatial variable, we assume

$$36 \quad (6) \quad (u(0, x), v(0, x)) = (u_0, v_0) = (\bar{u}_0, f(\bar{u}_0) - \lambda^2 \partial_x \bar{u}_0).$$

37 By using the change of variables (3), the BGK initial data read:

$$38 \quad (7) \quad (f_1^\varepsilon(0, x), f_2^\varepsilon(0, x)) = \left(M_1(\bar{u}_0) - \frac{\varepsilon\lambda}{2} \partial_x \bar{u}_0, M_2(\bar{u}_0) + \frac{\varepsilon\lambda}{2} \partial_x \bar{u}_0 \right),$$

39 where the fluctuations are given by $\pm \frac{\varepsilon\lambda}{2} \partial_x \bar{u}_0$.

System (1) is the parabolic scaled version of the hyperbolic relaxation approximation for systems of conservation laws, the Jin-Xin system, introduced in [10] in 1995. This model has been studied in [18, 6, 10], and the hyperbolic relaxation limit has been investigated. A complete review on hyperbolic conservation laws with relaxation, and a focus on the Jin-Xin system is presented in [16]. By means of the Chapman-Enskog expansion, local attractivity of diffusion waves for the Jin-Xin model was established in [6]. In [13], the authors showed that, under some assumptions on the initial data and the function $f(u)$, the first component of system (1) with $\varepsilon = 1$ decays asymptotically towards the fundamental solution to the Burgers equation, for the case of $f(u) = \alpha u^2/2$. Besides, [20] is a complete study of the long time behavior of this model for a more general class of functions $f(u) = |u|^{q-1}u$, with $q \geq 2$. The method developed in [20] can be also extended to the multidimensional case in space, and provides sharp decay rates. Here we study the parabolic scaled version of the system studied in [20], i.e. (1), and we consider a more general function $f(u) = au + h(u)$, where a is a constant, and $h(u)$ is a polynomial function whose terms are at least quadratic. We point out that only the case $a = 0$ has been handled in [20], and in many previous works as well. In accordance with the theory presented in [1] on partially dissipative hyperbolic systems, we are able to cover also the case $a \neq 0$. Furthermore, besides the asymptotic behavior of the solutions, here we are interested in studying the diffusion limit, for vanishing ε , of the Jin-Xin system, which is the main improvement of the present paper with respect to the results achieved in [1]. Indeed, because of the presence of the singular parameter, we cannot approximate the analysis of the Green function of the linearized problems, as the authors did in [1], and explicit calculations in that context are needed.

The diffusive Jin-Xin system has been already investigated in the following works below. In [9], initial data around a traveling wave were considered, while in [4] the authors write system (1) in terms of a BGK model, and the diffusion limit is studied by using monotonicity properties of the solution. In all these cases, u, v are scalar functions. For simplicity, here we also take scalar unknowns u, v . However, our approach, which takes its roots in [1], can be generalized to the case of vectorial functions $u, v \in \mathbb{R}^N$. As mentioned before, the novelty of the present paper consists in dealing with the singular approximation and, in the meantime, with the the large time asymptotic of system (1), which behaves like the limit parabolic equation (2), without using

monotonicity arguments. We obtain, indeed, sharp decay estimates in time to the solution to system (1) in the Sobolev spaces, which are uniform with respect to the singular parameter. This provides the convergence to the limit nonlinear parabolic equation (2) both asymptotically in time, and in the vanishing ε -limit. To this end, we perform a crucial change of variables that highlights the dissipative property of the Jin-Xin system, and provides a faster decay of the dissipative variable with respect to the conservative one, which allows to close the estimates. Next, a deep investigation on the Green function of the linearized system (1) and the related spectral analysis is provided, since explicit expressions are needed in order to deal with the singular parameter ε . The dissipative property of the diffusive Jin-Xin system, together with the uniform decay estimates discussed above, and the Green function analysis combined with the Duhamel formula provide our main results, stated in the following. We should mention that we intend to adapt the present argument to the BGK approximation to the Navier-Stokes equations considered in [2], in order to extend the local in time convergence of the approximating system to the solutions to the Navier-Stokes equations on the two dimensional torus, proved in [2], to the global in time result. Define

$$E_m = \max\{\|u_0\|_{L^1} + \varepsilon\|v_0 - au_0\|_{L^1}, \|u_0\|_m + \varepsilon\|v_0 - au_0\|_m\},$$

40 where $\|\cdot\|_m$ stands for the $H^m(\mathbb{R})$ Sobolev norm and $H^0(\mathbb{R}) = L^2(\mathbb{R})$. Precise as-
41 sumptions on $f(u)$ in (1) are stated here.

42 ASSUMPTION 1.1. $f(u) : \mathbb{R} \rightarrow \mathbb{R}$ is a Lipschitz function such that:

- 43 • $f(0) = 0$;
- 44 • $f'(0) = a$, which is a constant value independent of ε, λ ;
- 45 • $f(u) = ah + h(u)$, where $h(u)$ is a polynomial functions of order higher than
46 or equal to 2.

THEOREM 1.1. Under Assumptions 1.1, consider the Cauchy problem associated with system (1) and initial data (6). If E_2 is sufficiently small, then the unique solution

$$(u, v) \in C([0, \infty), H^2(\mathbb{R})) \cap C^1([0, \infty), H^1(\mathbb{R})).$$

Moreover, the following decay estimate holds:

$$\|u(t)\|_2 + \varepsilon\|v(t) - au(t)\|_2 \leq C \min\{1, t^{-1/4}\} E_2.$$

47 Now, consider the equation below,

$$48 \quad (8) \quad \partial_t w_p + a\partial_x w_p + \partial_x h(w_p) - \lambda^2 \partial_{xx} w_p = 0.$$

49 We state the following result.

THEOREM 1.2. Under Assumptions 1.1, let w_p be the solution to the nonlinear equation (8) with initial data

$$w_p(0) = u(0) = u_0 \in L^1(\mathbb{R}) \cap H^{\beta+4}(\mathbb{R}),$$

50 where $\beta > 0$, and u_0 in (6) is the initial datum for the Jin-Xin system (1). For any
51 $\mu \in [0, 1/2)$, if E_1 is sufficiently small with respect to $(1/2 - \mu)$, then we have the
52 following decay estimate:

$$53 \quad (9) \quad \|D^\beta(u(t) - w_p(t))\|_0 \leq C\varepsilon \min\{1, t^{-1/4 - \mu - \beta/2}\} E_{|\beta|+4},$$

54 with $C = C(E_{|\beta|+4})$.

55 Once the right scaled variables, i.e. $(u, \varepsilon^2 v)$, expressed at the beginning of Section
 56 2, have been identified, and the so-called *conservative-dissipative* (C-D) form in [1] for
 57 system (1) has been found, our approach essentially relies on the method developed
 58 in [1], with substantial differences listed here.

- 59 • We need an explicit Green function analysis of the linearized system rather
 60 than expansions and approximations, in order to deal with the singular pa-
 61 rameter ε . The analysis performed in first part of Section 4 is as precise as it
 62 is possible.
- 63 • Some estimates in [1] rely on the use of the Shizuta-Kawashima (SK) con-
 64 dition, explained in the following. Consider a linear first order system in
 65 compact form: $\partial_t \mathbf{u} + A \partial_x \mathbf{u} = G \mathbf{u}$. Passing to the Fourier transform, define
 66 $E(i\xi) = G - iA\xi$. The (SK) condition states that, if $\lambda(z)$ is an eigenvalue
 67 of $E(z)$, then $Re(\lambda(i\xi)) \leq -c \frac{|\xi|^2}{1+|\xi|^2}$, for some constant $c > 0$ and for every
 68 $\xi \in \mathbb{R} - \{0\}$. As it can be seen in (57), the eigenvalues associated with the
 69 compact linearized system in (C-D) form (22) of system (1) present different
 70 weights in ε . Thus, we cannot simply apply the (SK) condition to estimate
 71 the remainders in paragraph *Remainders in between* disregarding ε , as the
 72 authors did in [1], since the weights in ε play a key role to deal with the sin-
 73 gular nonlinear term in the Duhamel formula (66). Again, a further analysis
 74 is needed. We are not using the entropy dissipation and the (SK) condition
 75 in order to start the study of the asymptotic behavior on a given global in
 76 time solution, as the authors did in [1], Theorem 2.5. The semilinear nature
 77 of the Jin-Xin system allows us to use the estimates coming from the Green
 78 function analysis, not only to study the asymptotic behavior of the system for
 79 small ε and long times, but also, and first of all, to prove the global existence
 80 of smooth solutions, uniformly in ε , to our singular system. Such a result is
 81 stated in Theorem 1.1.
- 82 • The coupling between the convergence to the limit equation (2) for vanishing
 83 ε and for large time in the last section is the main novelty of the present
 84 paper, and new ideas are needed to get this result.

85 **2. General setting.** First of all, we write system (1) in the following form:

$$86 \quad (10) \quad \begin{cases} \partial_t u_1 + \frac{\partial_x u_2}{\varepsilon^2} = 0, \\ \partial_t u_2 + \lambda^2 \partial_x u_1 = f(u_1) - \frac{u_2}{\varepsilon^2}. \end{cases}$$

87 The unknown variable is $\mathbf{u} = (u_1, u_2) = (u, \varepsilon^2 v)$, in the spirit of the scaled variables
 88 introduced in [2], which are the right scaling to get the conservative-dissipative form
 89 discussed below. Here we write $f(u_1) = au_1 + h(u_1)$, where $a = f'(0)$, and system
 90 (10) reads

$$91 \quad (11) \quad \begin{cases} \partial_t u_1 + \frac{\partial_x u_2}{\varepsilon^2} = 0, \\ \partial_t u_2 + \lambda^2 \partial_x u_1 = au_1 + h(u_1) - \frac{u_2}{\varepsilon^2}. \end{cases}$$

92 Equations (11) can be written in compact form:

$$93 \quad (12) \quad \partial_t \mathbf{u} + A \partial_x \mathbf{u} = -B \mathbf{u} + N(u_1),$$

94 where

$$95 \quad (13) \quad A = \begin{pmatrix} 0 & \frac{1}{\varepsilon^2} \\ \lambda^2 & 0 \end{pmatrix}, \quad -B = \begin{pmatrix} 0 & 0 \\ a & -\frac{1}{\varepsilon^2} \end{pmatrix}, \quad N(u) = \begin{pmatrix} 0 \\ h(u) \end{pmatrix}.$$

96 In particular, $-B\mathbf{u}$ is the linear part of the source term, while $N(u)$ is the remaining
97 nonlinear one, which only depends on the first component of $\mathbf{u} = (u, \varepsilon^2 v)$. Now, we
98 look for a right constant symmetrizer Σ for system (12), which also highlights the
99 dissipative properties of the linear source term. Thus, we find

$$100 \quad (14) \quad \Sigma = \begin{pmatrix} 1 & a\varepsilon^2 \\ a\varepsilon^2 & \lambda^2\varepsilon^2 \end{pmatrix}.$$

101 Taking \mathbf{w} such that

$$\mathbf{u} = \begin{pmatrix} u_1 \\ u_2 \end{pmatrix} = \begin{pmatrix} u \\ \varepsilon^2 v \end{pmatrix} = \Sigma \mathbf{w} = \begin{pmatrix} (\Sigma \mathbf{w})_1 \\ (\Sigma \mathbf{w})_2 \end{pmatrix},$$

$$102 \quad (15) \quad \text{where } \mathbf{w} = \begin{pmatrix} w_1 \\ w_2 \end{pmatrix} = \begin{pmatrix} \frac{\lambda^2 u - a\varepsilon^2 v}{\lambda^2 - a^2\varepsilon^2} \\ \frac{v - au}{\lambda^2 - a^2\varepsilon^2} \end{pmatrix},$$

103 system (12) reads

$$104 \quad (16) \quad \Sigma \partial_t \mathbf{w} + A_1 \partial_x \mathbf{w} = -B_1 \mathbf{w} + N((\Sigma \mathbf{w})_1),$$

105 where

$$106 \quad A_1 = A_1^T = A\Sigma = \begin{pmatrix} a & \lambda^2 \\ \lambda^2 & a\lambda^2\varepsilon^2 \end{pmatrix}, \quad -B_1 = -B\Sigma = \begin{pmatrix} 0 & 0 \\ 0 & a^2\varepsilon^2 - \lambda^2 \end{pmatrix},$$

107

$$108 \quad (17) \quad N((\Sigma \mathbf{w})_1) = \begin{pmatrix} 0 \\ h(w_1 + a\varepsilon^2 w_2) \end{pmatrix}.$$

109 By using the Cauchy inequality we get the following lemma.

110 LEMMA 2.1. *The symmetrizer Σ is definite positive. Precisely*

$$111 \quad (18) \quad \frac{1}{2} \|w_1\|_0^2 + \varepsilon^2 \|w_2\|_0^2 (\lambda^2 - 2a^2\varepsilon^2) \leq (\Sigma \mathbf{w}, \mathbf{w})_0 \leq \|w_1\|_0^2 (1 + a\varepsilon^2) + \|w_2\|_0^2 (a + \lambda^2) \varepsilon^2.$$

112 Notice that from the theory on hyperbolic systems, [14], the Cauchy problem
113 for (16) with initial data \mathbf{w}_0 in $H^m(\mathbb{R})$, $m \geq 2$, has a unique local smooth solution
114 \mathbf{w}^ε for each fixed $\varepsilon > 0$. We denote by T^ε the maximum time of existence of this
115 local solution and, hereafter, we consider the time interval $[0, T^*]$, with $T^* \in [0, T^\varepsilon)$
116 for every ε . In the following, we study the Green function of system (16), and we
117 establish some uniform energy estimates and decay rates of the smooth solution to
118 system (16).

119 **2.1. The conservative-dissipative form.** In this section, we introduce a linear
 120 change of variable, so providing a particular structure for our system, the so-called
 121 *conservative-dissipative form* (C-D), defined in [1]. The (C-D) form allows to identify
 122 a conservative and a dissipative variable for system (1), such that a faster decay of the
 123 dissipative variable, playing a key role in the following, is observed. Thanks to this
 124 change of variables, we are able indeed to handle the case $a \neq 0$ in (11). Hereafter,
 125 (\cdot, \cdot) denotes the standard scalar product in $L^2(\mathbb{R})$, and $\|\cdot\|_m$ is the $H^m(\mathbb{R})$ -norm, for
 126 $m \in \mathbb{N}$, where $H^0(\mathbb{R}) = L^2(\mathbb{R})$.

127 **PROPOSITION 2.2.** *Given the right symmetrizer Σ in (14) for system (12), denot-*
 128 *ing by*

$$129 \quad (19) \quad \tilde{\mathbf{w}} = M\mathbf{u} = \begin{pmatrix} 1 & 0 \\ \frac{-a\varepsilon}{\sqrt{\lambda^2 - a^2\varepsilon^2}} & \frac{1}{\varepsilon\sqrt{\lambda^2 - a^2\varepsilon^2}} \end{pmatrix} \mathbf{u} = \begin{pmatrix} u \\ \frac{\varepsilon(v - au)}{\sqrt{\lambda^2 - a^2\varepsilon^2}} \end{pmatrix},$$

130 *system (12) can be written in (C-D) form defined in [1], i.e.*

$$131 \quad (20) \quad \partial_t \tilde{\mathbf{w}} + \tilde{A} \partial_x \tilde{\mathbf{w}} = -\tilde{B} \tilde{\mathbf{w}} + \tilde{N}(\tilde{w}_1),$$

132 *where*

$$133 \quad (21) \quad \tilde{A} = \begin{pmatrix} a & \frac{\sqrt{\lambda^2 - a^2\varepsilon^2}}{\varepsilon} \\ \frac{\sqrt{\lambda^2 - a^2\varepsilon^2}}{\varepsilon} & -a \end{pmatrix}, \quad \tilde{B} = \begin{pmatrix} 0 & 0 \\ 0 & \frac{1}{\varepsilon^2} \end{pmatrix},$$

$$\tilde{N}(\tilde{w}_1) = \begin{pmatrix} 0 \\ \frac{h(\tilde{w}_1)}{\varepsilon\sqrt{\lambda^2 - a^2\varepsilon^2}} \end{pmatrix}.$$

134 **3. The Green function of the linear partially dissipative system.** We
 135 consider the linear part of the (C-D) system (20)-(21) without the *tilde* for simplicity,

$$136 \quad (22) \quad \partial_t \mathbf{w} + A \partial_x \mathbf{w} = -B \mathbf{w}.$$

137 We want to apply the approach developed in [1], to study the singular approximation
 138 system above. The main difficulty here is to deal with the singular perturbation
 139 parameter ε . We consider the Green kernel $\Gamma(t, x)$ of (22), which satisfies

$$140 \quad (23) \quad \begin{cases} \partial_t \Gamma + A \partial_x \Gamma = -B \Gamma, \\ \Gamma(0, x) = \delta(x) I. \end{cases}$$

141 Taking the Fourier transform $\hat{\Gamma}$, we get

$$142 \quad (24) \quad \begin{cases} \frac{d}{dt} \hat{\Gamma} = (-B - i\xi A) \hat{\Gamma}, \\ \hat{\Gamma}(0, \xi) = I. \end{cases}$$

143 Consider the entire function

$$144 \quad (25) \quad E(z) = -B - zA = \begin{pmatrix} -az & -\frac{z\sqrt{\lambda^2 - a^2\varepsilon^2}}{\varepsilon} \\ -\frac{z\sqrt{\lambda^2 - a^2\varepsilon^2}}{\varepsilon} & az - \frac{1}{\varepsilon^2} \end{pmatrix}.$$

145 Formally, the solution to (24) is given by

$$146 \quad (26) \quad \hat{\Gamma}(t, \xi) = e^{E(i\xi)t} = \sum_{n=0}^{\infty} (-B - i\xi A)^n.$$

Since $E(z)$ in (25) is symmetric, if z is not exceptional we can write

$$E(z) = \lambda_1(z)P_1(z) + \lambda_2(z)P_2(z),$$

where $\lambda_1(z), \lambda_2(z)$ are the eigenvalues of $E(z)$, and $P_1(z), P_2(z)$ the related eigenprojections, given by

$$P_j(z) = -\frac{1}{2\pi i} \oint_{|\xi - \lambda_j(z)| < \epsilon} (E(z) - \xi I)^{-1} d\xi, \quad j = 1, 2.$$

147 Following [1], we study the low frequencies (case $z = 0$) and the high frequencies one
148 (case $z = \infty$) separately.

149 **Case** $z = 0$. The total projector for the eigenvalues near to 0 is

$$150 \quad (27) \quad P(z) = -\frac{1}{2\pi i} \oint_{|\xi| < \epsilon} (E(z) - \xi I)^{-1} d\xi.$$

151 Besides, the following expansion holds true, see [11],

$$152 \quad (28) \quad P(z) = Q_0 + \sum_{n \geq 1} z^n P^n(z),$$

153 where Q_0 is the eigenprojection for $E(0) - \xi I = -B - \xi I$, i.e.

$$154 \quad (29) \quad Q_0 = -\frac{1}{2\pi i} \oint_{|\xi| < \epsilon} (-B - \xi I)^{-1} d\xi = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix},$$

$$P^n(z) = -\frac{1}{2\pi i} \oint R^{(n)}(\xi) d\xi, \quad n \geq 1,$$

$R^{(n)}$ being the n -th term in the expansion of the resolvent (30). Here Q_0 is the projection onto the null space of the source term, while we denote by $Q_- = I - Q_0$ the complementing projection, and by L_-, L_0 and R_-, R_0 the related left and right eigenprojectors, see [11, 1], i.e.

$$L_- = R_-^T = \begin{pmatrix} 0 & 1 \end{pmatrix}, \quad L_0 = R_0^T = \begin{pmatrix} 1 & 0 \end{pmatrix}, \\ Q_- = R_- L_-, \quad Q_0 = R_0 L_0.$$

155 On the other hand, from [11],

$$156 \quad R(\xi, z) = (E(z) - \xi I)^{-1} = (-B - zA - \xi I)^{-1} = (-B - \xi I)^{-1} \sum_{n=0}^{\infty} (Az(-B - \xi I)^{-1})^n \\ 157 \quad = (-B - \xi I)^{-1} + \sum_{n \geq 1} (-B - \xi I)^{-1} z^n (A(-B - \xi I)^{-1})^n \\ 158 \quad = R_0(\xi) + \sum_{n \geq 1} R^{(n)}(\xi), \\ 159$$

160 i.e.

$$161 \quad (30) \quad R^{(n)} = z^n(-B - \xi I)^{-1}(A(-B - \xi I)^{-1})^n.$$

162 Since we are in a neighborhood of $z = 0$, at this point the authors in [1] consider
 163 only the first two terms of the asymptotic expansion of the total projector (28), so
 164 obtaining an expression with a remainder $O(z^2)$. The same approximation cannot be
 165 performed here, since we need to handle the singular terms in ε . Thus, we develop an
 166 explicit spectral analysis for the Green function of our problem. First of all,

$$167 \quad (31) \quad A(-B - \xi I)^{-1} = \begin{pmatrix} -\frac{a}{\xi} & -\frac{\varepsilon\sqrt{\lambda^2 - a^2\varepsilon^2}}{1 + \varepsilon^2\xi} \\ -\frac{\sqrt{\lambda^2 - a^2\varepsilon^2}}{\varepsilon\xi} & \frac{a\varepsilon^2}{1 + \varepsilon^2\xi} \end{pmatrix},$$

which is diagonalizable, i.e.

$$A(-B - \xi I)^{-1} = VDV^{-1},$$

168 where D is the diagonal matrix with entries given by the eigenvalues, and V is the
 169 matrix with the eigenvectors on the columns. Explicitly, setting

$$170 \quad (32) \quad \phi := a^2 + 4\lambda^2\xi + 4\varepsilon^2\lambda^2\xi^2,$$

171 we have

$$172 \quad (33) \quad D = \text{diag} \left\{ \frac{-a \pm \sqrt{\phi}}{2\xi(1 + \varepsilon^2\xi)} \right\},$$

$$V = \begin{pmatrix} \frac{\varepsilon(a + \sqrt{\phi} + 2a\varepsilon^2\xi)}{2(1 + \varepsilon^2\xi)\sqrt{\lambda^2 - a^2\varepsilon^2}} & \frac{\varepsilon(a - \sqrt{\phi} + 2a\varepsilon^2\xi)}{2(1 + \varepsilon^2\xi)\sqrt{\lambda^2 - a^2\varepsilon^2}} \\ 1 & 1 \end{pmatrix},$$

173

$$174 \quad (34) \quad V^{-1} = \begin{pmatrix} \frac{(1 + \varepsilon^2\xi)\sqrt{\lambda^2 - a^2\varepsilon^2}}{\varepsilon\sqrt{\phi}} & \frac{-a + \sqrt{\phi} - 2a\varepsilon^2\xi}{2\sqrt{\phi}} \\ -\frac{(1 + \varepsilon^2\xi)\sqrt{\lambda^2 - a^2\varepsilon^2}}{\varepsilon\sqrt{\phi}} & \frac{a + \sqrt{\phi} + 2a\varepsilon^2\xi}{2\sqrt{\phi}} \end{pmatrix}.$$

This way, denoting by

$$\theta_1 = a - \sqrt{\phi} + 2a\varepsilon^2\xi, \quad \theta_2 = a + \sqrt{\phi} + 2a\varepsilon^2\xi, \quad \psi_1 = -\frac{a + \sqrt{\phi}}{2\xi(1 + \varepsilon^2\xi)}, \quad \psi_2 = -\frac{a - \sqrt{\phi}}{2\xi(1 + \varepsilon^2\xi)},$$

with ϕ in (32), from (30) we have

$$R^{(n)} = z^n(-B - \xi I)^{-1}(A(-B - \xi I)^{-1})^n = z^n(-B - \xi I)^{-1}(VD^nV^{-1})$$

$$= z^n \begin{pmatrix} \frac{\theta_1 \psi_2^n - \theta_2 \psi_1^n}{2\sqrt{\phi\xi}} & -\varepsilon \sqrt{\frac{\lambda^2 - a^2 \varepsilon^2}{\phi}} (\psi_1^n - \psi_2^n) \\ -\varepsilon \sqrt{\frac{\lambda^2 - a^2 \varepsilon^2}{\phi}} (\psi_1^n - \psi_2^n) & -\frac{\varepsilon^2}{2(1 + \varepsilon^2 \xi) \sqrt{\phi}} (\theta_2 \psi_2^n - \theta_1 \psi_1^n) \end{pmatrix}.$$

175 The above matrix is bounded in ε , and so now we can approximate the expression
 176 of the total projector (28) up to the second order. To this end, we consider $R^{(n)}$ for
 177 $n = 0, 1, 2$, we apply the integral formula (29), and we obtain

$$178 \quad (35) \quad P(z) = \begin{pmatrix} 1 + O(z^2) & -\varepsilon z \sqrt{\lambda^2 - a^2 \varepsilon^2} + \varepsilon O(z^2) \\ -\varepsilon z \sqrt{\lambda^2 - a^2 \varepsilon^2} + \varepsilon O(z^2) & \varepsilon^2 z^2 (\lambda^2 - a^2 \varepsilon^2) + \varepsilon^2 O(z^3) \end{pmatrix}.$$

179 Now, let $L(z), R(z)$ be the left and the right eigenprojector of $P(z)$, i.e.

$$180 \quad \begin{aligned} P(z) &= R(z)L(z), & L(z)R(z) &= I \\ 181 \quad L(z)P(z) &= L(z), & P(z)R(z) &= R(z). \end{aligned}$$

183 According to (35), we consider the second order approximation, given by

$$184 \quad (36) \quad \tilde{P}(z) = \begin{pmatrix} 1 & -\varepsilon z \sqrt{\lambda^2 - a^2 \varepsilon^2} \\ -\varepsilon z \sqrt{\lambda^2 - a^2 \varepsilon^2} & \varepsilon^2 z^2 (\lambda^2 - a^2 \varepsilon^2) \end{pmatrix},$$

185 with

$$186 \quad \tilde{L}(z) = \begin{pmatrix} 1 & -\varepsilon z \sqrt{\lambda^2 - a^2 \varepsilon^2} \end{pmatrix}, \quad \tilde{R}(z) = \begin{pmatrix} 1 \\ -\varepsilon z \sqrt{\lambda^2 - a^2 \varepsilon^2} \end{pmatrix},$$

188 where

$$189 \quad (37) \quad \begin{aligned} \tilde{P}(z) &= \tilde{R}(z)\tilde{L}(z), & \tilde{L}(z)\tilde{R}(z) &= 1 + \varepsilon^2 O(z^2), \\ \tilde{P}(z)\tilde{R}(z) &= \tilde{R}(z) + \varepsilon^2 O(z^2), & \tilde{L}(z)\tilde{P}(z) &= \tilde{L}(z) + \varepsilon^2 O(z^2), \end{aligned}$$

and so

$$P(z) = \tilde{P}(z) + O(z^2), \quad R(z) = \tilde{R}(z) + O(z^2), \quad L(z) = \tilde{L}(z) + O(z^2).$$

Let us point out that further expansions of $L(z), R(z)$ are not singular in ε too, since the weights in ε of these vectors come from (35). Precisely, one can see that $L^\varepsilon(z)$ depends on ε as follows:

$$L^\varepsilon(\cdot) = \begin{pmatrix} 1 & O(\varepsilon) \end{pmatrix} = [R(\cdot)^\varepsilon]^T.$$

190 Now, by using the left and the right operators, we decompose $E(z)$, see [1],

$$191 \quad (38) \quad E(z) = R(z)F(z)L(z) + R_-(z)F_-(z)L_-(z),$$

where $L_-(z), R_-(z)$ are left and right eigenprojectors associated with

$$P_-(z) = I - P(z),$$

while

$$F(z) = L(z)E(z)R(z), \quad F_-(z) = L_-(z)E(z)R_-(z).$$

192 We use the previous approximations of $L(z), R(z)$, so obtaining

$$193 \quad (39) \quad F(z) = (\tilde{L}(z) + O(z^2))(-B - Az)(\tilde{R}(z) + O(z^2)) = -az + (\lambda^2 - a^2\varepsilon^2)z^2 + O(z^3).$$

194 Now, consider $F_-(z)$. Matrix (35) and the above definition imply that

$$195 \quad (40) \quad P_-(z) = \begin{pmatrix} O(z^2) & z\varepsilon\sqrt{\lambda^2 - a^2\varepsilon^2} + \varepsilon O(z^2) \\ z\varepsilon\sqrt{\lambda^2 - a^2\varepsilon^2} + \varepsilon O(z^2) & 1 + \varepsilon^2 O(z^2) \end{pmatrix},$$

and, approximating again,

$$L_-(z) = \tilde{L}_-(z) + O(z^2) = \begin{pmatrix} z\varepsilon\sqrt{\lambda^2 - a^2\varepsilon^2} & 1 \end{pmatrix} + O(z^2),$$

$$R_-(z) = \tilde{R}_-(z) + O(z^2) = \begin{pmatrix} z\varepsilon\sqrt{\lambda^2 - a^2\varepsilon^2} \\ 1 \end{pmatrix} + O(z^2).$$

196 Thus,

$$197 \quad (41) \quad F_-(z) = \tilde{L}_-(z)(-B - Az)\tilde{R}_-(z) + O(z^2) = -\frac{1}{\varepsilon^2} + az + O(z^2).$$

198 This yields the proposition below.

199 **PROPOSITION 3.1.** *We have the following decomposition near $z = 0$:*

$$200 \quad (42) \quad E(z) = F(z)P(z) + E_-(z),$$

201 with $F(z)$ in (39), $P(z)$ in (35), $E_-(z) = R_-(z)F_-(z)L_-(z)$, and $F_-(z)$ in (41).

Case $z = \infty$. We consider $E(z) = -B - Az = z(-B/z - A) = zE_1(1/z)$ and, setting $z = i\xi$ and $\zeta = 1/z = -i\eta$, with $\xi, \eta \in \mathbb{R}$, we have

$$\begin{aligned} E_1(\zeta) &= -A - \zeta B \\ &= \begin{pmatrix} -a & -\frac{\sqrt{\lambda^2 - a^2\varepsilon^2}}{\varepsilon} \\ -\frac{\sqrt{\lambda^2 - a^2\varepsilon^2}}{\varepsilon} & a - \frac{\zeta}{\varepsilon^2} \end{pmatrix} = \begin{pmatrix} -a & -\frac{\sqrt{\lambda^2 - a^2\varepsilon^2}}{\varepsilon} \\ -\frac{\sqrt{\lambda^2 - a^2\varepsilon^2}}{\varepsilon} & a + \frac{i\eta}{\varepsilon^2} \end{pmatrix}. \end{aligned}$$

Since $E_1(\zeta)$ is symmetric, we determine the eigenvalues and the right eigenprojectors,

$$-A - \zeta B = \lambda_1^{E_1}(\zeta)R_1(\zeta)R_1^T(\zeta) + \lambda_2^{E_1}(\zeta)R_2(\zeta)R_2^T(\zeta),$$

such that, for $j = 1, 2$, $R_j^T(\zeta)R_j(\zeta) = I$. The following expression for the eigenvalues of $E_1(i\eta)$ is provided

$$\lambda_{1,2}^{E_1}(z) = \frac{i\eta}{2\varepsilon^2} \pm \frac{\sqrt{4\varepsilon^2\lambda^2 + 4a\eta\varepsilon^2i - \eta^2}}{2\varepsilon^2},$$

and it is simple to prove that both the corresponding eigenvalues of $E(z)$, which can be obtained multiplying $\lambda_1^{E_1}(z)$ and $\lambda_2^{E_1}(z)$ by $z = i\xi = i/\eta$, have a strictly negative real part in the high frequencies regime ($|\zeta| = |\eta| \ll 1$) and in the vanishing ε

limit. Moreover, setting $\delta_{1,2} = \sqrt{8\varepsilon^2\lambda^2 + 2\zeta^2 - 8a\varepsilon^2\zeta \pm (-2\zeta\sqrt{\mu} + 4a\varepsilon^2\sqrt{\mu})}$, where $\mu = 4\varepsilon^2\lambda^2 + \zeta^2 - 4a\varepsilon^2\zeta$, the normalized right eigenprojectors are given by:

$$R_1(\zeta) = \frac{1}{\delta_1} \begin{pmatrix} (2a\varepsilon^2 - \zeta) + \sqrt{\mu} \\ 2\varepsilon\sqrt{\lambda^2 - a^2\varepsilon^2} \end{pmatrix}, \quad R_2(\zeta) = \frac{1}{\delta_2} \begin{pmatrix} (2a\varepsilon^2 - \zeta) - \sqrt{\mu} \\ 2\varepsilon\sqrt{\lambda^2 - a^2\varepsilon^2} \end{pmatrix}.$$

The eigenprojectors are bounded in ε , even for ζ near zero. Thus, we can approximate the total projector of $E_1(\zeta) = -A - \zeta B$ in a more convenient way, i.e. we decompose

$$A = \lambda_1 R_1 R_1^T + \lambda_2 R_2 R_2^T,$$

where $\lambda_1 = \lambda/\varepsilon$, $\lambda_2 = -\lambda/\varepsilon$, and the corresponding eigenprojectors read

$$R_1 = \frac{1}{\sqrt{2\lambda}} \begin{pmatrix} \sqrt{\lambda + a\varepsilon} \\ \sqrt{\lambda - a\varepsilon} \end{pmatrix}, \quad R_2 = \frac{1}{\sqrt{2\lambda}} \begin{pmatrix} -\sqrt{\lambda - a\varepsilon} \\ \sqrt{\lambda + a\varepsilon} \end{pmatrix}.$$

202 Now, by considering the total projector for the family of eigenvalues going to $\lambda_j =$
203 $\pm\lambda/\varepsilon$ as $\zeta \approx 0$, we obtain the following approximations:

$$204 \quad (43) \quad F_{1j}(\zeta) = -\lambda_j I + \zeta R_j^T (-B) R_j + O(\zeta^2).$$

205 Explicitly,

$$206 \quad (44) \quad F_{11}(\zeta) = -\frac{\lambda}{\varepsilon} - \frac{(\lambda - a\varepsilon)\zeta}{2\lambda\varepsilon^2} + O(\zeta^2), \quad F_{12}(\zeta) = \frac{\lambda}{\varepsilon} - \frac{(\lambda + a\varepsilon)\zeta}{2\lambda\varepsilon^2} + O(\zeta^2).$$

207 Since $E(z) = zE_1(1/z)$, we multiply $F_1(\zeta) = F_1(1/z)$ by z and, for $|z| \rightarrow +\infty$,

$$208 \quad (45) \quad \lambda_1(z) = -\frac{\lambda}{\varepsilon}z - \frac{\lambda - a\varepsilon}{2\lambda\varepsilon^2} + O(1/z), \quad \lambda_2(z) = \frac{\lambda}{\varepsilon}z - \frac{\lambda + a\varepsilon}{2\lambda\varepsilon^2} + O(1/z),$$

209 while the projectors are

$$210 \quad (46) \quad \mathcal{P}_j(z) = R_j R_j^T + O(1/z), \quad j = 1, 2.$$

211 **REMARK 3.1.** Notice that the term $O(1/z)$ in (45) could be singular in ε . How-
212 ever, from the previous discussion, the eigenvalues of $E(z)$ have a strictly negative
213 real part. This implies that the coefficients of the even powers of z in (45) have a
214 negative sign, while the others are imaginary terms. Thus, $e^{\lambda_{1,2}(z)}$ are bounded in ε .

215 **PROPOSITION 3.2.** We have the following decomposition near $z = \infty$:

$$216 \quad (47) \quad E(z) = \lambda_1(z)\mathcal{P}_1(z) + \lambda_2(z)\mathcal{P}_2(z),$$

217 with $\lambda_1(z), \lambda_2(z)$ in (45), and $\mathcal{P}_1(z), \mathcal{P}_2(z)$ in (46).

218 4. Green function estimates.

Green function estimates near $z = 0$. We associate to (39) the parabolic equation

$$\partial_t w + a\partial_x w = (\lambda^2 - a^2\varepsilon^2)\partial_{xx} w.$$

219 We can write the explicit solution

$$220 \quad (48) \quad g(t, x) = \frac{1}{2\sqrt{(\lambda^2 - a^2\varepsilon^2)\pi t}} \exp \left\{ -\frac{(x - at)^2}{4(\lambda^2 - a^2\varepsilon^2)t} \right\}.$$

221 This means that, for some $c_1, c_2 > 0$,

$$222 \quad (49) \quad |g(t, x)| \leq \frac{c_1}{\sqrt{t}} e^{-(x-at)^2/ct}, \quad (t, x) \in \mathbb{R}^+ \times \mathbb{R}, \quad \forall \varepsilon > 0.$$

Now, recalling Proposition 3.1 and considering the approximation $\tilde{P}(z)$ in (36) of the total projector $P(z)$ in (35),

$$e^{E(z)t} = \hat{g}(z)\tilde{P}(z) + R_-(z)e^{F_-(z)t}L_-(z) + \hat{R}_1(t, z),$$

223 where $\hat{g}(z) = -az - (\lambda^2 - \varepsilon^2 a^2)z^2$, and $R_1(t, x)$ is a remainder term, we take the
224 inverse of the Fourier transform of

$$225 \quad (50) \quad \hat{K}(z) = \hat{g}(z)\tilde{P}(z),$$

226 which yields the expression of the first part of the Green function near $z = 0$, i.e.

$$227 \quad (51) \quad K(t, x) = \begin{pmatrix} g(t, x) & \varepsilon\sqrt{\lambda^2 - a^2\varepsilon^2} \frac{dg(t, x)}{dx} \\ \varepsilon\sqrt{\lambda^2 - a^2\varepsilon^2} \frac{dg(t, x)}{dx} & \varepsilon^2(\lambda^2 - a^2\varepsilon^2) \frac{d^2g(t, x)}{d^2x} \end{pmatrix}.$$

228 Here, $\hat{K}(t, \xi)$ is the approximation of $\hat{\Gamma}(t, \xi)$ in (26) for $|\xi| \approx 0$. Thus, for $\xi \in [-\delta, \delta]$
229 with $\delta > 0$ sufficiently small, we consider the following remainder term

$$\begin{aligned} R_1(t, x) &= \frac{1}{2\pi} \int_{-\delta}^{\delta} (e^{E(i\xi)t} - e^{\hat{K}(t, \xi)t}) e^{i\xi x} d\xi \\ 230 \quad (52) \quad &= \frac{1}{2\pi} \int_{-\delta}^{\delta} e^{i\xi(x-at) - \xi^2(\lambda^2 - a^2\varepsilon^2)t} (e^{O(\xi^3 t)} P(i\xi) - \tilde{P}(i\xi)) d\xi \\ &+ \frac{1}{2\pi} \int_{-\delta}^{\delta} R_-(i\xi) e^{F_-(i\xi)t} L_-(i\xi) e^{i\xi x} d\xi. \end{aligned}$$

231 We need an estimate for the remainder above. First of all, from (41) and (40),

$$\begin{aligned} 232 \quad \left| \frac{1}{2\pi} \int_{-\delta}^{\delta} R_-(i\xi) e^{F_-(i\xi)t} L_-(i\xi) e^{i\xi x} d\xi \right| &\leq \left| \frac{1}{2\pi} \int_{-\delta}^{\delta} P_-(i\xi) e^{(-1/\varepsilon^2 + ai\xi + O(\xi^2))t} e^{i\xi x} d\xi \right| \\ 233 \quad & \\ 234 \quad &\leq C e^{-t/\varepsilon^2} \end{aligned}$$

for some constant C . Following [1],

$$|e^{O(\xi^3 t)} P(i\xi) - \tilde{P}(i\xi)| = |z^3| t e^{2\mu|z|^2 t} \begin{pmatrix} O(1) & O(\varepsilon)|z| \\ O(\varepsilon)|z| & O(\varepsilon^2)|z|^2 \end{pmatrix},$$

for a constant $\mu > 0$. This way,

$$R_1(t, x) = e^{-(x-at)^2/(ct)} \begin{pmatrix} O(1)(1+t)^{-1} & O(\varepsilon)(1+t)^{-3/2} \\ O(\varepsilon)(1+t)^{-3/2} & O(\varepsilon^2)(1+t)^{-2} \end{pmatrix}.$$

236 *Green function estimates near $z = \infty$.* We associate to (45) the following equa-
 237 tions:

$$238 \quad \partial_t w + \frac{\lambda}{\varepsilon} \partial_x w = -\frac{\lambda - a\varepsilon}{2\lambda\varepsilon^2} w, \quad \partial_t w - \frac{\lambda}{\varepsilon} \partial_x w = -\frac{\lambda + a\varepsilon}{2\lambda\varepsilon^2} w.$$

240 We can write explicitly the solutions

$$241 \quad g_1(t, x) = \delta(x - \lambda t/\varepsilon) e^{-(\lambda - a\varepsilon)t/(2\lambda\varepsilon^2)}, \quad g_2(t, x) = \delta(x + \lambda t/\varepsilon) e^{-(\lambda + a\varepsilon)t/(2\lambda\varepsilon^2)}.$$

Thus,

$$|g_j(t, x)| \leq C \delta(x \pm \lambda t/\varepsilon) e^{-ct/\varepsilon^2}, \quad j = 1, 2.$$

243 We determine the Fourier transform of the Green function for $|z|$ going to infinity,

$$244 \quad (53) \quad \hat{\mathcal{K}}(t, \xi) = \exp \left\{ -i \frac{\lambda t \xi}{\varepsilon} - \frac{(\lambda - a\varepsilon)t}{2\lambda\varepsilon^2} \right\} \mathcal{P}_1(\infty) + \exp \left\{ i \frac{\lambda t \xi}{\varepsilon} - \frac{(\lambda + a\varepsilon)t}{2\lambda\varepsilon^2} \right\} \mathcal{P}_2(\infty).$$

245 This way, from Proposition 3.2, the remainder term here is

$$246 \quad (54) \quad R_2(t, x) = \frac{1}{2\pi} \int_{|\xi| \geq N} (e^{E(i\xi)t} - \hat{\mathcal{K}}(t, \xi)) e^{i\xi x} d\xi, \quad \text{and}$$

247

$$248 \quad |R_2| \leq \frac{1}{2\pi} \left| \int_{|\xi| \geq N} e^{i\xi(x - \lambda t/\varepsilon) - (\lambda - a\varepsilon)t/(2\lambda\varepsilon^2)} \cdot (e^{O(1)t/(i\xi) + O(1)t/\xi^2} \mathcal{P}_1(i\xi) - \mathcal{P}_1(\infty)) d\xi \right|$$

$$249 \quad + \frac{1}{2\pi} \left| \int_{|\xi| \geq N} e^{i\xi(x + \lambda t/\varepsilon) - (\lambda + a\varepsilon)t/(2\lambda\varepsilon^2)} \cdot (e^{O(1)t/(i\xi) + O(1)t/\xi^2} \mathcal{P}_2(i\xi) - \mathcal{P}_2(\infty)) d\xi \right|.$$

251 Following [1] and thanks to Remark 3.1,

$$252 \quad |R_2(t, x)| \leq C e^{-ct/\varepsilon^2} \left[\left| \int_{|\xi| \geq N} \frac{e^{i\xi(x \pm \lambda t/\varepsilon)}}{\xi} d\xi \right| + \int_{|\xi| \geq N} \frac{1}{\xi^2} d\xi \right] \leq C e^{-ct/\varepsilon^2}.$$

253

254 *Remainders in between.* Until now, we studied the Green function of the linearized
 255 diffusive Jin-Xin system for $z \approx 0$, which yields the parabolic kernel \hat{K} in (50), and
 256 for $z \approx \infty$, so obtaining $\hat{\mathcal{K}}$ in (53). In these two cases, we also provided estimates for
 257 the remainder terms:

- 258 • R_1 in (52) for the parabolic kernel K for $|\xi| \leq \delta$, with δ sufficiently small;
- 259 • R_2 in (54) for the transport kernel \mathcal{K} for $|\xi| \geq N$, with N big enough.

260 It remains to estimate the last remainder terms, namely the parabolic kernel K for
 261 $|\xi| \geq \delta$, $t \geq 1$, the transport kernel \mathcal{K} for $|\xi| \leq N$, and the kernel $E(z)$ for $\delta \leq |\xi| \leq N$.

262 *Parabolic kernel $K(t, x)$ for $|\xi| \geq \delta$, $\delta \ll 1$.* Let us define

$$263 \quad (55) \quad R_3(t, x) = \frac{1}{2\pi} \int_{|\xi| \geq \delta} \hat{K}(t, \xi) e^{i\xi x} d\xi.$$

264 Thus, from (50), for $t \geq 1$,

$$265 \quad |R_3(t, x)| \leq C \left| \int_{|\xi| \geq \delta} e^{i\xi(x - at)} e^{-(\lambda^2 - a^2\varepsilon^2)\xi^2 t} \tilde{P}(i\xi) d\xi \right|$$

$$266 \quad \leq \frac{C e^{-t/C}}{\sqrt{t}} \begin{pmatrix} O(1) & O(\varepsilon) \\ O(\varepsilon) & O(\varepsilon^2) \end{pmatrix}.$$

267

268 *Transport kernel for $|\xi| \leq N$. Set*

$$269 \quad (56) \quad R_4(t, x) = \frac{1}{2\pi} \int_{|\xi| \leq N} \hat{\mathcal{K}}(t, \xi) e^{i\xi x} d\xi,$$

270 and, from (53),

$$271 \quad |R_4(t, x)| \leq C e^{-(\lambda + |a|\varepsilon)t/(2\lambda\varepsilon^2)} \sum \left| \int_{-N}^N e^{i\xi(x \pm \lambda t/\varepsilon)} d\xi \right|$$

$$272 \quad \leq C e^{-ct/\varepsilon^2} \min \left\{ N, \frac{1}{|x \pm \lambda t/\varepsilon|} \right\}.$$

273

Kernel $E(z)$ for $\delta \leq |\xi| \leq N$. Finally, we set

$$R_5(t, x) = \frac{1}{2\pi} \int_{\delta \leq |\xi| \leq N} e^{E(i\xi)t} e^{i\xi t} d\xi.$$

274 Unlike [1], here we cannot simply apply the (SK) condition, as mentioned in the
275 Introduction, and a further analysis is needed. The eigenvalues of $E(i\xi) = -i\xi A - B$
276 are expressed here:

$$277 \quad (57) \quad \lambda_{1/2} = \frac{1}{2\varepsilon^2} \left(-1 \pm \sqrt{1 - 4\varepsilon^2(ia\xi + \lambda^2\xi^2)} \right) = \frac{-2(ia\xi + \lambda^2\xi^2)}{1 \pm \sqrt{1 - 4\varepsilon^2(ia\xi + \lambda^2\xi^2)}}.$$

By using the Taylor expansion for $\varepsilon \approx 0$,

$$\lambda_1 = -\frac{ia\xi + \lambda^2\xi^2}{1 - \varepsilon^2(ia\xi + \lambda^2\xi^2)}, \quad \lambda_2 = -\frac{1}{\varepsilon^2}.$$

Explicitly, denoting by

$$\Delta = \sqrt{1 - 4\varepsilon^2\xi(\lambda^2\xi + ia)}, \quad \phi_1 = -1 + \Delta + 2ia\xi\varepsilon^2, \quad \phi_2 = 1 + \Delta - 2ia\xi\varepsilon^2,$$

278 one can find that

$$279 \quad e^{E(i\xi)t} = \begin{pmatrix} \frac{e^{\lambda_2 t}}{2\Delta} \phi_1 + \frac{e^{\lambda_1 t}}{2\Delta} \phi_2 & -\frac{i\xi\varepsilon\sqrt{\lambda^2 - a^2\varepsilon^2}(e^{\lambda_1 t} - e^{\lambda_2 t})}{\Delta} \\ -\frac{i\xi\varepsilon\sqrt{\lambda^2 - a^2\varepsilon^2}(e^{\lambda_1 t} - e^{\lambda_2 t})}{\Delta} & \frac{e^{\lambda_1 t}}{2\Delta} \phi_1 + \frac{e^{\lambda_2 t}}{2\Delta} \phi_2 \end{pmatrix},$$

where $\phi_1 = -1 + \Delta = -2\varepsilon^2\xi(\lambda^2\xi + ia) + O(\varepsilon^2) = O(\varepsilon^2)$, $\phi_2 = 1 + \Delta = O(1)$,

280 and, in terms of the singular parameter ε , this yields

$$281 \quad e^{E(i\xi)t} = \begin{pmatrix} O(1)(e^{\lambda_1 t} + e^{\lambda_2 t}) & O(\varepsilon)(e^{\lambda_1 t} - e^{\lambda_2 t}) \\ O(\varepsilon)(e^{\lambda_1 t} - e^{\lambda_2 t}) & e^{\lambda_1 t} O(\varepsilon^2) + O(1)e^{\lambda_2 t} \end{pmatrix}.$$

282 Putting the above calculations all together and integrating in space with respect to
283 the Fourier variable for $\delta \leq |\xi| \leq N$, we get

$$284 \quad (58) \quad |R_5(t, x)| \leq C \begin{pmatrix} O(1)e^{-t/C} & O(\varepsilon)e^{-t/C} \\ O(\varepsilon)e^{-t/C} & O(\varepsilon^2)e^{-t/C} + O(1)e^{-t/\varepsilon^2} \end{pmatrix}.$$

285 From (52), (54), (55), (56), (58), we denote the remainder by

$$286 \quad (59) \quad R(t) = R_1(t) + R_2(t) + R_3(t) + R_4(t) + R_5(t).$$

287 The previous estimates provide the following lemma.

LEMMA 4.1. *Let $\Gamma(t, x)$ be the Green function of the linear system (22). We have the following decomposition:*

$$\Gamma(t, x) = K(t, x) + \mathcal{K}(t, x) + R(t, x),$$

288 with $K(t, x), \mathcal{K}(t, x), R(t, x)$ in (51), (53) and (59) respectively. Moreover, for some
289 constants c, C ,

$$290 \quad \bullet \quad |K(t, x)| \leq e^{-(x-at)^2/(ct)} \begin{pmatrix} O(1)(1+t)^{-1} & O(\varepsilon)(1+t)^{-3/2} \\ O(\varepsilon)(1+t)^{-3/2} & O(\varepsilon^2)(1+t)^{-2} \end{pmatrix};$$

$$291 \quad \bullet \quad |\mathcal{K}(t, x)| \leq Ce^{-ct/\varepsilon^2};$$

$$292 \quad \bullet \quad |R(t)| \leq e^{-(x-at)^2/(ct)} \begin{pmatrix} O(1)(1+t)^{-1} & O(\varepsilon)(1+t)^{-3/2} \\ O(\varepsilon)(1+t)^{-3/2} & O(\varepsilon^2)(1+t)^{-2} \end{pmatrix} \\ + \begin{pmatrix} O(1) & O(\varepsilon) \\ O(\varepsilon) & O(\varepsilon^2) \end{pmatrix} e^{-ct} + Id e^{-ct/\varepsilon^2}.$$

Decay estimates. Let us consider the solution to the Cauchy problem associated with the linear system (22) and initial data \mathbf{w}_0 ,

$$\hat{\mathbf{w}}(t, \xi) = \hat{\Gamma}(t, \xi) \hat{\mathbf{w}}_0(\xi) = e^{E(i\xi)t} \hat{\mathbf{w}}_0(\xi).$$

293 By using the decomposition provided by Lemma 4.1, we get the following theorem.

THEOREM 4.2. *Consider the linear system in (22), i.e.*

$$\partial_t \mathbf{w} + A \partial_x \mathbf{w} = -B \mathbf{w},$$

and let $Q_0 = R_0 L_0$ and $Q_- = R_- L_-$ as before, i.e. the eigenprojectors onto the null space and the negative definite part of $-B$ respectively. Then, for any function $\mathbf{w}_0 \in L^1 \cap L^2(\mathbb{R}, \mathbb{R})$, the solution $\mathbf{w}(t) = \Gamma(t) \mathbf{w}_0$ to the related Cauchy problem can be decomposed as

$$\mathbf{w}(t) = \Gamma(t) \mathbf{w}_0 = K(t) \mathbf{w}_0 + \mathcal{K}(t) \mathbf{w}_0 + R(t) \mathbf{w}_0.$$

294 Moreover, for any index β , the following estimates hold:

$$295 \quad (60) \quad \|L_0 D^\beta K(t) \mathbf{w}_0\|_0 \leq C \min\{1, t^{-1/4-|\beta|/2}\} \|L_0 \mathbf{w}_0\|_{L^1} \\ + C\varepsilon \min\{1, t^{-3/4-|\beta|/2}\} \|L_- \mathbf{w}_0\|_{L^1},$$

296

$$297 \quad (61) \quad \|L_- D^\beta K(t) \mathbf{w}_0\|_0 \leq C\varepsilon \min\{1, t^{-3/4-|\beta|/2}\} \|L_0 \mathbf{w}_0\|_{L^1} \\ + C\varepsilon^2 \min\{1, t^{-5/4-|\beta|/2}\} \|L_- \mathbf{w}_0\|_{L^1},$$

298

$$299 \quad (62) \quad \|D^\beta \mathcal{K}(t) \mathbf{w}_0\|_0 \leq Ce^{-ct/\varepsilon^2} \|D^\beta \mathbf{w}_0\|_0,$$

300

$$301 \quad (63) \quad \|L_0 D^\beta R(t) \mathbf{w}_0\|_0 \leq C \min\{1, t^{-1/4-|\beta|/2}\} \|L_0 \mathbf{w}_0\|_{L^1} \\ + C\varepsilon \min\{1, t^{-3/4-|\beta|/2}\} \|L_- \mathbf{w}_0\|_{L^1} \\ + Ce^{-ct} \|L_0 \mathbf{w}_0\|_{L^1} + C\varepsilon e^{-ct} \|L_- \mathbf{w}_0\|_{L^1} + Ce^{-ct/\varepsilon^2} \|\mathbf{w}_0\|_{L^1},$$

302

$$\begin{aligned}
& \|L_- D^\beta R(t) \mathbf{w}_0\|_0 \leq C\varepsilon \min\{1, t^{-3/4-|\beta|/2}\} \|L_0 \mathbf{w}_0\|_{L^1} \\
& + C\varepsilon^2 \min\{1, t^{-5/4-|\beta|/2}\} \|L_- \mathbf{w}_0\|_{L^1} \\
& + C\varepsilon e^{-ct} \|L_0 \mathbf{w}_0\|_{L^1} + C\varepsilon^2 e^{-ct} \|L_- \mathbf{w}_0\|_{L^1} + C e^{-ct/\varepsilon^2} \|\mathbf{w}_0\|_{L^1}.
\end{aligned}
\tag{64}$$

304

Proof. From Lemma 4.1, for some constants $c, C > 0$, and for an index β , it holds

$$\|D^\beta \mathcal{K}(t) \mathbf{w}_0\|_0 \leq C e^{-ct/\varepsilon^2} \|D^\beta \mathbf{w}_0\|_0.
\tag{65}$$

306

On the other hand, the hyperbolic kernel (50) can be estimated as

$$\begin{aligned}
& |L_0 \widehat{K}(t) \mathbf{w}_0| \leq C e^{-c|\xi|^2 t} (|L_0 \hat{\mathbf{w}}_0| + \varepsilon |\xi| |L_- \hat{\mathbf{w}}_0|), \\
& |L_- \widehat{K}(t) \mathbf{w}_0| \leq C e^{-c|\xi|^2 t} (\varepsilon |\xi| |L_0 \hat{\mathbf{w}}_0| + \varepsilon^2 |\xi|^2 |L_- \hat{\mathbf{w}}_0|).
\end{aligned}$$

310 This yields

$$\begin{aligned}
& \|L_0 K(t) \mathbf{w}_0\|_0^2 \leq C \int_0^\infty \int_{S^0} e^{-2c|\xi|^2 t} (|L_0 \hat{\mathbf{w}}_0(\xi)|^2 + \varepsilon^2 |\xi|^2 |L_- \hat{\mathbf{w}}_0(\xi)|^2) d\zeta d\xi \\
& \leq C \min\{1, t^{-1/2}\} \|L_0 \hat{\mathbf{w}}_0\|_\infty^2 + C\varepsilon^2 \min\{1, t^{-3/2}\} \|L_- \hat{\mathbf{w}}_0\|_\infty^2 \\
& \leq C \min\{1, t^{-1/2}\} \|L_0 \mathbf{w}_0\|_{L^1}^2 + C\varepsilon^2 \min\{1, t^{-3/2}\} \|L_- \mathbf{w}_0\|_{L^1}^2,
\end{aligned}$$

315 and

$$\begin{aligned}
& \|L_- K(t) \mathbf{w}_0\|_0^2 \leq C \int_0^\infty \int_{S^0} e^{-2c|\xi|t} (\varepsilon^2 |\xi|^2 |L_0 \hat{\mathbf{w}}_0(\xi)|^2 + \varepsilon^4 |\xi|^2 |L_- \hat{\mathbf{w}}_0(\xi)|^2) d\zeta d\xi \\
& \leq C\varepsilon^2 \min\{1, t^{-3/2}\} \|L_0 \mathbf{w}_0\|_{L^1}^2 + C\varepsilon^4 \min\{1, t^{-5/2}\} \|L_- \mathbf{w}_0\|_{L^1}^2.
\end{aligned}$$

319 Besides, for every β we multiply by $\xi^{2\beta}$ the integrand and we get

$$\begin{aligned}
& \|L_0 D^\beta K(t) \mathbf{w}_0\|_0 \leq C \min\{1, t^{-1/4-|\beta|/2}\} \|L_0 \mathbf{w}_0\|_{L^1} \\
& + C\varepsilon \min\{1, t^{-3/4-|\beta|/2}\} \|L_- \mathbf{w}_0\|_{L^1}, \\
& \|L_- D^\beta K(t) \mathbf{w}_0\|_0 \leq C\varepsilon \min\{1, t^{-3/4-|\beta|/2}\} \|L_0 \mathbf{w}_0\|_{L^1} \\
& + C\varepsilon^2 \min\{1, t^{-5/4-|\beta|/2}\} \|L_- \mathbf{w}_0\|_{L^1}.
\end{aligned}$$

323 The estimates for $R(t)$ are obtained in a similar way. \square

5. Decay estimates and convergence. Consider the local solution \mathbf{w} to the Cauchy problem associated with (20), where we drop the *tilde*, and initial data \mathbf{w}_0 . The solution to the nonlinear problem (20) can be expressed by using the Duhamel formula

$$\begin{aligned}
& \mathbf{w}(t) = \Gamma(t) \mathbf{w}_0 + \int_0^t \Gamma(t-s) (N(w_1(s)) - DN(0)w_1(s)) ds \\
& = \Gamma(t) \mathbf{w}_0 + \int_0^t \Gamma(t-s) \left(\begin{array}{c} 0 \\ \frac{h(w_1(s))}{\varepsilon \sqrt{\lambda^2 - a^2 \varepsilon^2}} \end{array} \right) ds \quad t \in [0, T^*].
\end{aligned}
\tag{66}$$

329 From (29) and the formulas below, we recall that $w_1 = L_0 \mathbf{w} = (1 - L_-) \mathbf{w}$ is the
 330 conservative variable, while $w_2 = L_- \mathbf{w}$ is the dissipative one. We remind the Green
 331 function decomposition given by Lemma 4.1. For the β -derivative,

$$\begin{aligned}
 332 \quad D^\beta \mathbf{w}(t) &= D^\beta K(t) \mathbf{w}(0) + \mathcal{K}(t) D^\beta \mathbf{w}(0) + R(t) D^\beta \mathbf{w}(0) \\
 333 \quad &+ \int_0^{t/2} D^\beta K(t-s) R_- L_- \begin{pmatrix} 0 \\ \frac{h(w_1(s))}{\varepsilon \sqrt{\lambda^2 - a^2 \varepsilon^2}} \end{pmatrix} ds \\
 334 \quad &+ \int_{t/2}^t K(t-s) R_- D^\beta L_- \begin{pmatrix} 0 \\ \frac{h(w_1(s))}{\varepsilon \sqrt{\lambda^2 - a^2 \varepsilon^2}} \end{pmatrix} ds \\
 335 \quad &+ \int_0^t \mathcal{K}(t-s) D^\beta \begin{pmatrix} 0 \\ \frac{h(w_1(s))}{\varepsilon \sqrt{\lambda^2 - a^2 \varepsilon^2}} \end{pmatrix} ds \\
 336 \quad &+ \int_0^{t/2} D^\beta R(t-s) R_- L_- \begin{pmatrix} 0 \\ \frac{h(w_1(s))}{\varepsilon \sqrt{\lambda^2 - a^2 \varepsilon^2}} \end{pmatrix} ds \\
 337 \quad &+ \int_{t/2}^t R(t-s) R_- D^\beta L_- \begin{pmatrix} 0 \\ \frac{h(w_1(s))}{\varepsilon \sqrt{\lambda^2 - a^2 \varepsilon^2}} \end{pmatrix} ds \\
 338 \quad & \\
 339 \quad & \\
 340 \quad &= D^\beta K(t) \mathbf{w}(0) + \mathcal{K}(t) D^\beta \mathbf{w}(0) + R(t) D^\beta \mathbf{w}(0) \\
 341 \quad &+ \int_0^{t/2} \begin{pmatrix} D^\beta K_{12}(t-s) \\ D^\beta K_{22}(t-s) \end{pmatrix} \frac{h(w_1(s))}{\varepsilon \sqrt{\lambda^2 - a^2 \varepsilon^2}} ds \\
 342 \quad &+ \int_{t/2}^t \begin{pmatrix} K_{12}(t-s) \\ K_{22}(t-s) \end{pmatrix} D^\beta \frac{h(w_1(s))}{\varepsilon \sqrt{\lambda^2 - a^2 \varepsilon^2}} ds \\
 343 \quad &+ \int_0^t \mathcal{K}(t-s) D^\beta \begin{pmatrix} 0 \\ \frac{h(w_1(s))}{\varepsilon \sqrt{\lambda^2 - a^2 \varepsilon^2}} \end{pmatrix} ds \\
 344 \quad & \\
 345 \quad & \\
 346 \quad &+ \int_0^{t/2} \begin{pmatrix} D^\beta R_{12}(t-s) \\ D^\beta R_{22}(t-s) \end{pmatrix} \frac{h(w_1(s))}{\varepsilon \sqrt{\lambda^2 - a^2 \varepsilon^2}} ds \\
 347 \quad & \\
 348 \quad &+ \int_{t/2}^t \begin{pmatrix} R_{12}(t-s) \\ R_{22}(t-s) \end{pmatrix} D^\beta \frac{h(w_1(s))}{\varepsilon \sqrt{\lambda^2 - a^2 \varepsilon^2}} ds. \\
 349 \quad &
 \end{aligned}$$

350 Notice that, from (51), K_{12}, K_{22} are of order ε and ε^2 respectively, and the same holds
 351 for

$$\begin{aligned}
 352 \quad R_{12} &= O(\varepsilon)(1+t)^{-3/2} e^{-(x-at)^2/ct} + O(\varepsilon) e^{-ct} + O(1) e^{-ct/\varepsilon^2}, \\
 353 \quad R_{22} &= O(\varepsilon^2)(1+t)^{-2} e^{-(x-at)^2/ct} + O(\varepsilon^2) e^{-ct} + O(1) e^{-ct/\varepsilon^2}.
 \end{aligned}$$

355 From the previous assumptions, $f(u) = f(w_1) = aw_1 + h(w_1)$, where $h(w_1) = w_1^2 \tilde{h}(w_1)$
 356 for some function $\tilde{h}(w_1)$. Thus, by using the estimates of Theorem 4.2, and recalling

357 that $\|\cdot\|_m = \|\cdot\|_{H^m(\mathbb{R})}$, for $m = 0, 1, 2$, ($H^0 = L^2$), we have, for $j = 1, 2$,

$$\begin{aligned}
358 \quad & \|\mathbf{w}(t)\|_m \leq C \min\{1, t^{-1/4}\} \|\mathbf{w}_0\|_{L^1} + C e^{-ct/\varepsilon^2} \|\mathbf{w}_0\|_m \\
359 \quad & + C \int_0^t \min\{1, (t-s)^{-3/4}\} (\|w_1^2 \tilde{h}(w_1)\|_{L^1} + \|w_1^2 \tilde{h}(w_1)\|_m) ds \\
360 \quad & + C \int_0^t e^{-c(t-s)} \|w_1^2 \tilde{h}(w_1)\|_m ds \\
361 \quad & + C \int_0^t \frac{1}{\varepsilon} e^{-c(t-s)/\varepsilon^2} \|w_1^2 \tilde{h}(w_1)\|_m ds.
\end{aligned}$$

363 For m big enough,

$$\begin{aligned}
364 \quad & \|\mathbf{w}(t)\|_m \leq C \min\{1, t^{-1/4}\} \|\mathbf{w}_0\|_{L^1} + C e^{-ct/\varepsilon^2} \|\mathbf{w}_0\|_m \\
365 \quad & + \int_0^t \min\{1, (t-s)^{-3/4}\} C(|w_1|_\infty) \|w_1\|_m^2 ds \\
366 \quad & + \int_0^t e^{-c(t-s)} C(|w_1|_\infty) \|w_1\|_m^2 ds \\
367 \quad & + \int_0^t \frac{1}{\varepsilon} e^{-c(t-s)/\varepsilon^2} C(|w_1|_\infty) \|w_1\|_m^2 ds.
\end{aligned}$$

369 From (19), we recall that $\mathbf{w} = \begin{pmatrix} w_1 \\ w_2 \end{pmatrix} = \begin{pmatrix} u \\ \frac{\varepsilon(v - au)}{\sqrt{\lambda^2 - a^2\varepsilon^2}} \end{pmatrix}$, and so, for $m = 2$,

$$\begin{aligned}
370 \quad & \|u(t)\|_2 + c\varepsilon\|v(t) - au(t)\|_2 \leq C \min\{1, t^{-1/4}\} (\|u_0\|_{L^1} + c\varepsilon\|v_0 - au_0\|_{L^1}) \\
371 \quad & + e^{-ct/\varepsilon^2} (\|u_0\|_2 + c\varepsilon\|v_0 - au_0\|_2) \\
372 \quad & + \int_0^t \min\{1, (t-s)^{-3/4}\} C(|u|_\infty) \|u\|_2^2 ds \\
373 \quad & + \int_0^t e^{-c(t-s)} C(|u|_\infty) \|u\|_2^2 ds \\
374 \quad & + \int_0^t \frac{1}{\varepsilon} e^{-c(t-s)/\varepsilon^2} C(|u|_\infty) \|u\|_2^2 ds.
\end{aligned}$$

376 Let us denote by

$$377 \quad (67) \quad E_m = \max\{\|u_0\|_{L^1} + \varepsilon\|v_0 - au_0\|_{L^1}, \|u_0\|_m + \varepsilon\|v_0 - au_0\|_m\},$$

378 where, according to (6), $v_0 = f(u_0) - \lambda^2 \partial_x u_0$, and

$$379 \quad (68) \quad M_0(t) = \sup_{0 \leq \tau \leq t} \{\max\{1, \tau^{1/4}\} (\|u(\tau)\|_2 + \varepsilon\|v(\tau) - au(\tau)\|_2)\}.$$

380 The first term of the right hand side of the above estimate gives

$$\begin{aligned}
381 \quad & C \min\{1, t^{-1/4}\} (\|u_0\|_{L^1} + c\varepsilon\|v_0 - au_0\|_{L^1}) + C e^{-ct/\varepsilon^2} (\|u_0\|_2 + c\varepsilon\|v_0 - au_0\|_2) \\
382 \quad & \leq C \min\{1, t^{-1/4}\} E_2.
\end{aligned}$$

Besides,

$$C(|u|_\infty)\|u\|_2^2 \leq C(|u|_\infty) \min\{1, s^{-1/2}\} M_0^2(s).$$

384

Thus,

385

$$\|u(t)\|_2 + \varepsilon\|v(t) - au(t)\|_2 \leq C \min\{1, t^{-1/4}\} E_2$$

386

$$+ M_0^2(t) \int_0^t e^{-c(t-s)} c(|u|_\infty) \min\{1, s^{-1/2}\} ds$$

387

$$+ M_0^2(t) \int_0^t \frac{1}{\varepsilon} e^{-c(t-s)/\varepsilon^2} c(|u|_\infty) \min\{1, s^{-1/2}\} ds$$

388

389

$$+ M_0^2(t) \int_0^t c(|u|_\infty) \min\{1, (t-s)^{-3/4}\} \min\{1, s^{-1/2}\} ds.$$

From the Sobolev embedding theorem,

$$c(|u(s)|_\infty) \leq c(\|u(s)\|_2) \leq C \min\{1, s^{-1/4}\} M_0(s) \leq C M_0(s).$$

390

This way,

391

$$\|u(t)\|_2 + \varepsilon\|v(t) - au(t)\|_2 \leq C \min\{1, t^{-1/4}\} E_2$$

392

$$+ C M_0^3(t) \int_0^t e^{-c(t-s)} \min\{1, s^{-1/2}\} ds$$

393

$$+ C M_0^3(t) \int_0^t \frac{1}{\varepsilon} e^{-c(t-s)/\varepsilon^2} \min\{1, s^{-1/2}\} ds$$

394

395

$$+ C M_0^3(t) \int_0^t \min\{1, (t-s)^{-3/4}\} \min\{1, s^{-1/2}\} ds.$$

396

Notice that

397

$$\frac{1}{\varepsilon} \int_0^t e^{-c(t-s)/\varepsilon^2} \min\{1, s^{-1/2}\} ds = \varepsilon e^{-ct/\varepsilon^2} \int_0^{t/\varepsilon^2} e^{c\tau} \min\{1, \varepsilon\sqrt{\tau}\} d\tau$$

398

$$\leq \varepsilon e^{-ct/\varepsilon^2} \int_0^{t/\varepsilon^2} e^{c\tau} d\tau$$

399

$$= \frac{\varepsilon}{c} [1 - e^{-ct/\varepsilon^2}]$$

400

$$\leq C\varepsilon.$$

402

By using this inequality in the previous estimate,

403

$$\|u(t)\|_2 + \varepsilon\|v(t) - au(t)\|_2 \leq C \min\{1, t^{-1/4}\} E_2$$

404

$$+ C M_0^3(t) \int_0^t e^{-c(t-s)} \min\{1, s^{-1/2}\} ds$$

405

$$+ \varepsilon C M_0^3(t)$$

406

407

$$+ C M_0^3(t) \int_0^t \min\{1, (t-s)^{-3/4}\} \min\{1, s^{-1/2}\} ds.$$

408

409

Applying usual lemmas on integration, as Lemma 5.2 in [1], we get the following inequality

410

$$M_0(t) \leq C(E_2 + M_0^3(t)).$$

Then, if E_2 is small enough,

$$M_0(t) \leq CE_2,$$

412 i.e.

$$413 \quad (69) \quad \|u(t)\|_2 + \varepsilon\|v(t) - au(t)\|_2 \leq C \min\{1, t^{-1/4}\}E_2.$$

414 By arguing as before and following [1], we have the theorem below.

415 **PROPOSITION 5.1.** *The following estimates hold, with C a constant independent*
 416 *of ε ,*

$$417 \quad (70) \quad \|D^\beta \mathbf{w}(t)\|_0 \leq C \min\{1, t^{-1/4-|\beta|/2}\}E_{|\beta|+3/2},$$

418

$$419 \quad (71) \quad \|D^\beta w_2(t)\|_0 \leq C \min\{1, t^{-3/4-|\beta|/2}\}E_{|\beta|+3/2},$$

420

$$421 \quad (72) \quad \|D^\beta \partial_t \mathbf{w}(t)\|_0 \leq C \min\{1, t^{-3/4-|\beta|/2}\}E_{|\beta|+5/2},$$

422

$$423 \quad (73) \quad \|D^\beta \partial_t w_2(t)\|_0 \leq C \min\{1, t^{-5/4-|\beta|/2}\}E_{|\beta|+7/2}.$$

424 The last result and estimate (69) prove Theorem 1.1.

REMARK 5.1. *Note that the time derivative estimates of the solution (72), (73) can be obtained by applying again the Duhamel formula and so, similarly to (69), $\|D^\beta \partial_t \mathbf{w}(t)\|_0$ is bounded by a constant depending on $\|D^\beta \partial_t \mathbf{w}|_{t=0}\|$. The latter is not singular in ε , thanks to the particular form of the initial data, as it is shown below. The initial data satisfy (6), i.e.*

$$v_0 = f(u_0) - \lambda^2 \partial_x u_0 = au_0 + h(u_0) - \lambda^2 \partial_x u_0.$$

In terms of the (C-D) variable \mathbf{w} ,

$$\begin{cases} u = w_1, \\ v = aw_1 + \frac{\sqrt{\lambda^2 - a^2\varepsilon^2}}{\varepsilon}w_2, \end{cases}$$

425 this gives the following relation:

$$426 \quad (74) \quad \frac{\sqrt{\lambda^2 - a^2\varepsilon^2}}{\varepsilon}w_2^0 = h(w_1^0) - \lambda^2 \partial_x w_1^0.$$

427 Using (74) in system (20),

$$428 \quad (75) \quad \begin{cases} \partial_t w_1|_{t=0} = -a\partial_x w_1^0 - \frac{\sqrt{\lambda^2 - a^2\varepsilon^2}}{\varepsilon}\partial_x w_2^0, \\ \partial_t w_2|_{t=0} = -\frac{\sqrt{\lambda^2 - a^2\varepsilon^2}}{\varepsilon}\partial_x w_1^0 + a\partial_x w_2^0 + \frac{\lambda^2}{\varepsilon\sqrt{\lambda^2 - a^2\varepsilon^2}}\partial_x w_1^0 \\ = \frac{a^2\varepsilon}{\sqrt{\lambda^2 - a^2\varepsilon^2}}\partial_x w_1^0 + a\partial_x w_2^0. \end{cases}$$

In terms of the original variable,

$$\begin{cases} \partial_t w_1|_{t=0} = -\partial_x f(\bar{u}_0) + \lambda^2 \partial_{xx} \bar{u}_0, \\ \partial_t w_2|_{t=0} = \frac{a\varepsilon}{\sqrt{\lambda^2 - a^2\varepsilon^2}}(\partial_x f(\bar{u}_0) - \lambda^2 \partial_{xx} \bar{u}_0). \end{cases}$$

429 **REMARK 5.2.** *As shown in the previous remark, the well-prepared initial data (6)*
 430 *allow to bound the H^s norm of the time derivative of the solution to (20), uniformly*
 431 *in ε . The case of general initial data should be explored in future works. However,*
 432 *in this remark let us consider, at least formally, the case of Maxwellian initial data,*
 433 *meaning that, instead of (6), we take*

(76)

$$434 (u_0, v_0) = (\bar{u}_0, f(\bar{u}_0)) = (\bar{u}_0, a\bar{u}_0 + h(\bar{u}_0)), \quad (f_1^\varepsilon(0, x), f_2^\varepsilon(0, x)) = (M_1(\bar{u}_0), M_2(\bar{u}_0)).$$

435 *In terms of the (C-D) variables \mathbf{w} in (19),*

$$436 (77) \quad (w_1(0, x), w_2(0, x)) = \left(\bar{u}_0, \frac{\varepsilon h(\bar{u}_0)}{\sqrt{\lambda^2 - a^2 \varepsilon^2}} \right).$$

437 *Consider the difference $\mathbf{w} - \mathbf{w}^M$, where \mathbf{w} is the solution to (22) with well-prepared*
 438 *initial data in (6), while \mathbf{w}^M is the solution to (22) with Maxwellian initial data (77).*
 439 *The associated initial conditions are:*

$$440 (78) \quad \mathbf{w}(0, x) - \mathbf{w}^M(0, x) = \left(0, \frac{-\lambda^2 \varepsilon \partial_x h(\bar{u}_0)}{\sqrt{\lambda^2 - a^2 \varepsilon^2}} \right).$$

We apply again the Duhamel formula as in (66), in order to estimate the behavior of
the difference between the two solutions. Taking into account the decomposition of the
Green function in Lemma 4.1, it follows that, for $t > 0$,

$$\|(\mathbf{w} - \mathbf{w}^M)(t, x)\|_0 \leq \varepsilon^2 C \|\partial_x \bar{u}_0\|.$$

441 *Thus, after the initial layer, for $t > 0$, the difference between the solution \mathbf{w} , with*
 442 *well-prepared initial data, and \mathbf{w}^M , with Maxwellian initial data, is of order ε^2 .*

Convergence in the diffusion limit and asymptotic behavior. We perform the one
 dimensional Chapman-Enskog expansion. Recalling that

$$w_1 = u, \quad w_2 = u_d,$$

where u is the conservative variable and u_d is the dissipative one, system (20) is

$$\partial_t \begin{pmatrix} u \\ u_d \end{pmatrix} + A \partial_x \begin{pmatrix} u \\ u_d \end{pmatrix} = \begin{pmatrix} 0 \\ q(u) \end{pmatrix},$$

443 with A in (21) and $q(u) = -\frac{w_2}{\varepsilon^2} + \frac{h(w_1)}{\varepsilon \sqrt{\lambda^2 - a^2 \varepsilon^2}}$. We consider the following nonlinear
 444 parabolic equation

$$445 \quad \partial_t u + a \partial_x u + \partial_x h(u) - (\lambda^2 - a^2 \varepsilon^2) \partial_{xx} u = \partial_x S,$$

446 where

$$447 (79) \quad S = \varepsilon \sqrt{\lambda^2 - a^2 \varepsilon^2} \{ \partial_t u_d - a \partial_x u_d \}.$$

448 The homogeneous equation is

$$449 (80) \quad \partial_t w_p + a \partial_x w_p + \partial_x h(w_p) - (\lambda^2 - a^2 \varepsilon^2) \partial_{xx} w_p = 0,$$

and associated Green function is provided here

$$\Gamma_p(t) = K_{11}(t) + \tilde{K}(t) + \tilde{R}(t),$$

450 with K_{11} in (51). We take the difference between the conservative variable $u = w_1$
451 and w_p ,
(81)

$$\begin{aligned} D^\beta(u(t) - w_p(t)) &= \int_0^{t/2} D^\beta D(K_{11}(t-s) + \tilde{R}(t-s))(h(w_p(s)) - h(u(s))) ds \\ &\quad + \int_0^{t/2} D^\beta D(K_{11}(t-s) + \tilde{R}(t-s))S(s) ds \\ 452 &\quad + \int_{t/2}^t D(K_{11}(t-s) + \tilde{R}(t-s))D^\beta(h(w_p(s)) - h(u(s)) + S(s)) ds \\ &\quad + \int_0^t \tilde{K}(t-s)D^\beta D(h(w_p(s)) - h(u(s)) + S(s)) ds. \end{aligned}$$

By using (70), (71), (73), we have

$$\|D^\beta S\|_0 \leq C\varepsilon \min\{1, t^{-5/4-\beta/2}\}E_{\beta+1}.$$

453 Let us define, for $\mu \in [0, 1/2)$,

$$454 \quad (82) \quad m_0(t) = \sup_{\tau \in [0, t]} \{\max\{1, \tau^{1/4+\mu}\} \|u(\tau) - w_p(\tau)\|_0\}.$$

455 For $\beta = 0$,

$$\begin{aligned} 456 \quad \|u(t) - w_p(t)\|_0 &\leq CE_1 m_0(t) \int_0^t \min\{1, (t-s)^{-3/4}\} \min\{1, s^{-1/2-\mu}\} ds \\ 457 &\quad + C\varepsilon E_3 \int_0^t \min\{1, (t-s)^{-3/4}\} \min\{1, s^{-1}\} ds \\ 458 &\quad + C(E_1 m_0(t) + \varepsilon E_4) \int_0^t e^{-c(t-s)} \min\{1, s^{-5/4}\} ds \\ 459 &\leq C \min\{1, s^{-1/4-\mu}\} (E_1 m_0(t) + \varepsilon E_1 + \varepsilon E_4 + (1/2 - \mu)^{-1} E_1 m_0(t)), \end{aligned}$$

460 i.e., if E_1 is small enough,

$$461 \quad (83) \quad m_0(t) \leq C\varepsilon E_4.$$

462 Similarly, it can be proved by induction that, for $\gamma < \beta$, defining

$$463 \quad (84) \quad m_\beta(t) = \sup_{\tau \in [0, t]} \{\max\{1, \tau^{1/4+\mu+\beta/2}\} \|D^\beta(u(\tau) - w_p(\tau))\|_0\},$$

and assuming $m_\gamma(t) \leq C(\mu)\varepsilon E_{\gamma+4}$, then

$$\|D^\beta(h(u(s)) - h(w_p(s)))\|_0 \leq C \min\{1, t^{-1/2-\mu-\beta/2}\} (C(\mu)E_{\beta+1}E_{\beta+3} + E_1 m_\beta(t)).$$

Using the last inequality, (79) and (83) in (81), finally we get

$$m_\beta(t) \leq C(\mu)\varepsilon E_{\beta+4},$$

465 which ends the proof of Theorem 1.2.

466 **Acknowledgments.** The author is grateful to Roberto Natalini for some helpful
 467 advices, encouragements and useful discussions.

468

REFERENCES

- 469 [1] S. BIANCHINI, B. HANOZET, R. NATALINI, Asymptotic behavior of smooth solutions for partially
 470 dissipative hyperbolic systems with a convex entropy, *Comm. Pure Appl. Math.* **60** (2007),
 471 1559-1622.
- 472 [2] R. BIANCHINI, R. NATALINI, Convergence of a vector BGK approximation for the incompressible
 473 Navier-Stokes equations, *submitted* (2017).
- 474 [3] F. BOUCHUT, Construction of BGK Models with a Family of Kinetic Entropies for a Given
 475 System of Conservation Laws, *J. Stat. Phys.* **95** (2003).
- 476 [4] F. BOUCHUT, F. GUARGUAGLINI, R. NATALINI, Diffusive BGK Approximations for Nonlinear
 477 Multidimensional Parabolic Equations, *Indiana Univ. Math. J.* **49** (2000), 723-749.
- 478 [5] M. CARFORA, R. NATALINI, A discrete kinetic approximation for the incompressible Navier-
 479 Stokes equations, *ESAIM: Math. Modelling Numer. Anal.* **42** (2008), 93-112.
- 480 [6] I.L. CHERN, Long-time effect of relaxation for hyperbolic conservation laws, *Comm. Math. Phys.*
 481 **172** (1995), 39-55.
- 482 [7] F. GOLSE, L. SAINT-RAYMOND, The Navier-Stokes limit of the Boltzmann equation for bounded
 483 collision kernels, *Invent. math.* **155** (81) (2004).
- 484 [8] B. HANOZET, R. NATALINI, Global Existence of Smooth Solutions for Partially Dissipative
 485 Hyperbolic Systems with a Convex Entropy, *Arch. Rational Mech. Anal.* **169** (2003), 89-
 486 117.
- 487 [9] S. JIN, H. L. LIU, Diffusion limit of a hyperbolic system with relaxation, *Meth. and Appl. Anal.*
 488 **5** (1998), 317-334.
- 489 [10] S. JIN, Z. XIN, The relaxation schemes for system of conservation laws in arbitrary space
 490 dimensions, *Comm. Pure Appl. Math.* **48** (1995), 235-277.
- 491 [11] T. KATO, Perturbation theory for linear operator, 2nd ed. Grundlehren der Mathematischen
 492 Wissenschaften, 132. *Springer, New York* (1976).
- 493 [12] S. KAWASHIMA, Large-time behavior of solutions to hyperbolic-parabolic systems of conservation
 494 laws and applications, *Proc. Roy. Soc. Edinburgh Sect. A* **106** (1987), 169-194.
- 495 [13] H. LIU, R. NATALINI, Long-time diffusive behavior of solutions to a hyperbolic relaxation system,
 496 *Asymptot. Anal.* **25** (1) (2001), 21-38.
- 497 [14] A. MAJDA, Compressible Fluid Flow and Systems of Conservation Laws in Several Space
 498 Variables, *Springer-Verlag, New York* (1984).
- 499 [15] P. MARCATI, K. NISHIHARA, The LpLq estimates of solutions to one-dimensional damped wave
 500 equations and their application to the compressible flow through porous media, *J. Diff. Eqs.*
 501 **191** (2003), 445469.
- 502 [16] C. MASCIA, Twenty-eight years with Hyperbolic Conservation Laws with Relaxation, *Acta*
 503 *Math. Scientia* **35** (4) (2015), 807-831.
- 504 [17] R. NATALINI, A discrete kinetic approximation of entropy solutions to multidimensional scalar
 505 conservation laws, *J. Diff. Eqs.* **148** (1998), 292-317.
- 506 [18] R. NATALINI, Convergence to equilibrium for the relaxation approximations of conservation
 507 laws, *Comm. Pure Appl. Math.* **49** (8) (1996), 795-823.
- 508 [19] T. NISHIDA, Nonlinear hyperbolic equations and related topics in fluid dynamics, *Département*
 509 *de Mathématique, Université de Paris-Sud, Orsay, Publications Mathématiques d'Orsay*
 510 (1978), 78-02.
- 511 [20] R. ORIVE, E. ZUAZUA, Long-time behavior of solutions to a non-linear hyperbolic relaxation
 512 system, *J. Diff. Eq.* **228** (2006), 17-38.
- 513 [21] M. PAICU AND G. RAUGEL, A hyperbolic perturbation of the Navier-Stokes equations, (Une
 514 perturbation hyperbolique des équations de Navier-Stokes.). *ESAIM, Proc.* **21** (2007), 65-87.
- 515 [22] B. PERTHAME, Kinetic formulation of conservation laws, *Oxford Lecture Series in Mathematics*
 516 *and its Applications* 21, *Oxford University Press* (2000).
- 517 [23] L. SAINT-RAYMOND, From the BGK model to the NavierStokes equations, *Annales scientifiques*
 518 *de l'École Normale Supérieure* **36.2** (2003), 271-317.
- 519 [24] D. SERRE, Systems of conservation laws Vol. I, Hyperbolicity, entropies, shock waves, *Cambridge*
 520 *University Press* (2000).
- 521 [25] Y. SHIZUTA, S. KAWASHIMA, Systems of equations of hyperbolic-parabolic type with applications
 522 to the discrete Boltzmann equation, *Hokkaido Math. J.* **14** (1985) n.2, 249-275.
- 523 [26] T. C. SIDERIS, B. THOMASES, D. WANG, Long time behavior of solutions to the 3D compressible
 524 Euler equations with damping, *Comm. Partial Differential Equations* **28** (2003) n. 3-4, 795-

- 525 816.
526 [27] M. TAYLOR, Partial differential equations III, *Applied Mathematical Sciences 117*, Springer
527 (1996).