



HAL
open science

Isochronous centers of polynomial Hamiltonian systems and a conjecture of Jarque and Villadelprat

Jacky Cresson, Jordy Palafox

► **To cite this version:**

Jacky Cresson, Jordy Palafox. Isochronous centers of polynomial Hamiltonian systems and a conjecture of Jarque and Villadelprat. *Journal of Differential Equations*, 2019, 266 (9), pp.5713-5747. hal-02020432

HAL Id: hal-02020432

<https://hal.science/hal-02020432>

Submitted on 22 Oct 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

Isochronous centers of polynomial Hamiltonian systems and a conjecture of Jarque and Villadelprat

Jacky Cresson^{a,*}, Jordy Palafox^a

^a *Université de Pau et des Pays de l'Adour,
Laboratoire de Mathématiques et leurs applications,
UMR CNRS 5142,
Batiment IPRA,
avenue de l'université,
64000 Pau, France*

Abstract

We study the conjecture of Jarque and Villadelprat stating that every center of a planar polynomial Hamiltonian system of even degree is nonisochronous. This conjecture has already been proved for quadratic and quartic systems. Using the correction of a vector field to characterize isochronicity and explicit computations of this quantity for polynomial vector fields, we are able to describe a very large class of nonisochronous Hamiltonian systems of even arbitrarily large degree.

Keywords: `elsarticle.cls`, L^AT_EX, Elsevier, template

2010 MSC: 00-01, 99-00

1. Introduction and main results

1.1. The Jarque-Villadelprat conjecture

In this paper, we study centers of planar polynomial Hamiltonian systems in the real case. In particular we focus on **isochronous centers**. Our main concern is the following conjecture stated by Jarque and Villadelprat in [20]:

Let X be a real polynomial Hamiltonian vector field of the form:

$$X(x, y) = -\partial_y H(x, y)\partial_x + \partial_x H(x, y)\partial_y, \quad (x, y) \in \mathbb{R}^2, \quad (1)$$

where $H(x, y)$ is a real polynomial in the variables x and y . The maximum degree of the polynomials $\partial_x H$ and $\partial_y H$ is the degree of the Hamiltonian vector field.

10

*Corresponding author

Email address: `jordy.palafox@univ-pau.fr` (Jordy Palafox)

URL: `jacky.cresson@univ-pau.fr` (Jacky Cresson)

Conjecture : *Every center of a planar polynomial Hamiltonian system of even degree is non-isochronous.*

The conjecture is known to be true for quadratic systems thanks to a result of Loud in [22] and
15 in the quartic case by a result of Jarque-Villadelprat in [20]. The proof of Jarque and Villadelprat
is based on a careful study of the bifurcations set and seems difficult to extend to an arbitrary
degree. The conjecture is open for the other cases despite partial results in this direction obtain
by B. Schuman in [27, 28] using an explicit computation of the first coefficients of the Birkhoff
normal form and Chen and al. [3] proving what they call a *weak version* of the conjecture, i.e.
20 that any vector fields having only even components is nonisochronous.

Different strategies can be used to go further toward this conjecture. A first class of methods
can be called *geometric* and are related to some special features of Hamiltonian or isochronous
centers. We can mention for example the work of L. Gavrilov [19] and P. Mardesic, C. Rousseau
25 and B. Toni [23]. Up to now, these methods are unable to reproduce some special results obtained
by B. Schuman [27, 28] for classes of polynomial vector fields of arbitrary degree. Another class of
methods can be called *analytic* and are more or less all dealing with the computations of quantities
which can be obtained algorithmically like *period constants* [16] and coefficients of *normal forms* [8]
(see also [17]). However, such methods are usually assumed to be *intractable* when one is dealing
30 with a vector field of arbitrary large degree (see for example [20] p.337). This is indeed the case
when one has no informations on the algebraic structure of these coefficients. Then one is reduced
to compute Grobner bases or to use the elimination method. However, one is quickly limited by
the computational complexity and the memory size need to perform these computations. Existing
results are restricted to polynomials of order 5.

35 A natural problem is then to look for methods allowing us to bypass these technical limitations.
In the analytic setting, this can be done for (pre)normal forms or other local analytic objects
using the *mould formalism* introduced by J. Ecalle in the 70 ([10],[11], see also [5]) which allows an
efficient algorithmic construction of these objects but also a very precise analysis of the coefficients.
40 The idea is to separate in these coefficients what is *universal* and what is not.

1.2. Main results

In this paper, using the formalism of *moulds* (see [10], [11]) and a particular object attached
to a vector field called the *correction* defined by Ecalle and Vallet in [14]. In particular, we obtain
a partial answer to the conjecture for arbitrarily large degree.

45

It is well known that isochronicity of a real center is equivalent to its linearisability (see [2], theorem 3.3, p.12). A main property of the correction is that it gives a very useful criterion for linearisability. Indeed, a vector field is linearisable if and only if its correction is zero. As the correction possesses an algorithmic and explicit form which is easily calculable using mould calculus
 50 we are able to give more informations on the isochronous set. This strategy was already used by one of us in [6].

In the following, we use the classical complex representation of real vector fields (see [21]). Let us denote by $X_{lin} = i(x\partial_x - \bar{x}\partial_{\bar{x}})$ and $X_r = P_r(x, \bar{x})\partial_x + \overline{P_r(x, \bar{x})}\partial_{\bar{x}}$ with $x \in \mathbb{C}$, P_r is a
 55 homogeneous polynomial of degree r , $P_r(x, \bar{x}) = \sum_{j=0}^r p_{r-j-1,j} x^{r-j} \bar{x}^j$.

We formulate our main results (Theorem 1 to Theorem 4) whose proofs are postpone in Section 4.

Theorem 1. *Let X be a non trivial real Hamiltonian vector field of even degree $2n$ given by:*

$$X = X_{lin} + \sum_{r=2}^{2n} X_r$$

If X satisfies one of the following conditions :

- 60 a) *there exists $1 \leq r < n - 1$ such that $p_{i,i} = 0$ for $i = 1, \dots, r - 1$ and $\text{Im}(p_{r,r}) > 0$,*
 b) *$p_{i,i} = 0$ for $i = 1, \dots, n - 1$,*

then the vector field is nonisochronous.

As a consequence we deduce that:

$$X = X_{lin} + X_2,$$

$$X = X_{lin} + X_2 + X_3 + X_4 \text{ with } \text{Im}(p_{1,1}) \geq 0,$$

$$X = X_{lin} + X_2 + X_3 + X_4 + X_5 + X_6 \text{ with } p_{1,1} > 0 \text{ or } p_{1,1} = 0 \text{ and } \text{Im}(p_{2,2}) > 0,$$

are nonisochronous.

65 As a corollary, we obtain the *weak version* of the Jarque-Villadelprat conjecture proved by X. Chen and al. [3]:

Corollary 1 (weak Jarque-Villadelprat conjecture). *Let X be a non trivial real Hamiltonian vector field of even degree $2n$ given by*

$$X = X_{lin} + X_2 + X_4 + \dots + X_{2n},$$

then X is nonisochronous.

The proof follows easily from Theorem 1 as for all $i = 1, \dots, n-1$, we have $p_{i,i} = 0$ due to the fact that there exists no odd components.

Theorem 2. *Let X be a non trivial real Hamiltonian vector field of the form:*

$$X = X_{lin} + X_k + \dots + X_{2l},$$

70 for $k \geq 2$, $2l \geq k$ and $l \leq k-1$. Then X is nonisochronous.

Using this last theorem, *without any conditions* we have that:

$$X = X_{lin} + X_2,$$

$$X = X_{lin} + X_3 + X_4,$$

$$X = X_{lin} + X_4 + X_5 + X_6,$$

or more funny

$$X = X_{lin} + \sum_{i=47}^{92} X_i$$

are nonisochronous. We see that Theorem 1 and Theorem 2 are complementary to each other.

75 Mixing the proofs of Theorem 1 and Theorem 2 we obtain:

Theorem 3. *Let X be a non trivial real polynomial Hamiltonian vector field on the form:*

$$X = X_{lin} + X_k + \dots + X_{2l} + \sum_{n=1}^m \sum_{j=c_n}^{2(c_n-1)} X_j$$

where $k \geq 2$, $2l \geq k$, $l \leq k-1$, $m \geq 1$ and the sequence c_n is defined by :

$$c_1 = 4l \quad \text{and} \quad \forall n \geq 2, \quad c_n = 4(c_{n-1} - 1), \quad (2)$$

Then X is nonisochronous.

A first example of nonisochronous vector field given by the last theorem is:

$$X = X_{lin} + X_2 + X_4 + X_5 + X_6.$$

Theorem 4. *Let $k \geq 2$ and $l \leq k-1$, a real polynomial Hamiltonian vector field denoted by X on one of these two forms:*

$$i) X = X_{lin} + X_k + \dots + X_{2l} + X_{2l+1} + \sum_{m=r}^{r+n} X_m$$

where $r \geq 2l+2$ and $\text{Im}(p_{l,l}) > 0$ or

$$ii) X = X_{lin} X_k + \dots + X_{2l} + X_{4l-1} + \sum_{m=r}^{r+n} X_r$$

where X_{2l} is nontrivial, $r \geq 4l$, with $\text{Im}(p_{2l-1,2l-1}) > 0$, are nonisochronous.

Using Theorem 3 and Theorem 4 we easily deduce the classical result that homogeneous per-
80 turbations of a linear center are non isochronous (see [27]) as well its generalization (see [28, 6]).

1.3. Plan of the paper

In Section 2, we give following J. Ecalle and B. Vallet [14] the definition of the correction of a vector field and remind some of its properties. We then look more specifically to the correction of polynomial real vector fields. We derive explicit formula allowing us to analyse its structure.

85

In Section 3, we prove that the set of isochronous Hamiltonian centers is an affine variety which can be explicitly described. We also prove that this variety is invariant under a non trivial torus action.

90

In Section 4, we give the proofs of our main results and some technical Lemmas.

We then discuss several perspectives for this work.

2. Correction of vector fields and Hamiltonian systems

2.1. Correction of a vector field

95

In this section, we remind the definition of the correction of a vector field following the work of J. Ecalle and B. Vallet [14]. In particular, we give the mould expansion of the correction, which plays a central role in our approach to study the linearisability. It must be noted that all these computations can be made in arbitrary dimension.

2.1.1. The correction of a vector field

We denote by X an analytic vector field on \mathbb{C}^ν at 0:

$$X = \sum_{1 \leq j \leq \nu} X_j(x) \partial_{x_j}$$

100 with $X_j(0) = 0$ and $X_j(x) \in \mathbb{C}\{x\}$. We can write the vector field X in its *prepared form*:

Definition 1. A vector field X is said to be in **prepared form** if it is given by

$$X = X_{lin} + \sum_{n \in A(X)} B_n \tag{3}$$

where X_{lin} is the linear part of X in diagonal form

$$X_{lin} = \sum_{j=1}^{\nu} \lambda_j x_j \partial_{x_j}, \tag{4}$$

the B_n are homogeneous differential operators of degree $n = (n_1, \dots, n_\nu) \in \mathbb{Z}^\nu$, where all n_i are integer except one which can be -1 , satisfying for all monomial $x^m = x_1^{m_1} \dots x_\nu^{m_\nu}$, where

105 $x = (x_1, \dots, x_\nu) \in \mathbb{C}^\nu$, $m = (m_1, \dots, m_\nu) \in \mathbb{N}^\nu$, the equality

$$B_n(x^m) = \beta_{n,m} x^{n+m}, \quad \beta_{n,m} \in \mathbb{C}^\nu, \tag{5}$$

and $A(X) \in \mathbb{Z}^\nu$ is the set of the degree of the operators B_n in the decomposition of X .

From the point of view of Analysis, *homogeneous differential operators* are more tractable. An operator B_n is said to be homogeneous of degree $n = (n_1, n_2)$ if for all monomial $x^l y^k$ we have $B_n(x^l y^k) = \beta_n^{l,k} \cdot x^{n_1+l} y^{n_2+k}$ with $\beta_n^{l,k} \in \mathbb{C}$.

110

In [14], J.Ecalte and B.Vallet introduce the **correction** of a vector field following previous works of G.Gallavotti [18] and H.Eliasson [15] in the Hamiltonian case.

Let us consider a vector field in prepared form. The correction is defined as follows([14], p.258):

115 **Definition 2.** *Let X be a vector field in prepared form with linear part X_{lin} . The **correction** of X , denoted by $Carr(X)$ is the vector field solution of the following problem:*

Find a vector field Z such that

$$X - Z \sim X_{lin}, \quad \text{and} \quad [X_{lin}, Z] = 0, \quad (6)$$

120 *where for two vector fields A and B , we denote $A \sim B$ if the vectors fields A and B are formally conjugate.*

The second condition implies that $Carr(X)$ is only made of resonant terms.

In [14], Ecalte and Vallet prove that the correction of a vector field admits a **mould expansion**.

Precisely, let us denote by $A^*(X)$ the set of the words given by the letters in $A(X)$ using the concatenation morphism *conc* on letters :

$$\begin{aligned} conc : A(X)^p &\rightarrow A^*(X) \\ (n_1, \dots, n_p) &\mapsto n_1 \cdot n_2 \cdot \dots \cdot n_p. \end{aligned}$$

125 for any integer p .

In the following, a word is denoted by $n_1 \cdot n_2 \cdot \dots \cdot n_p$ or $n_1 n_2 \dots n_p$.

Remark 1. *The length of the word $n_1 \dots n_p$ is p . The word of length 0 is denoted by \emptyset .*

130 **Definition 3.** *The set $A^*(X)$ is composed by all the words of all lengths that is, if $\mathbf{n} \in A^*(X)$ there exists an integer $p \geq 0$ such that $\mathbf{n} = conc(n_1, \dots, n_p)$ where $n_j \in A(X)$ for $j = 1, \dots, p$. We denote $A^p(X)$ the set of words of length p .*

For all $\mathbf{n} = n_1 \cdot \dots \cdot n_r \in A^*(X)$, we denote:

$$B_{\mathbf{n}} = B_{n_1} \circ \dots \circ B_{n_r}. \quad (7)$$

The correction can be written as ([14], Lemma 3.2 p.267):

$$Carr(X) = \sum_{\mathbf{n} \in A^*(X)} Carr^{\mathbf{n}} B_{\mathbf{n}}, \quad (8)$$

where $Carr^{\mathbf{n}} \in \mathbb{C}$ for all $\mathbf{n} \in A^*(X)$, or simply

$$Carr(X) = \sum_{\bullet} Carr^{\bullet} B_{\bullet}. \quad (9)$$

135 following Ecalle's notations (see [10] and [5]).

The mapping

$$Carr^{\bullet} : \begin{array}{l} A(X)^* \longrightarrow \mathbb{C}, \\ \mathbf{n} \longmapsto Carr^{\mathbf{n}}, \end{array} \quad (10)$$

is called a **mould** (see [10, 11] and [5]).

140 The main point is that the mould $Carr^{\bullet}$ can be computed *algorithmically* using a recursive formula on the length of words.

Precisely for all $n \in A(X)$, let us denote by $\omega(n)$ the quantity:

$$\omega(n) = \langle n, \lambda \rangle,$$

where the $\langle \cdot, \cdot \rangle$ is the usual scalar product on \mathbb{C}^n and $\lambda = (\lambda_1, \dots, \lambda_\nu)$ are the eigenvalues of X_{lin} .

We can extend ω to a morphism from $(A^*(X), conc)$ to $(\mathbb{C}, +)$, i.e. for all $\mathbf{n} = n_1.n_2 \dots n_r \in A^*(X)$,

145 $n_i \in A(X)$, $i = 1, \dots, r$, we have

$$\omega(\mathbf{n}) = \omega(n_1) + \dots + \omega(n_r). \quad (11)$$

The quantity $\omega(n)$ is called the **weight** of the letter n .

We have the following theorem (formula 3.42 in [14]):

Theorem 5 (Variance formula). *The mould of the correction is given by the formula for any word $\mathbf{n} = n_1 \cdot \dots \cdot n_r$:*

$$\omega(n_1)Carr^{n_1, n_2, \dots, n_r} + Carr^{n_1+n_2, n_3, \dots, n_r} = \sum_{n_1 \mathbf{bc} = \mathbf{n}} Carr^{n_1 \mathbf{c}} Carr^{\mathbf{b}}. \quad (12)$$

150 The proof of this theorem is nontrivial. It follows from the *variance formula* for a vector field discussed in ([14], Prop 3.1 p.270). The variance of a vector field gives many different way to

compute the mould of the correction.

The main consequence of the previous Theorem is the *universal* character of the mould of the
 155 correction. Precisely, following the definition of universality used in [5], we have :

Theorem 6 (Universality of the Correction's mould). *There exists a one parameter family of complex functions $C_r : \mathcal{D}_r \subset \mathbb{C}^r \rightarrow \mathbb{C}$, $r \in \mathbb{N}$ such that for all X the correction's mould $Carr^\bullet$ defined on $A(X)^*$ is given for all $\mathbf{n} \in A(X)^*$ such that $l(\mathbf{n}) = r$, $r \in \mathbb{N}$ by*

$$Carr^{\mathbf{n}} = \begin{cases} C_r(\omega(n_1), \dots, \omega(n_r)), & \text{if } \omega(\mathbf{n}) = 0, \\ 0 & \text{otherwise.} \end{cases} \quad (13)$$

This property is fundamental concerning our problem as the computation of these coefficients
 160 is done once and for all and does not depends on the value of the coefficients entering the polynomials but on the alphabet generated by the vector field. Up to our knowledge only the mould formalism is able to produce such kind of coefficients allowing to write the correction (this is not the case for example dealing with the classical Lie framework).

165 As for prenormal forms (see [13]), a specific role is played by resonant terms:

Definition 4. *A word $\mathbf{n} \in A^*(X)$ is said to be resonant if $\omega(\mathbf{n}) = 0$.*

In the following, we give explicit expressions for C_1 , C_2 and C_3 .

2.1.2. The mould of the correction

The following theorem concerns precisely the length 1,2 and 3 :

170 **Theorem 7.** *The universal correction functions $C_r : \mathbb{C}^r \rightarrow \mathbb{C}$ of Theorem 6 are given for $r = 1, 2, 3$ by*

$$C_1(x) = \begin{cases} 1 & \text{if } z_1 = 0, \\ 0 & \text{otherwise.} \end{cases} \quad C_2(z_1, z_2) = \begin{cases} -\frac{1}{z_1} & \text{if } z_1 + z_2 = 0, z_1 \neq 0, \\ 0 & \text{otherwise.} \end{cases} \quad (14)$$

$$C_3(z_1, z_2, z_3) = \begin{cases} \frac{1}{z_1(z_1 + z_2)}, & \text{if } z_1 + z_2 + z_3 = 0, z_1 \neq 0, z_1 + z_2 \neq 0, \\ 0 & \text{otherwise.} \end{cases} \quad (15)$$

The proof is based on explicit computations which are summarized in the following Lemmas whose proof are given in Appendix.

175

We can remark that the values of the correction's mould depend on the weight of the letters.

Lemma 1. *The mould $Carr^\bullet$ verifies :*

- 1) $Carr^0 = 0$,
- 2) If \mathbf{n} is non resonant, $Carr^{\mathbf{n}} = 0$,
- 3) If $\mathbf{n} = n_1 \dots n_r$ is such that there exists j satisfying $\omega(n_j) = 0$ then $Carr^{\mathbf{n}} = 0$.

Moreover, we have:

Lemma 2. 1) If $\omega(n) = 0$, $Carr^n = 1$,

2) If $\omega(n_1 \cdot n_2) = 0$ with $\omega(n_1) = -\omega(n_2) \neq 0$, we have $Carr^{n_1 \cdot n_2} = -\frac{1}{\omega(n_1)}$,

3) If $\omega(n_1 \cdot n_2 \cdot n_3) = 0$ with $\omega(n_j) \neq 0$ $j = 1, 2, 3$, we have $Carr^{n_1 \cdot n_2 \cdot n_3} = \frac{1}{\omega(n_1)(\omega(n_1) + \omega(n_2))}$.

We refer to Appendix B.2 and Appendix B.3 for the proofs.

2.1.3. Some computations of the correction mould

Let us consider the quadratic case, i.e.

$$X = X_{lin} + X_2, \quad (16)$$

where X_{lin} is diagonal with eigenvalues $(i, -i)$ and X_2 is a homogeneous polynomial vector field of degree 2 given by

$$X_2 = (p_{1,0}x^2 + p_{0,1}xy + p_{-1}, 2y^2)\partial_x + (q_{2,-1}x^2 + q_{0,1}xy + q_{1,0})\partial_y. \quad (17)$$

We have:

Lemma 3. *The alphabet generated by X_2 is given by*

$$A(X) = \{n_1 = (1, 0), n_{-1} = (0, 1), n_3 = (2, -1), n_{-3} = (-1, 2)\}. \quad (18)$$

Proof. The alphabet generated by X_2 is related to the decomposition of X_2 in homogeneous differential operators. A simple rearrangement of terms proves that

$$X_2 = B_{(1,0)} + B_{(0,1)} + B_{(-1,2)} + B_{(2,-1)}, \quad (19)$$

where

$$\begin{aligned} B_{(1,0)} &= x(p_{1,0}x\partial_x + q_{1,0}y\partial_y), & B_{(0,1)} &= y(p_{0,1}x\partial_x + q_{0,1}y\partial_y), \\ B_{(-1,2)} &= p_{-1,2}y^2\partial_x, & B_{(2,-1)} &= q_{2,-1}x^2\partial_y. \end{aligned} \quad (20)$$

This concludes the proof. □

All the letters in $A(X)$ are non resonant. Indeed, we have

$$\omega(n_1) = i, \omega(n_{-1}) = -i, \omega(n_3) = 3i, \omega(n_{-3}) = -3i. \quad (21)$$

As a consequence, the correction mould is always zero in length 1.

In length 2 however, some resonant combinations are possible. We have

Word \mathbf{n}	$Carr^{\mathbf{n}}$
$n_1 \cdot n_{-1}$	i
$n_{-1} \cdot n_1$	$-i$
$n_3 \cdot n_{-3}$	$\frac{i}{3}$
$n_{-3} \cdot n_3$	$\frac{-i}{3}$

In length 4, the correction mould is given by :

Word \mathbf{n}	$Carr^{\mathbf{n}}$	Word \mathbf{n}	$Carr^{\mathbf{n}}$
$n_{-3} \cdot n_{-3} \cdot n_3 \cdot n_3$	$\frac{-i}{54}$	$n_1 \cdot n_{-3} \cdot n_{-1} \cdot n_3$	$\frac{-i}{6}$
$n_{-3} \cdot n_{-1} \cdot n_1 \cdot n_3$	$\frac{-i}{12}$	$n_1 \cdot n_{-3} \cdot n_1 \cdot n_1$	$\frac{-i}{2}$
$n_{-3} \cdot n_{-1} \cdot n_3 \cdot n_1$	$\frac{i}{12}$	$n_1 \cdot n_{-3} \cdot n_3 \cdot n_{-1}$	$\frac{i}{6}$
$n_{-3} \cdot n_1 \cdot n_{-1} \cdot n_3$	$\frac{i}{6}$	$n_1 \cdot n_{-1} \cdot n_{-3} \cdot n_3$	0
$n_{-3} \cdot n_1 \cdot n_1 \cdot n_1$	$\frac{i}{6}$	$n_1 \cdot n_{-1} \cdot n_{-1} \cdot n_1$	0
$n_{-3} \cdot n_1 \cdot n_3 \cdot n_{-1}$	$\frac{-i}{6}$	$n_1 \cdot n_{-1} \cdot n_1 \cdot n_{-1}$	$-i$
$n_{-3} \cdot n_3 \cdot n_{-3} \cdot n_3$	$\frac{i}{27}$	$n_1 \cdot n_{-1} \cdot n_3 \cdot n_{-3}$	0
$n_{-3} \cdot n_3 \cdot n_{-1} \cdot n_1$	0	$n_1 \cdot n_1 \cdot n_{-3} \cdot n_1$	$\frac{i}{2}$
$n_{-3} \cdot n_3 \cdot n_1 \cdot n_{-1}$	0	$n_1 \cdot n_1 \cdot n_{-1} \cdot n_{-1}$	$\frac{i}{2}$
$n_{-3} \cdot n_3 \cdot n_3 \cdot n_{-3}$	0	$n_1 \cdot n_1 \cdot n_1 \cdot n_{-3}$	$\frac{-i}{6}$
$n_{-1} \cdot n_{-3} \cdot n_1 \cdot n_3$	$\frac{i}{12}$	$n_1 \cdot n_3 \cdot n_{-3} \cdot n_{-1}$	$\frac{i}{12}$
$n_{-1} \cdot n_{-3} \cdot n_3 \cdot n_1$	$\frac{-i}{12}$	$n_1 \cdot n_3 \cdot n_{-1} \cdot n_{-3}$	$\frac{-i}{12}$
$n_{-1} \cdot n_{-1} \cdot n_{-1} \cdot n_3$	$\frac{i}{6}$	$n_3 \cdot n_{-3} \cdot n_{-3} \cdot n_3$	0
$n_{-1} \cdot n_{-1} \cdot n_1 \cdot n_1$	$\frac{-i}{2}$	$n_3 \cdot n_{-3} \cdot n_{-1} \cdot n_1$	0
$n_{-1} \cdot n_{-1} \cdot n_3 \cdot n_{-1}$	$\frac{-i}{2}$	$n_3 \cdot n_{-3} \cdot n_1 \cdot n_{-1}$	0
$n_{-1} \cdot n_1 \cdot n_{-3} \cdot n_3$	0	$n_3 \cdot n_{-3} \cdot n_3 \cdot n_{-3}$	$\frac{-i}{27}$
$n_{-1} \cdot n_1 \cdot n_{-1} \cdot n_1$	i	$n_3 \cdot n_{-1} \cdot n_{-3} \cdot n_1$	$\frac{i}{6}$
$n_{-1} \cdot n_1 \cdot n_1 \cdot n_{-1}$	0	$n_3 \cdot n_{-1} \cdot n_{-1} \cdot n_{-1}$	$\frac{-i}{6}$
$n_{-1} \cdot n_1 \cdot n_3 \cdot n_{-3}$	0	$n_3 \cdot n_{-1} \cdot n_1 \cdot n_{-3}$	$\frac{-i}{6}$
$n_{-1} \cdot n_3 \cdot n_{-3} \cdot n_1$	$\frac{-i}{6}$	$n_3 \cdot n_1 \cdot n_{-3} \cdot n_{-1}$	$\frac{-i}{12}$
$n_{-1} \cdot n_3 \cdot n_{-1} \cdot n_{-1}$	$\frac{i}{2}$	$n_3 \cdot n_1 \cdot n_{-1} \cdot n_{-3}$	$\frac{i}{12}$
$n_{-1} \cdot n_3 \cdot n_1 \cdot n_{-3}$	$\frac{i}{6}$	$n_3 \cdot n_3 \cdot n_{-3} \cdot n_{-3}$	$\frac{i}{54}$

2.2. Correction of a polynomial vector field

2.2.1. Prepared form and alphabet

205 Let X a polynomial vector field in \mathbb{C}^2 of the form:

$$X = X_{lin} + \sum_{r=2}^l X_r, \quad (22)$$

with

$$X_r = P_r(x, y)\partial_x + Q_r(x, y)\partial_y, \quad (23)$$

where P_r and Q_r are homogeneous polynomials of degree r such that:

$$P_r(x, y) = \sum_{k=0}^r p_{k-1, r-k} x^k y^{r-k}, \quad Q_r(x, y) = \sum_{l=0}^r q_{r-k, k-1} x^{r-k} y^k. \quad (24)$$

In the following, we describe **explicitly** the prepared form of X , the set $A(X)$ and the operators B_n .

Lemma 4. *For all $r \geq 2$, we can associate an alphabet to the vector field X_r , written $A(X_r)$ given by:*

$$A(X_r) = \{(r, -1), (-1, r), (k-1, r-k) \text{ with } k = 1, \dots, r\}.$$

Moreover, we get from $A(X_r)$ the set denoted $\mathcal{B}(X_r)$ of the homogeneous differential operators given by the decomposition :

$$\begin{aligned} B_{(-1, r)} &= p_{-1, r} y^r \partial_x, \\ B_{(r, -1)} &= q_{r, -1} x^r \partial_y, \\ B_{(k-1, r-k)} &= x^{k-1} y^{r-k} (p_{k-1, r-k} x \partial_x + q_{k-1, r-k} y \partial_y) \end{aligned}$$

210 with $k \in \{1, \dots, r\}$.

Proof. We first rewrite any vector field X_r , $r = 2, \dots, l$, in order to make apparent the order, in term of differential operators, of all its components. We have :

$$\begin{aligned} P_r(x, y)\partial_x + Q_r(x, y)\partial_y &= \sum_{k=1}^r (p_{k-1, r-k} x^{k-1} y^{r-k} x \partial_x + q_{r-k, k-1} x^{r-k} y^{k-1} y \partial_y) + p_{-1, r} y^r \partial_x + q_{r, -1} x^r \partial_y, \\ &= \sum_{k=1}^r (O_{(k-1, r-k)} + \tilde{O}_{(r-k, k-1)}) + O_{(-1, r)} + O_{(r, -1)}, \end{aligned}$$

with

$$\begin{aligned} O_{(k-1, r-k)} &= p_{k-1, r-k} x^{k-1} y^{r-k} x \partial_x, \\ \tilde{O}_{(r-k, k-1)} &= q_{r-k, k-1} x^{r-k} y^{k-1} y \partial_y, \\ O_{(-1, r)} &= p_{-1, r} y^r \partial_x, \\ O_{(r, -1)} &= q_{r, -1} x^r \partial_y. \end{aligned}$$

We want to know if there exists some operators of the same degree among the operators $O_{(k-1, r-k)}$ and $\tilde{O}_{(r-k, k-1)}$. If it is the case, we gather them in a same operator of the form $B_{(n^1, n^2)} = x^{n^1} y^{n^2} (p_{(n^1, n^2)} x \partial_x + q_{(n^1, n^2)} y \partial_y)$.

For that, we have to solve the equation

$$k - 1 = r - \tilde{k} \quad (25)$$

where $k, \tilde{k} \in \{1, \dots, r\}$. As $\tilde{k} \in \{1, \dots, r\}$, $r - \tilde{k} \in \{0, \dots, r - 1\}$ hence $r - \tilde{k} + 1 \in \{1, \dots, r\}$. So there always exists solutions. This concludes the proof. \square

Example 1. We consider the vector field $X = X_{lin} + X_2 + X_3$ where

$$X_2 = (p_{1,0}x^2 + p_{0,1}xy + p_{-1,2}y^2) \partial_x + (q_{-1,2}x^2 + q_{1,0}xy + q_{0,1}yr) \partial_y,$$

$$X_3 = (p_{2,0}x^3 + p_{1,1}x^2y + p_{0,2}xy^2 + p_{-1,3}y^3) \partial_x + (q_{3,-1}x^3 + q_{2,0}x^2y + q_{1,1}xy^2 + q_{0,2}y^3) \partial_y.$$

Hence we obtain the three following alphabets :

$$A(X_2) = \{(2, -1), (1, 0), (0, 1), (-1, 2)\},$$

$$A(X_3) = \{(3, -1), (2, 0), (1, 1), (0, 2), (-1, 3)\}.$$

For example, we also have the elements of the set $\mathcal{B}(X_3)$:

$$B_{(3,-1)} = q_{3,-1}x^3\partial_y,$$

$$B_{(2,0)} = x^2(p_{2,0}x\partial_x + q_{2,0}y\partial_y),$$

$$B_{(1,1)} = xy(xp_{1,1}x\partial_x + q_{1,1}y\partial_y),$$

$$B_{(0,2)} = y^2(p_{0,2}x\partial_x + q_{0,2}y\partial_y),$$

$$B_{(-1,3)} = p_{-1,3}y^3\partial_x.$$

Definition 5. We define the degree of a vector fields as the maximum of the degree of its defining polynomial.

220 In a same way, we define the degree of the homogeneous differential operators B_n or of a Lie bracket of B_n which appear in the decomposition of X . We denote by $\deg(B_n)$ (resp. $\deg([B_n])$) the degree of B_n (resp. $[B_n]$).

Lemma 5. Let X be a vector fields of the form $X = X_{lin} + \sum_{r=2}^m X_r$ then X admits the alphabet

$$A(X) = \bigcup_{r=2}^m A(X_r) \text{ and } \mathcal{B}(X), \text{ the set of homogeneous differential operators of } X, \text{ is given by}$$

$$225 \mathcal{B}(X) = \bigcup_{r=2}^m \mathcal{B}(X_r).$$

Proof. For all $n = (n^1, n^2) \in A(X)$, we define the application :

$$p : A(X) \rightarrow \mathbb{N}$$

$$n = (n^1, n^2) \mapsto n^1 + n^2.$$

For every $r \geq 2$, for all $n \in A(X_r)$, we have $p(n) = r - 1$, so $\forall r, r'$, such that $r \neq r'$, we have $A(X_r) \cap A(X_{r'}) = \emptyset$ because $p(A(X_r)) \neq p(A(X_{r'}))$. Moreover, as $Der(\mathbb{C}^2) = \bigoplus_{r \geq 1} Der_r(\mathbb{C}^2)$ and $\mathcal{B}(X_r) \subset Der_r(\mathbb{C}^2)$, then $\mathcal{B}(X_r) \cap \mathcal{B}(X_{r'}) = \emptyset$ if $r \neq r'$. \square

The elements of the alphabet $A(X)$ are named *letter*.

230 2.2.2. *Depth*

As we have a one-to-one correspondence between $(A^*(X), \text{conc})$ and $(\mathcal{B}^*(X), \circ)$:

$$\begin{aligned} A^*(X) &\rightarrow \mathcal{B}^*(X), \\ \mathbf{n} = n_1 \cdot \dots \cdot n_r &\mapsto B_{\mathbf{n}} = B_{n_1} \circ \dots \circ B_{n_r}, \end{aligned}$$

the degree of $[B_{\mathbf{n}}]$ gives a natural notion of *depth* for the words defined by:

Definition 6. We denote by $p : A^*(X) \rightarrow \mathbb{N}$ the mapping defined by:

$$p(\mathbf{n}) = \text{deg}([B_{\mathbf{n}}]) - 1.$$

Lemma 6. The mapping p is a morphism from $(A^*(X), \text{conc})$ in $(\mathbb{N}, +)$.

Proof. We prove it by induction on the length of the words. Let $n_1, n_2 \in A(X)$,

$$\begin{aligned} p(n_1 \cdot n_2) &= \text{deg}([B_{n_1} B_{n_2}]) - 1 \\ &= \text{deg}(B_{n_1}) + \text{deg}(B_{n_2}) - 1 - 1 \\ &= p(n_1) + p(n_2). \end{aligned}$$

Let $n_1 \in A(X)$ and $\mathbf{n} \in A^*(X)$, so:

$$\begin{aligned} p(n_1 \cdot \mathbf{n}) &= \text{deg}([B_{n_1}, B_{\mathbf{n}}]) - 1 \\ &= \text{deg}(B_{n_1}) + \text{deg}(B_{\mathbf{n}}) - 1 - 1 \\ &= p(n_1) + p(\mathbf{n}). \end{aligned}$$

□

2.2.3. *The projection Theorem and rewriting of a mould expansion*

235 Let $\mathcal{M}(X)$ be a **mould series**, i.e. an expression of the form

$$\mathcal{M}(X) = \sum_{\mathbf{n} \in A^*(X)} M^{\mathbf{n}} B_{\mathbf{n}}, \quad (26)$$

where M^{\bullet} is an **alternal mould** ([14]), i.e. $\mathcal{M}(X)$ is *primitive* ([29], p.17) and then a vector field. In this case, using the *projection theorem* ([29], p.28), the mould series $\mathcal{M}(X)$ can be expressed in the following form :

$$\mathcal{M}(X) = \sum_{r \geq 1} \frac{1}{r} \sum_{\substack{\mathbf{n} \in A^*(X) \\ l(\mathbf{n})=r}} M^{\mathbf{n}} [B_{\mathbf{n}}], \quad (27)$$

where

$$[B_{\mathbf{n}}] = [B_{n_1 \dots n_r}] = [\dots [[B_{n_1}, B_{n_2}], B_{n_3}], \dots], B_{n_{r-1}}, B_{n_r}]. \quad (28)$$

240 We can reorganize this sum using the depth as follows:

$$\mathcal{M}(X) = \sum_{d \geq 1} \mathcal{M}_d(X), \quad (29)$$

where

$$\mathcal{M}_d(X) = \sum_{\substack{\mathbf{n} \in A^*(X) \\ p(\mathbf{n})=d}} M^{\mathbf{n}} B_{\mathbf{n}}. \quad (30)$$

A useful consequence is that the equation $\mathcal{M}(X) = 0$ is equivalent to $\mathcal{M}_d(X) = 0$ for all $d \geq 1$.

2.2.4. Expression of the correction and criterion of linearisability

The main property of the correction is that it provides a useful and simple criterion of linearizability. Indeed, we have by definition of the correction (see [14], p.258) :

Lemma 7. *A vector field X is linearizable if and only if $Carr(X) = 0$.*

Using the above decomposition, we obtain an explicit criterion for linearizability writing $Carr(X)$ as :

$$Carr(X) = \sum_{p \geq 1} \left(\sum_{\substack{\mathbf{n} \in A^*(X) \\ p(\mathbf{n})=p}} Carr^{\mathbf{n}} B_{\mathbf{n}} \right) = \sum_{p \geq 1} Carr_p(X).$$

Theorem 8. *A vector fields X is linearizable if and only if $Carr_p(X) = 0 \forall p \geq 1$.*

In the following, we derive some properties of the quantities $Carr_p(X)$.

2.3. Correction of real polynomial Hamiltonian vector fields

250 2.3.1. General properties

An interesting property of the correction is that we just have to consider the even depth, indeed:

Theorem 9. *Let X be a real Hamiltonian vector fields as above. Its correction in odd depth is zero, i.e.*

$$Carr_{2p+1}(X) = 0, \quad (31)$$

255 for all integer p .

This theorem is a consequence of the following lemma :

Lemma 8. *For a resonant word \mathbf{n} , the related Lie bracket is on the form :*

$$[B_{\mathbf{n}}] = (xy)^{\frac{p(\mathbf{n})}{2}} (P_{\mathbf{n}} x \partial_x + Q_{\mathbf{n}} y \partial_y).$$

Proof. For all resonant word $\mathbf{n} = n_1 \cdot \dots \cdot n_r$, the related Lie bracket is :

$$[B_{\mathbf{n}}] = x^{\sum n_j^1} y^{\sum n_j^2} (P_{\mathbf{n}} x \partial_x + Q_{\mathbf{n}} y \partial_y).$$

where $n_j = (n_j^1, n_j^2)$. As \mathbf{n} is resonant, we have :

$$\begin{aligned} \omega(\mathbf{n}) &= \omega(n_1) + \dots + \omega(n_r) \\ &= i \left(\sum n_j^1 - \sum n_j^2 \right) = 0, \end{aligned}$$

then $\sum n_j^1 = \sum n_j^2 = \alpha \in \mathbb{N}$. We just have to remark that $p(\mathbf{n}) = p(n_1) + \dots + p(n_r) = \sum n_j^1 + \sum n_j^2 = 2\alpha$, then $\alpha = \frac{p(\mathbf{n})}{2}$. \square

As a consequence, we can restrict our attention to the even components of the correction. For a given integer p , terms in $Carr_{2p}$ can be decomposed with respect to the length of words. Precisely, we have

$$Carr_p(X) = Carr_{p,1}(X) + \dots + Carr_{p,p}(X), \quad (32)$$

where

$$Carr_{p,j}(X) = \sum_{\substack{\mathbf{n} \in A^*(X) \\ p(\mathbf{n})=p, l(\mathbf{n})=j}} Carr^{\mathbf{n}} B_{\mathbf{n}}, \quad (33)$$

for $j = 1, \dots, p$.

The main point is that of course, this is a *finite* sum. Indeed, as each differential operator entering in the definition are at least of depth one, we can not have more than a word of length p as for all $j \in \mathbb{N}^*$, $\mathbf{n} \in A^*(X)$ such that $l(\mathbf{n}) = j$, we have $p(\mathbf{n}) \geq j$.

Moreover, some of these terms are easily determined.

Lemma 9. *Let $p \in \mathbb{N}^*$, we have*

$$\begin{aligned} Carr_{2p,2p}(X) &= Carr_{2p,2p}(X_2), \\ Carr_{2p,2}(X) &= Carr_{2p,2}(X_{p+1}) + \sum_{r=2}^p Carr_{2p,2}(X_r, X_{2p-r+2}), \\ Carr_{2p,1}(X) &= Carr_{2p}(X_{2p+1}) = B_{p,p}. \end{aligned} \quad (34)$$

This Lemma has important implications on the following, as it gives the maximal degree of the homogeneous vector fields X_r entering in the computation of a given correction term. In particular, for $Carr_{2p}$ we have no terms coming from the X_r with $r \geq 2p + 2$.

The proof of Lemma 9 is based on the properties of the set of resonant words with respects to length and depth.

Proof. i) The first equality easily follows from the fact that an element of depth $2p$ and length $2p$ is necessarily made of elements of depth 1 corresponding to operators in X_2 .

280 ii) The second equality is a direct rewriting of the definition for a length 2 contribution to the correction term of depth $2p$. We denoted by $\mathcal{W}(X_r)$ the set of weights coming from the component X_r of X and given by :

$$\mathcal{W}(X_r) = \{\langle n, \lambda \rangle, n \in A(X_r)\}. \quad (35)$$

The first term comes from the following decomposition lemma:

Lemma 10. *For every $r \geq 2$, $\mathcal{W}(X_r)$ can be decomposed in the following way :*

$$\mathcal{W}(X_r) = \mathcal{W}^+(X_r) \cup \mathcal{W}^-(X_r) \cup \mathcal{W}^0(X_r),$$

285 *where $\mathcal{W}^+(X_r)$ is the set of positive weights coming from X_r , $\mathcal{W}^-(X_r) = -\mathcal{W}^+(X_r)$ and $\mathcal{W}^0(X)$ is the set of the zero weight.*

This decomposition shows the interaction between each homogeneous component X_l intervening in $\text{Carr}_{2p,2}(X)$. In length 2, as any weight as its symmetric counterpart, we always have a contribution of X_{p+1} .

290 iii) The third equality follows directly from a computation:

Lemma 11. *A component X_r , $r \geq 2$, produces a resonant letter in $A(X)$ if and only if r is odd. In this case, the letter is unique and given by $n_0 = (\frac{r-1}{2}, \frac{r-1}{2})$.*

Proof. Let $r \geq 2$ be fixed. By Lemma 4 we know the set of letters produced by X_r . The two letters $(-1, r)$ and $(r, -1)$ are never resonant. For the other ones given by $(k-1, r-k)$, one must
295 solve the equation of resonance

$$2k - r - 1 = 0, \quad (36)$$

for $k = 1, \dots, r$. This equation has a unique solution given by

$$k = \frac{r+1}{2}, \quad (37)$$

which is valid, as k must be an integer, only when r is odd. □

This concludes the proof of Lemma 9. □

2.3.2. Explicit computation and the fundamental Lemma

300 We now explicit the quantities $Carr_p(X)$ when X is a real Hamiltonian polynomial vector field.

Let X a polynomial vector field in \mathbb{C}^2 of the form:

$$X = X_{lin} + \sum_{r=2}^d (P_r(x, y)\partial_x + Q_r(x, y)\partial_y), \quad (38)$$

where P_r and Q_r are homogeneous polynomials of degree r .

305 **Lemma 12.** *The complex vector field (38) corresponds to a real vector field if for all $r = 2, \dots, d$, we have*

$$\overline{p_{i,j}} = q_{j,i}, \quad i = 0, \dots, r-1, \quad j = r-i. \quad (39)$$

The proof follows easily from the fact that $\bar{x} = y$ and $Q_r(x, y) = \overline{P_r(x, y)}$ which gives $Q_r(x, y) = \overline{P_r(y, x)}$ for all $r = 2, \dots, d$.

Real Hamiltonian systems satisfy moreover the following conditions:

310 **Lemma 13.** *The complex vector field (38) corresponds to a real Hamiltonian vector fields if conditions (39) are satisfied and moreover if for all $r = 2, \dots, d$, we have*

$$p_{i-1, r-i} = -\frac{r-i+1}{i} \overline{p_{r-i, i-1}}, \quad i = 1, \dots, r. \quad (40)$$

We give some examples of relations between the coefficients in X_2 and X_3 :

Example 2. *For the vector field X_2 defined above, we have :*

$$p_{1,0} = \frac{-1}{2} \overline{p_{0,1}},$$

$$p_{-1,2} = \overline{q_{2,-1}}.$$

For the vector field X_3 , we have :

$$p_{2,0} = \frac{-1}{3} \overline{p_{0,2}},$$

$$p_{1,1} = -\overline{p_{1,1}},$$

$$p_{-1,3} = \overline{q_{3,-1}}.$$

Under the two previous conditions on the coefficients we have:

Lemma 14 (Fundamental Lemma). *Let X be a real Hamiltonian vector fields of the form $X = i(x\partial_x - y\partial_y) + \sum_{j=r}^{2r-1} (P_j(x, y)\partial_x + \overline{P_j(x, y)}\partial_y)$ with $\bar{x} = y$, then :*

$$Carr_{2(r-1)}(X) = p_{r-1, r-1} + i \left(\sum_{k=\lfloor \frac{r+1}{2} \rfloor + 1}^r \frac{r(r+1)}{(r-k+1)^2} |p_{k-1, r-k}|^2 + \frac{r}{r+1} |p_{-1, r}|^2 \right).$$

Proof. The computation of the correction in depth $2(k-1)$ requires to know which operators appear in it and the alphabet. In this depth, there are just the homogeneous differential operators from the polynomial P_r and the resonant letter from the polynomial P_{2r-1} .

The alphabet given by P_r is $A(X_r) = \{(r, -1), (-1, r), (k-1, r-k) \text{ with } k = 1, \dots, r\}$.

We also have the homogeneous differential operators :

$$B_{(-1,r)} = p_{-1,r} y^r \partial_x, B_{(r,-1)} = q_{r,-1} x^r \partial_y, B_{(k-1,r-k)} = x^{k-1} y^{r-k} (p_{k-1,r-k} x \partial_x + q_{k-1,r-k} y \partial_y)$$

with $k \in \{1, \dots, r\}$.

Here the resonant words are :

$$\begin{aligned} &((k-1, r-k), (r-k, k-1)), \\ &((r-k, k-1), (k-1, r-k)), \\ &((-1, r), (r-1)), \\ &((r, -1), (-1, r)). \end{aligned}$$

So the Lie brackets associated, using the real and Hamiltonian conditions, are:

$$\begin{aligned} [B_{(k-1,r-k)}, B_{(r-k,k-1)}] &= \frac{r(r+1)(2k-r-1)}{(r-k+1)^2} |p_{k-1,r-k}|^2 (xy)^{r-1} (x\partial_x - y\partial_y) \\ &= -[B_{(r-k,k-1)}, B_{(k-1,r-k)}], \\ [B_{(-1,r)}, B_{(r,-1)}] &= -r |p_{-1,r}|^2 (xy)^{r-1} (x\partial_x - y\partial_y) = -[B_{(r,-1)}, B_{(-1,r)}]. \end{aligned}$$

The mould of the correction in this resonant word are:

$$\begin{aligned} Carr^{(k-1,r-k) \cdot (r-k,k-1)} &= \frac{i}{2k-r-1}, \\ Carr^{(-1,r) \cdot (r,-1)} &= \frac{-i}{r+1}. \end{aligned}$$

The only resonant letter in P_{2k-1} is $p_{r-1,r-1}$ with the operator $B_{(r-1,r-1)} = p_{r-1,r-1} (xy)^{r-1} (x\partial_x - y\partial_y)$. Using the alternality of the mould $Carr^\bullet$ and the skew-symmetric of the Lie brackets, we get the formula :

$$Carr_{2(k-1)}(X) = p_{r-1,r-1} + i \left(\sum_{k=\lfloor \frac{r+1}{2} \rfloor + 1}^r \frac{r(r+1)}{(r-k+1)^2} |p_{k-1,r-k}|^2 + \frac{r}{r+1} |p_{-1,r}|^2 \right).$$

□

315 2.3.3. Examples of computations

We give examples of computations of the correction.

The quadratic case. We consider the real Hamiltonian vector field :

$$X = X_{lin} + X_2$$

We remind that the alphabet is given by $A(X_2) = \{(1, 0), (0, 1), (2, -1), (-1, 2)\}$ (see Section 2.1.3) and is non resonant.

320 The first none trivial correction term is then in depth 2.

The correction in depth 2 is given by:

$$\begin{aligned} Carr_2(X) &= Carr_{2,2}(X) \\ &= \sum_{\substack{\mathbf{n} \in A^*(X) \\ p(\mathbf{n})=2 \\ \omega(\mathbf{n})=0}} Carr^{\mathbf{n}} B_{\mathbf{n}} \\ &= \sum_{\substack{\mathbf{n} \in A^*(X) \\ p(\mathbf{n})=2 \\ \omega(\mathbf{n})=0}} \frac{1}{\ell(\mathbf{n})} Carr^{\mathbf{n}} B_{[\mathbf{n}]} \\ &= \frac{1}{2} \left(Carr^{(1,0) \cdot (0,1)} B_{[(1,0) \cdot (0,1)]} + Carr^{(0,1) \cdot (1,0)} B_{[(0,1) \cdot (1,0)]} \right. \\ &\quad \left. + Carr^{(2,-1) \cdot (-1,2)} B_{[(2,-1) \cdot (-1,2)]} + Carr^{(-1,2) \cdot (2,-1)} B_{[(-1,2) \cdot (2,-1)]} \right). \end{aligned}$$

By the Fundamental Lemma, we finally have :

$$Carr_2(X) = i \left(6|p_{1,0}|^2 + \frac{2}{3}|p_{-1,2}|^2 \right).$$

The cubic case. We consider the real Hamiltonian vector field :

$$X = X_{lin} + X_2 + X_3$$

with $X_2 = (p_{1,0}x^2 + p_{0,1}xy + p - 1, 2y^2)\partial_x + (q_{2,-1}x^2 + q_{0,1}xy + q_{1,0})\partial_y$ and $X_3 = (p_{2,0}x^3 + p_{1,1}x^2y + p_{0,2}xy^2 + p_{-1,3}y^3)\partial_x + (q_{3,-1}x^3 + q_{2,0}x^2y + q_{1,1}xy^2 + q_{0,2}y^3)\partial_y$.

As above, the first none trivial correction is in depth 2 :

$$Carr_2(X) = Carr_{2,1}(X_3) + Carr_{2,2}(X_2).$$

The only operator which is of depth 2 from X_3 is given by its resonant letter $(1, 1)$ and the operator $B_{1,1} = xy(p_{1,1}x\partial_x + q_{1,1}y\partial_y)$. So, using the previous result on the quadratic case, we have :

$$Carr_2(X) = p_{1,1} + i \left(6|p_{1,0}|^2 + \frac{2}{3}|p_{-1,2}|^2 \right).$$

In depth 4, we have :

$$Carr_4(X) = Carr_{4,2}(X_3, X_3) + Carr_{4,3}(X_3, X_2, X_2) + Carr_{4,4}(X_2, X_2, X_2, X_2).$$

The different $Carr_{i,j}(X)$ are given by :

$$\begin{aligned} Carr_{4,2}(X_3, X_3) &= i(12|p_{2,0}|^2 + \frac{3}{4}|p_{-1,3}|^2), \\ Carr_{4,3}(X_3, X_2, X_2) &= -i(120Im(p_{2,0}\bar{p}_{1,0}^2) + \frac{26}{3}Im(\bar{p}_{-1,3}p_{-1,2}\bar{p}_{1,0}) + 40Im(p_{2,0}p_{-1,2}p_{1,0})) \\ Carr_{4,4}(X_2, X_2, X_2, X_2) &= i\left(-144|p_{1,0}|^4 + 12|p_{1,0}|^2|p_{-1,2}|^2 - \frac{8}{9}|p_{-1,2}|^4 + 40Re(p_{-1,2}p_{0,1}^3)\right). \end{aligned}$$

The quartic case. We consider the real Hamiltonian vector field :

$$X = X_{lin} + X_2 + X_3 + X_4$$

with X_2 and X_3 as above and $X_4 = (p_{3,0}x^4 + p_{2,1}x^3y + p_{1,2}x^2y^2 + p_{0,3}xy^3 + p_{-1,4}y^4)\partial_x + (q_{4,-1}x^4 + q_{3,0}x^3y + q_{2,1}x^2y^2 + q_{1,2}xy^3 + q_{0,3}y^4)\partial_y$.

The correction in depth 2 is the same as the cubic case.

In depth 4, we have :

$$\begin{aligned} Carr_4(X) &= Carr_{4,2}(X_4, X_2) + Carr_{4,2}(X_3, X_3) \\ &\quad + Carr_{4,3}(X_3, X_2, X_2) + Carr_{4,4}(X_2, X_2, X_2, X_2), \end{aligned}$$

where $Carr_{4,2}(X_4, X_2)$ is given by :

$$Carr_{4,2}(X_4, X_2) = i(12Re(p_{2,1}\bar{p}_{1,0}) + 8Re(p_{3,0}p_{-1,2})).$$

Maple program. For the interested readers, we can send some Maple program to compute the correction of a polynomial vector fields. The program can be found in [25].

3. The isochronous center affine variety

325 In this part, we prove that the set of isochronous center is a rational affine variety which is invariant under a non trivial \mathbb{C}^* action. This affine variety is moreover explicitly described. We also give estimates on the growth of the degree of each rational polynomials entering in this description as well as the growth of the rational coefficients.

3.1. Affine variety of isochronous center

330 We consider real vector fields written in complex form as $X = X_{lin} + P(x, y)\partial_x + Q(x, y)\partial_y$ where P and Q are polynomials with coefficients in \mathbb{C} such that $\overline{P(x, y)} = Q(y, x)$. We denote by $N(d)$ the number of independent coefficients defining P and by \mathbf{p} any element of this set. By the reality condition, the coefficient of Q can be deduced from those of P . We then identify the set of complex polynomials of a given degree d with $\mathbb{C}^{N(d)}$, where $N(d)$ is given by $N(d) = \frac{(d-1)(d+4)}{2}$.

335

We denote by \mathcal{L}_d the set of polynomial perturbations (P, Q) of degree d such that X is linearizable. The set \mathcal{L} can be seen as a subset of $\mathbb{C}^{N(d)}$. Precisely, we have :

Theorem 10 (Geometric structure). *For all $d \geq 2$, the set \mathcal{L}_d of isochronous centers is an affine variety over \mathbb{Q} in $\mathbb{C}^{N(d)}$.*

The proof is based on a precise description of the algebraic form of the correction. For all $\mathbf{n} \in A^*(X)$, let us denote by $P(\mathbf{n})$ and $Q(\mathbf{n})$ the coefficients given by Lemma 8 and satisfying

$$[B_{\mathbf{n}}] = (xy)^{\frac{p(\mathbf{n})}{2}} (P(\mathbf{n})x\partial_x + Q(\mathbf{n})y\partial_y).$$

340 We have :

Theorem 11 (Algebraic structure). *For all $p \in \mathbb{N}^*$ the correction term $Carr_{2p}(X)$ has the form*

$$Carr_{2p}(X) = (xy)^p \left[Ca_{2p}(\mathbf{p})x\partial_x + \overline{Ca_{2p}(\mathbf{p})}y\partial_y \right], \quad (41)$$

where

$$Ca_{2p}(\mathbf{p}) = \sum_{i=1}^{2p} \frac{1}{i!} Ca_{2p,i}(\mathbf{p}), \quad (42)$$

with

$$Ca_{2p,i}(\mathbf{p}) := \sum_{\substack{\mathbf{n} \in A^*(X) \\ p(\mathbf{n})=2p, l(\mathbf{n})=i}} Carr^{\mathbf{n}}P(\mathbf{n}), \quad i = 1, \dots, 2p. \quad (43)$$

345 The quantities $Ca_{2p,i}(\mathbf{p})$, $i = 1, \dots, 2p$, are explicit polynomials of degree i in the coefficients of P with coefficients in \mathbb{Q} if i is even and $i\mathbb{Q}$ otherwise. Moreover, these polynomials can be computed algorithmically using recursive formula.

The proof of this theorem is a consequence of two results. First, nested Lie brackets have a very special shapes which can be easily computed. Precisely, we have :

350 **Lemma 15.** *For all $\mathbf{n} \in A^*(X)$, the coefficients $P(\mathbf{n})$ and $Q(\mathbf{n})$ are polynomials in $\mathbb{Z}[\mathbb{C}^{N(d)}]$ of degree $l(\mathbf{n})$ and defined recursively on the length of \mathbf{n} by*

$$\begin{aligned} P(n\mathbf{n}) &= (|\mathbf{n}|^1 - n^1)p_n P(\mathbf{n}) + |\mathbf{n}|^2 q_n P(\mathbf{n}) - n^2 p_n Q(\mathbf{n}), \\ Q(n\mathbf{n}) &= (|\mathbf{n}|^2 - n^2)q_n Q(\mathbf{n}) + |\mathbf{n}|^1 p_n Q(\mathbf{n}) - n^1 q_n P(\mathbf{n}). \end{aligned} \quad (44)$$

where for $\mathbf{n} = n_1 \cdot \dots \cdot n_r \in A^*(X)$, we let $|n|^j = n_1^j + \dots + n_r^j$, $j = 1, 2$ with $n_i = (n_i^1, n_i^2)$.

Second, the correction mould can also be computed by a recursive formula from which we deduce :

355 **Lemma 16.** *For all $\mathbf{n} \in A^*(X)$, the mould $Carr^{\mathbf{n}}$ belongs to \mathbb{Q} if $l(\mathbf{n})$ is odd and $i\mathbb{Q}$ if $l(\mathbf{n})$ is even.*

The proof of Theorem 11 easily follows.

3.1.1. Proof of Theorem 10

Using Theorem 11 and the characterization of isochronous centers given by Theorem 8, the set \mathcal{L}_d is defined by the zero set of an infinite family of polynomials over $\mathbb{C}^{N(d)}$ given by

$$\mathcal{L}_d = \left\{ P \in \mathbb{C}^{N(d)}, Ca_{2p}(X) = 0, p \geq 1 \right\}. \quad (45)$$

360 We define the *ascending chain of ideals* I_k generated by $\langle Ca_{2k}(\mathbf{p}), \dots, Ca_{2k}(\mathbf{p}) \rangle$ in $\mathbb{C}[\mathbf{p}]$. By the *Hilbert Basis Theorem* (see [4], Theorem 4, p.77), there exists an $M(d) \in \mathbb{N}^*$ such that $I_M = I_{M+1} = \dots$. We denote by \mathcal{I}_d the resulting ideal. As a consequence, the set \mathcal{L}_d can be obtained as (see [4], Definition 8, p.81)

$$\mathcal{L}_d = \mathbf{V}(\mathcal{I}_d) = \left\{ \mathbf{p} \in \mathbb{C}^{N(d)} \mid f(\mathbf{p}) = 0 \text{ for all } f \in \mathcal{I}_d \right\}, \quad (46)$$

and corresponds to the affine variety (see [4], Proposition 9 p.81) defined by

$$\mathcal{L}_d = \mathbf{V}(f_1, \dots, f_{s(d)}) = \left\{ \mathbf{p} \in \mathbb{C}^{N(d)} \mid f_i(\mathbf{p}) = 0 \text{ for all } i = 1, \dots, s(d) \right\}, \quad (47)$$

365 where the finite family of polynomials $f_i, i = 1, \dots, s(d)$ is a generating set of \mathcal{I}_d . As the polynomials defining this variety have coefficients in \mathbb{Q} or $i\mathbb{Q}$ this concludes the proof.

Remark 2. *Theorem 10 together with Theorem 11 gives explicit informations on the degree as well as on the growth of the rational coefficients entering in the definition of the affine variety. A natural question is up to which extend these informations can be used to provide a natural upper*
370 *bound on the number of generators for the ideal generating \mathcal{L}_d thanks to a constructive version of the Hilbert basis theorem. This will be explored in another work.*

3.2. \mathbb{C}^* -invariance

The resonant character of the correction has an interesting consequence on the rational algebraic variety of isochronous center. Indeed, let us consider the following action of \mathbb{C}^* :

Definition 7. *Let $\lambda \in \mathbb{C}^*$, we denote by T_λ the map*

$$\begin{aligned} T_\lambda : \mathcal{C}^{N(d)} &\rightarrow \mathcal{C}^{N(d)} \\ p_\bullet &\mapsto \lambda^{\omega(\bullet)} p_\bullet \end{aligned}$$

375 where \bullet is an arbitrary letter.

We extend this action for all monomials $p_{\mathbf{n}} = p_1^{n_1} \dots p_r^{n_r}$, with \mathbf{n} is a word $n_1 \cdot \dots \cdot n_r$, we have:

$$\begin{aligned} T_\lambda(p_{\mathbf{n}}) &= T_\lambda(p_1^{n_1} \dots p_r^{n_r}), \\ &= \lambda^{\omega(n_1) + \dots + \omega(n_r)} p_{\mathbf{n}} \\ &= \lambda^{\omega(\mathbf{n})} p_{\mathbf{n}}. \end{aligned}$$

We denoted by $[B_{\mathbf{n}}] = [B_{n_1, \dots, n_r}] = (xy)^{\frac{p(\mathbf{n})}{2}} (P(\mathbf{n})x\partial_x + Q(\mathbf{n})y\partial_y)$ where, as we have just shown, $P(\mathbf{n})$ and $Q(\mathbf{n})$ are polynomial in the coefficient of B_{n_1}, \dots, B_{n_r} . We have the following lemma which show the \mathbb{C}^* -invariance:

Lemma 17. *For all resonant word \mathbf{n} , we have :*

$$T_\lambda(P(\mathbf{n})) = P(\mathbf{n}) \text{ and } T_\lambda(Q(\mathbf{n})) = Q(\mathbf{n}).$$

Proof. By definition of a resonant word, we have $\omega(\mathbf{n}) = 0$. So :

$$T_\lambda(p_{\mathbf{n}}) = \lambda^{\omega(\mathbf{n})} p_{\mathbf{n}} = \lambda^0 p_{\mathbf{n}} = p_{\mathbf{n}}.$$

□

380 Finally we can generalise this lemma in the following corollary :

Corollary 2. *For all $\lambda \in \mathbb{C}^*$, the algebraic variety \mathcal{L}_d of the isochronous centers is invariant under the action of T_λ .*

Proof. To prove this corollary we just have to remind that only the resonant word contribute to the linearisability. We can conclude by the above lemma. □

385 4. Proof of the main results

4.1. Proof of Theorem 1

Let X be a real Hamiltonian vector field of even degree $2n$ of the form :

$$X = X_{lin} + \sum_{r=2}^{2n} X_r.$$

For each X_r , we can associate its depth as follows :

X_r	Depth
X_2	1
X_3	2
X_3	3
...	...
X_{2n-1}	$2n - 2$
X_{2n}	$2n - 1$

By Theorem 7, we are only interested by the even depth. As a consequence, we look for all possible combinations of arbitrary length which give rise under Lie bracket to an even depth vector field. In the following, we denote by $[X_{k_1}, \dots, X_{k_r}]$ the set of operators that one obtain by nested Lie brackets of homogeneous differential operators B_{n_i} coming from X_{k_i} , $i = 1, \dots, r$. As an example, we have :

X_r and $[X_r, X_{r'}]$	Depth
X_3	2
$[X_2, X_2]$	2
X_5	4
$[X_4, X_2]$	4
$[X_3, X_3]$	4
$[X_3, X_2, X_2]$	4
X_7	6
$[X_6, X_2]$	6
$[X_5, X_3]$	6
$[X_4, X_4]$	6
$[X_5, X_2, X_2]$	6
$[X_4, X_3, X_2]$	6
$[X_4, X_2, X_2, X_2]$	6
$[X_3, X_2, X_2, X_2, X_2]$	6
$[X_2, X_2, X_2, X_2, X_2, X_2]$	6

The correction in depth 2 is given by :

$$Carr_2(X) = Carr_{2,1}(X_3) + Carr_{2,2}(X_2),$$

As the depth is a morphism we have the contribution of the Lie bracket of X_2 , we also have the contribution of the resonant letter of X_3 . By the Fundamental Lemma, the correction is given by

$$Carr_2(X) = p_{1,1} + i \left(6|p_{1,0}|^2 + \frac{2}{3}|p_{-1,2}| \right).$$

395 By the linearisation criterion, we must have $Carr_2(X) = 0$. This implies that

$$\begin{cases} \operatorname{Re}(p_{1,1}) &= 0, \\ -\operatorname{Im}(p_{1,1}) &= 6|p_{1,0}|^2 + \frac{2}{3}|p_{-1,2}|. \end{cases} \quad (48)$$

As X is real and Hamiltonian, the first equation is always satisfied. The second one has only a non trivial solution if and only if $\operatorname{Im}(p_{1,1}) < 0$. The situation when $\operatorname{Im}(p_{1,1}) \geq 0$ leads to two distinct cases. When $\operatorname{Im}(p_{1,1}) = 0$, the Birkhoff sphere reduce to 0 and we obtain $X_2 = 0$. When $\operatorname{Im}(p_{1,1}) > 0$, the equation can not be satisfied and the vector field is then nonisochronous.

400

Assume that $p_{1,1} = 0$ then $X_2 = 0$ and we are reduce to the case

$$X = X_{lin} + X_3 + \cdots + X_r. \quad (49)$$

As $p_{1,1} = 0$, the first non zero term of the correction is $Carr_4(X)$. By the fundamental Lemma, the term $Carr_4(X)$ has exactly the same algebraic structure than the preceding $Carr_2(X) = 0$

case. The role of X_2 is played by X_3 and the role of the resonant term of X_3 is played by the
 405 resonant term of X_5 . Here again, we recover the same dichotomy between the case $Im(p_{2,2}) = 0$
 and $Im(p_{2,2}) > 0$. In the first case, we obtain that $X_3 = 0$ and we are led to the same sit-
 uation as before. Otherwise if $Im(p_{2,2}) > 0$ the term $Carr_4(X)$ can not be zero and we have a
 nonisochronous center.

410 The preceding discussion is representative of the general strategy of proof. Let us assume that
 $p_{j,j} = 0$ for $j = 1, \dots, r - 1$. Then, we prove by induction that $X_2 = \dots = X_{r-1} = 0$. In order to
 finish the proof, two cases must be discussed depending the value of r .

Case 1 : $r < n - 1$. The component X_{2r+1} is non trivial due to the condition $Im(p_{r,r}) > 0$.
 415 By the Fundamental Lemma and the linearisability criterion, we must have

$$Carr_{2r}(X) = p_{r,r} + i \left(\sum_{k=\lfloor \frac{r+1}{2} \rfloor + 1}^r \frac{r(r+1)}{(r-k+1)^2} |p_{k-1,r-k}|^2 + \frac{r}{r-1} |p_{-1,r}|^2 \right) = 0 \quad (50)$$

As $Im(p_{r,r}) > 0$, this equality can not be satisfied and X is nonisochronous.

Case 2 : $r = n - 1$. In this case, we are reduced to an homogeneous perturbation of degree
 $2n$ and the correction is given by

$$Carr_{2n-2}(X) = \sum_{k=\lfloor \frac{r+1}{2} \rfloor + 1}^r \frac{r(r+1)}{(r-k+1)^2} |p_{k-1,r-k}|^2 + \frac{r}{r-1} |p_{-1,r}|^2 = 0$$

As X_{2n} is nontrivial, this equation can not be satisfied and X is nonisochronous.

420 This concludes the proof of the Theorem.

4.2. Proof of Theorem 2

The proof follows the lines of those of Theorem 1. We have to distinguish two cases : k is even
 or odd.

425 **Case 1: k is even.** The vector field X_k does not contain resonant terms. As a consequence,
 its first contribution to the correction appears in depth $2(k - 1)$ corresponding to resonant Lie
 brackets of homogeneous differential operators in X_k of length two. As $l \leq k - 1$, this implies
 that $2l - 1 < 2l \leq 2(k - 1)$. The even component between X_k and X_{2l} will come into play in the
 correction only with a greater depth as $2(k - 1)$ by Lie brackets of length at least two. In the
 430 same way, for odd components, the resonant term will intervene in the correction with a strictly
 smaller depth in length one and the other terms in depth greater than $2(k - 1)$ by a Lie brackets

of length at least two.

As a consequence, the correction term coming from an even k is given by :

$$Carr_{2(k-1)}(X) = i \left(\sum_{k=\lfloor \frac{r+1}{2} \rfloor + 1}^r \frac{r(r+1)}{(r-k+1)^2} |p_{k-1,r-k}|^2 + \frac{r}{r-1} |p_{-1,r}|^2 \right). \quad (51)$$

435 In order to satisfy the linearisability criterion, we must have $Carr_{2(k-1)} = 0$. If the component X_k is non trivial then the system is already non isochronous. Otherwise, we have $X_k = 0$ and we are led to the same problem but with an odd component.

Case 2 : k is odd. In this case the vector field X_k contain a resonant homogeneous operator.

440 Let us write $k = 2m + 1$ then $B_{m,m}$ is of depth $2m$ and weight zero. We have

$$Carr_{k-1}(X) = Carr_{2m}(X) = B_{m,m}. \quad (52)$$

By the linearisability criterion, $Carr_{k-1}(X) = 0$ and the resonant term $B_{m,m}$ in X_k is zero. The contribution of X_k in length 2 follows the same argument as for the even case and we deduce that finally $X_k = 0$.

445 As a consequence, we can prove by induction that in order to be linearisable the components X_k, \dots, X_{2l} must be zero. But, by assumption, we have that X_{2l} is non trivial. As a consequence, the vector field X is necessarily nonisochronous.

4.3. Proof of Theorem 3

The strategy of proof follows those of Theorem 2. The main observation is that there exists
 450 no interactions between each family of vector fields $\{X_k, \dots, X_{2l}\}$ and $\{X_{c_n}, \dots, X_{2(c_n-1)}\}$, $n = 1, \dots, m$. Indeed, let us first analyse the depth of all these objects. We have :

X_r	Depth
X_k	$k - 1$
X_{k+1}	k
\vdots	\vdots
X_{2l-1}	$2l - 2$
X_{2l}	$2l - 1$
0	0
X_{4l}	$4l - 1$
\vdots	\vdots
X_{8l-2}	$8l - 3$
\dots	\dots
X_{c_m}	$c_m - 1$
\vdots	\vdots
$X_{2(c_m-1)}$	$2(c_m - 1) - 1$

Following the same lines as for Theorem 2, we see that the arguments based on the contributions of a given component belonging to $\{X_k, \dots, X_{2l}\}$ are valid. In other words, we easily proved
455 that in order to be linearisable, then one must have $X_k = \dots = X_{2l-2} = 0$. The last argument concerning X_{2l} is also satisfies because the first contribution of X_{2l} to the correction is of length two and depth $4l - 2$ which is not disturbed by terms of the remaining family $\{X_{c_n}, \dots, X_{2(c_n-1)}\}$, $n = 1, \dots, m$ as the minimal contribution of these terms to the correction is in depth $4l - 1$.

460 As a consequence, a vector fields of this type will be linearisable if $X_k = \dots = X_{2l} = 0$.

By the same argument, we see that there exists no interaction between the family $\{X_{c_1}, \dots, X_{2(c_1-1)}\}$ and the remaining one $\{X_{c_n}, \dots, X_{2(c_n-1)}\}$, $n = 2, \dots, m$. We deduce that a linearisable vector field of this type must satisfy $X_{c_1} = \dots = X_{2(c_1-1)}$.

465

By induction, we easily deduce that a vector field of this type is linearisable if and only if all the components vanish. As by assumption we are considering a non trivial vector field, we are leaded to a contradiction and the vector field is necessarily nonisochronous.

5. Conclusion and perspectives

470 5.1. Toward a complete proof of the Jarque-Villadelprat conjecture

Our main results give a very strong support to the Jarque-Villadelprat conjecture. The remaining cases always deal with the role of the resonant term in the deformation of the Birkhoff's

spheres. However, the phenomenon which is working for a quartic perturbation, which was precisely studied by Jarque and Villadelprat using geometrical methods, must applies in the same way
 475 for arbitrary degrees. Indeed, as we have seem in our derivation of the main results, the algebraic structure of the correction during the cancellation process does not change and can be closely investigated. We have then some directions in order to solve completely the Jarque-Villadelprat conjecture :

- Can we prove using an extension of our method the quartic case ?
- 480 • Can we extend the geometrical method of Jarque-Villadelprat in the cases which are let open by our work ?
- Can we prove the remaining cases using other methods ?

We believe that a better understanding of the algebraic structure of the correction will be of importance in order to go further.

485 *5.2. Effective Hilbert basis and the isochronous centers affine variety*

A second aspect of our work is the explicit and algorithmic description of the isochronous center affine variety. As already said, we have informations about the growth of the degree and the coefficients entering in the description of this variety. A natural question is then to look for effective version of the Hilbert basis theorem in order to get some informations about the minimal
 490 number of generators of the ideal. The isochronous centers seem to be more tractable than the usual center. However, it is clear that any advance in this direction will have consequences on the local 16th Hilbert problem. Indeed, the same kind of combinatoric and tools can be used to obtain analogous information for centers of polynomial vector fields (see [8]).

5.3. Isochronicity for complex Hamiltonian systems

495 In [21], the authors study isochronicity of complex Hamiltonian systems when the linear part has for spectrum $(1, -1)$. Our method and results extend naturally to this case and give an explicit and algorithmic description of the isochronous centers affine variety. This will be the subject of a forthcoming work.

Appendix A. Notations

- 500 • $A(X)$ alphabet associated to a vector field X ;
- $A^*(X)$ set of words given by the alphabet $A(X)$;
- $\mathcal{B}(X)$ set of homogeneous differential operator associated to a vector fields X ;

- $Carr(X)$ the correction associated to a vector fields X ;
- $Carr^\bullet$ the mould of the correction;
- 505 • n element of $A(X)$;
- \mathbf{n} element of $A^*(X)$;
- B_n element of $\mathcal{B}(X)$;
- $B_{\mathbf{n}}$ element of $(\mathcal{B}(X), \circ)$;
- ω the weight application;
- 510 • p the depth application;
- $ping$ the application which inverts the two components of a letter;
- ret the application on word which inverts the order of the letters;

Appendix B. Properties of the Correction's mould

The proofs of Lemma 1 and 2 use the definition of the mould of the correction using the
 515 variance formula (12) and also an alternative one based on the knowledge of the mould of a given
 prenormal form that we remind in the next Section.

Appendix B.1. An alternative definition of the correction

We first remind the **composition** of two moulds M^\bullet and N^\bullet (see [5], Definition III.44 p.347):

$$(M^\bullet \circ N^\bullet)^{\mathbf{n}} = \sum_{1 \leq k \leq \ell(\mathbf{n})} \sum_{\omega_1 \dots \omega_k}^* M^{\|\omega_1\| \dots \|\omega_k\|} \times N^{\omega_1} \times \dots \times N^{\omega_k} \quad (\text{B.1})$$

520 where $\sum_{\omega_1 \dots \omega_k}^*$ means the sum on all the decomposition of the word \mathbf{n} in k words. Moreover, $\|w_j\|$
 is a letter obtained by the word ω_j summing all its letter if the alphabet is provided with a law of
 semi-group.

The neutral element for the composition of moulds is denoted by I^\bullet and is defined by $I^{\mathbf{n}} = 1$
 525 if $\ell(\mathbf{n}) = 1$ and $I^{\mathbf{n}} = 0$ for $\ell(\mathbf{n}) \neq 1$.

To prove the different results about the correction, we use another equivalent definition of the
 mould of the correction related to the choice of a given **prenormal form** (see [14], p.267 **Lemma**
3.2) an

$$I^{\mathbf{n}} - Carr^{\mathbf{n}} = \lim_{n \rightarrow +\infty} ((I^\bullet - M^\bullet) \circ r)^{\mathbf{n}} \quad (\text{B.2})$$

530 where $\mathbf{n} = n_1 \cdot \dots \cdot n_r$ and M^\bullet is a mould associated to a given prenormal form.

In the following, we often use the mould $Tram^\bullet$ associated to the **trimmed prenormal form** (see [13], [5] and [24]) and defined by

$$Tram^\bullet = 0, \quad Tram^\mathbf{n} = (Sam^\bullet)^{l(\mathbf{n})}, \quad (\text{B.3})$$

where Sam^\bullet is a mould defined as follows: Let P_r be the family of **Poincaré functions** $\mathcal{P} = \{P_r\}_{r \in \mathbb{N}^*}$ defined by G. Morin in ([24], §.7.9.1 p.114-115) by

$$P_1(z) = \begin{cases} 1 & \text{if } z = 0, \\ 0 & \text{otherwise.} \end{cases} \quad (\text{B.4})$$

and for $r \geq 2$ by

$$P_r(z_1, \dots, z_r) = \begin{cases} \frac{1}{z_1 \dots z_r} \sum_{k=1}^r (-1)^{r-k} \frac{z_k (r-k) - z_{k+1} - \dots - z_r}{(k-1)!(r-k+1)!}, & \text{if } z \in (\mathbb{C}^*)^r, \\ \frac{(-1)^{r-1}}{(i-1)!(r-i)!} \frac{1}{z_1 \dots z_{i-1} z_{i+1} \dots z_r}, & \text{if } z \in S_{r,i}, \\ 0 & \text{otherwise,} \end{cases} \quad (\text{B.5})$$

where $S_{r,i} = (\mathbb{C}^*)^{i-1} \times \{0\} \times (\mathbb{C}^*)^{r-i}$.

The mould Sam^\bullet is defined by (see [24], Lemma 7.9.9., p.115):

$$Sam^\bullet = \begin{cases} 1 & \text{if } n = \emptyset, \\ P_r(\omega(n_1), \dots, \omega(n_r)) & \text{if } \mathbf{n} = n_1 \dots n_r, r \geq 1. \end{cases} \quad (\text{B.6})$$

540 Moreover we have :

$$I^\mathbf{n} - Carr^\mathbf{n} = ((I^\bullet - M^\bullet)^\circ r)^\mathbf{n} = ((I^\bullet - M^\bullet)^\circ r+k)^\mathbf{n} \quad (\text{B.7})$$

where $k \in \mathbb{N}$.

Now, we can prove the Lemma 1 and Lemma 2.

Appendix B.2. Proof of Lemma 1

1) Using the formula (B.2) for the correction via a given prenormal form mould M^\bullet , we obtain for the word $n = \emptyset$:

$$\begin{aligned} I^\emptyset - Carr^\emptyset &= ((I^\bullet - M^\bullet)^{\circ 0})^\emptyset \\ &= I^\emptyset - M^\emptyset, \end{aligned}$$

545 as $I^\emptyset = M^\emptyset = 0$, we deduce that $Carr^\emptyset = 0$.

2) Let $\mathbf{n} = n_1 \cdot \dots \cdot n_r$ be a non resonant word of length r , i.e. $\omega(\mathbf{n}) \neq 0$, then we have :

$$I^{\mathbf{n}} - Carr^{\mathbf{n}} = ((I^{\bullet} - Sam^{\bullet})^{\circ r})^{\mathbf{n}}, \quad (\text{B.8})$$

If \mathbf{n} is a letter i.e $\ell(\mathbf{n}) = 1$, such that $\omega(n) \neq 0$ then

$$\begin{aligned} I^{\mathbf{n}} - Carr^{\mathbf{n}} &= ((I^{\bullet} - M^{\bullet})^{\circ 1})^{\mathbf{n}} \\ &= (I^{\bullet} - M^{\bullet})^{\|\mathbf{n}\|} (I^{\bullet} - M^{\bullet})^{\mathbf{n}}, \end{aligned} \quad (\text{B.9})$$

where $M^{\|\mathbf{n}\|} = M^{\mathbf{n}} = 0$ and $I^{\mathbf{n}} = I^{\|\mathbf{n}\|} = 1$ hence $1 - Carr^{\mathbf{n}} = 1$ and $Carr^{\mathbf{n}} = 0$.

550

If the length of \mathbf{n} is greater than 2, the variance formula gives:

$$\begin{aligned} I^{\mathbf{n}} - Carr^{\mathbf{n}} &= ((I^{\bullet} - M^{\bullet})^{\circ r})^{\mathbf{n}} \\ -Carr^{\mathbf{n}} &= ((I^{\bullet} - M^{\bullet})^{\circ r+1})^{\mathbf{n}} \\ &= ((I^{\bullet} - M^{\bullet}) \circ (I^{\bullet} - M^{\bullet})^{\circ r})^{\mathbf{n}} \\ &= ((I^{\bullet} - M^{\bullet}) \circ (I^{\bullet} - Carr^{\bullet}))^{\mathbf{n}} \\ &= \sum_{1 \leq k \leq r} \sum_{w_1 \dots w_k = \mathbf{n}} (I^{\bullet} - M^{\bullet})^{\|w_1\| \dots \|w_k\|} (I^{\bullet} - Carr^{\bullet})^{w_1} \dots (I^{\bullet} - Carr^{\bullet})^{w_k}. \end{aligned}$$

By induction, we assume the result is true in length $r - 1$. As \mathbf{n} is non resonant, for any given decomposition in k sub-words $\mathbf{n} = w_1 \dots w_k$, there exists at least one of these sub-words which is non resonant, let says w_j of length $l(w_j) < l(\mathbf{n})$ such that $I^{w_j} - Carr^{w_j} = 0$ as long as $l(w_j) \geq 2$.

555

If this sub-word w_j is a letter, then the decomposition is made of at least two terms, i.e. $k \geq 2$, and all the remaining sub-words are either resonant or non resonant. and the quantity $(I^{\bullet} - M^{\bullet})^{\|w_1\| \dots \|w_j\| \dots \|w_k\|}$ reduces to $-M^{\bullet})^{\|w_1\| \dots \|w_j\| \dots \|w_k\|}$. However, the mould M^{\bullet} is zero over non resonant words as it corresponds to a prenormal form, so that $-M^{\bullet})^{\|w_1\| \dots \|w_j\| \dots \|w_k\|} = 0$ and $Carr^{\mathbf{n}} = 0$.

560

3) We prove this result by induction on the length. If $\ell(\mathbf{n}) = 2$, $\mathbf{n} = n_1 \cdot n_2$. If n_1 is resonant, as \mathbf{n} is resonant, n_2 is also resonant. So we have :

$$\begin{aligned} Carr^{\mathbf{n}} &= -(I^{\|\mathbf{n}\|} - Carr^{\|\mathbf{n}\|})(I^{\mathbf{n}} - Carr^{\mathbf{n}}) \\ &\quad - (I^{\|n_1\| \cdot \|n_2\|} - Carr^{\|n_1\| \cdot \|n_2\|})(I^{n_1} - Carr^{n_1})(I^{n_2} - Carr^{n_2}). \end{aligned} \quad (\text{B.10})$$

As $I^{\|\mathbf{n}\|} - Carr^{\|\mathbf{n}\|} = 1 - 1 = 0$ and $I^{n_1} - Carr^{n_1} = 0$, we have $Carr^{\mathbf{n}} = 0$ if $\ell(\mathbf{n}) = 2$.

565

Now we assume that the result is satisfied for all words of length $\leq r - 1$, $r \geq 3$. We consider $\mathbf{n} = n_1 \cdot \dots \cdot n_r$ such that $\omega(\mathbf{n}) = 0$, $l(\mathbf{n}) = r \geq 3$ and at least one letter, let says n_j is resonant.

We have to study the following equality,

$$\begin{aligned}
I^{\mathbf{n}} - Carr^{\mathbf{n}} &= [(I^\bullet - M^\bullet) \circ r]^{\mathbf{n}}, \\
&= \sum_{1 \leq k \leq r} \sum_{w_1 \dots w_k = \mathbf{n}} (I^\bullet - M^\bullet)^{\|w_1\| \dots \|w_k\|} (I^\bullet - Carr^\bullet)^{w_1} \dots (I^\bullet - Carr^\bullet)^{w_k}.
\end{aligned} \tag{B.11}$$

There exists an integer l such that n_j appears in the decomposition of one w_l for $1 \leq l \leq k$.

570 Either $\ell(w_l) = 1$, so $w_l = n_j$ and $I^{w_l} - Carr^{w_l} = 1 - 1 = 0$, or $\ell(w_l) \geq 2$, so by the induction hypothesis $Carr^{w_l} = 0$ and $I^{w_l} = 0$ by definition. As a consequence, we obtain that $Carr^{\mathbf{n}} = 0$.

Appendix B.3. Proof of Lemma 2

Appendix B.3.1. The mould $Carr^\bullet$ in length 1

Let n a letter such that $\omega(n) = 0$. By the above definition,

$$I^n - Carr^n = I^n - M^n,$$

so $Carr^n = M^n$, where M^\bullet is a prenormal form. We can take for example the mould $Tram^\bullet$ (see 575 [5]), which is the mould of the Poincaré-Dulac normal form. So $Carr^n = Tram^n = 1$.

Appendix B.3.2. The mould $Carr^\bullet$ in length 2

Lemma 18. *If $\mathbf{n} \in A(X)^*$ is such that $l(\mathbf{n}) = 2$, $\omega(\mathbf{n}) = 0$, we have the following possibilities:*

1. *If all the letters have zero weight, i.e. $\omega(n_i) = 0$ for all $i = 1, 2, 3$, then $Carr^{\mathbf{n}} = 0$.*
2. *If all the letters have a non zero weight then $Carr^{\mathbf{n}} = 1/\omega(n_1)$.*

580 *Proof.* If $\omega(n_1 n_2) = 0$, we have $\omega(n_1) = -\omega(n_2)$ as ω is a morphism. Using the variance formula, we have :

$$\omega(n_1)Carr^{n_1 \cdot n_2} + Carr^{n_1 + n_2} = Carr^{n_1}Carr^{n_2}. \tag{B.12}$$

As $\omega(n_i) \neq 0$, we have $Carr^{n_1} = Carr^{n_2} = 0$ so that

$$\omega(n_1)Carr^{n_1 \cdot n_2} + Carr^{n_1 + n_2} = 0. \tag{B.13}$$

We then obtain

$$\omega(n_1)Carr^{n_1 \cdot n_2} = -Carr^{n_1 + n_2} = -1. \tag{B.14}$$

As a consequence, we have

$$Carr^{n_1 \cdot n_2} = \frac{-1}{\omega(n_1)}. \tag{B.15}$$

We can also prove this result using the alternative definition :

$$\begin{aligned}
I^{n_1 \cdot n_2} - Carr^{n_1 \cdot n_2} &= ((I^\bullet - M^\bullet) \circ^2)^{n_1 \cdot n_2} \\
&= ((I^\bullet - M^\bullet) \circ (I^\bullet - Carr^\bullet))^{n_1 \cdot n_2} \\
&= (I^\bullet - M^\bullet)^{\|n_1 \cdot n_2\|} (I^\bullet - Carr^\bullet)^{n_1 \cdot n_2} + \\
&+ (I^\bullet - M^\bullet)^{\|n_1\| \cdot \|n_2\|} (I^\bullet - Carr^\bullet)^{n_1} (I^\bullet - Carr^\bullet)^{n_2}.
\end{aligned}$$

As above we will use the mould $Tram^\bullet$ in length 2, $Carr^{n_1 \cdot n_2} = Tram^{n_1 \cdot n_2} = \frac{\omega(n_2) - \omega(n_1)}{\omega(n_1)\omega(n_2)}$. As $\omega(n_1) = -\omega(n_2)$, we finally have :

$$Carr^{n_1 \cdot n_2} = \frac{1}{\omega(n_1)}.$$

As we can see the result differs by a multiplication by -1 . It is due to the fact in [13] and [14], the nested Lie brackets are taken in this form $[B_{n_1 \dots n_r}] = [B_{n_r}, [B_{n_{r-1}}, [\dots [B_{n_2}, B_{n_1}] \dots]]$ whereas in [7] and [5] we consider $[B_{n_1 \dots n_r}] = [\dots [B_{n_1}, B_{n_2}], \dots], B_{n_{r-1}}, B_{n_r}]$. And we have the relation:

$$[B_{n_r}, [B_{n_{r-1}}, [\dots [B_{n_2}, B_{n_1}] \dots]] = (-1)^{r+1} [B_{n_1}, B_{n_2}], \dots], B_{n_{r-1}}, B_{n_r}].$$

585

□

Appendix B.3.3. The mould $Carr^\bullet$ in length 3

Using the variance formula we obtain for a word $\mathbf{n} = n_1 n_2 n_3$ of length 3 the following equality:

$$\omega(n_1)Carr^{n_1 \cdot n_2 \cdot n_3} + Carr^{(n_1+n_2) \cdot n_3} = Carr^{n_1 \cdot n_3}Carr^{n_2} + Carr^{n_1}Carr^{n_2 \cdot n_3}. \quad (\text{B.16})$$

One can prove the following lemma:

590 **Lemma 19.** *If $\mathbf{n} \in A(X)^*$ is such that $l(\mathbf{n}) = 3$, $\omega(\mathbf{n}) = 0$, we have the following possibilities:*

1. *If all the letters have zero weight, i.e. $\omega(n_i) = 0$ for all $i = 1, 2, 3$, then $Carr^{\mathbf{n}} = 0$.*
2. *If there exists a letter $a \in \{n_1, n_2, n_3\}$ such that $\omega(a) = 0$ and $\omega(b) \neq 0$ for $b \in \{n_1, n_2, n_3\} \setminus \{a\}$, then $Carr^{\mathbf{n}} = 0$.*
3. *If all the letters have a non zero weight, i.e. $\omega(n_i) \neq 0$, $i = 1, 2, 3$, then*

$$Carr^{n_1 \cdot n_2 \cdot n_3} = \frac{1}{\omega(n_1)(\omega(n_1) + \omega(n_2))}. \quad (\text{B.17})$$

595 *Proof.* 1) This case follows from Lemma ??.

2) Let us take for example $a = n_2$ such that $\omega(n_2) = 0$, then $\omega(n_1 n_3) = 0$ and $\omega(n_3) = -\omega(n_1)$. As a consequence, using (B.16) and the fact that $Carr^{n_2} = 1$, we obtain

$$\omega(n_1)Carr^{n_1 \cdot n_2 \cdot n_3} + Carr^{(n_1+n_2) \cdot n_3} = Carr^{n_1 \cdot n_3} + Carr^{n_1}Carr^{n_2 \cdot n_3}. \quad (\text{B.18})$$

Let us assume that n_1 is such that $\omega(n_1) \neq 0$, then $Carr^{n_1} = 0$ and the previous equality reduces
600 to

$$\omega(n_1)Carr^{n_1 \cdot n_2 \cdot n_3} = Carr^{n_1 \cdot n_3} - Carr^{(n_1+n_2) \cdot n_3}. \quad (\text{B.19})$$

Using the expression of the mould $Carr^\bullet$ in length 2, we have

$$Carr^{n_1 \cdot n_3} = -1/\omega(n_1), \quad Carr^{(n_1+n_2) \cdot n_3} = -1/\omega(n_1 + n_2) = -1/\omega(n_1). \quad (\text{B.20})$$

We deduce easily that

$$\omega(n_1)Carr^{n_1 \cdot n_2 \cdot n_3} = 0, \quad (\text{B.21})$$

and as a consequence $Carr^{n_1 \cdot n_2 \cdot n_3} = 0$.

605 The same computations prove that $Carr^{\mathbf{n}} = 0$ if $\omega(n_3) = 0$ and $\omega(n_1) \neq 0$.

The remaining case, i.e. $\omega(n_1) = 0$ and $\omega(n_2) \neq 0$, can be deduced from the alternality of the mould $Carr^\bullet$ which induces the following equality (see [5], §.4.1 p.331):

$$Carr^{n_1 n_2 n_3} + Carr^{n_2 n_1 n_3} + Carr^{n_2 n_3 n_1} = 0. \quad (\text{B.22})$$

610 The second and third terms are zero thanks to our previous computations. As a consequence, we obtain $Carr^{n_1 n_2 n_3} = 0$ if $\omega(\mathbf{n}) = 0$, $\omega(n_1) = 0$ and $\omega(n_2) \neq 0$.

3) From now on, let us assume that all the $\omega(n_i)$ are non zero. Then using formula (B.16) and the fact that $Carr^{n_1} = 0$ and $Carr^{n_2} = 0$, we obtain

$$\omega(n_1)Carr^{n_1 \cdot n_2 \cdot n_3} + Carr^{(n_1+n_2) \cdot n_3} = 0. \quad (\text{B.23})$$

Using the expression of $Carr^{(n_1+n_2) \cdot n_3}$ we deduce that

$$Carr^{n_1 \cdot n_2 \cdot n_3} = \frac{1}{\omega(n_1)(\omega(n_1) + \omega(n_2))}. \quad (\text{B.24})$$

615 As in length 2, we can give an alternative proof using the mould $Tram^\bullet$ to compute the correction :

$$\begin{aligned} I^{n_1 \cdot n_2 \cdot n_3} - Carr^{n_1 \cdot n_2 \cdot n_3} &= ((I^\bullet - Tram^\bullet)^\circ 3)^{n_1 \cdot n_2 \cdot n_3} \\ &= ((I^\bullet - Tram^\bullet)^\circ 4)^{n_1 \cdot n_2 \cdot n_3} \\ &= (I^\bullet - Tram^\bullet) \circ (I^\bullet - Carr^\bullet)^{n_1 \cdot n_2 \cdot n_3} \\ &= -Tram^{n_1 \cdot n_2 \cdot n_3}. \end{aligned} \quad (\text{B.25})$$

So $Carr^{n_1 \cdot n_2 \cdot n_3} = Tram^{n_1 \cdot n_2 \cdot n_3} = \frac{1}{\omega(n_1)(\omega(n_1) + \omega(n_2))}$. □

Appendix C. Technical results

Appendix C.1. Proof of Theorem 11

620 For all $i = 1, \dots, 2p$, we have using the Theorem of projection

$$\begin{aligned} Carr_{2p,i}(X) &= \frac{1}{i!} \sum_{\substack{\mathbf{n} \in A^*(X) \\ p(\mathbf{n})=2p, l(\mathbf{n})=i}} Carr^{\mathbf{n}}[B_{\mathbf{n}}], \\ &= \frac{1}{i!} \sum_{\substack{\mathbf{n} \in A^*(X) \\ p(\mathbf{n})=2p, l(\mathbf{n})=i}} Carr^{\mathbf{n}}(xy)^{\frac{p(\mathbf{n})}{2}} (P(\mathbf{n})x\partial_x + Q(\mathbf{n})y\partial_y), \\ &= \frac{1}{i!} (xy)^p \sum_{\substack{\mathbf{n} \in A^*(X) \\ p(\mathbf{n})=2p, l(\mathbf{n})=i}} Carr^{\mathbf{n}}(P(\mathbf{n})x\partial_x + Q(\mathbf{n})y\partial_y). \end{aligned} \quad (\text{C.1})$$

By definition of $Ca_{2p,i}(X)$ we obtain

$$Carr_{2p,i}(X) = \frac{1}{i!}(xy)^p \left[Ca_{2p,i}(X)x\partial_x + \overline{Ca_{2p,i}(X)}y\partial_y \right]. \quad (C.2)$$

Appendix C.2. Proof of Lemma 15

Let $B_n = B_{(n^1, n^2)} = x^{n^1}y^{n^2}(p_n x\partial_x + q_n y\partial_y)$ and $B_m = B_{(m^1, m^2)} = x^{m^1}y^{m^2}(p_m x\partial_x + q_m y\partial_y)$. Then, the Lie brackets of B_n and B_m is :

$$\begin{aligned} [B_{nm}] &:= [B_n, B_m] = x^{n^1+m^1}y^{n^2+m^2}(P_{n,m}x\partial_x + Q_{n,m}y\partial_y), \\ &= x^{|nm|^1}y^{|nm|^2}(P_{n,m}x\partial_x + Q_{n,m}y\partial_y) \end{aligned}$$

where $P_{n,m}$ and $Q_{n,m}$ are polynomials in the coefficients of B_n and B_m , precisely :

$$\begin{aligned} P_{n,m} &= (m^1 - n^1)p_n p_m + m^2 q_n p_m - n^2 p_n q_m, \\ Q_{n,m} &= (m^2 - n^2)q_n q_m + m^1 p_n q_m - n^1 q_n p_m. \end{aligned}$$

We easily prove by induction that all the Lie bracket in any length are of the above form.

References

625 [1] A. Cima, F. Maosas, J. Villadelprat, Isochronicity for several classes of Hamiltonian systems. J. Differential Equations 157 (1999), no. 2, 373-413.

[2] J. Chavarriga, M. Sabatini, A survey of isochronous centers, Qual. Theory Dyn. Syst. 1 (1999), 1-70.

[3] Y. Chen, V. G. Romanovski, W. Zhang, Non-isochronicity of the center for polynomial Hamil-
630 tonian systems with even degree nonlinearities, Nonlinear Analysis 68 (2008), p. 2769-2778.

[4] D.A. Cox, J. Little, D. O’Shea, *Ideals, Varieties, and Algorithms*, An introduction to computational algebraic geometry and commutative algebra, Fourth Edition, Springer, 2015.

[5] J. Cresson, *Calcul Moulien*, Annales de la Faculté des Sciences de Toulouse Mathématiques Vol. XVIII, no. 2, 2009, pp. 307-395.

635 [6] J. Cresson, Obstruction à la linéarisation des champs de vecteurs polynomiaux. Canad. Math. Bull. 45 (2002), no. 3, 355-363.

[7] J. Cresson, J. Raissy, About the Trimmed and the Poincaré-Dulac Normal Form of Diffeomorphisms, Bollettino U.M.I (9) V (2012), 55-80.

[8] J. Cresson, B. Schuman, Formes normales et problème du centre, Bulletin des Sciences Math-
640 ématiques 125, no.3, 235-252, 2001.

- [9] Ecalle J., Singularités non abordables par la géométrie, Ann. Inst. Fourier, 42 (1-2), 1992, 73-164.
- [10] Ecalle J., Les fonctions résurgentes, Vol.1, Les algèbres de fonctions résurgentes, Publications Mathématiques d'Orsay, (1981).
- 645 [11] Ecalle J., Les fonctions résurgentes, Vol.3, L'équation du pont et la classification analytique des objets locaux, Publications Mathématiques d'Orsay, (1985).
- [12] Ecalle J., Schlomiuk D., The nilpotent and distinguished form of resonant vector fields or diffeomorphisms, Ann. Inst. Fourier 43 5 (1993) 1407-1483.
- [13] J. Ecalle, B. Vallet, Prenormalization, correction, and linearization of resonant vector fields
650 or diffeomorphisms, Prepublication d'Orsay (1995) 101.p.
- [14] J. Ecalle, B. Vallet, Correction an linearization of resonant vector fields and diffeomorphisms, Math. Z. 229 (1998) 249-318.
- [15] L. H. Eliasson, Hamiltonian systems with linear normal form near an invariant torus, in Nonlinear Dynamics Turchetti Ed., World Scientific,, Singapore, 1989.
- 655 [16] J-P. Francoise, R. Pons, Les conditions du centre pour un champ de vecteurs quadratiques, CRAS 318 (1994), p. 909-912.
- [17] J-P. Francoise, *Géométrie analytique et systèmes dynamiques*, PUF, 1995.
- [18] G.. Gallavotti, A criterion of integrability for perturbed nonresonant harmonic oscillators, "wick ordering" of the perturbations in classical mechanics and invariance of the frequency
660 spectrum, Communication in Mathematical Physics Vol. 87, p.365-383, 1982.
- [19] L. Gavrilov, Isochronicity of plane polynomial Hamiltonian systems, Nonlinearity 10 (1997) 433-448.
- [20] X. Jarque, J. Villadelprat, Nonexistence of isochronous centers in planar polynomial Hamiltonian systems of degree four. J. Differential Equations 180 (2002), no. 2, 334-373.
- 665 [21] J. Llibre, V. G. Romanovski. Isochronicity and linearizability of planar polynomial Hamiltonian systems. J. Differential Equations 259 (2015), no. 5, 1649-1662.
- [22] W.S. Loud, Behaviour of the period of solutions of certain plane autonomous systems near centers, Contrib. Differential Equations 3 (1964) 21-36.
- [23] P. Mardesic, C. Rousseau, B. Toni, Linearisation of isochronous centers, J. Differential equa-
670 tions 121 (1995), 67-108.

- [24] G. Morin, *Calcul moulien et théorie des formes normales classiques et renormalisés*, Ph.D., Observatoire de Paris, 2010.
- [25] J. Palafox, *Calcul moulien, arborification, symétries et applications*, Ph.D. Université de Pau et des Pays de l'Adour, Juin 2018.
- 675 [26] C. Reutenauer, *Free Lie algebras*, London Math. Soc. Monographs, new series 7 (1993).
- [27] B. Schuman, Sur la forme normale de Birkhoff et les centres isochrones. C. R. Acad. Sci. Paris Sér. I Math. 322 (1996), no. 1, 21-24.
- [28] B. Schuman, Une classe d'hamiltoniens polynomiaux isochrones. Canad. Math. Bull. 44 (2001), no. 3, 323-334.
- 680 [29] J-P. Serre, *Lie algebras and Lie groups*, W.C. Benjamin Inc (1965).
- [30] M. Urabe, Potential forces which yield periodic motions of a fixed period, J. Math. Mech. 10 (1961) 569-578.