



HAL
open science

Convergence rate of inertial proximal algorithms with general extrapolation and proximal coefficients

Hedy Attouch, Zaki Chbani, Hassan Riahi

► **To cite this version:**

Hedy Attouch, Zaki Chbani, Hassan Riahi. Convergence rate of inertial proximal algorithms with general extrapolation and proximal coefficients. 2019. hal-02021322

HAL Id: hal-02021322

<https://hal.science/hal-02021322v1>

Preprint submitted on 15 Feb 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

CONVERGENCE RATE OF INERTIAL PROXIMAL ALGORITHMS WITH GENERAL EXTRAPOLATION AND PROXIMAL COEFFICIENTS

HEDY ATTOUCH, ZAKI CHBANI, AND HASSAN RIAHI

Abstract. In a Hilbert space setting, with the aim of developing rapid methods for nonsmooth convex optimization, we analyze the convergence rate of the inertial proximal algorithms. These algorithms involve both extrapolation coefficients (including Nesterov acceleration method) and proximal coefficients in a general form. They can be interpreted as the discrete time version of inertial continuous gradient systems with general damping and time scale coefficients. Based on the proper setting of these parameters, we show the fast convergence of values and the convergence of iterates. In doing so, we provide an overview of this class of algorithms. Our study complements the previous Attouch-Cabot paper (SIOPT, 2018) by introducing into the algorithm time scaling aspects, and sheds new light on the Güler seminal papers on the convergence rate of the accelerated proximal methods for convex optimization.

Key words: Inertial proximal algorithms; general extrapolation coefficient; Lyapunov analysis; Nesterov accelerated gradient method; nonsmooth convex optimization; time rescaling.

AMS subject classification. 37N40, 46N10, 49M30, 65B99, 65K05, 65K10, 90B50, 90C25.

1. INTRODUCTION

Throughout the paper, \mathcal{H} is a real Hilbert space endowed with the scalar product $\langle \cdot, \cdot \rangle$ and the corresponding norm $\|\cdot\|$. The function $\Phi : \mathcal{H} \rightarrow \mathbb{R} \cup \{+\infty\}$, which is to minimize, is supposed to be convex lower semicontinuous and proper, and such that $\operatorname{argmin}_{\mathcal{H}} \Phi \neq \emptyset$. We will analyze the convergence properties of the Inertial Proximal algorithm

$$(1) \quad (\text{IP})_{\alpha_k, \beta_k} \quad \begin{cases} y_k = x_k + \alpha_k(x_k - x_{k-1}) \\ x_{k+1} = \operatorname{prox}_{\beta_k \Phi}(y_k), \end{cases}$$

where the sequences (α_k) and (β_k) play the role of parameters. As a standing assumption, we assume that, for every $k \geq 1$, α_k is non-negative and β_k is positive. For a judicious choice of α_k and β_k , we will obtain fast convergence of the values and convergence of the iterates. In doing so, we will extend previous studies to the case of inertial proximal methods with general coefficients, including large proximal steps β_k . We recall that, for $\beta > 0$, the proximal mapping $\operatorname{prox}_{\beta \Phi} : \mathcal{H} \rightarrow \mathcal{H}$ is defined by: for every $x \in \mathcal{H}$,

$$\operatorname{prox}_{\beta \Phi}(x) = \operatorname{argmin}_{\xi \in \mathcal{H}} \left\{ \beta \Phi(\xi) + \frac{1}{2} \|x - \xi\|^2 \right\}.$$

One can consult [15, 28, 29, 32], for a recent account on the proximal methods, that play a central role in non-smooth optimization as a basic block of many splitting algorithms. Proximal methods have a close relationship with gradient-type continuous evolution systems, from which they can be derived by implicit discretization. As such, they retain their convergence properties (see [30] for first-order evolution systems). Precisely, to guide our study, when Φ is continuously differentiable, we will use the link between the $(\text{IP})_{\alpha_k, \beta_k}$ proximal algorithm and the continuous second-order evolution equation

$$(\text{IGS})_{\gamma, \beta} \quad \ddot{x}(t) + \gamma(t)\dot{x}(t) + \beta(t)\nabla\Phi(x(t)) = 0.$$

In $(\text{IGS})_{\gamma, \beta}$, $\gamma(t)$ is a positive damping coefficient, and $\beta(t)$ is a time scale coefficient. The inertial proximal algorithm $(\text{IP})_{\alpha_k, \beta_k}$ can be interpreted as an implicit discretization of $(\text{IGS})_{\gamma, \beta}$, which gives the following dynamic interpretation of the parameters (α_k) and (β_k) :

- The sequence (α_k) of non-negative extrapolation coefficients captures the inertial effect. The property $\alpha_k \rightarrow 1$, which plays a central role in the Nesterov acceleration method, is associated with the vanishing damping property $\gamma(t) \rightarrow 0$, as $t \rightarrow +\infty$.

- The sequence (β_k) of positive proximal coefficients takes into account the temporal scale effects. The coefficient β_k in $(\text{IP})_{\alpha_k, \beta_k}$ is the discrete temporal version of $\beta(t)$. The fast time parametrization of the orbits (respectively iterates) results from the property $\beta(t) \rightarrow +\infty$ as $t \rightarrow +\infty$ (respectively $\beta_k \rightarrow +\infty$ as $k \rightarrow +\infty$). It plays a key role in obtaining fast optimization properties.

1.1. Presentation of the results. Historical aspects. Recall two important instances of $(\text{IP})_{\alpha_k, \beta_k}$ already studied, and which constitute special cases of our study:

1. The case $\beta_k \equiv \beta > 0$ where the proximal parameter is fixed, and α_k is general, is written as follows

$$(2) \quad (\text{IP})_{\alpha_k, \beta} \quad \begin{cases} y_k = x_k + \alpha_k(x_k - x_{k-1}) \\ x_{k+1} = \text{prox}_{\beta\Phi}(y_k). \end{cases}$$

It was analyzed by Attouch-Cabot [5], then by Attouch-Cabot-Chbani-Riahi [6] in the presence of perturbations, errors. The following result gives the convergence rate of values in this case. It is expressed with the help of the sequence (t_k) that is related to (α_k) by the formula $t_k := 1 + \sum_{i=k}^{+\infty} \prod_{j=k}^i \alpha_j$. The inverse mapping takes the simpler form $\alpha_k = \frac{t_k - 1}{t_{k+1}}$.

Theorem 1.1 (Attouch-Cabot [5]). *Suppose that the sequence (α_k) satisfies the following properties (K_0) and (K_1) .*

$$(K_0) \quad \forall k \geq 1, \quad \sum_{i=k}^{+\infty} \prod_{j=k}^i \alpha_j < +\infty,$$

$$(K_1) \quad \forall k \geq 1, \quad t_{k+1}^2 - t_k^2 \leq t_{k+1}.$$

Then, for any sequence (x_k) generated by the algorithm $(\text{IP})_{\alpha_k, \beta}$

$$\Phi(x_k) - \min_{\mathcal{H}} \Phi = \mathcal{O}\left(\frac{1}{t_k^2}\right) \quad \text{as } k \rightarrow +\infty.$$

This result comes as a result of a rich history. In a seminal paper [26], Nesterov proposed the extrapolation coefficient $\alpha_k = \frac{t_k - 1}{t_{k+1}}$ with $t_1 = 1$ and $t_{k+1} = \frac{\sqrt{4t_k^2 + 1} + 1}{2}$. This choice corresponds to taking equality in (K_1) , and leads to an increasing sequence (α_k) which behaves like $1 - \frac{3}{k}$ as $k \rightarrow +\infty$. Then t_k is of order k , and the corresponding convergence rate for values is $\mathcal{O}(\frac{1}{k^2})$. This result has been extended by Beck-Teboulle to structured convex minimization problems in [16], that's the FISTA inertial proximal-gradient algorithm. Since the introduction of Nesterov's scheme, much progress has been made in the development of first-order accelerated methods, to name a few [15, 16, 18, 19, 22, 23, 24, 28, 32, 37, 35]. Recently, a special attention has been devoted to the case $\alpha_k = 1 - \frac{\alpha}{k}$, where $\alpha > 0$ (or an equivalent expression), see [8, 11, 18, 37]. Given $x_0, x_1 \in \mathcal{H}$, for $k \geq 1$ the algorithm writes

$$(3) \quad \begin{cases} y_k = x_k + \left(1 - \frac{\alpha}{k}\right)(x_k - x_{k-1}) \\ x_{k+1} = \text{prox}_{\beta\Phi}(y_k). \end{cases}$$

For $\alpha = 3$ we recover a first-order approximation of the original choice of Nesterov. The great novelty of the algorithm (3) is that, while keeping the same computational complexity as in the case $\alpha = 3$, taking $\alpha > 3$ offers many advantages. First, it ensures the convergence of the sequences (x_k) , as proved by Chambolle-Dossal [18], see [8] for complementary results. Let us recall that the convergence of the sequences generated by FISTA has not been established so far. This is a puzzling question in the study of numerical optimization methods. Second, as proved by Attouch-Peypouquet in [11], it provides the better convergence rate of values

$$\Phi(x_k) - \min_{\mathcal{H}} \Phi = o\left(\frac{1}{k^2}\right) \quad \text{as } k \rightarrow +\infty.$$

When taking $\alpha_k = 1 - \frac{\alpha}{k}$, $\alpha = 3$ appears as a critical value. The subcritical case $\alpha < 3$ has been recently considered by Apidopoulos-Aujol-Dossal [3] and Attouch-Chbani-Riahi [9] with the convergence rate of values $\mathcal{O}\left(\frac{1}{k^{\frac{2\alpha}{3}}}\right)$. Theorem 1.1 allows to cover all these situations (except the subcritical case), and many others including the case $\alpha_k = 1 - \frac{c}{k^r}$ with $0 \leq r \leq 1$.

2. The case β_k general, and $\alpha_k = 1 - \frac{\alpha}{k}$, is written as follows

$$(4) \quad (\text{IP})_{\alpha, \beta_k} \quad \begin{cases} y_k = x_k + \left(1 - \frac{\alpha}{k}\right)(x_k - x_{k-1}) \\ x_{k+1} = \text{prox}_{\beta_k\Phi}(y_k). \end{cases}$$

The following result gives the convergence rate of values for the algorithm (4). It was obtained by the authors in [7, Theorem 7.1].

Theorem 1.2 (Attouch-Chbani-Riahi [7]). *Take $\alpha \geq 1$. Suppose that the sequence (β_k) satisfies the growth condition: there exists $k_1 \in \mathbb{N}$ such that for all $k \geq k_1$*

$$(H_\beta) \quad \beta_{k+1} \leq \frac{k^2}{(k+1)(k+2-\alpha)} \beta_k.$$

Then, for any sequence (x_k) generated by the algorithm $(\text{IP})_{\alpha, \beta_k}$, we have

$$\Phi(x_k) - \min_{\mathcal{H}} \Phi = \mathcal{O}\left(\frac{1}{k^2 \beta_k}\right) \quad \text{as } k \rightarrow +\infty.$$

In this result, the introduction of the scaling factor β_k permits to pass from the convergence rate $\mathcal{O}\left(\frac{1}{k^2}\right)$ to $\mathcal{O}\left(\frac{1}{k^2 \beta_k}\right)$. When taking $\beta_k = k^\delta$, condition (H_β) is satisfied for $\delta < \alpha - 3$. In this case, for a given $\delta > 0$, by taking α large enough, namely $\alpha > 3 + \delta$, we can improve the convergence rate of values from $\mathcal{O}\left(\frac{1}{k^2}\right)$ to $\mathcal{O}\left(\frac{1}{k^{2+\delta}}\right)$. This is a clear acceleration effect.

In Theorem 2.3 we will prove the following result, which includes the two studies above (it is formulated below in a condensed form).

Theorem 1.3. *Suppose that the sequence (α_k) satisfies (K_0) . Let (t_k) be the sequence defined by $\alpha_k = \frac{t_k - 1}{t_{k+1}}$. Suppose that the sequences (α_k) and (β_k) satisfy jointly*

$$(K_{1, \alpha_k, \beta_k}) \quad t_{k+1}^2 \beta_k - t_k^2 \beta_{k-1} - t_{k+1} \beta_k \leq 0 \quad \text{for every } k \geq 1.$$

Let (x_k) be a sequence generated by algorithm $(\text{IP})_{\alpha_k, \beta_k}$. Then,

$$\Phi(x_k) - \min_{\mathcal{H}} \Phi = \mathcal{O}\left(\frac{1}{t_k^2 \beta_{k-1}}\right) \quad \text{as } k \rightarrow +\infty.$$

This result permits to better understand the joint tuning of the extrapolation parameter α_k and the scaling parameter β_k in $(\text{IP})_{\alpha_k, \beta_k}$ which provides fast convergence of the values. Our study includes Güler's accelerated proximal algorithm, and provides new insight on this result. In Güler's algorithm, the damping coefficient is written $\alpha_k = g_k \left(\frac{1}{g_{k-1}} - 1\right)$, which allows to formulate the condition $(K_{1, \alpha_k, \beta_k})$ in a simpler form, directly on the data. Based on this approach, and considering various temporal discretizations, our dynamic approach with general damping and scaling coefficients provides a family of inertial proximal algorithms with fast convergence properties.

1.2. Link with inertial gradient systems and time scaling.

1.2.1. *Link with inertial gradient systems.* First verify that the inertial proximal algorithm $(\text{IP})_{\alpha_k, \beta_k}$ can be interpreted as a discretized version in time of the damped inertial dynamic

$$(\text{IGS})_{\gamma, \beta} \quad \ddot{x}(t) + \gamma(t)\dot{x}(t) + \beta(t)\nabla\Phi(x(t)) = 0.$$

Take a fixed time step $h > 0$, and set $\tau_k = kh$, $x_k = x(\tau_k)$. Implicit discretization of $(\text{IGS})_{\gamma, \beta}$ with centered second-order variation, gives

$$(5) \quad \frac{1}{h^2}(x_{k+1} - 2x_k + x_{k-1}) + \frac{\gamma(kh)}{h}(x_k - x_{k-1}) + \beta(kh)\nabla\Phi(x_{k+1}) = 0.$$

Hence, $x_{k+1} + h^2\beta(kh)\nabla\Phi(x_{k+1}) = x_k + (1 - h\gamma(kh))(x_k - x_{k-1})$, which gives the algorithm $(\text{IP})_{\alpha_k, \beta_k}$

$$\begin{cases} y_k &= x_k + (1 - h\gamma(kh))(x_k - x_{k-1}) \\ x_{k+1} &= \text{prox}_{h^2\beta(kh)\Phi}(y_k), \end{cases}$$

with extrapolation coefficient $\alpha_k = 1 - h\gamma(kh)$, and prox. parameter $h^2\beta(kh)$.

1.2.2. *From Polyak heavy ball to vanishing damping.* Take $\beta(\cdot) \equiv 1$ and discuss the role of the time-dependent parameter $\gamma(t)$ in the continuous dynamic $(\text{IGS})_{\gamma, 1}$. In the context of mechanics, $\gamma(t)$ is a viscous friction coefficient. Taking $\gamma(t)$ positive causes the dissipation of the global energy of the system. This plays a central role in its stabilization and optimization properties. The introduction of inertial dynamics (second-order in time) in optimization comes with B. Polyak [31] who, in 1964, introduced the so-called heavy ball with friction method, which corresponds to a constant friction coefficient $\gamma(t) \equiv \gamma_0 > 0$ in $(\text{IGS})_{\gamma, 1}$. It turns out that in this model, the friction is important and neutralizes the inertial effect, which prevents obtaining fast optimization methods. Indeed, taking the damping parameter $\gamma(t)$ which tends to zero as $t \rightarrow +\infty$ (vanishing damping) is a key property for obtaining fast optimization methods. A decisive step in this direction was obtained by Su, Boyd, and Candès [37], who showed that the Nesterov acceleration method can be obtained as a discretization

of the dynamical system $(\text{IGS})_{\gamma,1}$ with $\gamma(t) = \frac{3}{t}$. Recently, Attouch-Chbani-Peypouquet-Redont [8] and May [25] showed convergence of the trajectories of the $(\text{IGS})_{\gamma}$ system with $\gamma(t) = \frac{\alpha}{t}$ and $\alpha > 3$

$$(6) \quad (\text{AVD})_{\alpha} \quad \ddot{x}(t) + \frac{\alpha}{t}\dot{x}(t) + \nabla\Phi(x(t)) = 0.$$

They also obtained the improved convergence rate $\Phi(x(t)) - \min_{\mathcal{H}} \Phi = o(\frac{1}{t^2})$ as $t \rightarrow +\infty$. On the basis of the correspondence $\alpha_k = 1 - h\gamma_k$ (where h is the discrete time step, and $\gamma_k = \gamma(kh)$), the discrete version of the vanishing damping property is that the sequence (α_k) tends to one from below as $k \rightarrow +\infty$. This makes the link with the historical aspects and the recent developments based on Nesterov acceleration method described in the previous section. For recent developments concerning the rich relations between inertial gradient systems with time-dependent friction and the inertial proximal-based algorithms, one can consult [2], [5], [7], [8], [14], [17], [37], [36].

1.2.3. *Link with time scaling.* The time-dependent parameter $\beta(t)$ comes from the time scaling of the dynamic. Let's illustrate it in the following model situation. Start from the $(\text{AVD})_{\alpha}$ system with $\gamma(t) = \frac{\alpha}{t}$ and $\alpha \geq 3$. Given a trajectory $x(\cdot)$ of $(\text{AVD})_{\alpha}$, we know that (see [4], [8], [37])

$$(7) \quad \Phi(x(t)) - \min_{\mathcal{H}} \Phi = \mathcal{O}\left(\frac{1}{t^2}\right).$$

Let's make the following change of time variable in $(\text{AVD})_{\alpha}$: $t = s^p$, where p is a positive parameter. Set $y(s) := x(s^p)$. By the derivation chain rule, we have

$$(8) \quad \ddot{y}(s) + \frac{\alpha p}{s}\dot{y}(s) + p^2 s^{2(p-1)}\nabla\Phi(y(s)) = 0,$$

where $\alpha_p = 1 + (\alpha - 1)p$. The convergence rate of values becomes

$$(9) \quad \Phi(y(s)) - \min_{\mathcal{H}} \Phi = \mathcal{O}\left(\frac{1}{s^{2p}}\right).$$

For $p > 1$, we have $\alpha_p > \alpha$, so the damping parameters for (8) are similar to those of $(\text{AVD})_{\alpha}$. The only major difference is the coefficient $s^{2(p-1)}$ in front of $\nabla\Phi(y(s))$, which explodes when $s \rightarrow +\infty$. From (9) we observe that the convergence rate of values can be made arbitrarily fast (in the scale of powers of $\frac{1}{s}$) with p large. The physical intuition is clear. Fast optimization is associated with the fast parameterization of the trajectories of the $(\text{AVD})_{\alpha}$ system. Our goal is to transpose these results to the discrete case, taking advantage of the fact that proximal algorithms usually inherit the properties of the continuous dynamics from which they come.

1.2.4. *Geometrical damping.* The optimization properties of the continuous dynamic $(\text{IGS})_{\gamma,\beta}$ come from its viscous damping term, which makes the system stabilize asymptotically. According to this mechanical and control point of view, it is natural to consider other types of damping. An important case is the damping driven by the Hessian which contains a rich geometric information, as considered in [2], [10], [12]. The temporal discretization of this dynamic gives the Ravine method initially introduced by Gelfand, Tsetlin 1961, then developed by B. Polyak. The exact link was obtained in [36] using high-resolution. Another interesting damping is the dry friction, taken account in an optimization framework in [1], and which gives finite time stabilization. Developing our Lyapunov analysis in these situations is an interesting subject for further studies.

1.3. **Organization of the paper.** Based on Lyapunov analysis, our main convergence result is established in Section 2. Depending on the behavior of the sequences (α_k) and (β_k) , we give the convergence rate of the values of the sequences (x_k) generated by algorithm $(\text{IP})_{\alpha_k, \beta_k}$. In Section 3, we analyze the convergence rate of the velocities. In Section 4, we show how to pass from \mathcal{O} to o estimates. In Section 5, we give general conditions on the sequences (α_k) and (β_k) which guarantee the weak convergence of the iterates. In Section 6, we study the link with the accelerated proximal algorithm of Güler. In Section 7 we study a parametrized family of rescaled proximal algorithms which extends the previous study. In Section 8, we study the stability with respect to perturbations. The paper is completed by some auxiliary technical lemmas contained in the Appendix.

2. CONVERGENCE RATE OF THE VALUES

The algorithm $(\text{IP})_{\alpha_k, \beta_k}$

$$(10) \quad (\text{IP})_{\alpha_k, \beta_k} \quad \begin{cases} y_k = x_k + \alpha_k(x_k - x_{k-1}) \\ x_{k+1} = \text{prox}_{\beta_k \Phi}(y_k), \end{cases}$$

can be equivalently formulated as

$$(11) \quad x_{k+1} - 2x_k + x_{k-1} + (1 - \alpha_k)(x_k - x_{k-1}) + \beta_k \partial\Phi(x_{k+1}) \ni 0.$$

This formulation can also be obtained from the implicit discretization of the inertial dynamic (IGS) $_{\gamma,\beta}$, see (5). Following [4], let us introduce the sequence (t_k) that plays a central role in the analysis of algorithm (IP) $_{\alpha_k,\beta_k}$.

2.1. **The sequence (t_k) .** Given $k \geq 1$, assume that $\sum_{i=k}^{+\infty} \prod_{j=k}^i \alpha_j < +\infty$ and set $t_k = 1 + \sum_{i=k}^{+\infty} \prod_{j=k}^i \alpha_j$. We use the convention $\prod_{j=k}^{k-1} \alpha_j = 1$, which gives the compact form of t_k

$$(12) \quad t_k = \sum_{i=k-1}^{+\infty} \prod_{j=k}^i \alpha_j.$$

If the series defining t_{k+1} is convergent, the series defining t_k is also convergent, and we have

$$1 + \alpha_k t_{k+1} = 1 + \alpha_k \left(\sum_{i=k}^{+\infty} \prod_{j=k+1}^i \alpha_j \right) = 1 + \sum_{i=k}^{+\infty} \prod_{j=k}^i \alpha_j = t_k.$$

Conversely, if $\alpha_k \neq 0$ and if t_k is well-defined, then the series defining t_{k+1} is convergent and the above equalities hold true. From now on, we assume that

$$(K_0) \quad \sum_{i=k}^{+\infty} \prod_{j=k}^i \alpha_j < +\infty \quad \text{for every } k \geq 1.$$

Let us summarize the above results.

Lemma 2.1. *Assume that the non-negative sequence (α_k) satisfies (K_0) . Then the sequence (t_k) is well defined and satisfies for every $k \geq 1$*

$$(13) \quad 1 + \alpha_k t_{k+1} = t_k.$$

The sequence (t_k) allows us to formulate the conditions on the sequence (α_k) in a dense form. Although the direct formulation (12) of (t_k) in terms of (α_k) is complicated, the inverse formulation is simple: $\alpha_k = \frac{t_k - 1}{t_{k+1}}$.

2.2. **Convergence rates of the values: Lyapunov analysis.** Let us introduce the ingredients that will serve for the Lyapunov analysis. For $k \geq 1$, the global energy W_k is defined by

$$(14) \quad W_k := \beta_{k-1} \left(\Phi(x_k) - \min_{\mathcal{H}} \Phi \right) + \frac{1}{2} \|x_k - x_{k-1}\|^2.$$

Its importance is due to the fact that, for damped inertial gradient systems, it is the global energy that decreases. Given $x^* \in \operatorname{argmin} \Phi$, the anchor sequence (h_k) is defined by: for $k \geq 1$

$$(15) \quad h_k := \frac{1}{2} \|x_k - x^*\|^2.$$

Let us define the sequence (\mathcal{E}_k) by: for every $k \geq 1$

$$(16) \quad \mathcal{E}_k := t_k^2 \beta_{k-1} \left(\Phi(x_k) - \min_{\mathcal{H}} \Phi \right) + \frac{1}{2} \|x_{k-1} + t_k(x_k - x_{k-1}) - x^*\|^2.$$

Note that

$$\mathcal{E}_k = t_k^2 W_k + \frac{1}{2} \|x_{k-1} - x^*\|^2 + t_k \langle x_k - x_{k-1}, x_{k-1} - x^* \rangle.$$

This formulation shows \mathcal{E}_k as a weighted sum of the global energy, the anchor function, and the discrete time derivative of the anchor function. For our next calculation, we will use the formulation (16), which, as a main advantage, involves only non-negative terms. The following result shows that the sequence (\mathcal{E}_k) is non-increasing, under the condition (K_{1,α_k,β_k}) , that involves the two sequences (α_k) , and (β_k) together.

Proposition 2.2. *Assume that the non-negative sequence (α_k) satisfies (K_0) . Let (x_k) be a sequence generated by algorithm (IP) $_{\alpha_k,\beta_k}$, and let (\mathcal{E}_k) be the sequence defined by (16). Then we have: for every $k \geq 1$*

$$(17) \quad \mathcal{E}_{k+1} - \mathcal{E}_k \leq (t_{k+1}^2 \beta_k - t_k^2 \beta_{k-1} - t_{k+1} \beta_k) \left(\Phi(x_k) - \min_{\mathcal{H}} \Phi \right).$$

Under the assumption

$$(K_{1,\alpha_k,\beta_k}) \quad t_{k+1}^2 \beta_k - t_k^2 \beta_{k-1} - t_{k+1} \beta_k \leq 0 \quad \text{for every } k \geq 1,$$

then the sequence (\mathcal{E}_k) is non-increasing.

Proof. By definition of the proximal operator, the iteration at step k of the algorithm $(IP)_{\alpha_k, \beta_k}$ writes

$$\frac{1}{\beta_k}(y_k - x_{k+1}) \in \partial\Phi(x_{k+1}).$$

Equivalently, we have the following subdifferential inequalities: for any $x \in \mathcal{H}$

$$(18) \quad \Phi(x) \geq \Phi(x_{k+1}) + \frac{1}{\beta_k} \langle x - x_{k+1}, y_k - x_{k+1} \rangle.$$

Let us write successively inequality (18) at $x = x_k$ and $x = x^* \in \operatorname{argmin} \Phi$. We obtain the two inequalities

$$(19) \quad \Phi(x_k) \geq \Phi(x_{k+1}) + \frac{1}{\beta_k} \langle x_k - x_{k+1}, y_k - x_{k+1} \rangle,$$

$$(20) \quad \Phi(x^*) \geq \Phi(x_{k+1}) + \frac{1}{\beta_k} \langle x^* - x_{k+1}, y_k - x_{k+1} \rangle.$$

Using $x_k - x_{k+1} = x_k - y_k + y_k - x_{k+1}$ in (19) and $x^* - x_{k+1} = x^* - y_k + y_k - x_{k+1}$ in (20) we obtain

$$(21) \quad \Phi(x_k) \geq \Phi(x_{k+1}) + \frac{1}{\beta_k} \langle x_k - y_k, y_k - x_{k+1} \rangle + \frac{1}{\beta_k} \|y_k - x_{k+1}\|^2,$$

$$(22) \quad \Phi(x^*) \geq \Phi(x_{k+1}) + \frac{1}{\beta_k} \langle x^* - y_k, y_k - x_{k+1} \rangle + \frac{1}{\beta_k} \|y_k - x_{k+1}\|^2.$$

Multiplying (21) by $t_{k+1} - 1 \geq 0$, then adding (22), we derive that

$$\begin{aligned} (t_{k+1} - 1)(\Phi(x_k) - \Phi(x^*)) &\geq t_{k+1}(\Phi(x_{k+1}) - \Phi(x^*)) + \frac{1}{\beta_k} \langle x_{k+1} - y_k, (t_{k+1} - 1)(y_k - x_k) + y_k - x^* \rangle \\ &\quad + \frac{t_{k+1}}{\beta_k} \|y_k - x_{k+1}\|^2. \end{aligned}$$

Observe that

$$\begin{aligned} (t_{k+1} - 1)(y_k - x_k) + y_k &= t_{k+1} y_k - (t_{k+1} - 1)x_k \\ &= x_k + t_{k+1} \alpha_k (x_k - x_{k-1}) \\ &= x_{k-1} + (1 + t_{k+1} \alpha_k)(x_k - x_{k-1}) \\ &= x_{k-1} + t_k (x_k - x_{k-1}) \quad \text{in view of (13)}. \end{aligned}$$

Setting $z_k = x_{k-1} + t_k(x_k - x_{k-1})$, we then deduce from (23) that

$$(23) \quad \begin{aligned} (t_{k+1} - 1)(\Phi(x_k) - \Phi(x^*)) &\geq t_{k+1}(\Phi(x_{k+1}) - \Phi(x^*)) \\ &\quad + \frac{1}{\beta_k} \langle x_{k+1} - y_k, z_k - x^* \rangle + \frac{t_{k+1}}{\beta_k} \|y_k - x_{k+1}\|^2. \end{aligned}$$

Equivalently, after multiplication by β_k

$$(24) \quad \begin{aligned} \beta_k(t_{k+1} - 1)(\Phi(x_k) - \Phi(x^*)) &\geq \beta_k t_{k+1}(\Phi(x_{k+1}) - \Phi(x^*)) \\ &\quad + \langle x_{k+1} - y_k, z_k - x^* \rangle + t_{k+1} \|y_k - x_{k+1}\|^2. \end{aligned}$$

To write (24) in a recursive form, observe that

$$\begin{aligned} z_{k+1} - z_k &= x_k + t_{k+1}(x_{k+1} - x_k) - x_{k-1} - t_k(x_k - x_{k-1}) \\ &= t_{k+1}(x_{k+1} - x_k) - (t_k - 1)(x_k - x_{k-1}) \\ &= t_{k+1}(x_{k+1} - x_k - \alpha_k(x_k - x_{k-1})) \quad \text{in view of (13)} \\ &= t_{k+1}(x_{k+1} - y_k). \end{aligned}$$

It ensues that

$$\|z_{k+1} - x^*\|^2 = \|z_k - x^*\|^2 + 2t_{k+1} \langle x_{k+1} - y_k, z_k - x^* \rangle + t_{k+1}^2 \|x_{k+1} - y_k\|^2,$$

which gives

$$\langle x_{k+1} - y_k, z_k - x^* \rangle = \frac{1}{2t_{k+1}} (\|z_{k+1} - x^*\|^2 - \|z_k - x^*\|^2) - \frac{t_{k+1}}{2} \|x_{k+1} - y_k\|^2.$$

Using this equality in (24), we find

$$\begin{aligned} \beta_k(t_{k+1} - 1)(\Phi(x_k) - \Phi(x^*)) &\geq \beta_k t_{k+1}(\Phi(x_{k+1}) - \Phi(x^*)) + \frac{t_{k+1}}{2} \|y_k - x_{k+1}\|^2 \\ &\quad + \frac{1}{2t_{k+1}} (\|z_{k+1} - x^*\|^2 - \|z_k - x^*\|^2). \end{aligned}$$

After multiplication by t_{k+1} , and neglecting the non-negative term $\frac{t_{k+1}}{2}\|y_k - x_{k+1}\|^2$, we obtain

$$\beta_k(t_{k+1}^2 - t_{k+1})(\Phi(x_k) - \Phi(x^*)) + \frac{1}{2}\|z_k - x^*\|^2 \geq \beta_k t_{k+1}^2(\Phi(x_{k+1}) - \Phi(x^*)) + \frac{1}{2}\|z_{k+1} - x^*\|^2.$$

Note that the sequence (\mathcal{E}_k) writes equivalently $\mathcal{E}_k = t_k^2 \beta_{k-1}(\Phi(x_k) - \min_{\mathcal{H}} \Phi) + \frac{1}{2}\|z_k - x^*\|^2$. Thus, we have obtained the following inequality

$$\mathcal{E}_k \geq \mathcal{E}_{k+1} + (t_k^2 \beta_{k-1} - \beta_k(t_{k+1}^2 - t_{k+1}))(\Phi(x_k) - \min_{\mathcal{H}} \Phi).$$

Under condition (K_{1, β_k}) we have $t_k^2 \beta_{k-1} - \beta_k(t_{k+1}^2 - t_{k+1}) \geq 0$. As a consequence $\mathcal{E}_k \geq \mathcal{E}_{k+1}$, and the sequence (\mathcal{E}_k) is non-increasing. \square

As a result of Proposition 2.2, we obtain the convergence rate of values as shown below.

Theorem 2.3. *Suppose that the sequence (α_k) satisfies (K_0) . Suppose that (α_k) and (β_k) satisfy $(K_{1, \alpha_k, \beta_k})$. Let (x_k) be a sequence generated by algorithm $(\text{IP})_{\alpha_k, \beta_k}$. Then we have*

(i) For every $k \geq 1$,

$$\Phi(x_k) - \min_{\mathcal{H}} \Phi \leq \frac{C}{t_k^2 \beta_{k-1}},$$

with $C = t_1^2 \beta_0(\Phi(x_1) - \min_{\mathcal{H}} \Phi) + \frac{1}{2}(d(x_0, \operatorname{argmin}_{\mathcal{H}} \Phi)^2 + t_1^2 \|x_1 - x_0\|^2)$.

(ii) Assume moreover that there exists $m < 1$ such that

$$(K_{1, \alpha_k, \beta_k}^+) \quad t_{k+1}^2 \beta_k - t_k^2 \beta_{k-1} \leq m t_{k+1} \beta_k \quad \text{for every } k \geq 1.$$

Then we have

$$\sum_{k=1}^{+\infty} t_{k+1} \beta_k (\Phi(x_k) - \min_{\mathcal{H}} \Phi) < +\infty.$$

Proof. (i) From Proposition 2.2, the sequence (\mathcal{E}_k) is non-increasing. It ensues that $\mathcal{E}_k \leq \mathcal{E}_1$ for every $k \geq 1$. Recalling the expression of \mathcal{E}_k , we deduce that

$$\begin{aligned} t_k^2 \beta_{k-1} (\Phi(x_k) - \min_{\mathcal{H}} \Phi) &\leq t_1^2 \beta_0 [\Phi(x_1) - \min_{\mathcal{H}} \Phi] + \frac{1}{2} \|x_0 - x^* + t_1(x_1 - x_0)\|^2 \\ &\leq t_1^2 \beta_0 [\Phi(x_1) - \min_{\mathcal{H}} \Phi] + \frac{1}{2} \|x_0 - x^*\|^2 + \frac{t_1^2}{2} \|x_1 - x_0\|^2 \end{aligned}$$

Since x^* can be taken arbitrarily in $\operatorname{argmin}_{\mathcal{H}} \Phi$, we finally obtain

$$t_k^2 \beta_{k-1} (\Phi(x_k) - \min_{\mathcal{H}} \Phi) \leq C,$$

with

$$C = t_1^2 \beta_0 (\Phi(x_1) - \min_{\mathcal{H}} \Phi) + \frac{1}{2} (d(x_0, \operatorname{argmin}_{\mathcal{H}} \Phi)^2 + t_1^2 \|x_1 - x_0\|^2).$$

(ii) By summing inequality (17) from $k = 1$ to n , we find

$$\mathcal{E}_{n+1} + \sum_{k=1}^n (t_{k+1} \beta_k - t_k^2 \beta_{k-1} + t_k^2 \beta_{k-1}) (\Phi(x_k) - \min_{\mathcal{H}} \Phi) \leq \mathcal{E}_1.$$

Since $\mathcal{E}_{n+1} \geq 0$ and since $t_{k+1}^2 \beta_k - t_k^2 \beta_{k-1} \leq m t_{k+1} \beta_k$, this implies that

$$(1 - m) \sum_{k=1}^n t_{k+1} \beta_k (\Phi(x_k) - \min_{\mathcal{H}} \Phi) \leq \mathcal{E}_1.$$

By letting n tend to infinity, we obtain

$$\sum_{k=1}^{\infty} t_{k+1} \beta_k (\Phi(x_k) - \min_{\mathcal{H}} \Phi) \leq \frac{\mathcal{E}_1}{1 - m},$$

which gives the claim. \square

2.3. Particular cases.

2.3.1. *Case $\beta_k \equiv s > 0$, α_k general.* This is the case without scaling. The algorithm $(\text{IP})_{\alpha_k, \beta_k}$ writes

$$(25) \quad \begin{cases} y_k = x_k + \alpha_k(x_k - x_{k-1}) \\ x_{k+1} = \text{prox}_{s\Phi}(y_k), \end{cases}$$

Condition $(K_{1, \alpha_k, \beta_k})$ becomes $t_{k+1}^2 - t_k^2 - t_{k+1} \leq 0$ for every $k \geq 1$. Under this condition, we obtain the convergence rate of values

$$\Phi(x_k) - \min_{\mathcal{H}} \Phi \leq \frac{C}{t_k^2}.$$

Thus, we recover the convergence rate of values of Attouch-Cabot [4].

2.3.2. *Case $\alpha_k = 1 - \frac{\alpha}{k}$, β_k general.* The algorithm $(\text{IP})_{\alpha_k, \beta_k}$ writes

$$(26) \quad \begin{cases} y_k = x_k + \left(1 - \frac{\alpha}{k}\right)(x_k - x_{k-1}) \\ x_{k+1} = \text{prox}_{\beta_k \Phi}(y_k), \end{cases}$$

Then, $t_k = \frac{k-1}{\alpha-1}$ (see [4]), and condition (K_{1, β_k}) becomes

$$\beta_{k+1} \leq \beta_k \frac{k^2}{(k+1)(k+2-\alpha)}.$$

Under this condition, we obtain the convergence rate of values

$$\Phi(x_k) - \min_{\mathcal{H}} \Phi \leq \frac{C}{k^2 \beta_k}.$$

Thus, we recover the result of Attouch-Chbani-Riahi [7, section 7].

3. CONVERGENCE RATE OF THE VELOCITIES

Let (x_k) be a sequence generated by the algorithm $(\text{IP})_{\alpha_k, \beta_k}$. We will analyze the convergence rate to zero of the velocity $\|x_k - x_{k-1}\|$. Our analysis is based on the global energy decay. Recall that the friction effect, and thus the dissipation of energy, is related to $\alpha_k \leq 1$.

Proposition 3.1. *Let (x_k) be a sequence generated by algorithm $(\text{IP})_{\alpha_k, \beta_k}$. Then, for every $k \geq 1$,*

$$(27) \quad \beta_k(\Phi(x_k) - \min_{\mathcal{H}} \Phi) + \frac{1}{2}\|x_k - x_{k-1}\|^2 \geq \beta_k(\Phi(x_{k+1}) - \mathcal{H}\Phi) + \frac{1}{2}\|x_{k+1} - x_k\|^2 + \frac{1 - \alpha_k^2}{2}\|x_k - x_{k-1}\|^2.$$

Proof. The iteration at step k of the algorithm $(\text{IP})_{\alpha_k, \beta_k}$ writes $\frac{1}{\beta_k}(y_k - x_{k+1}) \in \partial\Phi(x_{k+1})$, which gives:

$$(28) \quad \Phi(x_k) - \min_{\mathcal{H}} \Phi \geq \Phi(x_{k+1}) - \min_{\mathcal{H}} \Phi + \frac{1}{\beta_k} \langle y_k - x_{k+1}, x_k - x_{k+1} \rangle.$$

According to the extrapolation formula, $y_k = x_k + \alpha_k(x_k - x_{k-1})$, and after multiplication by β_k we get

$$\begin{aligned} \beta_k(\Phi(x_k) - \min_{\mathcal{H}} \Phi) &\geq \beta_k(\Phi(x_{k+1}) - \min_{\mathcal{H}} \Phi) + \langle x_k - x_{k+1} + \alpha_k(x_k - x_{k-1}), x_k - x_{k+1} \rangle \\ &= \beta_k(\Phi(x_{k+1}) - \min_{\mathcal{H}} \Phi) + \|x_k - x_{k+1}\|^2 + \langle \alpha_k(x_k - x_{k-1}), x_k - x_{k+1} \rangle. \end{aligned}$$

By using the algebraic inequality

$$|\langle \alpha_k(x_k - x_{k-1}), x_k - x_{k+1} \rangle| \leq \frac{1}{2}\|x_k - x_{k+1}\|^2 + \frac{\alpha_k^2}{2}\|x_k - x_{k-1}\|^2,$$

we deduce that

$$\beta_k(\Phi(x_k) - \min_{\mathcal{H}} \Phi) \geq \beta_k(\Phi(x_{k+1}) - \min_{\mathcal{H}} \Phi) + \|x_k - x_{k+1}\|^2 - \frac{1}{2}\|x_k - x_{k+1}\|^2 - \frac{\alpha_k^2}{2}\|x_k - x_{k-1}\|^2.$$

Equivalently

$$\beta_k(\Phi(x_k) - \min_{\mathcal{H}} \Phi) + \frac{1}{2}\|x_k - x_{k-1}\|^2 \geq \beta_k(\Phi(x_{k+1}) - \min_{\mathcal{H}} \Phi) + \frac{1}{2}\|x_{k+1} - x_k\|^2 + \frac{1 - \alpha_k^2}{2}\|x_k - x_{k-1}\|^2,$$

which gives the claim. \square

Theorem 3.2. *Suppose that (α_k) and (β_k) satisfy $(K_{1, \alpha_k, \beta_k}^+)$. Then, we have*

$$\sum_{k=1}^{+\infty} t_k \|x_k - x_{k-1}\|^2 < +\infty.$$

Proof. Consider, for $k \geq 1$, the global energy W_k , where we set $m = \min_{\mathcal{H}} \Phi$:

$$W_k := \beta_{k-1} (\Phi(x_k) - m) + \frac{1}{2} \|x_k - x_{k-1}\|^2.$$

Let us multiply the inequality (27) obtained in Proposition 3.1 by t_{k+1}^2

$$\begin{aligned} t_{k+1}^2 \left(\beta_k (\Phi(x_k) - m) + \frac{1}{2} \|x_k - x_{k-1}\|^2 \right) &\geq t_{k+1}^2 \beta_k (\Phi(x_{k+1}) - m) \\ &\quad + \frac{t_{k+1}^2}{2} \|x_{k+1} - x_k\|^2 + \frac{t_{k+1}^2 (1 - \alpha_k^2)}{2} \|x_k - x_{k-1}\|^2. \end{aligned}$$

Let us reformulate this inequality in terms of the sequence (W_k) . We have

$$\begin{aligned} t_{k+1}^2 \left(\beta_{k-1} (\Phi(x_k) - m) + (\beta_k - \beta_{k-1}) (\Phi(x_k) - m) + \frac{1}{2} \|x_k - x_{k-1}\|^2 \right) \\ \geq t_{k+1}^2 W_{k+1} + \frac{t_{k+1}^2 (1 - \alpha_k^2)}{2} \|x_k - x_{k-1}\|^2, \end{aligned}$$

which gives

$$t_{k+1}^2 W_k + t_{k+1}^2 (\beta_k - \beta_{k-1}) (\Phi(x_k) - m) \geq t_{k+1}^2 W_{k+1} + \frac{t_{k+1}^2 (1 - \alpha_k^2)}{2} \|x_k - x_{k-1}\|^2.$$

Equivalently

$$t_{k+1}^2 (W_{k+1} - W_k) + \frac{t_{k+1}^2 (1 - \alpha_k^2)}{2} \|x_k - x_{k-1}\|^2 \leq t_{k+1}^2 (\beta_k - \beta_{k-1}) (\Phi(x_k) - m).$$

Let us rewrite the above expression as

$$\begin{aligned} t_{k+1}^2 W_{k+1} - t_k^2 W_k + (t_k^2 - t_{k+1}^2) W_k + \frac{t_{k+1}^2 (1 - \alpha_k^2)}{2} \|x_k - x_{k-1}\|^2 \\ \leq t_{k+1}^2 (\beta_k - \beta_{k-1}) (\Phi(x_k) - m). \end{aligned}$$

Recalling the expression of W_k , we deduce that

$$\begin{aligned} t_{k+1}^2 W_{k+1} - t_k^2 W_k + \frac{1}{2} [t_k^2 - t_{k+1}^2 + t_{k+1}^2 (1 - \alpha_k^2)] \|x_k - x_{k-1}\|^2 \\ \leq [(t_{k+1}^2 - t_k^2) \beta_{k-1} + t_{k+1}^2 (\beta_k - \beta_{k-1})] (\Phi(x_k) - m). \end{aligned}$$

On the one hand, by (13), and since $t_k \geq 1$ we have

$$t_k^2 - t_{k+1}^2 + t_{k+1}^2 (1 - \alpha_k^2) = t_k^2 - t_{k+1}^2 \alpha_k^2 = t_k^2 - (t_k - 1)^2 = 2t_k - 1 \geq t_k.$$

On the other hand, by using condition $(K_{1, \alpha_k, \beta_k}^+)$ we have

$$(t_{k+1}^2 - t_k^2) \beta_{k-1} + t_{k+1}^2 (\beta_k - \beta_{k-1}) = t_{k+1}^2 \beta_k - t_k^2 \beta_{k-1} \leq t_{k+1} \beta_k.$$

Combining the above results we obtain

$$(29) \quad t_{k+1}^2 W_{k+1} - t_k^2 W_k + \frac{1}{2} t_k \|x_k - x_{k-1}\|^2 \leq t_{k+1} \beta_k (\Phi(x_k) - m).$$

By Theorem 2.3, under the assumption $(K_{1, \alpha_k, \beta_k}^+)$ we have

$$\sum_{k=1}^{\infty} t_{k+1} \beta_k (\Phi(x_k) - m) < +\infty.$$

As a consequence, by summing the above inequalities, we obtain

$$\frac{1}{2} \sum_{k=1}^{\infty} t_k \|x_k - x_{k-1}\|^2 \leq t_1^2 W_1 + \sum_{k=1}^{\infty} t_{k+1} \beta_k (\Phi(x_k) - m) < +\infty,$$

which gives the claim. \square

Corollary 3.3. *Suppose that the sequences (α_k) and (β_k) satisfy $(K_{1, \alpha_k, \beta_k}^+)$. Suppose that $0 \leq \alpha_k \leq 1$ for every $k \geq 1$, and that the sequence (β_k) satisfies: there exists some positive constant C such that, for all $k \geq 1$*

$$(30) \quad \beta_k \leq C \beta_{k+1}.$$

Then, for any sequence (x_k) generated by the algorithm $(\text{IP})_{\alpha_k, \beta_k}$, we have

$$\sum_{k=1}^{+\infty} t_k W_k < +\infty, \text{ where } W_k := \beta_{k-1} (\Phi(x_k) - \min_{\mathcal{H}} \Phi) + \frac{1}{2} \|x_k - x_{k-1}\|^2.$$

Proof. By Theorem 2.3 (ii), under the assumption $(K_{1,\alpha_k,\beta_k}^+)$ we have

$$(31) \quad \sum_{k=1}^{+\infty} t_{k+1}\beta_k(\Phi(x_k) - \min_{\mathcal{H}} \Phi) < +\infty.$$

By assumption (30) we have $t_k\beta_{k-1} \leq Ct_k\beta_k$. By Lemma 2.1, we have $t_k = 1 + \alpha_k t_{k+1}$. Since $0 \leq \alpha_k \leq 1$, and according to $1 \leq t_{k+1}$, this implies

$$t_k \leq 1 + t_{k+1} \leq 2t_{k+1}.$$

Combining the two above inequalities, we obtain $t_k\beta_{k-1} \leq 2Ct_{k+1}\beta_k$. According to (31) this implies

$$(32) \quad \sum_{k=1}^{+\infty} t_k\beta_{k-1}(\Phi(x_k) - \min_{\mathcal{H}} \Phi) \leq 2C \sum_{k=1}^{+\infty} t_{k+1}\beta_k(\Phi(x_k) - \min_{\mathcal{H}} \Phi) < +\infty.$$

Combining this inequality with the estimate of the velocities $\sum_{k=1}^{+\infty} t_k \|x_k - x_{k-1}\|^2 < +\infty$, obtained in Theorem 3.2, we obtain $\sum_{k=1}^{+\infty} t_k W_k < +\infty$. \square

Remark 3.4. In Corollary 3.3, the assumption on the sequence (β_k) is satisfied in almost all practical situations. Clearly, it is satisfied if the sequence (β_k) is non-decreasing. It may be also decreasing like $\frac{1}{k^r}$ for any $r \geq 0$, or $\exp(-k)$. A limiting case in which it is not satisfied is $\frac{1}{k!}$. Moreover, the assumption on the sequence (α_k) can be weakened, just assuming that it is bounded from above.

4. FROM \mathcal{O} TO o ESTIMATES

In line with Attouch-Peypouquet [11] and Attouch-Cabot [5], we have the following convergence rate of values, where \mathcal{O} is replaced by o .

Theorem 4.1. *Let us assume that the non-negative sequence (α_k) is bounded from above, (α_k) and (β_k) satisfy the condition $(K_{1,\alpha_k,\beta_k}^+)$, and there exists some positive constant C such that $\beta_k \leq C\beta_{k+1}$ for all $k \geq 1$.*

Then, for any sequence (x_k) generated by the algorithm $(\text{IP})_{\alpha_k,\beta_k}$, we have:

i) *If $\sum \frac{1}{t_k} = +\infty$, then*

$$\Phi(x_k) - \min_{\mathcal{H}} \Phi = o\left(\frac{1}{t_k^2\beta_k}\right) \quad \text{and} \quad \|x_k - x_{k-1}\|^2 = o\left(\frac{1}{t_k^2}\right).$$

ii) *If $\sum t_k\beta_{k-1} = +\infty$, $0 \leq \alpha_k \leq 1$, and (β_k) is non-decreasing, then*

$$\Phi(x_k) - \min_{\mathcal{H}} \Phi = o\left(\frac{1}{\sum_{i=1}^k t_i\beta_{i-1}}\right) \quad \text{and} \quad \|x_k - x_{k-1}\|^2 = o\left(\frac{\beta_{k-1}}{\sum_{i=1}^k t_i\beta_{i-1}}\right).$$

In particular, $\Phi(x_k) \rightarrow \min_{\mathcal{H}} \Phi$ as $k \rightarrow +\infty$.

Proof. Let's consider the sequence of global energies (W_k) introduced in the proof of Theorem 3.2

$$W_k := \beta_{k-1}(\Phi(x_k) - m) + \frac{1}{2}\|x_k - x_{k-1}\|^2.$$

By Corollary 3.3 it satisfies

$$\sum_{k=1}^{\infty} t_k W_k < +\infty.$$

i) Returning to (29) we have

$$t_{k+1}^2 W_{k+1} - t_k^2 W_k + \frac{1}{2}t_k \|x_k - x_{k-1}\|^2 \leq t_{k+1}\beta_k(\Phi(x_k) - m).$$

The non-negative sequence (a_k) with $a_k := t_k^2 W_k$ satisfies the relation

$$a_{k+1} - a_k \leq \omega_k$$

with $\omega_k = t_{k+1}\beta_k(\Phi(x_k) - m)$. According to $\sum_{k \geq 1} t_{k+1}\beta_k(\Phi(x_k) - m) < +\infty$ (see Theorem 2.3 (ii)) we have $(\omega_k) \in l^1(\mathbb{N})$. By a standard argument, we deduce that the limit of the sequence (a_k) exists, that is

$$\lim_{k \rightarrow +\infty} t_k^2 W_k \quad \text{exists.}$$

Let $c := \lim_{k \rightarrow +\infty} t_k^2 W_k$. Let's show that $c = 0$. Let's argue by contradiction, and suppose that $c > 0$. As a result, $t_k W_k \sim \frac{c}{t_k}$. According to $\sum_{k=1}^{\infty} t_k W_k < +\infty$, this implies $c \sum \frac{1}{t_k} < +\infty$, a clear contradiction with the hypothesis $\sum \frac{1}{t_k} = +\infty$. Hence, $c = 0$. So, $\lim_{k \rightarrow +\infty} t_k^2 W_k = 0$, which gives the claim.

ii) The argument is based on the non-increasing property of the sequence $(\frac{1}{\beta_{k-1}}W_k)$. Let us return to Proposition 3.1. After dividing (27) by β_k , we get

$$(33) \quad \begin{aligned} (\Phi(x_k) - \min_{\mathcal{H}} \Phi) + \frac{1}{2\beta_k} \|x_k - x_{k-1}\|^2 &\geq (\Phi(x_{k+1}) - \min_{\mathcal{H}} \Phi) \\ &+ \frac{1}{2\beta_k} \|x_{k+1} - x_k\|^2 + \frac{1 - \alpha_k^2}{2\beta_k} \|x_k - x_{k-1}\|^2. \end{aligned}$$

Let's formulate the above inequality in terms of the sequence (W_k) . We obtain

$$(34) \quad \frac{1}{\beta_{k-1}}W_k + \left(\frac{1}{2\beta_k} - \frac{1}{2\beta_{k-1}}\right) \|x_k - x_{k-1}\|^2 \geq \frac{1}{\beta_k}W_{k+1} + \frac{1 - \alpha_k^2}{2\beta_k} \|x_k - x_{k-1}\|^2.$$

Since $0 \leq \alpha_k \leq 1$, and the sequence (β_k) is non-decreasing, we deduce that

$$(35) \quad \frac{1}{\beta_{k-1}}W_k \geq \frac{1}{\beta_k}W_{k+1}.$$

Hence, the sequence $(\frac{1}{\beta_{k-1}}W_k)$ is non-increasing. Let us rewrite the estimate $\sum_{k=1}^{\infty} t_k W_k < +\infty$ as

$$\sum_{k=1}^{\infty} t_k \beta_{k-1} \left(\frac{1}{\beta_{k-1}} W_k \right) < +\infty.$$

We are now in position to apply Lemma A.2 in the appendix, with the sequences $(t_k \beta_{k-1})$ and $(\frac{1}{\beta_{k-1}}W_k)$, respectively in place of (τ_k) and (ε_k) . We obtain that

$$\frac{1}{\beta_{k-1}}W_k = o\left(\frac{1}{\sum_{i=1}^k t_i \beta_{i-1}}\right) \quad \text{as } k \rightarrow +\infty.$$

Hence

$$W_k = o\left(\frac{\beta_{k-1}}{\sum_{i=1}^k t_i \beta_{i-1}}\right) \quad \text{as } k \rightarrow +\infty,$$

which gives the claim. \square

As a direct consequence of Theorem 4.1, in the case without inertia, i.e. $\alpha_k \equiv 0$ (which gives $t_k \equiv 1$) we obtain the following result.

Corollary 4.2. *Suppose that the sequence (β_k) is non-decreasing. Then for any sequence (x_k) generated by the proximal algorithm $x_{k+1} = \text{prox}_{\beta_k \Phi}(x_k)$ we have the following estimates:*

$$\Phi(x_k) - \min_{\mathcal{H}} \Phi = o\left(\frac{1}{\sum_{i=1}^k \beta_{i-1}}\right) \quad \text{and} \quad \|x_k - x_{k-1}\|^2 = o\left(\frac{\beta_{k-1}}{\sum_{i=1}^k \beta_{i-1}}\right).$$

When (β_k) is non-decreasing, this notably improves the results obtained by Güler in [20, Theorem 3.1].

5. CONVERGENCE OF THE ITERATES

Let us now fix $x^* \in \text{argmin} \Phi$, and define $h_k := \frac{1}{2} \|x_k - x^*\|^2$. The sequence (h_k) takes account of the anchoring of the sequence of iterates (x_k) to the solution set. The next result will be useful for establishing the convergence of the iterates of the algorithm $(\text{IP})_{\alpha_k, \beta_k}$.

Proposition 5.1. *For any $k \geq 1$, the following inequality holds*

$$(36) \quad h_{k+1} - h_k - \alpha_k (h_k - h_{k-1}) \leq \frac{1}{2} (\alpha_k^2 + \alpha_k) \|x_k - x_{k-1}\|^2 - \beta_k (\Phi(x_{k+1}) - \min_{\mathcal{H}} \Phi).$$

Proof. Observe that

$$\begin{aligned} \|y_k - x^*\|^2 &= \|x_k + \alpha_k(x_k - x_{k-1}) - x^*\|^2 \\ &= \|x_k - x^*\|^2 + \alpha_k^2 \|x_k - x_{k-1}\|^2 + 2\alpha_k \langle x_k - x^*, x_k - x_{k-1} \rangle \\ &= \|x_k - x^*\|^2 + \alpha_k^2 \|x_k - x_{k-1}\|^2 + \alpha_k \|x_k - x^*\|^2 + \alpha_k \|x_k - x_{k-1}\|^2 - \alpha_k \|x_{k-1} - x^*\|^2 \\ &= \|x_k - x^*\|^2 + \alpha_k (\|x_k - x^*\|^2 - \|x_{k-1} - x^*\|^2) + (\alpha_k^2 + \alpha_k) \|x_k - x_{k-1}\|^2 \\ &= 2[h_k + \alpha_k (h_k - h_{k-1})] + (\alpha_k^2 + \alpha_k) \|x_k - x_{k-1}\|^2. \end{aligned}$$

Set briefly $A_k = h_{k+1} - h_k - \alpha_k(h_k - h_{k-1}) = \frac{1}{2}\|x_{k+1} - x^*\|^2 - [h_k + \alpha_k(h_k - h_{k-1})]$. According to the above inequality, we get

$$\begin{aligned} A_k &= \frac{1}{2}\|x_{k+1} - x^*\|^2 - \frac{1}{2}\|y_k - x^*\|^2 + \frac{1}{2}(\alpha_k^2 + \alpha_k)\|x_k - x_{k-1}\|^2 \\ &= \left\langle x_{k+1} - y_k, \frac{1}{2}(x_{k+1} + y_k) - x^* \right\rangle + \frac{1}{2}(\alpha_k^2 + \alpha_k)\|x_k - x_{k-1}\|^2 \\ &= \langle x_{k+1} - y_k, y_k - x^* \rangle + \frac{1}{2}\|x_{k+1} - y_k\|^2 + \frac{1}{2}(\alpha_k^2 + \alpha_k)\|x_k - x_{k-1}\|^2. \end{aligned}$$

So, we obtain the equality

$$(37) \quad h_{k+1} - h_k - \alpha_k(h_k - h_{k-1}) = \frac{1}{2}(\alpha_k^2 + \alpha_k)\|x_k - x_{k-1}\|^2 + \langle y_k - x_{k+1}, x^* - y_k \rangle + \frac{1}{2}\|y_k - x_{k+1}\|^2.$$

Now use that $x^* \in \operatorname{argmin} \Phi$. By definition of the proximal operator, $x_{k+1} = \operatorname{prox}_{\beta_k \Phi}(y_k)$ equivalently gives $\frac{1}{\beta_k}(y_k - x_{k+1}) \in \partial \Phi(x_{k+1})$. The convex subdifferential inequality applied at x_{k+1} gives

$$\Phi(x^*) \geq \Phi(x_{k+1}) + \left\langle \frac{1}{\beta_k}(y_k - x_{k+1}), x^* - x_{k+1} \right\rangle.$$

Since $\Phi(x^*) = \min_{\mathcal{H}} \Phi$, we infer that

$$\langle y_k - x_{k+1}, x^* - x_{k+1} \rangle \leq -\beta_k(\Phi(x_{k+1}) - \min_{\mathcal{H}} \Phi).$$

Hence

$$\|y_k - x_{k+1}\|^2 + \langle y_k - x_{k+1}, x^* - y_k \rangle \leq -\beta_k(\Phi(x_{k+1}) - \min_{\mathcal{H}} \Phi).$$

Combining this inequality with (37) gives

$$h_{k+1} - h_k - \alpha_k(h_k - h_{k-1}) \leq \frac{1}{2}(\alpha_k^2 + \alpha_k)\|x_k - x_{k-1}\|^2 - \beta_k(\Phi(x_{k+1}) - \min_{\mathcal{H}} \Phi),$$

which completes the proof of Proposition 5.1. \square

We will also need the following estimate for the velocities which completes Theorem 3.2.

Proposition 5.2. *Suppose that the sequences (α_k) and (β_k) satisfy the condition $(K_{1, \alpha_k, \beta_k}^+)$ and that the sequence (β_k) is non-decreasing. Then, we have*

$$\sum_{k=1}^{+\infty} t_{k+1} \|x_k - x_{k-1}\|^2 < +\infty.$$

Proof. Condition $(K_{1, \alpha_k, \beta_k}^+)$ can be equivalently written as

$$(t_{k+1}^2 - t_k^2)\beta_k + t_k^2(\beta_k - \beta_{k-1}) \leq t_{k+1}\beta_k$$

Since the sequence (β_k) is non-decreasing, we infer $t_{k+1}^2 - t_k^2 \leq t_{k+1}$. Hence

$$t_{k+1} - t_k \leq \frac{t_{k+1}}{t_{k+1} + t_k} \leq 1.$$

Since $1 \leq t_k$, we immediately obtain $t_{k+1} \leq 2t_k$. So

$$\sum_{k=1}^{+\infty} t_{k+1} \|x_k - x_{k-1}\|^2 \leq 2 \sum_{k=1}^{+\infty} t_k \|x_k - x_{k-1}\|^2 < +\infty,$$

where the last inequality comes from Theorem 3.2. \square

Theorem 5.3. *Assume that*

- i) *the sequences (α_k) and (β_k) satisfy the condition $(K_{1, \alpha_k, \beta_k}^+)$;*
- ii) *$\alpha_k \in [0, 1]$ for every $k \geq 1$;*
- iii) *the sequence (β_k) is non-decreasing;*
- iv) *$\sum_{k=1}^{\infty} t_k \beta_{k-1} = +\infty$.*

Then, any sequence (x_k) generated by the algorithm $(\text{IP})_{\alpha_k, \beta_k}$ converges weakly, and its limit belongs to $\operatorname{argmin} \Phi$.

Proof. We apply the Opial lemma, see Lemma A.1. Assume that there exist $\bar{x} \in H$ and a sequence (k_n) such that $k_n \rightarrow +\infty$, and $x_{k_n} \rightharpoonup \bar{x}$ weakly as $n \rightarrow +\infty$. Since the convex function Φ is lower semicontinuous, it is lower semicontinuous for the weak topology, hence satisfies

$$\Phi(\bar{x}) \leq \liminf_{n \rightarrow +\infty} \Phi(x_{k_n}) = \lim_{k \rightarrow +\infty} \Phi(x_k) = \min_{\mathcal{H}} \Phi,$$

where the last equality comes from Theorem 4.1. It ensues that $\bar{x} \in \operatorname{argmin} \Phi$, which shows the first point. Let us now fix $x^* \in \operatorname{argmin} \Phi$, and show that $\lim_{k \rightarrow +\infty} \|x_k - x^*\|$ exists. Set $h_k = \frac{1}{2} \|x_k - x^*\|^2$. From Proposition 5.1, (h_k) satisfies the following inequalities

$$h_{k+1} - h_k - \alpha_k(h_k - h_{k-1}) \leq \frac{1}{2}(\alpha_k^2 + \alpha_k)\|x_k - x_{k-1}\|^2 \leq \|x_k - x_{k-1}\|^2 \quad \text{since } \alpha_k \in [0, 1].$$

Taking the positive part, we find

$$(h_{k+1} - h_k)_+ \leq \alpha_k(h_k - h_{k-1})_+ + \|x_k - x_{k-1}\|^2.$$

By Proposition 5.2 we have $\sum_{k=1}^{+\infty} t_{k+1} \|x_k - x_{k-1}\|^2 < +\infty$. By applying Lemma A.3 (appendix) with $a_k = (h_k - h_{k-1})_+$ and $\omega_k = \|x_k - x_{k-1}\|^2$, we obtain $\sum_{k=1}^{+\infty} (h_k - h_{k-1})_+ < +\infty$. Since (h_k) is nonnegative, this implies that $\lim_{k \rightarrow +\infty} h_k$ exists. The second point of the Opial lemma is shown, which ends the proof. \square

6. LINK WITH GÜLER'S ACCELERATED PROXIMAL ALGORITHM

Let's recall Güler's accelerated proximal algorithm, see [21]. We modify the notations to adapt them to our framework. The proximal parameter noted λ_k in Güler's article is noted here β_k , and the inertial parameter is denoted by g_k in place of γ_k .

a) Initialization of ν_0 and A_0 .

b) Step k :

- Choose $\beta_k > 0$, and calculate $g_k > 0$ by solving
$$g_k^2 + g_k A_k \beta_k - A_k \beta_k = 0.$$
- Define
 - i) $y_k = (1 - g_k)x_k + g_k \nu_k$;
 - ii) $x_{k+1} = \operatorname{prox}_{\beta_k \Phi}(y_k)$;
 - iii) $\nu_{k+1} = \nu_k + \frac{1}{g_k}(x_{k+1} - y_k)$;
 - iv) $A_{k+1} = (1 - g_k)A_k$.

(38)

6.1. **Güler's accelerated proximal algorithm as $(\text{IP})_{\alpha_k, \beta_k}$.** Let us show that Güler's proximal algorithm can be written as an inertial proximal algorithm $(\text{IP})_{\alpha_k, \beta_k}$. First verify that, for all $k \geq 1$

$$(39) \quad \nu_k = x_{k-1} + \frac{1}{g_{k-1}}(x_k - x_{k-1}).$$

For this, we use an induction argument. Suppose (39) is satisfied at step k . Using successively (38) iii), (39), (38) i), and (39) again, we obtain

$$\begin{aligned} \nu_{k+1} &= \nu_k + \frac{1}{g_k}(x_{k+1} - y_k) = x_{k-1} + \frac{1}{g_{k-1}}(x_k - x_{k-1}) + \frac{1}{g_k}(x_{k+1} - y_k) \\ &= \frac{1}{g_k}x_{k+1} + x_{k-1} + \frac{1}{g_{k-1}}(x_k - x_{k-1}) - \frac{1}{g_k}((1 - g_k)x_k + g_k \nu_k) \\ &= \frac{1}{g_k}x_{k+1} - \frac{1 - g_k}{g_k}x_k = x_k + \frac{1}{g_k}(x_{k+1} - x_k), \end{aligned}$$

which shows that (39) is satisfied at step $k + 1$. Then, combining (38) i) with (39), we obtain

$$\begin{aligned} y_k &= (1 - g_k)x_k + g_k \nu_k = (1 - g_k)x_k + g_k \left(x_{k-1} + \frac{1}{g_{k-1}}(x_k - x_{k-1}) \right) \\ &= x_k + \left(\frac{g_k}{g_{k-1}} - g_k \right) (x_k - x_{k-1}). \end{aligned}$$

Hence, Güler's proximal algorithm can be written as

$$(40) \quad \begin{cases} y_k = x_k + \alpha_k(x_k - x_{k-1}) \\ x_{k+1} = \operatorname{prox}_{\beta_k \Phi}(y_k), \end{cases}$$

with

$$(41) \quad \alpha_k = g_k \left(\frac{1}{g_{k-1}} - 1 \right).$$

By construction of g_k , we have $g_k = \frac{1}{2} \left(-A_k \beta_k + \sqrt{(A_k \beta_k)^2 + 4A_k \beta_k} \right)$, which gives $0 \leq g_k < 1$. According to (41), we deduce that $\alpha_k > 0$. As a result, the first formula of (40) defines an extrapolation operation. This

makes Güler's algorithm an inertial proximal algorithm, within the framework of the algorithm $(IP)_{\alpha_k, \beta_k}$. From item *iv*) in Güler's algorithm, we immediately get:

$$(42) \quad A_k = A_0 \prod_{j=0}^{k-1} (1 - g_j).$$

From the second-order algebraic equation that defines g_k and (38) *iv*), we have

$$(43) \quad g_k^2 = A_k \beta_k (1 - g_k) = \beta_k A_{k+1}.$$

Combining (42) with (43), we obtain the following relation:

$$(44) \quad \beta_k = \frac{g_k^2}{A_0 \prod_{j=0}^k (1 - g_j)}.$$

According to (41)-(42)-(44), we have obtained that all the parameters entering into Güler's algorithm can be expressed according to the single parameter g_k .

6.2. Convergence rate of Güler's accelerated proximal algorithm. Let us show that the convergence rate of Güler's accelerated proximal algorithm falls within the framework of Theorem 2.3. For this, we must make the link between the parameters used in Güler's algorithm and the parameter t_k used in Theorem 2.3. We start from

$$(45) \quad \alpha_k = g_k \left(\frac{1}{g_{k-1}} - 1 \right).$$

Let us show how to invert this relation. Set $\gamma_k = 1 - \alpha_k$. According to (45)

$$\gamma_k = 1 - g_k \left(\frac{1}{g_{k-1}} - 1 \right) = g_k \left(1 + \frac{1}{g_k} - \frac{1}{g_{k-1}} \right).$$

Set $w_k := \frac{1}{g_k}$. The sequence (w_k) satisfies the finite-difference equation

$$1 + w_k - w_{k-1} = \gamma_k w_k,$$

which gives, for all $k \geq 1$,

$$w_{k-1} = 1 + \alpha_k w_k.$$

Comparing with the relation (13) that defines the sequence (t_k) in Lemma 2.1, we obtain $w_k = t_{k+1}$. Hence, for all $k \geq 1$

$$(46) \quad g_k = \frac{1}{t_{k+1}}.$$

We have now all the ingredients to verify that the condition

$$(K_{1, \alpha_k, \beta_k}) \quad t_{k+1}^2 \beta_k - t_k^2 \beta_{k-1} - t_{k+1} \beta_k \leq 0 \quad \text{for every } k \geq 1,$$

is satisfied by Güler's algorithm. We have

$$\begin{aligned} t_{k+1}^2 \beta_k - t_k^2 \beta_{k-1} - t_{k+1} \beta_k &= t_{k+1} \beta_k (t_{k+1} - 1) - t_k^2 \beta_{k-1} \\ &= \frac{\beta_k}{g_k} \left(\frac{1 - g_k}{g_k} \right) - \frac{\beta_{k-1}}{g_{k-1}^2} \\ &= \frac{g_k^2}{A_0 \prod_{j=0}^k (1 - g_j)} \frac{1}{g_k} \left(\frac{1 - g_k}{g_k} \right) - \frac{1}{g_{k-1}^2} \frac{g_{k-1}^2}{A_0 \prod_{j=0}^{k-1} (1 - g_j)} \\ &= \frac{1}{A_0 \prod_{j=0}^{k-1} (1 - g_j)} - \frac{1}{A_0 \prod_{j=0}^{k-1} (1 - g_j)} = 0. \end{aligned}$$

So, Güler's accelerated proximal algorithm corresponds to taking equality in the condition $(K_{1, \alpha_k, \beta_k})$. Let us summarize the above results.

Proposition 6.1. *Güler's accelerated proximal algorithm is within the framework of algorithm $(IP)_{\alpha_k, \beta_k}$ with $\alpha_k = g_k \left(\frac{1}{g_{k-1}} - 1 \right)$. It satisfies the condition $(K_{1, \alpha_k, \beta_k})$ with equality instead of inequality, i.e., for all $k \geq 1$*

$$t_{k+1}^2 \beta_k - t_k^2 \beta_{k-1} - t_{k+1} \beta_k = 0.$$

The following rate of convergence is satisfied: For every $k \geq 1$,

$$(47) \quad \Phi(x_k) - \min_{\mathcal{H}} \Phi \leq \frac{C}{t_k^2 \beta_{k-1}},$$

with $C = t_1^2 \beta_0 (\Phi(x_1) - \min_{\mathcal{H}} \Phi) + \frac{1}{2} (d(x_0, \operatorname{argmin}_{\mathcal{H}} \Phi))^2 + t_1^2 \|x_1 - x_0\|^2$.

Moreover, for every $k \geq 1$,

$$(48) \quad \Phi(x_k) - \min_{\mathcal{H}} \Phi = \mathcal{O} \left(\frac{1}{\sum_{j=1}^k t_j \beta_{j-1}} \right).$$

The above formulas can be expressed equivalently in terms of Güler's parameters thanks to the relation $t_k = \frac{1}{g_{k-1}}$:

$$\Phi(x_k) - \min_{\mathcal{H}} \Phi = \mathcal{O} \left(\frac{g_{k-1}^2}{\beta_{k-1}} \right) \quad \Phi(x_k) - \min_{\mathcal{H}} \Phi = \mathcal{O} \left(\frac{1}{\sum_{j=1}^k \frac{\beta_{j-1}}{g_{j-1}}} \right)$$

Proof. The convergence rate given in (47) results from Theorem 2.3. Let us now exploit the fact that in the case of Güler's algorithm, the condition $(K_{1, \alpha_k, \beta_k})$ is satisfied, with equality. By adding the equalities

$$t_j^2 \beta_{j-1} - t_{j-1}^2 \beta_{j-2} = t_j \beta_{j-1}$$

from $j = 1$ to k , we get

$$t_k^2 \beta_{k-1} = \sum_{j=1}^k t_j \beta_{j-1} + t_1 \beta_0.$$

This gives (48), by just assuming that these quantities are infinite. This is a minimal assumption to insure convergence of the algorithm. \square

Let's now compare the convergence rate given as above, which results from Theorem 2.3, with the convergence rate obtained by Güler in [21]

$$(49) \quad \Phi(x_k) - \min_{\mathcal{H}} \Phi = \mathcal{O} \left(\frac{1}{\left(\sum_{j=1}^{k-1} \sqrt{\beta_j} \right)^2} \right).$$

The two above formulas give the same rate of convergence in most practical situations. Let's give a simple result where we can deduce the convergence rate obtained by Güler from the Proposition 6.1. Suppose that (β_k) is a non-decreasing sequence, and that $t_k \geq Ck$ for some positive constant C , and k large enough. This last assumption is satisfied for the Nesterov type acceleration formula with $\alpha_k = 1 - \frac{\alpha}{k}$ and $\alpha \geq 3$, see [5]. Since (β_k) is non-decreasing

$$\left(\sum_{j=1}^{k-1} \sqrt{\beta_j} \right)^2 \leq (k-1)^2 \beta_{k-1}.$$

Since $t_k \geq Ck$ for k large enough, we deduce that (for some other constant C)

$$\frac{1}{\left(\sum_{j=1}^{k-1} \sqrt{\beta_j} \right)^2} \geq \frac{C}{t_k^2 \beta_{k-1}},$$

which gives the claim.

Remark 6.2. Güler's accelerated proximal algorithm corresponds to taking equality in $(K_{1, \alpha_k, \beta_k})$. As such, it fits into the critical case, introduced historically by Nesterov, and for which the convergence of the iterates is still an open question. By contrast, as shown in Theorem 5.3, taking strict inequality (precisely assuming the condition $(K_{1, \alpha_k, \beta_k}^+)$) gives the convergence of iterates, and makes it possible to pass from the capital \mathcal{O} to small o convergence rates. See [18] and [11] who initiated this type of result.

6.3. Dynamic interpretation of Güler's algorithm. Let's come with the dynamic interpretation of Güler's algorithm, as formulated in (40-41). According to the formulation (41) of α_k we get

$$x_{k+1} + \beta_k \partial \Phi(x_{k+1}) \ni y_k = x_k + g_k \left(\frac{1}{g_{k-1}} - 1 \right) (x_k - x_{k-1}).$$

Equivalently,

$$(50) \quad x_{k+1} - 2x_k + x_{k-1} + \left(g_k - \frac{g_k - g_{k-1}}{g_{k-1}} \right) (x_k - x_{k-1}) + \beta_k \partial \Phi(x_{k+1}) = 0.$$

This can be interpreted as a time discretization of the second-order evolution equation (when Φ is smooth)

$$(51) \quad \ddot{x}(t) + \left(g(t) - \frac{\dot{g}(t)}{g(t)} \right) \dot{x}(t) + \beta(t) \nabla \Phi(x(t)) = 0.$$

So, Güler's proximal algorithm is based on the damped inertial dynamic $(IGS)_{\gamma,\beta}$ where the damping coefficient is expressed as

$$\gamma(t) = g(t) - \frac{\dot{g}(t)}{g(t)} = g(t) \left(1 + \frac{d}{dt} \left(\frac{1}{g} \right) (t) \right).$$

This makes the link with the next section, where we show that, in this context, the corresponding dynamic and proximal algorithms can be studied more directly: the conditions ensuring the rapid convergence of the algorithm can be expressed directly on the parameters g_k and β_k . This avoids the use of the coefficients t_k whose study requires additional analysis, see [5] for more details.

7. A CLASS OF INERTIAL PROXIMAL ALGORITHMS

Motivated by the results above, we consider the proximal algorithms that can be obtained (when Φ is smooth) by various temporal discretizations of the second-order evolution equation

$$(52) \quad \ddot{x}(t) + g(t) \left(1 + \frac{d}{dt} \left(\frac{1}{g} \right) (t) \right) \dot{x}(t) + \lambda(t) \nabla \Phi(x(t)) = 0.$$

We will see that, without loss of generality, the convergence analysis for the Inertial Proximal algorithm $(IP)_{\alpha_k, \beta_k}$ can be developed within this setting. In doing so, we will introduce a class of inertial proximal algorithms, including the one studied in the previous sections, and which share similar convergence properties. Moreover, the convergence results can now be expressed directly on the parameters describing the algorithms.

7.1. Link between the two descriptions.

7.1.1. *Continuous case.* Let's show that taking $g(t) \left(1 + \frac{d}{dt} \left(\frac{1}{g} \right) (t) \right)$ as the damping coefficient $\gamma(t)$ is not a restrictive assumption. To see this, given a continuous function $\gamma(\cdot)$, we have to integrate the differential equation

$$(53) \quad g(t) \left(1 + \frac{d}{dt} \left(\frac{1}{g} \right) (t) \right) = \gamma(t).$$

Take as a new unknown function $w := \frac{1}{g}$. Then (53) is equivalent to solve the non-autonomous linear differential equation

$$\dot{w}(t) - \gamma(t)w(t) = -1.$$

Set $p(t) := \exp \left(\int_{t_0}^t \gamma(\tau) d\tau \right)$ for $t \geq t_0$, then, following [4] and assuming that $\int_{t_0}^{+\infty} \frac{ds}{p(s)} < +\infty$, we obtain as a solution

$$w(t) = p(t) \int_t^{+\infty} \frac{1}{p(s)} ds.$$

This is the function $\Gamma(\cdot)$ introduced in [4]. Hence $g(t) = \frac{1}{\Gamma(t)}$ is solution of (53). The general solution is $g(t) = \frac{1}{C p(t) + \Gamma(t)}$ with $\frac{1}{C} = \lim_{t \rightarrow +\infty} p(t)g(t)$. Let's summarize the above results in the following lemma.

Lemma 7.1. *Let $\gamma : [t_0, +\infty[\rightarrow \mathbb{R}$ be a positive continuous function. Set $p(t) = \exp \left(\int_{t_0}^t \gamma(\tau) d\tau \right)$ and assume $\int_{t_0}^{+\infty} \frac{ds}{p(s)} < +\infty$. Consider the following differential equation, where the unknown function is $g(\cdot)$:*

$$(54) \quad g(t) \left(1 + \frac{d}{dt} \left(\frac{1}{g} \right) (t) \right) = \gamma(t).$$

Then, the following non-negative continuously differentiable function $g(\cdot)$ is solution of (54):

$$(55) \quad g(t) = \frac{1}{\Gamma(t)} \quad \text{where} \quad \Gamma(t) = p(t) \int_t^{+\infty} \frac{1}{p(s)} ds.$$

Precisely, it is the solution of (54) that satisfies $\lim_{t \rightarrow +\infty} \frac{1}{p(t)g(t)} = 0$.

Note that the above definition of Γ does not depend on the initial time t_0 . When $\beta(t) \equiv \beta > 0$ is fixed, fast convergence of the values is obtained in [4] under the condition $\gamma(t)\Gamma(t) \leq \frac{3}{2}$. According to (54) and (55), in terms of $g(t)$, this condition takes the equivalent form

$$g(t) \left(1 + \frac{d}{dt} \left(\frac{1}{g} \right) (t) \right) \frac{1}{g(t)} = 1 + \frac{d}{dt} \left(\frac{1}{g} \right) (t) \leq \frac{3}{2}.$$

So, the condition $\gamma(t)\Gamma(t) \leq \frac{3}{2}$ becomes

$$(56) \quad \frac{d}{dt} \left(\frac{1}{g} \right) (t) \leq \frac{1}{2},$$

with the corresponding convergence rate of the values

$$(57) \quad \Phi(x(t)) - \min_{\mathcal{H}} \Phi = \mathcal{O}(g(t)^2).$$

This shows the obvious interest in formulating the damping coefficient in the form $g(t) \left(1 + \frac{d}{dt} \left(\frac{1}{g}\right)(t)\right)$. There is no loss of generality, and the conditions for obtaining rapid convergence results can be formulated directly on the data g . For example, let's start with $g(t) = \frac{\alpha-1}{t}$. From (54), we immediately obtain $\gamma(t) = \frac{\alpha}{t}$. Then, (56)-(57) shows that the well-known condition $\alpha \geq 3$ provides the $\mathcal{O}\left(\frac{1}{t^2}\right)$ convergence rate of values (see [4], [14], [37]).

7.1.2. *Discrete case.* Similar results occur in the discrete case. Recall that the implicit temporal discretization of (52) gives

$$(58) \quad x_{k+1} - 2x_k + x_{k-1} + g_k \left(1 + \frac{1}{g_k} - \frac{1}{g_{k-1}}\right) (x_k - x_{k-1}) + \lambda_k \partial \Phi(x_{k+1}) \ni 0.$$

Equivalently

$$(59) \quad \begin{cases} y_k = x_k + \alpha_k(x_k - x_{k-1}) \\ x_{k+1} = \text{prox}_{\lambda_k \Phi}(y_k), \end{cases}$$

where

$$(60) \quad \alpha_k = g_k \left(\frac{1}{g_{k-1}} - 1\right).$$

Set $\gamma_k = 1 - \alpha_k$. This gives $\gamma_k = g_k \left(1 + \frac{1}{g_k} - \frac{1}{g_{k-1}}\right)$. As shown in the previous section, inverting this relation gives, for all $k \geq 1$, $g_k = \frac{1}{t_{k+1}}$.

Let's summarize the above results in the following lemma.

Lemma 7.2. *Let (γ_k) be a sequence of non-negative numbers. Consider the following finite-difference equation, where the unknown sequence is (g_k)*

$$(61) \quad g_k \left(1 + \frac{1}{g_k} - \frac{1}{g_{k-1}}\right) = \gamma_k.$$

Then, the following sequence of non-negative numbers is solution of (61):

$$(62) \quad g_k = \frac{1}{t_{k+1}} \quad \text{where} \quad t_k = 1 + \sum_{i=k}^{+\infty} \prod_{j=k}^i \alpha_j.$$

Following Theorem 1.1 recalled in the introduction, when $\beta_k \equiv \beta > 0$ is fixed, we obtain a fast convergence of the values (see [5]) under the condition

$$(K_1) \quad \forall k \geq 1, \quad t_{k+1}^2 - t_k^2 \leq t_{k+1}.$$

According to the above lemma this condition can be equivalently formulated in terms of g_k as

$$(63) \quad g_k \geq 1 - \left(\frac{g_k}{g_{k-1}}\right)^2,$$

with the corresponding convergence rate of the values

$$(64) \quad \Phi(x_k) - \min_{\mathcal{H}} \Phi = \mathcal{O}(g_k^2).$$

Starting from $g_k = \frac{\alpha-1}{k}$, (61) immediately gives $\gamma_k = \frac{\alpha}{k}$. Then (63)-(64) shows that the well-known condition $\alpha \geq 3$ provides the $\mathcal{O}\left(\frac{1}{k^2}\right)$ convergence rate of values, a classical result (see [5], [8], [11], [18], [37]).

7.2. **A parametrized family of proximal inertial algorithms.** Let us start from the second-order evolution equation (52) and introduce various temporal discretizations. When considering the implicit discretization for the potential term, which gives proximal algorithms, we can take a general convex lower semicontinuous proper function Φ . As novelty, we introduce a parameter $\theta \in [0, 1]$ which takes into account different discretizations of the damping term: for $k \geq 1$,

$$\begin{aligned} (x_{k+1} - 2x_k + x_{k-1}) &+ g_k(1 - \theta) \left(1 + \frac{1}{g_{k+1}} - \frac{1}{g_k}\right) (x_{k+1} - x_k) \\ &+ g_k \theta \left(1 + \frac{1}{g_k} - \frac{1}{g_{k-1}}\right) (x_k - x_{k-1}) + \lambda_k \partial \Phi(x_{k+1}) \ni 0. \end{aligned}$$

For $\theta = 1$, we recover the algorithm considered in the previous sections. After dividing by g_k , we obtain

$$\begin{aligned} & \left(\frac{1}{g_k} + (1 - \theta) \left(1 + \frac{1}{g_{k+1}} - \frac{1}{g_k} \right) \right) (x_{k+1} - x_k) \\ & - \left(\frac{1}{g_k} + \theta \left(\frac{1}{g_{k-1}} - \frac{1}{g_k} - 1 \right) \right) (x_k - x_{k-1}) + \frac{\lambda_k}{g_k} \partial\Phi(x_{k+1}) \ni 0. \end{aligned}$$

Set, for $k \geq 1$, $\theta_k := \frac{1}{g_k} + \theta \left(\frac{1}{g_{k-1}} - \frac{1}{g_k} - 1 \right)$. Note that

$$\frac{1}{g_k} + (1 - \theta) \left(1 + \frac{1}{g_{k+1}} - \frac{1}{g_k} \right) = 1 + \theta_{k+1}.$$

So, we can reformulate the above algorithm in the condensed form

$$(65) \quad x_{k+1} + \frac{\lambda_k}{g_k(1 + \theta_{k+1})} \partial\Phi(x_{k+1}) \ni x_k + \frac{\theta_k}{1 + \theta_{k+1}} (x_k - x_{k-1}).$$

This gives the Inertial Proximal algorithm with parameter θ :

Inertial Proximal algorithm with parameter θ .

$$(66) \quad (\text{IP})_{\alpha_k, \beta_k, \theta} \begin{cases} y_k = x_k + \alpha_k (x_k - x_{k-1}) \\ x_{k+1} = \text{prox}_{\beta_k \Phi}(y_k), \\ \alpha_k := \frac{\theta_k}{1 + \theta_{k+1}}; \\ \beta_k := \frac{\lambda_k}{g_k(1 + \theta_{k+1})}; \\ \theta_k := \frac{1}{g_k} + \theta \left(\frac{1}{g_{k-1}} - \frac{1}{g_k} - 1 \right). \end{cases}$$

$(\text{IP})_{\alpha_k, \beta_k, \theta}$ fits into the general framework of $(\text{IP})_{\alpha_k, \beta_k}$, with extrapolation parameters and proximal parameters written in a specific form.

7.3. Convergence rates. An analysis parallel to that developed in the previous sections can be developed. We only detail the study of the rapid convergence of values.

Theorem 7.3. Consider the inertial proximal algorithm $(\text{IP})_{\alpha_k, \beta_k, \theta}$ where $\alpha_k := \frac{\theta_k}{1 + \theta_{k+1}}$, $\beta_k := \frac{\lambda_k}{g_k(1 + \theta_{k+1})}$, and $\theta_k := \frac{1}{g_k} + \theta \left(\frac{1}{g_{k-1}} - \frac{1}{g_k} - 1 \right)$. Suppose that $0 < g_k \leq 1$, $0 \leq \theta \leq 1$, and the parameters (g_k) , (λ_k) and θ satisfy the growth condition: there exists $k_1 \in \mathbb{N}$ such that for all $k \geq k_1$

$$(K_{1, g_k, \lambda_k, \theta}) \quad \lambda_{k+1} \leq \frac{g_{k+1}}{g_k} \frac{\theta_{k+1} + 1}{\theta_{k+2}} \lambda_k.$$

Then, for any sequence (x_k) generated by the algorithm $(\text{IP})_{\alpha_k, \beta_k, \theta}$, we have

$$\left\{ \begin{array}{l} (i) \quad \Phi(x_k) - \min_{\mathcal{H}} \Phi = \mathcal{O} \left(\frac{g_{k-1}}{\lambda_{k-1}(1 + \theta_k)} \right), \text{ as } k \rightarrow +\infty; \\ (ii) \quad \sum_{k \geq 1} \beta_{k, \theta} (\Phi(x_k) - \min_{\mathcal{H}} \Phi) < +\infty \\ \text{where } \beta_{k, \theta} := \frac{\lambda_{k-1}}{g_{k-1}} (1 + \theta_k) - \frac{\lambda_k}{g_k} \theta_{k+1} \text{ is non-negative by } (K_{1, g_k, \lambda_k, \theta}). \end{array} \right.$$

The proof is similar to that of the more general perturbed case (Theorem 8.1).

7.4. Some examples. Depending on the choice of the parameter θ , we obtain a specific algorithm, with its convergence rate. The classical situation (considered below) corresponds to the explicit discretization ($\theta = 1$) of the damping term. Let's consider the following cases of particular interest:

a) *Case $\theta = 1$:* it corresponds to the explicit discretization of the damping term

$$(67) \quad (x_{k+1} - 2x_k + x_{k-1}) + g_k \left(1 + \frac{1}{g_k} - \frac{1}{g_{k-1}} \right) (x_k - x_{k-1}) + \beta_k \partial\Phi(x_{k+1}) \ni 0.$$

This is precisely the algorithm studied in the previous section with $\alpha_k = g_k \left(\frac{1}{g_{k-1}} - 1 \right)$. Let us compare the result obtained by applying respectively Theorem 2.3 (using the t_k) and Theorem 7.3 (using the g_k in an explicit form). On the one hand, according to $g_k = \frac{1}{t_{k+1}}$, the growth condition (K_{1,α_k,β_k}) of Theorem 2.3 can be written

$$\beta_{k+1} \leq \frac{g_{k+1}}{g_k^2 \left(\frac{1}{g_{k+1}} - 1 \right)} \beta_k.$$

Making $\theta = 1$ in the formula giving the parameters, we have

$$\theta_k := \frac{1}{g_k} + \left(\frac{1}{g_{k-1}} - \frac{1}{g_k} - 1 \right) = \frac{1}{g_{k-1}} - 1.$$

Hence $\theta_{k+1} + 1 = \frac{1}{g_k}$, which gives $\beta_k = \lambda_k$. The formula $(K_{1,g_k,\lambda_k,\theta})$ in Theorem 7.3 becomes

$$(68) \quad \beta_{k+1} \leq \frac{g_{k+1}}{g_k} \frac{\frac{1}{g_k}}{\frac{1}{g_{k+1}} - 1} \beta_k = \frac{g_{k+1}}{g_k^2 \left(\frac{1}{g_{k+1}} - 1 \right)} \beta_k.$$

So we recover the same growth condition. Let us now compare the convergence rates. Theorem 2.3 gives

$$\Phi(x_k) - \min_{\mathcal{H}} \Phi = \mathcal{O} \left(\frac{1}{t_k^2 \beta_{k-1}} \right).$$

Theorem 7.3 gives

$$\Phi(x_k) - \min_{\mathcal{H}} \Phi = \mathcal{O} \left(\frac{g_{k-1}}{\lambda_{k-1} (1 + \theta_k)} \right).$$

From $\theta_k + 1 = \frac{1}{g_k}$ we get

$$\frac{g_{k-1}}{\lambda_{k-1} (1 + \theta_k)} = \frac{g_{k-1}^2}{\lambda_{k-1}} = \frac{1}{t_k^2 \beta_{k-1}}.$$

So, we recover the same convergence rate.

b) *Case $\theta = 0$* : it corresponds to the implicit discretization of the damping term

$$(69) \quad (x_{k+1} - 2x_k + x_{k-1}) + g_k \left(1 + \frac{1}{g_{k+1}} - \frac{1}{\alpha_k} \right) (x_{k+1} - x_k) + \beta_k \partial \Phi(x_{k+1}) \ni 0.$$

We have $\theta_k = \frac{1}{g_k}$, which gives algorithm (IP) $_{\alpha_k,\beta_k,\theta}$ with $\alpha_k = \frac{g_{k+1}}{g_k(1+g_{k+1})}$ and $\beta_k = \frac{\lambda_k g_{k+1}}{g_k(1+g_{k+1})}$. Theorem 7.3 gives that, under the condition $(K_{1,g_k,\lambda_k,\theta})$

$$\lambda_{k+1} \leq \frac{(1+g_{k+1})g_{k+2}}{g_k} \lambda_k,$$

we have

$$\Phi(x_k) - \min_{\mathcal{H}} \Phi = \mathcal{O} \left(\frac{g_k g_{k-1}}{\lambda_{k-1} (1 + g_k)} \right), \text{ as } k \rightarrow +\infty;$$

Consider the particular case $g_k = \frac{\alpha-1}{k-1}$. This gives $\alpha_k = \frac{k-1}{k+\alpha-1}$, which corresponds to a variant of the Nesterov acceleration scheme considered in [8], [18], [37]. An elementary calculation shows that the growth condition above and the corresponding convergence rate give results comparable to those of the explicit case.

8. STABILITY WITH RESPECT TO PERTURBATIONS, ERRORS

Consider the perturbed version of the evolution equation (52)

$$(70) \quad \ddot{x}(t) + g(t) \left(1 + \frac{d}{dt} \left(\frac{1}{g} \right) (t) \right) \dot{x}(t) + \lambda(t) \nabla \Phi(x(t)) = e(t)$$

where the second member of (70), denoted by $e(\cdot)$, can be interpreted as an external action on the system, a perturbation, or a control term. By following a parallel approach to the time discretization procedure described in section 7.2, we obtain : for $k \geq 1$,

$$\begin{aligned} (x_{k+1} - 2x_k + x_{k-1}) &+ g_k(1-\theta) \left(1 + \frac{1}{g_{k+1}} - \frac{1}{g_k} \right) (x_{k+1} - x_k) \\ &+ g_k \theta \left(1 + \frac{1}{g_k} - \frac{1}{g_{k-1}} \right) (x_k - x_{k-1}) + \lambda_k \partial \Phi(x_{k+1}) \ni e_k \end{aligned}$$

From the algorithmic point of view, the sequence (e_k) of elements of \mathcal{H} takes into account the presence of perturbations, approximations, or errors. After dividing by g_k , we obtain

$$\begin{aligned} & \left(\frac{1}{g_k} + (1-\theta) \left(1 + \frac{1}{g_{k+1}} - \frac{1}{g_k} \right) \right) (x_{k+1} - x_k) \\ & - \left(\frac{1}{g_k} + \theta \left(\frac{1}{g_{k-1}} - \frac{1}{g_k} - 1 \right) \right) (x_k - x_{k-1}) + \frac{\lambda_k}{g_k} \partial\Phi(x_{k+1}) \ni \frac{1}{g_k} e_k. \end{aligned}$$

Set, for $k \geq 1$

$$\theta_k := \frac{1}{g_k} + \theta \left(\frac{1}{g_{k-1}} - \frac{1}{g_k} - 1 \right).$$

Note that

$$\frac{1}{g_k} + (1-\theta) \left(1 + \frac{1}{g_{k+1}} - \frac{1}{g_k} \right) = 1 + \theta_{k+1}.$$

So, we can reformulate the above algorithm in the condensed form

$$(71) \quad (1 + \theta_{k+1})(x_{k+1} - x_k) + \frac{\lambda_k}{g_k} \partial\Phi(x_{k+1}) \ni \theta_k(x_k - x_{k-1}) + \frac{1}{g_k} e_k.$$

Equivalently,

$$(72) \quad x_{k+1} + \frac{\lambda_k}{g_k(1 + \theta_{k+1})} \partial\Phi(x_{k+1}) \ni x_k + \frac{\theta_k}{1 + \theta_{k+1}}(x_k - x_{k-1}) + \frac{1}{g_k(1 + \theta_{k+1})} e_k.$$

This gives the Inertial Proximal algorithm with parameter θ and error r_k , $(\text{IP})_{\alpha_k, \beta_k, r_k, \theta}$ for short

Inertial Proximal Algorithm with parameter θ and error r_k .

$$(73) \quad (\text{IP})_{\alpha_k, \beta_k, r_k, \theta} \begin{cases} y_k = x_k + \alpha_k(x_k - x_{k-1}) \\ x_{k+1} = \text{prox}_{\beta_k \Phi}(y_k - r_k), \\ \alpha_k := \frac{\theta_k}{1 + \theta_{k+1}}; \\ \beta_k := \frac{\lambda_k}{g_k(1 + \theta_{k+1})}; \\ r_k = -\frac{1}{g_k(1 + \theta_{k+1})} e_k \\ \theta_k := \frac{1}{g_k} + \theta \left(\frac{1}{g_{k-1}} - \frac{1}{g_k} - 1 \right). \end{cases}$$

The following result extends Theorem 2.3 to the perturbed case.

Theorem 8.1. *Consider the inertial proximal algorithm $(\text{IP})_{\alpha_k, \beta_k, r_k, \theta}$. Suppose that $0 < g_k \leq 1$, $0 \leq \theta \leq 1$, and the parameters (g_k) , (λ_k) and θ satisfy the growth condition: there exists $k_1 \in \mathbb{N}$ such that for all $k \geq k_1$*

$$(K_{1, g_k, \lambda_k, \theta}) \quad \lambda_{k+1} \leq \frac{g_{k+1}}{g_k} \frac{\theta_{k+1} + 1}{\theta_{k+2}} \lambda_k.$$

Suppose that the sequence (r_k) satisfies the summability property

$$(74) \quad \sum_{k \geq 1} (1 + \theta_{k+1}) \|r_k\| < +\infty.$$

Then, for any sequence (x_k) generated by the algorithm $(\text{IP})_{\alpha_k, \beta_k, r_k, \theta}$, we have

$$\left\{ \begin{array}{l} (i) \quad \Phi(x_k) - \min_{\mathcal{H}} \Phi = \mathcal{O} \left(\frac{g_{k-1}}{\lambda_{k-1}(1 + \theta_k)} \right), \text{ as } k \rightarrow +\infty; \\ (ii) \quad \sum_{k \geq 1} \beta_{k, \theta} (\Phi(x_k) - \min_{\mathcal{H}} \Phi) < +\infty \\ \text{where } \beta_{k, \theta} := \frac{\lambda_{k-1}}{g_{k-1}} (1 + \theta_k) - \frac{\lambda_k}{g_k} \theta_{k+1} \text{ is non-negative by } (K_{1, g_k, \lambda_k, \theta}). \end{array} \right.$$

Proof. To make the presentation simpler, without loss of generality, we take $k_1 = 1$. By definition of the proximal operator, the iteration at step k of the algorithm $(\text{IP})_{\alpha_k, \beta_k, \theta}$ is written

$$\frac{1}{\beta_k} (y_k - x_{k+1} - r_k) \in \partial\Phi(x_{k+1}).$$

Equivalently, we have the following subdifferential inequalities: for any $x \in \mathcal{H}$

$$(75) \quad \Phi(x) \geq \Phi(x_{k+1}) + \frac{1}{\beta_k} \langle x - x_{k+1}, y_k - x_{k+1} \rangle - \frac{1}{\beta_k} \langle x - x_{k+1}, r_k \rangle.$$

Let us write successively inequality (75) at $x = x_k$ and $x = x^* \in \operatorname{argmin} \Phi$. We obtain the two inequalities

$$(76) \quad \Phi(x_k) \geq \Phi(x_{k+1}) + \frac{1}{\beta_k} \langle x_k - x_{k+1}, y_k - x_{k+1} \rangle - \frac{1}{\beta_k} \langle x_k - x_{k+1}, r_k \rangle,$$

$$(77) \quad \Phi(x^*) \geq \Phi(x_{k+1}) + \frac{1}{\beta_k} \langle x^* - x_{k+1}, y_k - x_{k+1} \rangle - \frac{1}{\beta_k} \langle x^* - x_{k+1}, r_k \rangle.$$

Using $x_k - x_{k+1} = x_k - y_k + y_k - x_{k+1}$ in (76) and $x^* - x_{k+1} = x^* - y_k + y_k - x_{k+1}$ in (77) we obtain

$$(78) \quad \Phi(x_k) \geq \Phi(x_{k+1}) + \frac{1}{\beta_k} \langle x_k - y_k, y_k - x_{k+1} \rangle - \frac{1}{\beta_k} \langle x_k - x_{k+1}, r_k \rangle + \frac{1}{\beta_k} \|y_k - x_{k+1}\|^2,$$

$$(79) \quad \Phi(x^*) \geq \Phi(x_{k+1}) + \frac{1}{\beta_k} \langle x^* - y_k, y_k - x_{k+1} \rangle - \frac{1}{\beta_k} \langle x^* - x_{k+1}, r_k \rangle + \frac{1}{\beta_k} \|y_k - x_{k+1}\|^2.$$

Multiplying (78) by $\frac{\theta_k}{\alpha_k} - 1 \geq 0$, then adding (79), we derive that

$$(80) \quad \begin{aligned} \left(\frac{\theta_k}{\alpha_k} - 1\right) (\Phi(x_k) - \Phi(x^*)) &\geq \frac{\theta_k}{\alpha_k} (\Phi(x_{k+1}) - \Phi(x^*)) + \frac{\theta_k}{\alpha_k \beta_k} \|y_k - x_{k+1}\|^2 \\ &+ \frac{1}{\beta_k} \left\langle x_{k+1} - y_k, \left(\frac{\theta_k}{\alpha_k} - 1\right) (y_k - x_k) + y_k - x^* \right\rangle \\ &+ \frac{1}{\beta_k} \left\langle \left(\frac{\theta_k}{\alpha_k} - 1\right) (x_{k+1} - x_k) + x_{k+1} - x^*, r_k \right\rangle. \end{aligned}$$

By definition of y_k we have

$$\begin{aligned} \left(\frac{\theta_k}{\alpha_k} - 1\right) (y_k - x_k) + y_k &= \left(\frac{\theta_k}{\alpha_k} - 1\right) \alpha_k (x_k - x_{k-1}) + x_k + \alpha_k (x_k - x_{k-1}) \\ &= x_k + \theta_k (x_k - x_{k-1}) = z_k \end{aligned}$$

where $z_k := x_k + \theta_k (x_k - x_{k-1})$. Moreover

$$\left(\frac{\theta_k}{\alpha_k} - 1\right) (x_{k+1} - x_k) + x_{k+1} = \theta_{k+1} (x_{k+1} - x_k) + x_{k+1} = z_{k+1}.$$

We then deduce from (80) that

$$\begin{aligned} \left(\frac{\theta_k}{\alpha_k} - 1\right) (\Phi(x_k) - \Phi(x^*)) &\geq \frac{\theta_k}{\alpha_k} (\Phi(x_{k+1}) - \Phi(x^*)) + \frac{1}{\beta_k} \langle x_{k+1} - y_k, z_k - x^* \rangle \\ &+ \frac{1}{\beta_k} \langle r_k, z_{k+1} - x^* \rangle + \frac{\theta_k}{\alpha_k \beta_k} \|y_k - x_{k+1}\|^2. \end{aligned}$$

Equivalently, after multiplication by β_k

$$(81) \quad \begin{aligned} \beta_k \left(\frac{\theta_k}{\alpha_k} - 1\right) (\Phi(x_k) - \Phi(x^*)) &\geq \frac{\beta_k \theta_k}{\alpha_k} (\Phi(x_{k+1}) - \Phi(x^*)) \\ &+ \langle x_{k+1} - y_k, z_k - x^* \rangle + \langle r_k, z_{k+1} - x^* \rangle + \frac{\theta_k}{\alpha_k} \|y_k - x_{k+1}\|^2. \end{aligned}$$

To write (81) in a recursive form, we use $z_{k+1} - z_k = (1 + \theta_{k+1}) (x_{k+1} - y_k)$. It ensues that

$$\|z_{k+1} - x^*\|^2 = \|z_k - x^*\|^2 + 2(1 + \theta_{k+1}) \langle x_{k+1} - y_k, z_k - x^* \rangle + (1 + \theta_{k+1})^2 \|x_{k+1} - y_k\|^2,$$

which gives

$$\langle x_{k+1} - y_k, z_k - x^* \rangle = \frac{1}{2(1 + \theta_{k+1})} (\|z_{k+1} - x^*\|^2 - \|z_k - x^*\|^2) - \frac{(1 + \theta_{k+1})}{2} \|x_{k+1} - y_k\|^2.$$

Using this equality in (81), we find

$$\begin{aligned} \beta_k \left(\frac{\theta_k}{\alpha_k} - 1\right) (\Phi(x_k) - \Phi(x^*)) &\geq \frac{\beta_k \theta_k}{\alpha_k} (\Phi(x_{k+1}) - \Phi(x^*)) + \langle r_k, z_{k+1} - x^* \rangle \\ &+ \frac{1}{2(1 + \theta_{k+1})} (\|z_{k+1} - x^*\|^2 - \|z_k - x^*\|^2) + \frac{(1 + \theta_{k+1})}{2} \|x_{k+1} - y_k\|^2, \end{aligned}$$

where we have used $\frac{\theta_k}{\alpha_k} - \frac{(1+\theta_{k+1})}{2} = \frac{(1+\theta_{k+1})}{2}$ (a consequence of the definition of α_k). After multiplication by $(1 + \theta_{k+1})$, and neglecting the non-negative term $\frac{(1+\theta_{k+1})}{2}\|x_{k+1} - y_k\|^2$, we obtain

$$(82) \quad \begin{aligned} & \beta_k(1 + \theta_{k+1}) \left(\frac{\theta_k}{\alpha_k} - 1 \right) (\Phi(x_k) - \Phi(x^*)) + \frac{1}{2}\|z_k - x^*\|^2 \\ & \geq \frac{\beta_k(1 + \theta_{k+1})\theta_k}{\alpha_k} (\Phi(x_{k+1}) - \Phi(x^*)) + \frac{1}{2}\|z_{k+1} - x^*\|^2 + (1 + \theta_{k+1})\langle r_k, z_{k+1} - x^* \rangle. \end{aligned}$$

According to $\beta_k(1 + \theta_{k+1}) = \frac{\lambda_k}{g_k}$, and $\frac{\theta_k}{\alpha_k} - 1 = \theta_{k+1}$ we have

$$\beta_k(1 + \theta_{k+1}) \left(\frac{\theta_k}{\alpha_k} - 1 \right) = \frac{\lambda_k}{g_k} \theta_{k+1}.$$

Hence, (82) can be equivalently written as

$$\begin{aligned} \frac{\lambda_k}{g_k} \theta_{k+1} (\Phi(x_k) - \Phi(x^*)) + \frac{1}{2}\|z_k - x^*\|^2 & \geq \frac{\lambda_k}{g_k} (1 + \theta_{k+1}) (\Phi(x_{k+1}) - \Phi(x^*)) \\ & + \frac{1}{2}\|z_{k+1} - x^*\|^2 + (1 + \theta_{k+1}) \langle r_k, z_{k+1} - x^* \rangle. \end{aligned}$$

This naturally leads us to introduce the sequence (\mathcal{E}_k)

$$(83) \quad \mathcal{E}_k = \frac{\lambda_{k-1}}{g_{k-1}} (1 + \theta_k) (\Phi(x_k) - \Phi(x^*)) + \frac{1}{2}\|z_k - x^*\|^2.$$

Thus, we have obtained the following inequality

$$(84) \quad \mathcal{E}_k \geq \mathcal{E}_{k+1} + \left(\frac{\lambda_{k-1}}{g_{k-1}} (1 + \theta_k) - \frac{\lambda_k}{g_k} \theta_{k+1} \right) (\Phi(x_k) - \min_{\mathcal{H}} \Phi) + (1 + \theta_{k+1}) \langle r_k, z_{k+1} - x^* \rangle.$$

Under condition $K_{1,g_k,\lambda_k,\theta}$ we have $\frac{\lambda_{k-1}}{g_{k-1}} (1 + \theta_k) - \frac{\lambda_k}{g_k} \theta_{k+1} \geq 0$. Hence,

$$(85) \quad (1 + \theta_{k+1}) \|r_k\| \times \|z_{k+1} - x^*\| + \mathcal{E}_k \geq \mathcal{E}_{k+1}.$$

By summing inequalities (85) from $j = 1$ to $k - 1$, we obtain

$$(86) \quad \mathcal{E}_k \leq \mathcal{E}_1 + \sum_{j=2}^k (1 + \theta_j) \|r_{j-1}\| \times \|z_j - x^*\|.$$

Since $\mathcal{E}_k \geq \frac{1}{2}\|z_k - x^*\|^2$, we deduce that

$$(87) \quad \|z_k - x^*\|^2 \leq 2\mathcal{E}_1 + \sum_{j=1}^k 2(1 + \theta_j) \|r_{j-1}\| \times \|z_j - x^*\|.$$

Let us apply the Gronwall lemma A.4 with $a_j = \|z_j - x^*\|$, $b_j = 2(1 + \theta_j) \|r_{j-1}\|$, and $c = \sqrt{2\mathcal{E}_1}$. We obtain

$$\|z_k - x^*\| \leq \sqrt{2\mathcal{E}_1} + \sum_{j=1}^{\infty} 2(1 + \theta_j) \|r_{j-1}\|.$$

By assumption (74), $\sqrt{2\mathcal{E}_1} + \sum_{j=1}^{\infty} 2(1 + \theta_j) \|r_{j-1}\|$ is a positive finite real number. Returning to (86) we obtain

$$(88) \quad \mathcal{E}_k \leq C := \mathcal{E}_1 + \sum_{j=2}^k (1 + \theta_j) \|r_{j-1}\| \left(\sqrt{2\mathcal{E}_1} + \sum_{j=1}^{\infty} 2(1 + \theta_j) \|r_{j-1}\| \right).$$

By definition of \mathcal{E}_k , we obtain, for all $k \geq k_1$

$$\frac{\lambda_{k-1}}{g_{k-1}} (1 + \theta_k) (\Phi(x_k) - \Phi(x^*)) \leq \mathcal{E}_k \leq C,$$

which gives the claim. The last item follows directly by summing (84). \square

Remark 8.2. According to the relations between the parameters, the summability assumption on the errors can be equivalently formulated in terms of e_k as

$$\sum_{k \geq 1} \frac{1}{g_k} \|e_k\| < +\infty.$$

When $\theta = 1$ and $\alpha_k = 1 - \frac{\alpha}{k}$, using $\frac{1}{g_k} = t_{k+1}$, we recover the condition $\sum_{k \geq 1} k \|e_k\| < +\infty$ considered in [8].

Remark 8.3. Because of its numerical importance, several papers have been devoted to the study of perturbation, errors in accelerated proximal-gradient methods. One can consult Aujol-Dossal [13], Schmidt-Le Roux-Bach [34], and Villa-Salzo-Baldassarres-Verri [35].

APPENDIX A. SOME AUXILIARY RESULTS

The following auxiliary lemmas are used throughout the paper. To establish the weak convergence of the iterates of $(\text{IP})_{\alpha_k, \beta_k}$, we apply Opial's Lemma [27], that we recall in its discrete form.

Lemma A.1. *Let S be a nonempty subset of \mathcal{H} , and (x_k) a sequence in \mathcal{H} . Assume that*

(i) *every sequential weak cluster point of (x_k) , as $k \rightarrow +\infty$, belongs to S ;*

(ii) *for every $z \in S$, $\lim_{k \rightarrow +\infty} \|x_k - z\|$ exists.*

Then (x_k) converges weakly as $k \rightarrow +\infty$ to a point in S .

Owing to the next lemma, we are able to estimate the rate of convergence of a sequence (ε_k) supposed to be non-increasing and summable with respect to weight coefficients, see [5, Lemma 21] for the proof.

Lemma A.2. *Let (τ_k) be a nonnegative sequence such that $\sum_{k=1}^{+\infty} \tau_k = +\infty$. Assume that (ε_k) is a non-negative and non-increasing sequence satisfying $\sum_{k=1}^{+\infty} \tau_k \varepsilon_k < +\infty$. Then we have*

$$\varepsilon_k = o\left(\frac{1}{\sum_{i=1}^k \tau_i}\right) \quad \text{as } k \rightarrow +\infty.$$

The following result shows the summability of a sequence (a_k) satisfying some suitable inequality.

Lemma A.3. *Given a non-negative sequence (α_k) satisfying (K_0) , let (t_k) be the sequence defined by $t_k = 1 + \sum_{i=k}^{+\infty} \prod_{j=k}^i \alpha_j$. Let (a_k) and (ω_k) be two sequences of nonnegative numbers such that*

$$(89) \quad a_{k+1} \leq \alpha_k a_k + \omega_k,$$

for all $k \geq 0$. If $\sum_{k=0}^{+\infty} t_{k+1} \omega_k < +\infty$, then $\sum_{k=0}^{+\infty} a_k < +\infty$.

Proof. By Lemma 2.1, we have $t_{k+1} \alpha_k = t_k - 1$. Multiplying inequality (89) by t_{k+1} gives

$$t_{k+1} a_{k+1} \leq (t_k - 1) a_k + t_{k+1} \omega_k,$$

or equivalently $a_k \leq (t_k a_k - t_{k+1} a_{k+1}) + t_{k+1} \omega_k$. By summing from $k = 0$ to n , we obtain

$$\begin{aligned} \sum_{k=0}^n a_k &\leq t_0 a_0 - t_{n+1} a_{n+1} + \sum_{k=0}^n t_{k+1} \omega_k \\ &\leq t_0 a_0 + \sum_{k=0}^{+\infty} t_{k+1} \omega_k < +\infty \quad \text{by assumption.} \end{aligned}$$

The conclusion follows by letting n tend to $+\infty$. □

Lemma A.4 ([8, Lemma 5.14]). *Let (a_k) be a sequence of nonnegative numbers such that, for all $k \in \mathbb{N}$, $a_k^2 \leq c^2 + \sum_{j=1}^k b_j a_j$, where (b_j) is a summable sequence of nonnegative numbers, and $c \geq 0$. Then, for all $k \in \mathbb{N}$, $a_k \leq c + \sum_{j=1}^{\infty} b_j$.*

REFERENCES

- [1] S. ADLY, H. ATTOUCH, A. CABOT, *Finite time stabilization of nonlinear oscillators subject to dry friction*, *Nonsmooth Mechanics and Analysis*, Adv. Mech. Math., 12 (2006), Springer, New York, pp. 289–304.
- [2] F. ÁLVAREZ, H. ATTOUCH, J. BOLTE, P. REDONT, *A second-order gradient-like dissipative dynamical system with Hessian-driven damping. Application to optimization and mechanics*, *J. Math. Pures Appl.*, 81 (2002), No. 8, pp. 747–779.
- [3] V. APIDOPOULOS, J.-F. AUJOL, CH. DOSSAL, *Convergence rate of inertial Forward-Backward algorithm beyond Nesterov's rule*, HAL-01551873, (2017), to appear in *Mathematical Programming*.
- [4] H. ATTOUCH, A. CABOT, *Asymptotic stabilization of inertial gradient dynamics with time-dependent viscosity*, *J. Differential Equations*, 263 (2017), pp. 5412–5458.
- [5] H. ATTOUCH, A. CABOT, *Convergence rates of inertial forward-backward algorithms*, *SIAM J. Optim.*, 28 (1) (2018), pp. 849–874.
- [6] H. ATTOUCH, A. CABOT, Z. CHBANI, H. RIAHI, *Accelerated forward-backward algorithms with perturbations. Application to Tikhonov regularization*, *Journal of Optimization Theory and Applications*, 179 (2018), pp. 1–36.
- [7] H. ATTOUCH, Z. CHBANI, H. RIAHI, *Fast proximal methods via time scaling of damped inertial dynamics*, HAL-01939292, 2018.
- [8] H. ATTOUCH, Z. CHBANI, J. PEYPOUQUET, P. REDONT, *Fast convergence of inertial dynamics and algorithms with asymptotic vanishing viscosity*, *Math. Program. Ser. B* 168 (2018), pp. 123–175.
- [9] H. ATTOUCH, Z. CHBANI, H. RIAHI, *Rate of convergence of the Nesterov accelerated gradient method in the subcritical case $\alpha \leq 3$* , arXiv:1706.05671v1 [math.OC] 2017, to appear in *ESAIM-COCV*.

- [10] H. ATTOUCH, P.E. MAINGÉ, P. REDONT, *A second-order differential system with Hessian-driven damping; Application to non-elastic shock laws*, Differential Equations and Applications, 4 (2012), No. 1, pp. 27–65.
- [11] H. ATTOUCH, J. PEYPOUQUET, *The rate of convergence of Nesterov’s accelerated forward-backward method is actually faster than $1/k^2$* , SIAM J. Optim., 26 (2016), No. 3, pp. 1824–1834.
- [12] H. ATTOUCH, J. PEYPOUQUET, P. REDONT, *Fast convex minimization via inertial dynamics with Hessian driven damping*, J. Differential Equations, 261, Issue 10, (2016), pp. 5734–5783.
- [13] J.-F. AUJOL, CH. DOSSAL, *Stability of over-relaxations for the Forward-Backward algorithm, application to FISTA*, SIAM J. Optim., 25 (2015), No. 4, pp. 2408–2433.
- [14] J.-F. AUJOL, CH. DOSSAL, *Optimal rate of convergence of an ODE associated to the Fast Gradient Descent schemes for $b > 0$* , 2017, <https://hal.inria.fr/hal-01547251v2>.
- [15] H. BAUSCHKE, P. L. COMBETTES, *Convex Analysis and Monotone Operator Theory in Hilbert spaces*, CMS Books in Mathematics, Springer, (2011).
- [16] A. BECK, M. TEBoulLE, *A fast iterative shrinkage-thresholding algorithm for linear inverse problems*, SIAM J. Imaging Sci., 2 (2009), No. 1, pp. 183–202.
- [17] R. I. BOT, E. R. CSETNEK, S.C. LÁSZLÓ, *A second order dynamical approach with variable damping to nonconvex smooth minimization*, to appear in Applicable Analysis, (2018).
- [18] A. CHAMBOLLE, CH. DOSSAL, *On the convergence of the iterates of the Fast Iterative Shrinkage Thresholding Algorithm*, Journal of Optimization Theory and Applications, 166 (2015), pp. 968–982.
- [19] P.L. COMBETTES, V.R. WAJS, *Signal recovery by proximal forward-backward splitting*, Multiscale Model. Simul., 4 (2005), pp. 1168–1200.
- [20] O. GÜLER, *On the convergence of the proximal point algorithm for convex optimization*, SIAM J. Control Optim., 29 (1991), pp. 403–419.
- [21] O. GÜLER, *New proximal point algorithms for convex minimization*, SIAM Journal on Optimization, 2 (1992), No. 4, pp. 649–664.
- [22] D. KIM, J.A. FESSLER, *Optimized first-order methods for smooth convex minimization*, Math. Program. 159 (2016), No. 1, pp. 81–107.
- [23] J. LIANG, J. FADILI, G. PEYRÉ, *Local linear convergence of forward-backward under partial smoothness*, Advances in Neural Information Processing Systems, 2014, pp. 1970–1978.
- [24] D.A. LORENZ, TH. POCK, *An inertial forward-backward algorithm for monotone inclusions*, J. Math. Imaging and Vision, 51 (2015), pp. 311–325.
- [25] R. MAY, *Asymptotic for a second-order evolution equation with convex potential and vanishing damping term*, Turkish Journal of Mathematics, 41 (2017), No. 3, pp. 681–685.
- [26] Y. NESTEROV, *A method of solving a convex programming problem with convergence rate $O(1/k^2)$* , Soviet Mathematics Doklady, 27 (1983), pp. 372–376.
- [27] Z. OPIAL, *Weak convergence of the sequence of successive approximations for nonexpansive mappings*, Bull. Amer. Math. Soc., 73 (1967), pp. 591–597.
- [28] N. PARIKH, S. BOYD, *Proximal algorithms*, Foundations and trends in optimization, volume 1, (2013), pp. 123–231.
- [29] J. PEYPOUQUET, *Convex optimization in normed spaces: theory, methods and examples*. Springer, 2015.
- [30] J. PEYPOUQUET, S. SORIN, *Evolution equations for maximal monotone operators: asymptotic analysis in continuous and discrete time*, J. Convex Anal., 17 (2010), No. 3-4, pp. 1113–1163.
- [31] B.T. POLYAK, *Some methods of speeding up the convergence of iteration methods*, U.S.S.R. Comput. Math. Math. Phys., 4 (1964), pp. 1–17.
- [32] B.T. POLYAK, *Introduction to optimization*. New York: Optimization Software. (1987).
- [33] R.T. ROCKAFELLAR, *Monotone operators and the proximal point algorithm*, SIAM J. Control Optim., 14 (1976), No. 5, pp. 877–898.
- [34] M. SCHMIDT, N. LE ROUX, F. BACH, *Convergence rates of inexact proximal-gradient methods for convex optimization*, NIPS’11 - 25 th Annual Conference on Neural Information Processing Systems, Dec 2011, Grenada, Spain. (2011) HAL inria-00618152v3.
- [35] S. VILLA, S. SALZO, L. BALDASSARRES, A. VERRI, *Accelerated and inexact forward-backward*, SIAM J. Optim., 23 (2013), No. 3, pp. 1607–1633 .
- [36] B. SHI, S. S. DU, M. I. JORDAN, W. J. SU, *Understanding the acceleration phenomenon via high-resolution differential equations*, arXiv:submit/2440124[cs.LG] 21 Oct 2018.
- [37] W. J. SU, S. BOYD, E. J. CANDÈS, *A differential equation for modeling Nesterov’s accelerated gradient method: theory and insights*. Neural Information Processing Systems 27 (2014), pp. 2510–2518.

IMAG, UNIV. MONTPELLIER, CNRS, MONTPELLIER, FRANCE

Email address: hedy.attouch@umontpellier.fr

CADI AYYAD UNIVERSITY, FACULTY OF SCIENCES SEMLALIA, MATHEMATICS, 40000 MARRAKECH, MORROCO

Email address: chbaniz@uca.ac.ma

CADI AYYAD UNIVERSITY, FACULTY OF SCIENCES SEMLALIA, MATHEMATICS, 40000 MARRAKECH, MORROCO

Email address: h-riahi@uca.ac.ma