



# Why Should We Stand for Geothermal Energy ? Example of the Negative Impacts of Oil and Gas Exploration Activities over the Marine Environment

Polina Lemenkova

## ► To cite this version:

Polina Lemenkova. Why Should We Stand for Geothermal Energy ? Example of the Negative Impacts of Oil and Gas Exploration Activities over the Marine Environment. *Geoenergy*, Grozny State Oil Technical University n.a. M.D. Millionshtchikov, Jun 2015, Grozny, Russia. pp.264-273, 10.6084/m9.figshare.7435490 . hal-02021981

**HAL Id: hal-02021981**

**<https://hal.science/hal-02021981>**

Submitted on 22 Mar 2019

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons CC0 - Public Domain Dedication 4.0 International License

11. Керимов И.А., Гайсумов М.Я., Гацаева Л.С. Эффективность использования геотермальных ресурсов Чеченской Республики // Актуальные проблемы геологии, геофизики и геоэкологии Северного Кавказа. Материалы III Всероссийской научно-технической конференции. Грозный: Академия наук Чеченской Республики, 2014. С. 147-151.

12. Маммаев О.А., Маммаев Б.О. Теплоэнергетические геотермальные ресурсы альтернативный источник теплоэнергоснабжения предгорно-равнинных городов и районов Дагестана // Труды Института геологии Дагестанского научного центра РАН. 2009, №33. С. 121-124.

13. Заурбеков Ш.Ш., Минцаев М.Ш., Шайпов А.А., Маммаев Ф.И., Черкасов С.В. Термальные воды Чеченской Республики и перспективы их использования // Материалы Международной научно-практической конференции «Актуальные проблемы защиты окружающей среды и техносферной безопасности в меняющихся антропогенных условиях». «Белые ночи - 2014». Грозный, 2014. С. 253-259.

УДК 550.8

# **WHY SHOULD WE STAND FOR GEOTHERMAL ENERGY? EXAMPLE OF THE NEGATIVE IMPACTS OF OIL AND GAS EXPLORATION ACTIVITIES OVER THE MARINE ENVIRONMENT**

©Lemenkova P.A.

Charles University in Prague, Institute for Environmental Studies, Prague

*Geothermal energy is a clean, environmentally friendly, renewable resource that provides energy around the world. Heat flowing constantly from the interior of the Earth ensure to be an inexhaustible supply of energy. However, existing traditional sources of energy, such as oil and gas are still popular nowadays. Current paper gives a remarkable example of how eco - non friendly and environmentally dangerous these sources of energy are. The given example of oil and gas operations within the Timan-Pechora basin in the shelf and the coast of the Pechora Sea causes contamination of waters by phenol and its accumulation in the bottom sediments and life tissues of the marine habitants. At the same time, the ecosystems of the south-eastern part of the Barents sea, the Pechora Sea, are character-*

*ized by their high biodiversity and high level of primary production. The last one is the fundamental biological characteristics for the marine ecosystems meaning the formation of the organic substance in the water by the chlorophyll-contains organisms – phytoplankton. The primary production plays an exceptional role in the functionality of the marine ecosystem's components as the level of primary production is a basic element in the ecological food-chain hierarchy and therefore, the ecosystem with damaged phytoplankton operation and low level of primary production dooms to die.*

*Therefore, the broad objective of the study was to identify location of main carbon deposits as sources of phenol contaminations accordingly to the nature features of the Pechora Sea ecosystem and to compare them with the distribution of the areas with high biological productivity and higher level of the primary production within the Pechora Sea. The results of the research demonstrate that both are located in the south-eastern part of the Pechora Sea. Thus, the major deposits of oil in Pechora Sea and amongst the Prirazlomnoe oil-field with large reserves of hydrocarbons are situated exactly in the eastern part of the Pechora Sea, some of them are located in immediate proximity to the coast. At the same time, shelf zone of the Pechora Sea is distinguished for its high level of the primary production, high bioproductivity and uniqueness of its natural resources, extremely hardly renewable under the conditions of the polar environment.*

*This paper details question of the ecological interrelation of the natural ecosystems and man's activities in this region, where any negative industrial human impacts can have an un-compensable consequences for the unique environment of this little sea. Also, the capability of aquatic ecosystem to resist towards human negative effects was examined, depending on the structure and character of its fundamental ecological components (e.g. types of bottom sediments, relief of the sea floor, and the direction of currents transferring).*

**Introduction.** At present, among all the types of the sea water contaminations in the Arctic Ocean, the most vital part plays phenol and hydrocarbon contamination, what could be taken up as an example of global technogene contamination. Annual entrance of oil products from different sources into the marine environment of World Ocean in general estimates of about 1.7–8.7 mil tons, what exceeds natural confluent of oil [5]. Main causes of the seawater pollution by phenol and other contaminants are consequences of the oil spilling during the mining works, the damages and the

accidents at the pipelines, shipwrecks, oil drilling and rigs. Besides phenol contamination, petroleum extraction brings others 'satellite' contaminants, e.g. heavy metals, nitric oxides, benz(a)pyrene etc. Phenol contamination is closely related with more and more increasing oil-drilling activities within the marine aquasystems of Barents, Pechora, Norway Seas, its transportation as a fuel of lubricating oils and other secondary chemical products.

Determinative factor in formation of modern ecological conditions of the hydrochemistry in the Pechora Sea has anthropogenic impact, namely, geological activities in the shelf and coastal regions [11]. It is closely connected with the intensity of the exploitation of the Timan-Pechora carbon basin and the Pechora coal-basin – major parts of industrial structure of the Nenets autonomous region. The Timan-Pechora oil-gas carbon basin is the largest source of raw fuel-energy materials in the northern part of Russia. Within the Nenets, autonomous region 76 deposits of oil and gas are discovered, including 64 oil-fields, 6 oil-gas, 4 gas-fields and 1 gas-oil deposit [16].

**Conditions for oil contamination.** Climatically, eastern part of the Pechora Sea differs a lot from its western part, being similar to this of Kara Sea. The temperature of the eastern part of the Pechora region is extreme, very likely to Kara Sea, whereas the air temperatures of the western part of the Pechora are generally continue the Barents Sea climatic conditions. The Pechora Sea has one of the mildest wind climate among other Arctic seas. The magnitude of the wind speed in the neighbouring seas (Barents and Kara) is quite similar, and this of Pechora is differs from them. Meanwhile, the characteristics of the wind (its magnitude and directions) are responsible for different climate, currents and certain ice conditions in the Pechora Sea [6]. Western part of the Barents Sea has the most severe wave climate [6]. The Gulf Stream continues here, thus making western Barents Sea turbulent [12]. In contrast to it, the Pechora Sea is sheltered by consequently lowest waves, compared to the western Barents Sea. A constant current of the water in the Pechora Sea is mostly induced by Kanin, Kolguev and Litke through the Kara Gates. The average speed of the Pechora's currents is about 0.02–0.05 m/s, which is lower than in the Barents Sea with its 0.05–0.5 m/s. Significantly higher speeds of water motion are induced only by the drift and tides. Generally, rather passive tempo of currents in the Pechora Sea calls forth certain stagnation and makes favorable conditions for the holding of the oil spills and contaminants within the waters.

**Table 2**  
**Monthly mean air temperature and extreme minimum  $t^{\circ}\text{C}$  [8].**

	January		February		March		April	
	$T_{\text{mean}}$	$T_{\text{min}}$	$T_{\text{mean}}$	$T_{\text{min}}$	$T_{\text{mean}}$	$T_{\text{min}}$	$T_{\text{mean}}$	$T_{\text{min}}$
Barents Sea (central)	-5.0	-24	-7	-25	-6	-24	-3	-22
Western Pechora Sea	-10.6	-40	-12.6	-40	-11.6	-42	-6.7	-32
<b>Eastern Pechora Sea</b>	<b>-17.5</b>	<b>-48</b>	<b>-18.3</b>	<b>-48</b>	<b>-17</b>	<b>-46</b>	<b>-9.8</b>	<b>-37</b>
Kara Sea	-18.3	-50	-20.1	-50	-20.7	-48	-12.4	-38

**Oil and gas activities in the region of Timan-Pechora.** Actually, four deposits of the oil-fields are being exploited at the territory of Nenets autonomous region with the total size of 3.5 mil ton annually. At present, the main activities that are been carrying out are the as follows: exploration of the Varandey oil gas field (under construction), the exploratory boring and preparation of the Pirazlomnoe deposit for further exploitation and others. Taking into account the oil drilling and production from Pirazlomnoe and Varandey deposits, the volume of carbon extraction will be increased until the 15–20 mil ton annually [16]. Maximal content of the phenols and oil products in the bottom sediments of the Pechora Sea is fixed up in the samples, taken in the region Bolvanskij Nos and also at the sandbar of the Pechora river, where the bottom sediments are represented by the silty sands. High levels of the phenols are fixed in the samples of the bottom sediments, taken by the shelf zone of the cape Konstantinovskiy, which can be explained by the drain of rivers Khylychiya and Dresvyanka where test oil-drilling is being practiced. The content of the phenols in the samples, taken by cape Bolvanovskiy Nos (0.8–0.9 mg/kg) exceeds 8–10 times those in the bottom sediments of the border zone 'bay-sea'. Higher content of the phenols are located in the zones with the siltstone-silty sediments [16].

**Sources of the oil contamination in the Pechora Sea.** One of the main sources of oil contamination in the Pechora Sea is maritime transportation. At present, due to the exploration of the oil-fields in the Pechora and Barents seas, discharge of the oil products is essentially increased both in the process of exploitation of the deposits, and by the transportation. What is more alerting, the intensity of the navigation, shipping and the oil-loading navy fleet is increased recently as well, which brings additional

sources of the oil contamination in the Pechora Sea [12]. During the oil transportation, the oil spilling do take place occasionally, including the wrecking (by its extraction), which often becomes a disaster. Additional source of the oil products into the Barents Sea is presented by the Gulf Stream currents system, which brings its waters into the Barents basin. The Gulf Stream transfers annually ca. 1-1.5 mil tons of phenols and of oil products within the World Ocean. Separate streams of the Gulf Stream, containing phenols and oil products have different places of discharge, namely, at the coasts of the North America and Europe, among which the Sargasso Sea, the Norway Sea and the Barents Sea.

**Distribution of the phenol contaminants within the Pechora Sea.** Unequal geographical distribution of the phenol contaminants within the space of the ecosystem is specified and conditional by the different geographic location of the sources of the contaminations, by their chemical features (e.g. forms of their existing in the water), and what is more important, by the particularities of the circulation of the seawater: its vertical structure and currents.

Maximal concentration of the contaminants in the Pechora Sea is detected in the coastal shelf waters, relatively calm and without intensive mixing, where the damping may occur and where they are being accumulated constantly. Some concentrations are occasionally take place (but definitely lesser) in the vast open regions of the sea, where they are carried out from the rest parts of the Gulf Stream and from the North-Atlantic currents systems, being generally transferred from the Barents Sea [7]. High concentrations of the oil aggregation are fixed along the main transport roots, in the shelf zones of the separated seas, and also in the waters coming from the Atlantic Ocean, carrying contaminations from the western Europe and north-western part of Africa [3].

Near the oil drilling stations and along the main ship transport roots the content of phenols and other oil products is higher than those in the open parts of the sea. In the estuaries where the ecological situation is worse, the concentration of oil in the bottom sediments reaches 5-160 mg/kg, coming down to the 0.1-1 mg/kg [2] in the sediments of the open deep sea regions. Distribution of the oil hydrocarbons in the shallow waters is rather homogeneous (0.05—0.15 mg/l) with maximal concentrations (0.4 mg/l) in the south-eastern part of the sea [9]. Vertical distribution of the pollutants within the water: the maximal concentrations of the oil hydrocarbons in the Pechora sea are located in the shelf waters, zone of interaction 'hydrosphere-atmosphere' [5]. At the same time, this layer of the

water is very important for the biological activities of the aquatic ecosystem and for the functionality of the marine environment: the process of photosynthesis takes place constantly here [1]. Hence, it becomes clear that the contamination of the most important layer of waters may have extreme consequences for the life existing in the ecosystem of this small sea.

**Data processing, methods and results.** The distribution of oil and gas deposits in bottom sediments of Pechora Sea and the contents of the chlorophyll in the shallow waters of the Pechora Sea were visualized on the maps. After digitizing the relief of Pechora basin using Autotrace software, the .shp-file was added to the map as a base layer for the thematic layers [13]. Using maps created by other authors and additional thematic information from the cited literature (containing geological and geomorphological characteristics, features of coastal and land relief, morfostructures of the Pechora submarine sea floor relief etc) the thematic maps were designed with the framework of this research [10]. Data containing various characteristics of objects (concentrations of contaminants) and spatial characteristics of objects were added to the current mapping project. References Maps: [1-27].

For mapping of oil and gas deposits within the Pechora basin its geographical natural specifics were studied, especially its geological, geomorphological conditions and tectonics that have impact on the environmental situation as such and distribution of its carbon deposits within the marine basin. For understanding the sustainability of the aquatic systems to resist the anthropogenic impacts, its climatic and meteorological conditions (current directions) were studied as well. Using information and data from the existing thematic maps and tables (References, Literature and Maps [1-27]) both ecological and geographical data of Pechora Sea were compared: first, its geographic aspects (geological, geomorphological and climatic factors) and second, the distribution of the main oil and gas fields within the Pechora basin, making a negative impact to the marine environment of unique Pechora ecosystem. Thematic information layers were stored in the different topological layers (.shp – files of ArcView system) and some mapping routines were executed for creating the final maps: data editing, overlay of different thematic layers thus making the synthesis of different information, creating of the legend, visualization of the layout. Maps visualize distribution of the main oil and gas deposits and the contents of the chlorophyll in the shallow waters of the Pechora Sea. Both maps show the element of the environment in the basin of the Pechora, demonstrating negative impact and effects of the oil and gas drilling activities under the condi-

tions of Arctic vulnerable tundra ecosystems. Two main zones of phenol concentrations are detected in the bottom sediments of the Pechora Sea: in the central part and in its eastern part (the most polluted area), containing Pirazlomnoe oil-field. In the western region the source of water contamination is not only caused by the geological activities, but also by the inflow of waters from the Barents and the Norway Sea, polluted by phenol and oil. In the south-western part of the Pechora Sea the inflow of the contaminated waters from the narrow entrance of the White and Barents seas can be considered as additional source for the phenol pollution of the Pechora Sea waters. Besides, zone of the high level of contaminants exists in the Kola bay in the Barents sea and in the neighboring aquatory. In the eastern part of the Pechora Sea recent geological activities concerning developing of the Pirazlomnoe oil-field and other smaller oil deposits could give rise to the ecological crisis in the aquatic ecosystems of the Pechora Sea.

**Conclusions.** As demonstrated in this paper, disastrous impacts on the environment is caused by the oil and gas drilling activities in one of the most environmentally precious regions of our world – Arctic. So, what can we do with it, how can we handle such a problem? Alternative sources of energy, such as geothermal energy, or wind energy can be a perfect replacement for the traditional sources. Moreover, geothermal energy can be effectively used as a supplement for the integrity of the power grid, enhancing the efficiency of the entire system while providing clean, reliable power for the humanity. Geothermal energy is also capable of achieving high capacity factors in results without damaging the ecosystems and the world. Current paper contributed to the popularization of the effective replacement of traditional (oil and gas) energy source by new-generation "green" energy, such as geothermal or wind.

**Acknowledgements:** Current research has been generously supported by the DAAD-Program (Referat 325-syp, Kennziffer A), at Alfred Wegener-Institut für Polar- und Meeresforschung Fachbereich: Geowissenschaften Arbeitsbereiche Bathymetrie und Geodäsie, Bremerhaven, Germany.

#### References:

1. Arild Moe ET Peter Johan Schei. (2006): Le Grand Nord. Debris & Potentials. <http://www.fni.no/doc&pdf/nordiques-2006.PDF>
2. Aibulatov N.A. (2001): Geocology of shelf and coasts of seas in Russia. - Noosphere, Moscow: 1-428.

3. Ole Gunnar Austvik (2006): Oil and gas in the High North - A perspective from Norway.
4. 5Bikmametova M., Bakalov V. (1997): Ecological problems of World Ocean.- MEO Chelyabinsk MGUK, Chelyabinsk: 13-14.
5. Dmitriev V.V. (1997): Diagnostics of ecological control and assessment of sustainability of water ecosystems towards anthropogenic impacts. Oceanology at St-Petersburg University, St-Petersburg: 186- 211, 269.
6. Golovin, N., Koptev, N., Maiorov, and O., Khvedynych (1996): Ice investigations in the Pechora and Kara Seas in the connection with a shelf development. Proc. Polar Tech. Conference, Moscow: 4, 38-39.
7. Gorshkov S.G. (1980): Atlas of the Oceans. Arctic Ocean, Arctic and Antarctic Research Institute, Ministry of defence USSR. Main Department of the Navigation and Oceanography, Moscow, USSR.
8. Gurevich V.I. (2002): Modern sedimentogenesis and geocology of West-Arctic shelf of Eurasia. - Scientific World, Moscow: 1-218.
9. Ilyin G.V., Petrov V.S. (1994): Research of the contents of the heavy metals in the grounds. Ecological researches of the industrial development in the zone of Shtokmanovskoe gas condensate deposits on the Barents sea shelf.- MMBI RAS, Apatiti: 44-48.
10. Kasimov N.S (1997): Complex ecological mapping (geographical aspect).- Moscow State University, ISBN 5-211-036654, Moscow: 1-146
11. Klenova M.V. (1967): Geology of Barents Sea- RAS USSR, Moscow: 1-367
12. Mennobart van Eerden, Pechora River Basin Integrated System Management.
13. Preobragensky V.S. (1990): Ecological maps (contents, requirements).- News RAS USSR. Series Geography (6): 51-56.
14. 15Romankevich E.A., A.P.Lisitzin, M.E.Vinogradov. (2003): Pechora Sea: integrated research.- Sea, Moscow: 1-352

#### References: Maps

1. AARI Team (1985): Distribution of waters from the Atlantic and Pacific Oceans in the Arctic (Scale 1:20000000), Atlas of Arctic. - Arctic and Antarctic Research Institute, Ministry of defence USSR, Moscow.
2. AARI Team (1985): Geological map of the Arctic ocean (Scale 1:10000000).- Atlas of Arctic, Arctic and Antarctic Research Institute SDGC Council of Ministers, USSR, Moscow
3. AARI Team (1985): Geomorphological map of the Arctic, (Scale 1:20000000).- State Scientific Center of the RF the Arctic and Antarctic Research Institute, Council of Ministers USSR, Moscow

4. AARI Team (1980): Partition of the bottom in the Arctic Ocean (Scale 1:15000000).- Arctic and Antarctic Scientific Research Institute (AARI). Ministry of defence USSR, Moscow.

5. AARI Team (1985): Prospect of the oil-gas production (1:20000000). Atlas of Arctic. AARI SDGC Council of Ministers USSR, Moscow

6. Central Intelligence Agency Team (1978): Mining (Scale 1:20000000).- Atlas of Polar Regions, Central Intelligence Agency, Washington, USA.

7. Central Intelligence Agency Team (1978): Oil and Gas in Arctic Region, Mackenzie Delta/Beaufort Sea, and Lower Ob' basin, Pechora (1:50000000). - Atlas of Polar Regions. Central Intelligence Agency, Washington, DC.

8. Geophysical Atlas of the Barents Sea, [http://www2.tgsnopec.com/whatsnew/Barents\\_Sea\\_geo\\_atlas.htm](http://www2.tgsnopec.com/whatsnew/Barents_Sea_geo_atlas.htm)

9. Gorshkov S.G. (1980): Types of the coasts in the Arctic, projection asimuthal equal-distance (1:15000000). Atlas of the Oceans. Arctic Ocean, AARI, Ministry of Defence USSR, Moscow, USSR.

10. La Direction Norvegienne Du Petrole (2006): Découvertes et exploitation en mer de Barents.- <http://www.fni.no/doc&pdf/nordiques-2006.PDF>

11. Matishov G.G. & Veslavsky S. (1995): Ecology of Novaya Zemlya region, (1:2000000). Institute of Oceanology, MMBI, RAS.

12. Oil and Gas development in Northern Russia, <http://www.kaldor.no/energy/berlin20060315-barents.html>

13. Okulich A.V., Lopatin B.G. & Jackson H.R. (1983): Carte géologique circumpolaire de l'Arctique, Carte 1765A, (Scale 1:6000000), Projection stéréographique polaire a 75 de Latitude N.- Commission géologique du Canada; Department of Energy, Mines and Resources; Institut de Sedimentaire and Petroleum Geology, Montreal, Quebec, Canada.

14. Pennwell Team (1985): Polar Oil and Gas. Arctic Ocean (Scale 1:10000000).- Pennwell Maps, Pennwell Publishing

15. Petroleum reserves and borders in the Barents Sea <http://www.kaldor.no/energy/berlin20060315-barents.html>

16. Rekacewicz Ph. (1998): Environmental threats in the Barents Region (Scale 1:20000000). - UNEP/GRID-Agenda, BarentsWatch Atlas, Oslo, Norway.

17. Rekacewicz Ph. (1998): Oil and gas development and seabird's colonies in the Barents Region (Scale 1:20000000). - BarentsWatch Atlas, Norwegian Polar Institute, UNEP/GRID-Agenda, Oslo.

18. Scott R. A., Natapov L.M., Anderson L. M., S.C.Jamie, W.C.F.Hanni (1995): Tectonic elements of the Arctic (Scale 1:10000000) - Cambridge University, Cambridge Arctic Shelf Program, Cambridge, UK

19. Sigmund Ellen M.O. (2002): Geological map. Land and Sea Areas of Northern Europe (Scale 1:4000000).- Geological Survey of Norway, Oslo

20. Team Commission de la Carte Géologique (1989): Carte tectonique internationale de l'Europe, Feuille 3. (Échelle 1:6000000) - Congrès Géologique International Commission de la Carte Géologique du Monde Sous-Commission de la Carte tectonique du Monde, Canada.

21. Team Ministry of Geology (1969): Tectonic map of polar regions of the Earth, 4 sheets with explanatory note, (Scale 1:10000000).- Ministry of Geology USSR, Scientific Research Institute of Arctic' Geology, Moscow

22. Team PennwellMaps (2001): Main oil deposits exploitation in the Barents Sea <http://images.pennwellnet.com/ogj/images/off/56071201.gif>

23. Tkachenko B.V. & Egiasarov B.Kh. (1970): Observed geologic map of islands of Soviet Arctic and adjoining areas (1:7500000). Council of Ministers USSR, "Geology of USSR", v. XXVI. USSR, Moscow.

24. U.S. Navy Hydrographic Department Team (1958): Bottom sediments in Arctic Ocean (Scale 1:20000000). Oceanographic Atlas of the Polar Seas. Part II. Arctic Ocean. U.S. Navy Hydrographic Office Washington, D.C.

25. U.S. Navy Hydrographic Department Team (1958): Circulation of Atlantic water in the Arctic Ocean and Adjacent seas (Scale 1:20000000).- Oceanographic Atlas of the Polar Seas. Part II. Arctic Ocean. U.S. Navy Hydrographic Office Washington, D.C.

26. U.S. Navy Hydrographic Department Team (1958): Generalized mayor rock types in Arctic Ocean (Scale 1:20 mln) Oceanographic Atlas of the Polar Seas. Part II. Arctic Ocean. U.S. Navy Hydrographic Office Washington D.C.

27. U.S. Navy Hydrographic Department Team (1958): Geologic structures in Arctic Ocean (Scale 1:20000000). - Oceanographic Atlas of the Polar Seas. Part II. Arctic Ocean. U.S. Navy Hydrographic Office Washington, D.C.