



**HAL**  
open science

## **BRANCHED1: A Key Hub of Shoot Branching**

Ming Wang, Marie-Anne Le Moigne, Jessica Bertheloot, Laurent Crespel,  
Maria Dolores Perez-Garcia, Laurent Ogé, Sabine Demotes-Mainard, Latifa  
Hamama, Jean-Michel Daviere, Soulayman Sakr

► **To cite this version:**

Ming Wang, Marie-Anne Le Moigne, Jessica Bertheloot, Laurent Crespel, Maria Dolores Perez-Garcia, et al.. BRANCHED1: A Key Hub of Shoot Branching. *Frontiers in Plant Science*, 2019, 10, pp.1-12. 10.3389/fpls.2019.00076 . hal-02025844

**HAL Id: hal-02025844**

**<https://hal.science/hal-02025844v1>**

Submitted on 19 Feb 2019

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



# BRANCHED1: A Key Hub of Shoot Branching

Ming Wang<sup>1</sup>, Marie-Anne Le Moigne<sup>1</sup>, Jessica Bertheloot<sup>1</sup>, Laurent Crespel<sup>1</sup>, Maria-Dolores Perez-Garcia<sup>1</sup>, Laurent Ogé<sup>1</sup>, Sabine Demotes-Mainard<sup>1</sup>, Latifa Hamama<sup>1</sup>, Jean-Michel Davière<sup>2</sup> and Soulayman Sakr<sup>1\*</sup>

<sup>1</sup> Institut de Recherche en Horticulture et Semences, Agrocampus-Ouest, INRA, SFR 4207 QUASAV, Université d'Angers, Beaucauzé, France, <sup>2</sup> Institut de Biologie Moléculaire des Plantes, UPR2357, Université de Strasbourg, Strasbourg, France

Shoot branching is a key process for plant growth and fitness. Newly produced axes result from axillary bud outgrowth, which is at least partly mediated through the regulation of *BRANCHED1* gene expression (BRC1/TB1/FC1). *BRC1* encodes a pivotal bud-outgrowth-inhibiting transcription factor belonging to the TCP family. As the regulation of *BRC1* expression is a hub for many shoot-branching-related mechanisms, it is influenced by endogenous (phytohormones and nutrients) and exogenous (light) inputs, which involve so-far only partly identified molecular networks. This review highlights the central role of BRC1 in shoot branching and its responsiveness to different stimuli, and emphasizes the different knowledge gaps that should be addressed in the near future.

## OPEN ACCESS

### Edited by:

Junko Kyoizuka,  
Tohoku University, Japan

### Reviewed by:

Philip B. Brewer,  
The University of Adelaide, Australia  
Laura Rossini,  
University of Milan, Italy

### \*Correspondence:

Soulayman Sakr  
soulayman.sakr@agrocampus-ouest.fr

### Specialty section:

This article was submitted to  
Plant Physiology,  
a section of the journal  
Frontiers in Plant Science

**Received:** 12 October 2018

**Accepted:** 17 January 2019

**Published:** 12 February 2019

### Citation:

Wang M, Le Moigne M-A, Bertheloot J, Crespel L, Perez-Garcia M-D, Ogé L, Demotes-Mainard S, Hamama L, Davière J-M and Sakr S (2019) BRANCHED1: A Key Hub of Shoot Branching. *Front. Plant Sci.* 10:76. doi: 10.3389/fpls.2019.00076

**Keywords:** TCP transcription factors, hormones, nutrients, light, regulation, shoot branching

## INTRODUCTION

Plants are sessile organisms that need to adjust their shape to suit the diversity of the changing environmental conditions in which they are growing. The regulation of shoot branching is a relevant strategy for plant survival and space occupancy, and involves an intricate regulatory network. Shoot branching depends on the status of bud dormancy, which is a temporary and reversible state (Shimizu and Mori, 1998). Shoot branching patterns, considered here as the distribution of branches along a parent stem, are generated during plant postembryonic development (Domagalska and Leyser, 2011). They depend on the ability of axillary vegetative buds located at the axil of each leaf to remain inactive or to produce a new branch in response to variable stimuli (Shinohara et al., 2013; Rameau et al., 2015; Wang and Jiao, 2018).

Shoot branching is an important feature of plant architecture that determines the interface between the plant and the surrounding environment. Shoot branching contributes to essential processes such as the establishment of leaf area and distribution that determine light interception and photosynthesis, which in turn influence the number of flowers and fruits, fruit filling and yield (Jiang and Egli, 1993; Richards, 2000). Branching also influences the plant competitiveness against weeds or the propagation of pests (Lemerle et al., 1996; Zhao et al., 2006; Simon et al., 2011). In ornamental plants, branching also determines plant visual quality, which drives consumers' preferences (Ta et al., 1987; Boumaza et al., 2009, 2010; Garbez et al., 2015).

Extensive studies have been undertaken for several decades to find out the mechanisms involved in branching. The currently accepted idea supports that endogenous, developmental, and environmental inputs converge into bud-located integrators, which are at the head of a network of mechanisms governing the ability of buds to grow out. Among these inputs, hormones, sugar,

nitrogen, light, and water play a determining role in shoot branching regulation (McSteen, 2009; González-Grandío et al., 2013; Niwa et al., 2013; Li-Marchetti et al., 2015; Rameau et al., 2015; Teichmann and Muhr, 2015; Corot et al., 2017; Le Moigne et al., 2018). Those factors may influence shoot branching via various physiological and molecular mechanisms, targeting different branching-related genes and acting synergistically or antagonistically. *BRC1* (*BRANCHED 1*) is well known to act locally in buds and is considered to be an important hub of different signals controlling the ability of a bud to grow out in many species (Aguilar-Martínez et al., 2007; Dun et al., 2009; Leyser, 2009; Beveridge and Kozuka, 2010; Rameau et al., 2015). *Arabidopsis thaliana* harbors two *BRANCHED* genes, namely *BRANCHED 1* (*BRC1*) and *BRANCHED 2* (*BRC2*); they encode TCP transcription factors closely related to *TEOSINTE BRANCHED1* (*TB1*) in maize and *FINE CULM 1* (*FC1*) in rice. In addition, they are conserved in many species of the plant kingdom (Table 1). The corresponding mutants show an altered branching phenotype as compared to the wild type (Aguilar-Martínez et al., 2007; González-Grandío et al., 2013). This review addresses the molecular identity of *BRC1*, its involvement in shoot branching, and its regulation in response to endogenous inputs (hormones and nutrients) and exogenous cues (light). We also discuss how *BRC1* can mechanistically govern bud outgrowth, and raise a few questions about future investigations.

## BRC1 BELONGS TO THE TCP TRANSCRIPTION FACTOR FAMILY

*AtBRC1* (also called *AtTCP18*) contains an open reading frame (ORF) made of ca.1,290-bp that encodes a protein with a TCP domain and an R domain. It belongs to the TCP gene family, an evolutionarily conserved family that first appeared in freshwater algae of the Charophyta family (Navaud et al., 2007). The TCP gene family was first described by Cubas et al. (1999) and is represented by four ‘founding members’: *TEOSINTE BRANCHED1* (*TB1*), *CYCLOIDEA* (*CYC*), *PROLIFERATING CELL NUCLEAR ANTIGEN FACTOR1* (*PCF1*), and *PCF2*, all identified on the basis of their functions in plant development or their DNA-binding capacities (for a review see Li, 2015; Danisman, 2016). In *Arabidopsis*, the TCP family comprises 24 genes encoding predicted proteins with a TCP domain (Cubas et al., 1999; Kosugi and Ohashi, 2002; Palatnik et al., 2003; Cubas, 2004) and categorized into two classes: class I (also known as PCF or TCP-P) is made up of 13 predicted proteins related to the PCF rice factors (Kosugi and Ohashi, 1997), and class II (also known as TCP-C) is made up of 11 predicted proteins related to the *Antirrhinum* *CYC* and *CIN* genes and to the *Zea mays* *TB1* gene (Luo et al., 1996; Doebley et al., 1997; Nath et al., 2003; Palatnik et al., 2003). All these transcription factors have the so-called TCP domain, a 59-amino-acid basic helix–loop–helix (bHLH), in common (Martín-Trillo and Cubas, 2010). Such a motif allows for DNA binding and protein–protein interactions in cells. The TCP domain is also necessary for nuclear localization (Kosugi and Ohashi, 1997; Cubas et al., 1999), and some TCP proteins

can be targeted to the nucleus in heterologous systems (Suzuki et al., 2001; Qin et al., 2004).

Besides the TCP domain, a few class-II TCPs, including *BRC1*, display a functionally unknown arginine-rich motif, the R-domain, which is predicted to mediate protein interactions (Lupas et al., 1991; Cubas et al., 1999). The R domain may involve the phosphorylation process of *BRC1* by a cAMP-dependent protein kinase (Dulhanty and Riordan, 1994; Martín-Trillo and Cubas, 2010). Additionally, most members of the *CYC/TB1* subclass, to which *BRC1* belongs, contain a conserved ECE (glutamic acid-cysteine-glutamic acid) motif that remains functionally uncharacterized and is located between their TCP and R domains (Howarth and Donoghue, 2006).

The TCP proteins of various species regulate many biological processes, including seed germination, plant branching, lateral organ development, floral asymmetry, gametophyte development, leaf senescence, circadian rhythms, and defense responses (for a review see Li, 2015; Danisman, 2016). These TCP-dependent regulations could occur directly through their binding to the promoter of target genes or indirectly *via* their interactions with plant hormones (Schommer et al., 2008; Guo et al., 2010; Danisman et al., 2012; Li and Zachgo, 2013; Nicolas and Cubas, 2016). In *Arabidopsis*, the *CYC/TB1* clade consists of *AtBRC1*, *AtBRC2* (also called *AtTCP12*) and *AtTCP1*, and is mainly involved in the development of axillary meristems, giving rise to either flowers or lateral shoots (Martín-Trillo and Cubas, 2010).

## BRC1 IS A CENTRAL ACTOR OF SHOOT BRANCHING

The shoot axillary meristem produces a branch when the appropriate endogenous and exogenous inputs occur, so as to adapt plant architecture to environmental conditions. In monocots, *TB1* from *Z. mays* (Doebley et al., 1997) and homologs of *TB1* in *Oryza sativa* (*OsTB1/FC1*, Takeda et al., 2003) and

**TABLE 1** | The publication of *BRC1* homolog genes in different species.

Species	Name of the gene	Reference	
Monocots	<i>Zea mays</i>	<i>TB1</i>	Doebley et al., 1997
	<i>Oryza sativa</i>	<i>Ostb1/FC1</i>	Takeda et al., 2003
	<i>Sorghum bicolor</i>	<i>SbTB1</i>	Kebrom et al., 2006
	<i>Hordeum vulgare</i>	<i>INTERMEDIUM-C</i>	Ramsay et al., 2011
	<i>Triticum aestivum</i>	<i>TB-D1</i>	Dixon et al., 2018
Eudicots	<i>Solanum tuberosum</i>	<i>StBRC1</i>	Nicolas et al., 2015
	<i>Pisum sativum</i>	<i>PsBRC1</i>	Braun et al., 2012
	<i>Dendranthema grandiflora</i>	<i>DgBRC1</i>	Chen et al., 2013
	<i>Arabidopsis thaliana</i>	<i>AtBRC1</i>	Aguilar-Martínez et al., 2007
	<i>Solanum lycopersicum</i>	<i>SIBRC1</i>	Martín-Trillo et al., 2011
	<i>Rosa hybrida</i>	<i>RhBRC1</i>	Barbier et al., 2015
	<i>Nicotiana tabacum</i>	<i>NtBRC1a; NtBRC1b; NtBRC1c; NtBRC1d</i>	Chen et al., 2016
	<i>Populus canescens</i>	<i>PcBRC1</i>	Muhr et al., 2018

*Sorghum bicolor* (SbTB1, Kebrom et al., 2006) promote bud arrest locally, without affecting the number of buds, and thus lead to reduced tillering. Consistently, *TB1* and *OsTB1* are mainly expressed in axillary bud meristems (Hubbard et al., 2002; Takeda et al., 2003), and their mutants *tb1* and *fc1* exhibit over-tillering phenotypes (Doebley et al., 1997; Wang et al., 1999; Takeda et al., 2003). The barley *TB1* ortholog, *INT-C*, has been shown to act mainly in the control of spike architecture, with a minor role in tillering (Ramsay et al., 2011). Moreover, modern maize displays less branching than the wild teosinte ancestor due to increased *TB1* expression (Studer et al., 2011; Zhou et al., 2011). However, the *int-c* loss-of-function mutant showed less tillers in barley, whose phenotype is opposite to the recessive *tb1* mutant in maize (Liller et al., 2015; Dong et al., 2019).

In dicots, genes closely related to *TB1* have been studied in a variety of species. In *Arabidopsis*, *AtBRC1* and *AtBRC2* both negatively regulate the branching process (Aguilar-Martínez et al., 2007; Poza-Carrión et al., 2007). However, *AtBRC1* seems to play a more pivotal role in axillary bud development than *AtBRC2*. The *AtBRC1* gene is predominantly expressed during the development of axillary buds (axillary meristems, bud leaf primordia and subtending vascular tissue). *AtBRC1* expression is inversely correlated with bud outgrowth and *brc1* mutant phenotypes are non-pleiotropic, while constitutive overexpression of *AtBRC1* reduces the growth of the whole plant (Aguilar-Martínez et al., 2007). Moreover, many *AtBRC1*-homologous genes have also been found to be involved in shoot branching suppression (Table 1). In addition, repressed buds in pea have been found to be as metabolically active as growing buds, so *BRC1* growth repression may not involve metabolism (Stafstrom and Sussex, 1988). Recent data demonstrate that *AtBRC1* is not always necessary for the complete inhibition of all buds in *Arabidopsis* (Seale et al., 2017).

Genomic sequences of *Solanum* species, including potato and tomato, also contain the *BRC1*-like gene, where it occurs under two forms (Brewer, 2015). More interestingly, in *Solanum tuberosum*, the *BRANCHED1a* (*StBRC1a*) gene encompasses an alternative splice site leading to the generation of two *BRC1a* protein isoforms, *BRC1a<sup>Long</sup>* and *BRC1a<sup>Short</sup>*, with distinct C-terminal regions (Martín-Trillo et al., 2011; Nicolas et al., 2015). The *BRC1a<sup>Long</sup>* C-terminal region has a strong activation domain and moves to the nucleus, whereas the *BRC1a<sup>Short</sup>* C-terminal region lacks an activation domain, which prevents the nuclear targeting of the protein (Nicolas et al., 2015). These different splice variants of *AtBRC1* have also been found in *Arabidopsis* (data not shown), but whether the mechanism mentioned above exists in *Arabidopsis* is still unknown. A central role of *BRC1* in shoot branching has also been revealed in pea (*PsBRC1*, Braun et al., 2012), *Chrysanthemum* (*DgBRC1*, Chen et al., 2013), and poplar (*PcBRC1*, Muhr et al., 2016 and 2018). In *Rosa* sp., Li-Marchetti et al. (2017) carried out a Quantitative Trait Loci (QTL) analysis of the plant architecture, using a segregating, recurrent blooming population called 'The Fairy' × 'Old Blush'. They showed that the branching angle of order 2 long axes, the number of short axes (the type of axis that comprises one to four internodes), and stem elongation were correlated, with QTL located in the genomic region of *RhBRC1*,

and assumed a pleiotropic role of *RhBRC1* in the establishment of the bushy shape of *Rosa* sp. Further work will be required to more accurately define the role of *BRC1* in the establishment of the plant complex architecture.


## BRC1 IS AN INTEGRATOR OF DIVERSE HORMONAL SIGNALING NETWORKS

Auxin, cytokinins (CK), and strigolactones (SL) are implicated in the hormonal regulation of *BRC1* expression. In this regulation network, auxin and SL act as inducers while CK act as repressors (Rameau et al., 2015; Teichmann and Muhr, 2015). According to Ferguson and Beveridge (2009), this kind of regulation could be involved in various metabolism pathways such as feedback regulation, long-distance hormone transport, and the interplay of plant hormone metabolism and signaling.

In apical dominance, the polar auxin transport (PAT) stream in the main stem, which is mediated by the PIN (PIN-FORMED) auxin-efflux facilitators located in xylem-associated cells (Petrášek and Friml, 2009), inhibits axillary bud outgrowth (Morris, 1977; Li and Bangerth, 1999; Zhang et al., 2007; Balla et al., 2011). Auxin cannot directly regulate *BRC1* expression because it is not transported from the stem to the buds in great enough amounts (Hall and Hillman, 1975). It is hypothesized that PAT prevents the establishment of auxin canalization from axillary buds to the stem, and that this might be necessary for the buds to grow out (Li and Bangerth, 1999; Domagalska and Leyser, 2011; Chabikwa et al., 2019). The characterization of the auxin-resistant *Arabidopsis* mutant *axr1* indicated that such an auxin effect occurred after axillary meristem initiation through the inhibition of bud outgrowth (Stirnberg et al., 1999).

Auxin can indirectly promote *BRC1* expression in the bud (Aguilar-Martínez et al., 2007). Furthermore, auxin-mediated *BRC1* regulation through the control of two antagonistic factors, CK and SL, fine-tunes *BRC1* expression inside buds (Rameau et al., 2015). The role of CK, a collection of adenine-related compounds, in bud outgrowth was evidenced decades ago, when CK application to dormant buds was shown to promote bud outgrowth (Wickson and Thimann, 1958; Sachs and Thimann, 1967; Bangerth, 1994; Tanaka et al., 2006). In parallel, auxin indirectly inhibits bud outgrowth by decreasing systemic and local CK levels, which determines the CK supply to the buds (Miyawaki et al., 2004; Nordström et al., 2004; Tanaka et al., 2006; Müller and Leyser, 2011). CK can act to promote branching partly by promoting PIN<sub>3,4,7</sub> cross-stem auxin transport between the bud and the adjoining stem, thereby potentially acting partly independently of *AtBRC1* repression directly in the bud (Waldie and Leyser, 2018). High CK levels in axillary buds lead to the activation of axillary buds through downregulation of *BRC1* expression (Braun et al., 2012), although *Psbrc1* (a pea *BRC1* mutant) remained sensitive to CK application. These findings might indicate that the branch-promoting hormone CK partly controls shoot branching by negatively regulating *BRC1* at the transcriptional level. In rice, transcript levels of *OsTB1/FC1* also decreased in a CK-dose-dependent manner (Minakuchi et al., 2010), and similar down-regulation of *DgBRC1* was reported





in *Chrysanthemum* (Dierck et al., 2016). This CK-dependent *BRC1* regulation can be part of the light intensity-dependent bud outgrowth regulation in *Rosa* sp. (Roman et al., 2016; Corot et al., 2017). The *Arabidopsis altered meristem program1* (*amp1*) mutants are characterized by higher levels of CK, more bud outgrowth, more axillary meristems, and reduced *BRC1* expression (Helliwell et al., 2001). Although CK are a powerful repressor of *BRC1/TB1/FC1* expression, the molecular mechanisms driven by this CK-dependent regulation still remain an open question (Figure 1).

Strigolactones (SL), a group of carotenoids derived from terpenoid lactones (Lin et al., 2009; Alder et al., 2012), act as endogenous shoot branching inhibitors (Gomez-Roldan et al., 2008; Leyser, 2008; Wang et al., 2013). Direct application of GR24 – an SL analog – on buds inhibited outgrowth on intact and decapitated plants (Brewer et al., 2009), and auxin application elevated the transcription levels of SL biosynthesis genes (Sorefan et al., 2003; Foo et al., 2005; Johnson et al., 2006; Zou et al., 2006; Arite et al., 2007, 2009; Hayward et al., 2009). These findings support that auxin-mediated bud outgrowth inhibition involves the promotion of systemic and local SL synthesis in the stem and thereby of SL levels inside buds. Consistently, different SL mutants exhibited a highly branched phenotype in pea [*ramosus* (*rms*)], petunia [*decreased apical dominance* (*dad*)], and *Arabidopsis* [*more axillary growth* (*max*)] (Crawford et al., 2010). A role for *BRC1* downstream of SL was first reported in *Arabidopsis* and pea, where *BRC1* expression was upregulated by

SL, and shoot branching in the *brc1* mutant was insensitive to SL (Aguilar-Martínez et al., 2007; Dun et al., 2012; Revel et al., 2015). However, SL application did not change the transcriptional activation of *OsTB1/FC1* expression in rice (Minakuchi et al., 2010). Recent investigations showed that *DWARF 53* (*D53*)/*SUPPRESSOR OF MAX2 1-LIKE* genes (*SMXL6*, 7, 8) acted downstream of SL as repressors of SL-dependent *BRC1* upregulation and thereby promoted shoot branching (Jiang et al., 2013; Zhou et al., 2013; Kong et al., 2014; Wang et al., 2015). Mutants deficient in *D53*-like genes indeed displayed constitutive *BRC1* upregulation (Soundappan et al., 2015; Wang et al., 2015; Seale et al., 2017). Moreover, SL perception by D14 ( $\alpha/\beta$  hydrolase) and the recruitment of the SCF complex resulted in the polyubiquitination and 26S-proteasome-mediated degradation of D53 (Kerr and Beveridge, 2017; Waters et al., 2017). D53 physically interacts with IPA1 (IDEAL PLANT ARCHITECTURE1), a repressor of shoot branching, and prevents it from upregulating *TB1* expression (Figure 1) (Song et al., 2017). IPA1, also named OsSPL14, is a member of the SQUAMOSA PROMOTER BINDING PROTEIN-LIKE (SPL) family of plant-specific transcription factors (Miura et al., 2010) that directly binds to the *TB1* promoter in rice and activates *TB1* transcriptional activity (Figure 1; Jiao et al., 2010; Lu et al., 2013). Further support for the relevance of the “IPA-1-related genes and *TB1*” module in shoot branching comes from a study in wheat, where TaD53 physically interacted with TaSPL3 and prevented TaSPL3 upregulation of *TaTB1* gene expression

(Liu et al., 2017). Although the *Arabidopsis* homologs of *IPA1* have been identified as being *SPL9/15*, further work will be required to confirm whether this mechanism is involved in the SL-dependent regulation of *AtBRC1*.

Besides auxin, CK, and SL, gibberellin (GA) might also be involved in the regulation of *BRC1* expression, even if the mechanism is still unknown (Lantzouni et al., 2017). GAs (diterpenoid tetracyclin molecules) are plant hormones that regulate various developmental processes, including stem elongation, germination, dormancy, flowering, flower development, and leaf and fruit senescence (Hedden and Spensel, 2015). In *Rosa* sp., GA biosynthesis strongly increases during bud outgrowth (Choubane et al., 2012). In the perennial woody plant *Jatropha curcas*, GA and CK synergistically promote lateral bud outgrowth, and both hormones negatively influence *BRC1* and *BRC2* expression (Ni et al., 2015). Simultaneously altered GA and SL levels positively influenced the expression of the *GA2 OXIDASE2* gene which encodes a GA-catabolic enzyme, and the expression of *BRC1* (Figure 1) (Lantzouni et al., 2017). Furthermore, GA is required for CK-mediated axillary bud outgrowth in *A. thaliana* (Jasinski et al., 2005; Lo et al., 2008).

## BRC1 EXPRESSION IS REGULATED BY LIGHT

Shoot branching is negatively affected by low light intensity and low ratios of red/far red (R:FR) light in many species (Kebrom et al., 2006; Finlayson et al., 2010; Su et al., 2011; Revel et al., 2015). In this process, light acts both as a driver of photosynthesis for the supply of sugars to axillary buds and as a photomorphogenic signal (Su et al., 2011). The signaling role of light in plant branching was first unraveled by Kebrom et al. (2006). In 2006 and 2010, these authors showed that active PHYB suppressed the expression of the *SbTB1* gene in sorghum, leading to high plant branching, whereas environmental conditions that inactivate phyB (low R/FR ratio) increased *SbTB1* expression and in turn repressed bud outgrowth. Additional experiments carried out in *Arabidopsis* confirmed these findings: a low R/FR ratio favored *AtBRC1* upregulation through the PHYB pathway, which is required for shoot branching reduction (Figure 1; González-Grandío et al., 2013). This effect seems to be reversible, as evidenced by the rapid and local downregulation of *AtBRC1* after increasing the R/FR ratio (Holalu and Finlayson, 2017). Such a response may contribute to the rapid adaptation of plants to fluctuations in the R/FR light ratio.

Besides light quality, a slight decrease of the photosynthetic leaf area is associated with a stimulation of *TBI* expression in sorghum seedlings and consequently a lower propensity of tiller buds to grow out (Kebrom and Mullet, 2015). In addition, darkness-exposed *Rosa* sp. exhibited no bud outgrowth and higher levels of *RhBRC1* transcripts than plants placed under light (Roman et al., 2016). All these findings indicate that *BRC1* expression is very sensitive to both light intensity and quality. However, this regulation may involve distinct mechanisms (Kebrom et al., 2010).

## BRC1 IS REGULATED BY NUTRIENTS

Sugars are well known to promote bud outgrowth in many species (Leduc et al., 2014; Rameau et al., 2015; Kebrom, 2017; Tarancón et al., 2017; Ferreira et al., 2018), and the relationship between sugars and bud outgrowth has been investigated for years (Maurel et al., 2004; Chao et al., 2007; Girault et al., 2010; Kebrom et al., 2010, 2012; Henry et al., 2011; Rabot et al., 2012; Mason et al., 2014; Barbier et al., 2015; Fichtner et al., 2017). Sugar effects are seemingly dependent on environmental conditions (Corot et al., 2017). Sugars not only serve as a carbon source for plant metabolism, but also as an important signaling entity that affects many developmental processes including *BRC1* gene expression (Price et al., 2004; Hellmann and Smeekens, 2014; Barbier et al., 2015; Sakr et al., 2018). In an interesting study, Mason et al. (2014) demonstrated that the initial signal responsible for the release of bud outgrowth after decapitation in pea was an increase in sugar availability rather than a decrease in apically supplied auxin, as traditionally thought. This is in line with the earlier proposal by Morris and collaborators (Morris et al., 2005), who assumed the existence of an auxin-independent “fast-decapitation signal” leading to bud outgrowth initiation after decapitation. Furthermore, Mason et al. (2014) also reported that the timing of the increase of the sugar flux inside buds and bud outgrowth tightly coincided with the downregulation of *BRC1* expression. In this process, sugar acts more likely as a signaling entity, because many non-metabolizable sugar analogs can trigger bud outgrowth (Rabot et al., 2012) and repress *BRC1* expression (Barbier et al., 2015). In addition, this effect of sugar on *BRC1* transcription could be mediated indirectly *via* sugar regulation of CK biosynthesis and SL signaling (Barbier et al., 2015) and/or directly (irrespective of hormonal action). Decapitation led to a rapid and sustained rise in trehalose-6 phosphate (T6P) levels in axillary buds and a decreased expression level of *BRC1*, which supports that T6P could partly mediate the sugar-dependent down-regulation of *BRC1* (Figure 1) (Fichtner et al., 2017). Further works are required to further unravel this molecular regulatory network. In the present state of knowledge, we cannot rule out that the transcriptional regulation of *BRC1* in response to sugars could involve many sugar-signaling pathways and also that *BRC1* expression is sensitive to the plant carbon status and/or energy levels (Martín-Fontecha et al., 2018).

Mineral nutrition influences tiller bud outgrowth in barley (Fletcher and Dale, 1974). In wheat, phosphorus deficiency directly altered the normal pattern of tiller emergence by reducing the rate of tiller emergence for each tiller (Rodríguez et al., 1999). Although several links exist between phosphate and the branching-related hormones (auxin, SL and CK), no direct effect of the phosphate status on *BRC1/TBI/FCI* gene expression is documented. Low-phosphate growth conditions enhance SL production in many species (Yoneyama et al., 2007; López-Ráez et al., 2008; Umehara et al., 2008; Domagalska and Leyser, 2011; Kohlen et al., 2011; Yamada et al., 2014). This situation leads to the repression of shoot branching (Umehara et al., 2008; Kohlen et al., 2011), but also to the stimulation of lateral root formation for soil

foraging (Yoneyama et al., 2007; Ruyter-Spira et al., 2011). In contrast to SL, low levels of inorganic phosphate reduce CK production, which correlates with a reduced number of branches (Horgan and Wareing, 1980).

In herbaceous and woody plants, high levels of nitrogen fertilization (nitrate and/or ammonium) result in (i) a large number of outgrowing buds (Lortie and Aarssen, 1997; Médiène et al., 2002; Cline et al., 2006; Emarat-Pardaz et al., 2013; Pal et al., 2013; Furet et al., 2014), and (ii) improved secondary axis elongation (Thitithanakul, 2012; Thitithanakul et al., 2012). Luo et al. (2017) confirmed that nitrogen deficiency did not affect the initiation of tiller buds, but suppressed tiller bud outgrowth in *O. sativa*. In *Arabidopsis*, low nitrate delayed axillary bud activation, and this process involved an effect of the plant nitrogen status rather than a direct nitrate-signaling pathway (De Jong et al., 2014). Recent results demonstrated a relationship between nitrogen fertilization and *BRC1* expression in rice (Li et al., 2016). They showed that high ammonium nitrate intake in the root environment induced a reduction of apical dominance through overexpression of miRNA393 in the buds; miRNA393 inhibits the expression of the genes involved in auxin synthesis and signaling (*OsTIR1*, *OsAFB2*, and *OsIAA6*) as well as *OsTBI*. In *Arabidopsis*, the *brc1-2/brc2-1* double mutant exhibited a higher number of branches than the wild type, but low availability of nitrate reduced this effect (Seale et al., 2017). As root nitrate is widely known to induce CK biosynthesis and signaling events in the whole plant (Crawford, 1995; Sakakibara et al., 1998; Takei et al., 2001, 2002; Forde, 2002a,b), and CK repress *BRC1* expression, we cannot exclude that nitrate may affect *BRC1* expression through CK modulation. In rice, the supply of a CK analog (BAP) or ammonium nitrate regulated SL amounts in the stem and the bud within 3 h after treatment, but nothing has been reported regarding *BRC1* expression (Xu et al., 2015).

In Rosaceae as in many woody plants, nitrate is reduced and assimilated into amino acids directly in the roots; consequently, asparagine, arginine, aspartate, and glutamine are the main forms of nitrogen translocated to the buds via the xylem sap (Millard et al., 1998; Malaguti et al., 2001; Grassi et al., 2002; Guak et al., 2003; Le Moigne et al., 2018). In rose, asparagine is a major nitrogen form involved in bud outgrowth (Le Moigne et al., 2018); this is in accordance with previous data showing that application of asparagine on the soil of olive trees or on the leaves of poplar trees contributed to enhance bud outgrowth and secondary axis elongation (Proietti and Tombesi, 1996; Cline et al., 2006). In rice, a lack of cytosolic glutamine synthetase1;2 in the vascular tissues of axillary buds severely reduced their outgrowth (Funayama et al., 2013; Ohashi et al., 2015) independently of the SL level (Ohashi et al., 2015). In rose bush, sucrose, glucose, and fructose had to be associated to asparagine to allow for the buds to grow out *in vitro* (Le Moigne et al., 2018). This effect involved the upregulation of *IPT3* gene expression in the stem and in the vicinity of the bud (Le Moigne et al., 2018) and the downregulation of *BRC1* (Barbier et al., 2015). In addition to a nutritional role, asparagine might also be a signal representing the nitrogen status of the plant, so as to counteract *BRC1* expression through CK stimulation.

## A BRC1-RELATED REGULATORY MECHANISM

Many studies ascribe an inhibitory function of mitotic cell activity to *BRC1* (Poza-Carrión et al., 2007; Kieffer et al., 2011). This is because early results of EMSA (Electrophoresis Mobility Shift Assay) revealed the capacity of the TCP domain to associate specifically with the promoter element of the rice proliferating cell nuclear antigen (PCNA) gene (Kosugi and Ohashi, 1997, 2002). These *cis*-regulatory modules are indispensable for the transcriptional activation of the PCNA gene in rice meristem tissues (Kosugi and Ohashi, 1997), which seems to be an ancient and prevalent role of TCP transcription factors (Ortiz-Ramírez et al., 2016).

*BRC1*-mediated branching is repressed by the regulation of abscisic acid (ABA) metabolism (Figure 1). ABA is a plant hormone that plays important roles in many phases of the plant life cycle (Seo and Koshida, 2002; Hayes, 2018; Wang et al., 2018). Evidence for a role of ABA in regulating bud growth comes from the positive correlation between a reduction of ABA levels in buds and their release from dormancy (Cline, 1991). Moreover, the *Arabidopsis era1* (*ENHANCED RESPONSE TO ABA 1*) mutant exhibited high sensitivity to ABA and reduced branching (Pei et al., 1998). In *brc1 Arabidopsis* mutants, the ABA-signaling pathway showed a significantly reduced response as compared to the wild type. Additional data revealed that the expression levels of two ABA markers, *ABA-RESPONSE PROTEIN (ABR)* and *UDP-GLUCOSYL TRANSFERASE 74D1 (UGT74D1)*, were significantly upregulated in the wild type but not in *brc1* mutants treated with low R:FR light (González-Grandío et al., 2013). González-Grandío and Cubas (2014) support a model in which ABA acts rather downstream of *BRC1*, because *ABRE-BINDING FACTOR 3 (ABF3)* and *ABA INSENSITIVE 5 (ABI5)*, two key regulators of the ABA response that contain TCP-binding sites in their promoters (Finkelstein and Lynch, 2000; Yoshida et al., 2010; González-Grandío et al., 2013; Nicolas and Cubas, 2016), are upregulated in axillary buds upon *BRC1* induction (González-Grandío and Cubas, 2014). They also indicated that *BRC1* bound to and positively regulated three transcription factors: *HOMEODOMAIN PROTEIN 21 (HB21)*, *HOMEODOMAIN PROTEIN 40 (HB40)*, and *HOMEODOMAIN PROTEIN 53 (HB53)*. These three proteins, together with *BRC1*, enhanced *NINE-CIS-EPOXICAROTENOID DIOXIGENASE 3 (NCED3)* expression, the main ABA-biosynthesis enzyme, leading in turn to ABA accumulation in buds (González-Grandío et al., 2017). This finding demonstrates a direct relationship between *BRC1* and ABA signaling, and places ABA downstream of *BRC1*. Consistently, *BRC1* expression was found to be insensitive to exogenous ABA application (Yao and Scott, 2015).

The TCP genes generally act by positively or negatively regulating the cell cycle (Sarvepalli and Nath, 2018). As a transcription factor, *BRC1* could bind to the promoter region of various genes to regulate the branching process and participate to many regulatory mechanisms (González-Grandío et al., 2013). In maize, *TBI* can directly activate the *tassels replace upper ears1 (tru1)* gene that encodes an ankyrin-repeat-domain protein by binding to the promoter



region of *tru1* (Dong et al., 2017). In *Arabidopsis*, bioinformatic analysis indicates that the promoter sequences of 1,950 genes expressed in the shoot bear the TCP-*cis* regulatory motif (5'-RRVMMV-3') and could be putatively regulated by *AtBRC1*. Based on Gene Ontology (GO) enrichment analysis, these putative target genes are thought to be mainly involved in metabolic processes, including amino acid metabolism [e.g., *ALANINE-2-OXOGLUTARATE AMINOTRANSFERASE 1* (*AOAT1*); *HYDROXYPYRUVATE REDUCTASE* (*ATHPR1*)] and sulfur (e.g., *sulfate transmembrane transporter* (*MOT2*), *sulfate transporter 1;2* (*SULTR1;2*)] (data not shown). We can therefore speculate that *BRC1/TB1* might control bud outgrowth via various pathways, such as stimulating the ABA-signaling pathway and inhibiting cell division and cell metabolism.

## CONCLUSION AND PERSPECTIVES

*BRC1/TB1/FC1* is an integrator gene involved in shoot branching, which fits well with the ability of *BRC1/TB1/FC1* expression to integrate many endogenous and exogenous inputs (Figure 1). However, the detailed mechanism whereby these stimuli regulate *BRC1* expression is still puzzling, and many mechanistic scenarios are plausible. Many questions are thus still open and include how CK and SL, the main two branching-related hormones, antagonistically regulate *BRC1* expression, and which molecular actors could be involved. Similar questions concern the sugar-mediated downregulation of *BRC1*, and the molecular mechanism behind the combined effect of nutrients and hormones on *BRC1* expression (Sakr et al., 2018). In addition, the regulation of gene expression includes many aspects, such as epigenetic regulation, transcriptional regulation, post-transcriptional regulation, translational and post-translational regulation. The relevance of these mechanisms in the regulation of *BRC1* expression deserves to be investigated in different biological contexts. Recent data showed that the protein interaction process also influences *BRC1* expression. For example, the florigen proteins FLOWERING LOCUS T (FT) and TWIN SISTER OF FT (TSF) influence axillary meristem development via their interaction with *AtBRC1* (Niwa et al.,

2013); TIE1 (TCP interactor containing EAR motif protein 1), a transcriptional repressor identified as involved in the control of leaf development, controls shoot branching by interacting with *BRC1* (Yang et al., 2018). Additional protein partners may also interact with *BRC1*, including those related to the energy and nutrient statuses [Sucrose non-fermenting-related kinase (SnRK1)/Target of rapamycin (TOR kinase)] (Martín-Fontecha et al., 2018). Meanwhile, our knowledge about the molecular network governing the *BRC1*-dependent reduction of plant branching is still limited, and the only available data report that *BRC1* action could be related to different biological functions such as cell proliferation, cell metabolism, hormone biosynthesis, ribosome biosynthesis, etc. All these findings indicate that further work is required to fully investigate the regulatory network behind the regulation and function of *BRC1* in shoot branching.

## DATA AVAILABILITY STATEMENT

Publicly available datasets were analyzed in this study. This data can be found here: <https://www.rosaceae.org>.

## AUTHOR CONTRIBUTIONS

All the authors listed have contributed significantly to this manuscript. MW, J-MD, and SS managed this review and contributed to different sections. M-DP-G and LO contributed to the TCP transcription factor section. JB, SD-M, LH, and LC contributed to both hormonal and environmental factors sections, and M-ALM contributed to the nutrient section.

## FUNDING

This work was supported by the program of China Scholarships Council (No. 201506320203) and by the ANR (Agence Nationale de la Recherche) project Labcom, called ESTIM (Evaluation de stimulateurs de vitalité des plantes).

## REFERENCES

- Aguilar-Martínez, J. A., Poza-Carrión, C., and Cubas, P. (2007). *Arabidopsis* *BRANCHED1* acts as an integrator of branching signals within axillary buds. *Plant Cell* 19, 458–472. doi: 10.1105/tpc.106.048934
- Alder, A., Jamil, M., Marzorati, M., Bruno, M., Vermathen, M., Bigler, P., et al. (2012). The path from  $\beta$ -carotene to carlactone, a strigolactone-like plant hormone. *Science* 335, 1348–1351. doi: 10.1126/science.1218094
- Arite, T., Iwata, H., Ohshima, K., Maekawa, M., Nakajima, M., Kojima, M., et al. (2007). *DWARF10*, an RMS1/MAX4/DAD1 ortholog, controls lateral bud outgrowth in rice. *Plant J.* 51, 1019–1029. doi: 10.1111/j.1365-313X.2007.03210.x
- Arite, T., Umehara, M., Ishikawa, S., Hanada, A., Maekawa, M., Yamaguchi, S., et al. (2009). *d14*, a strigolactone-insensitive mutant of rice, shows an accelerated outgrowth of tillers. *Plant Cell Physiol.* 50, 1416–1424. doi: 10.1093/pcp/pcp091
- Balla, J., Kalousek, P., Reinöhl, V., Friml, J., and Procházka, S. (2011). Competitive canalization of PIN-dependent auxin flow from axillary buds controls pea bud outgrowth. *Plant J.* 65, 571–577. doi: 10.1111/j.1365-313X.2010.04443.x
- Bangerth, F. (1994). Response of cytokinin concentration in the xylem exudate of bean (*Phaseolus vulgaris* L.) plants to decapitation and auxin treatment, and relationship to apical dominance. *Planta* 194, 439–442. doi: 10.1007/BF00197546
- Barbier, F., Peron, T., Lecerf, M., Perez-Garcia, M.-D., Barriere, Q., Rolčík, J., et al. (2015). Sucrose is an early modulator of the key hormonal mechanisms controlling bud outgrowth in *Rosa hybrida*. *J. Exp. Bot.* 66, 2569–2582. doi: 10.1093/jxb/erv047
- Beveridge, C. A., and Kyozuka, J. (2010). New genes in the strigolactone-related shoot branching pathway. *Curr. Opin. Plant Biol.* 13, 34–39. doi: 10.1016/j.pbi.2009.10.003
- Boumazza, R., Demotes-Mainard, S., Huché-Thellier, L., and Guérin, V. (2009). Visual characterization of the esthetic quality of the rosebush. *J. Sens. Stud.* 24, 774–796. doi: 10.1111/j.1745-459X.2009.00238.x
- Boumazza, R., Huché-Thellier, L., Demotes-Mainard, S., Le Coz, E., Leduc, N., Pelleschi-Travier, S., et al. (2010). Sensory profiles and preference analysis in ornamental horticulture: the case of the rosebush. *Food Qual. Prefer.* 21, 987–997. doi: 10.1016/j.foodqual.2010.05.003



- Braun, N., de Saint Germain, A., Pillot, J. P., Boutet-Mercey, S., Dalmais, M., Antoniadis, I., et al. (2012). The pea TCP transcription factor PsBRC1 acts downstream of strigolactones to control shoot branching. *Plant Physiol.* 158, 225–238. doi: 10.1104/pp.111.182725
- Brewer, P. B. (2015). Plant architecture: the long and the short of branching in potato. *Curr. Biol.* 25, R724–R725. doi: 10.1016/j.cub.2015.06.066
- Brewer, P. B., Dun, E. A., Ferguson, B. J., Rameau, C., and Beveridge, C. A. (2009). Strigolactone acts downstream of auxin to regulate bud outgrowth in pea and Arabidopsis. *Plant Physiol.* 150, 482–493. doi: 10.1104/pp.108.134783
- Chabikwa, T. G., Brewer, P. B., and Beveridge, C. A. (2019). Initial bud outgrowth occurs independent of auxin flow out of buds. *Plant Physiol.* 179, 55–65. doi: 10.1104/pp.18.00519
- Chao, W. S., Foley, M. E., Horvath, D. P., and Anderson, J. V. (2007). Signals regulating dormancy in vegetative buds. *Int. J. Plant Dev. Biol.* 1, 49–56.
- Chen, X., Zhou, X., Xi, L., Li, J., Zhao, R., Ma, N., et al. (2013). Roles of *DgBRC1* in regulation of lateral branching in chrysanthemum (*Dendranthema × grandiflora* cv. Jinba). *PLoS One* 8:e61717. doi: 10.1371/journal.pone.0061717
- Chen, X. J., Xia, X. J., Guo, X., Zhou, Y. H., Shi, K., Zhou, J., et al. (2016). Apoplastic H<sub>2</sub>O<sub>2</sub> plays a critical role in axillary bud outgrowth by altering auxin and cytokinin homeostasis in tomato plants. *New Phytol.* 211, 1266–1278. doi: 10.1111/nph.14015
- Choubane, D., Rabot, A., Mortreau, E., Legourrierec, J., Péron, T., Foucher, F., et al. (2012). Photocontrol of bud burst involves gibberellin biosynthesis in *Rosa* sp. *J. Plant Physiol.* 169, 1271–1280. doi: 10.1016/j.jplph.2012.04.014
- Cline, M. G. (1991). Apical dominance. *Bot. Rev.* 57, 318–358. doi: 10.1007/BF02858771
- Cline, M. G., Thangavelu, M., and Dong-Il, K. (2006). A possible role of cytokinin in mediating long-distance nitrogen signaling in the promotion of sylleptic branching in hybrid poplar. *J. Plant Physiol.* 163, 684–688. doi: 10.1016/j.jplph.2005.06.005
- Corot, A., Roman, H., Douillet, O., Autret, H., Perez-Garcia, M. D., Citerne, S., et al. (2017). Cytokinins and abscisic acid act antagonistically in the regulation of the bud outgrowth pattern by light intensity. *Front. Plant Sci.* 8:1724. doi: 10.3389/fpls.2017.01724
- Crawford, N. M. (1995). Nitrate: nutrient and signal for plant growth. *Plant Cell* 7, 859–868. doi: 10.1105/tpc.7.7.859
- Crawford, S., Shinohara, N., Sieberer, T., Williamson, L., George, G., Hepworth, J., et al. (2010). Strigolactones enhance competition between shoot branches by dampening auxin transport. *Development* 137, 2905–2913. doi: 10.1242/dev.051987
- Cubas, P. (2004). “Role of TCP genes in the evolution of morphological characters in angiosperms,” in *Developmental Genetics and Plant Evolution*, eds Q. C. B. Cronk, R. M. Bateman, and J. A. Hawkins (Boca Raton, FL: CRC Press), 262–281.
- Cubas, P., Lauter, N., Doebley, J., and Coen, E. (1999). The TCP domain: a motif found in proteins regulating plant growth and development. *Plant J.* 18, 215–222. doi: 10.1046/j.1365-313X.1999.00444.x
- Danisman, S. (2016). TCP transcription factors at the interface between environmental challenges and the plant's growth responses. *Front. Plant Sci.* 7:1930. doi: 10.3389/fpls.2016.01930
- Danisman, S., Van der Wal, F., Dhondt, S., Waites, R., de Folter, S., Bimbo, A., et al. (2012). Arabidopsis class I and class II TCP transcription factors regulate jasmonic acid metabolism and leaf development antagonistically. *Plant Physiol.* 159, 1511–1523. doi: 10.1104/pp.112.200303
- De Jong, F., Thodey, K., Lejay, L. V., and Bevan, M. W. (2014). Glucose elevates NITRATE TRANSPORTER2.1 protein levels and nitrate transport activity independently of its HEXOKINASE1-mediated stimulation of NITRATE TRANSPORTER2.1 expression. *Plant Physiol.* 164, 308–320. doi: 10.1104/pp.113.230599
- Dierck, R., De Keyser, E., De Riek, J., Dhooghe, E., Van Huylenbroeck, J., Prinsen, E., et al. (2016). Change in auxin and cytokinin levels coincides with altered expression of branching genes during axillary bud outgrowth in Chrysanthemum. *PLoS One* 11:e0161732. doi: 10.1371/journal.pone.0161732
- Dixon, L. E., Greenwood, J. R., Bencivenga, S., Zhang, P., Cockram, J., Mellers, G., et al. (2018). *TEOSINTE BRANCHED1* regulates inflorescence architecture and development in bread wheat (*Triticum aestivum* L.). *Plant Cell* 30, 563–581. doi: 10.1105/tpc.17.00961
- Doebley, J., Stec, A., and Hubbard, L. (1997). The evolution of apical dominance in maize. *Nature* 386, 485–488. doi: 10.1038/386485a0
- Domagalska, M. A., and Leyser, O. (2011). Signal integration in the control of shoot branching. *Nat. Rev. Mol. Cell Biol.* 12, 211–221. doi: 10.1038/nrm3088
- Dong, Z., Alexander, M., and Chuck, G. (2019). Understanding grass domestication through maize mutants. *Trends Genet.* 35, 118–128. doi: 10.1016/j.tig.2018.10.007
- Dong, Z., Li, W., Unger-Wallace, E., Yang, J., Vollbrecht, E., and Chuck, G. (2017). Ideal crop plant architecture is mediated by *tassels replace upper ears1*, a BTB/POZ ankyrin repeat gene directly targeted by *TEOSINTE BRANCHED1*. *Proc. Natl. Acad. Sci. U.S.A.* 114, E8656–E8664. doi: 10.1073/pnas.1714960114
- Dulhanty, A. M., and Riordan, J. R. (1994). Phosphorylation by cAMP-dependent protein kinase causes a conformational change in the R domain of the cystic fibrosis transmembrane conductance regulator. *Biochemistry* 33, 4072–4079. doi: 10.1021/bi00179a036
- Dun, E. A., Brewer, P. B., and Beveridge, C. A. (2009). Strigolactones: discovery of the elusive shoot branching hormone. *Trends Plant Sci.* 14, 364–372. doi: 10.1016/j.tplants.2009.04.003
- Dun, E. A., de Saint Germain, A., Rameau, C., and Beveridge, C. A. (2012). Antagonistic action of strigolactone and cytokinin in bud outgrowth control. *Plant Physiol.* 158, 487–498. doi: 10.1104/pp.111.186783
- Emarat-Pardaz, J., Shakiba, M. R., Toorchi, M., and Mohammadinassab, A. D. (2013). The influence of light intensities and nitrogen on growth of *Hypericum perforatum* L. *Int. J. Agric.* 3, 775–781.
- Ferguson, B. J., and Beveridge, C. A. (2009). Roles for auxin, cytokinin, and strigolactone in regulating shoot branching. *Plant Physiol.* 149, 1929–1944. doi: 10.1104/pp.109.135475
- Ferreira, D. A., Soldi, M. C. M., Cheavegatti Gianotto, A., Carneiro, M. S., Amadeu, R. R., Aricetti, J. A., et al. (2018). Metabolite profiles of sugarcane culm reveal the relationship among metabolism and axillary bud outgrowth in genetically related sugarcane commercial cultivars. *Front. Plant Sci.* 9:857. doi: 10.3389/fpls.2018.00857
- Fichtner, F., Barbier, F. F., Feil, R., Watanabe, M., Annunziata, M. G., Chabikwa, T. G., et al. (2017). Trehalose 6-phosphate is involved in triggering axillary bud outgrowth in garden pea (*Pisum sativum* L.). *Plant J.* 92, 611–623. doi: 10.1111/tpj.13705
- Finkelstein, R. R., and Lynch, T. J. (2000). The Arabidopsis abscisic acid response gene *ABI5* encodes a basic leucine zipper transcription factor. *Plant Cell* 12, 599–609. doi: 10.1105/tpc.12.4.599
- Finlayson, S. A., Krishnareddy, S. R., Kebrom, T. H., and Casal, J. J. (2010). Phytochrome regulation of branching in Arabidopsis. *Plant Physiol.* 152, 1914–1927. doi: 10.1104/pp.109.148833
- Fletcher, G. M., and Dale, J. E. (1974). Growth of tiller buds in barley: effects of shade treatment and mineral nutrition. *Ann. Bot.* 38, 63–76. doi: 10.1093/oxfordjournals.aob.a084802
- Foo, E., Bullier, E., Goussot, M., Foucher, F., Rameau, C., and Beveridge, C. A. (2005). The branching gene *RAMOSUS1* mediates interactions among two novel signals and auxin in pea. *Plant Cell* 17, 464–474. doi: 10.1105/tpc.104.026716
- Forde, B. G. (2002a). Local and long-range signaling pathways regulating plant responses to nitrate. *Annu. Rev. Plant Biol.* 53, 203–224. doi: 10.1146/annurev.arplant.53.100301.135256
- Forde, B. G. (2002b). The role of long-distance signalling in plant responses to nitrate and other nutrients. *J. Exp. Bot.* 53, 39–43. doi: 10.1093/jxbbot/53.366.39
- Funayama, K., Kojima, S., Tabuchi-Kobayashi, M., Sawa, Y., Nakayama, Y., Hayakawa, T., et al. (2013). Cytosolic glutamine synthetase1;2 is responsible for the primary assimilation of ammonium in rice roots. *Plant Cell Physiol.* 54, 934–943. doi: 10.1093/pcp/pct046
- Furet, P. M., Lothier, J., Demotes-Mainard, S., Travier, S., Henry, C., Guérin, V., et al. (2014). Light and nitrogen nutrition regulate apical control in *Rosa hybrida* L. *J. Plant Physiol.* 171, 7–13. doi: 10.1016/j.jplph.2013.10.008
- Garbez, M., Galopin, G., Sigogne, M., Favre, P., Demotes-Mainard, S., and Symoneaux, R. (2015). Assessing the visual aspect of rotating virtual rose bushes by a labeled sorting task. *Food Qual. Prefer.* 40(Part B), 287–295. doi: 10.1016/j.foodqual.2014.06.008

- Girault, T., Abidi, F., Sigogne, M., Pelleschi-Travier, S. A. N. D. R. I. N. E., Boumaza, R., Sakr, S., et al. (2010). Sugars are under light control during bud burst in *Rosa* sp. *Plant Cell Environ.* 33, 1339–1350. doi: 10.1111/j.1365-3040.2010.02152.x
- Gomez-Roldan, V., Femas, S., Brewer, P. B., Puech-Pagès, V., Dun, E. A., Pillot, J. P., et al. (2008). Strigolactone inhibition of shoot branching. *Nature* 455, 189–194. doi: 10.1038/nature07271
- González-Grandío, E., and Cubas, P. (2014). Identification of gene functions associated to active and dormant buds in *Arabidopsis*. *Plant Signal. Behav.* 9:e27994. doi: 10.4161/psb.27994
- González-Grandío, E., Pajoro, A., Franco-Zorrilla, J. M., Tarancón, C., Immink, R. G., and Cubas, P. (2017). Abscisic acid signaling is controlled by a *BRANCHED1/HD-ZIP I* cascade in *Arabidopsis* axillary buds. *Proc. Natl. Acad. Sci. U.S.A.* 114, E245–E254. doi: 10.1073/pnas.1613199114
- González-Grandío, E., Poza-Carrión, C., Sorzano, C. O. S., and Cubas, P. (2013). *BRANCHED1* promotes axillary bud dormancy in response to shade in *Arabidopsis*. *Plant Cell* 25, 834–850. doi: 10.1105/tpc.112.108480
- Grassi, G., Millard, P., Wendler, R., Minotta, G., and Tagliavini, M. (2002). Measurement of xylem sap amino acid concentrations in conjunction with whole tree transpiration estimates spring N remobilization by cherry (*Prunus avium* L.) trees. *Plant Cell Environ.* 25, 1689–1699. doi: 10.1046/j.1365-3040.2002.00949.x
- Guak, S., Neilsen, D., Millard, P., Wendler, R., and Neilsen, G. H. (2003). Determining the role of N remobilization for growth of apple (*Malus domestica* Borkh.) trees by measuring xylem-sap N flux. *J. Exp. Bot.* 54, 2121–2131. doi: 10.1093/jxb/erg228
- Guo, Z., Fujioka, S., Blancaflor, E. B., Miao, S., Gou, X., and Li, J. (2010). TCP1 modulates brassinosteroid biosynthesis by regulating the expression of the key biosynthetic gene *DWARF4* in *Arabidopsis thaliana*. *Plant Cell* 22, 1161–1173. doi: 10.1105/tpc.109.069203
- Hall, S. M., and Hillman, J. R. (1975). Correlative inhibition of lateral bud growth in *Phaseolus vulgaris* L. Timing of bud growth following decapitation. *Planta* 123, 137–143. doi: 10.1007/BF00383862
- Hayes, S. (2018). Revealing the invisible: a synthetic reporter for ABA. *Plant Physiol.* 177, 1346–1347. doi: 10.1104/pp.18.00646
- Hayward, A., Stirnberg, P., Beveridge, C., and Leyser, O. (2009). Interactions between auxin and strigolactone in shoot branching control. *Plant Physiol.* 151, 400–412. doi: 10.1104/pp.109.137646
- Hedden, P., and Sponsel, V. (2015). A century of gibberellin research. *J. Plant Growth Regul.* 34, 740–760. doi: 10.1007/s00344-015-9546-1
- Helliwell, C. A., Chin-Atkins, A. N., Wilson, I. W., Chapple, R., Dennis, E. S., and Chaudhury, A. (2001). The *Arabidopsis* *AMP1* gene encodes a putative glutamate carboxypeptidase. *Plant Cell* 13, 2115–2125. doi: 10.1105/TPC.010146
- Hellmann, H. A., and Smeekens, S. (2014). Sugar sensing and signaling in plants. *Front. Plant Sci.* 5:113. doi: 10.3389/fpls.2014.00113
- Henry, C., Rabot, A., Laloi, M., Mortreau, E., Sigogne, M., Leduc, N., et al. (2011). Regulation of *RhSUC2*, a sucrose transporter, is correlated with the light control of bud burst in *Rosa* sp. *Plant Cell Environ.* 34, 1776–1789. doi: 10.1111/j.1365-3040.2011.02374.x
- Holalu, S. V., and Finlayson, S. A. (2017). The ratio of red light to far red light alters *Arabidopsis* axillary bud growth and abscisic acid signalling before stem auxin changes. *J. Exp. Bot.* 68, 943–952. doi: 10.1093/jxb/erw479
- Horgan, J. M., and Wareing, P. F. (1980). Cytokinins and the growth responses of seedlings of *Betula pendula* Roth. and *Acer pseudoplatanus* L. to nitrogen and phosphorus deficiency. *J. Exp. Bot.* 31, 525–532. doi: 10.1093/jxb/31.2.525
- Howarth, D. G., and Donoghue, M. J. (2006). Phylogenetic analysis of the “ECE”(CYC/TB1) clade reveals duplications predating the core eudicots. *Proc. Natl. Acad. Sci. U.S.A.* 103, 9101–9106. doi: 10.1073/pnas.0602827103
- Hubbard, L., McSteen, P., Doebley, J., and Hake, S. (2002). Expression patterns and mutant phenotype of teosinte branched1 correlate with growth suppression in maize and teosinte. *Genetics* 162, 1927–1935.
- Jasinski, S., Piazza, P., Craft, J., Hay, A., Woolley, L., Rieu, I., et al. (2005). KNOX action in *Arabidopsis* is mediated by coordinate regulation of cytokinin and gibberellin activities. *Curr. Biol.* 15, 1560–1565. doi: 10.1016/j.cub.2005.07.023
- Jiang, H., and Egli, D. B. (1993). Shade induced changes in flower and pod number and flower and fruit abscission in soybean. *Agron. J.* 85, 221–225. doi: 10.2134/agronj1993.00021962008500020011x
- Jiang, L., Liu, X., Xiong, G., Liu, H., Chen, F., Wang, L., et al. (2013). *DWARF53* acts as a repressor of strigolactone signalling in rice. *Nature* 504, 401–405. doi: 10.1038/nature12870
- Jiao, Y., Wang, Y., Xue, D., Wang, J., Yan, M., Liu, G., et al. (2010). Regulation of *OsSPL14* by *OsmiR156* defines ideal plant architecture in rice. *Nat. Genet.* 42, 541–544. doi: 10.1038/ng.591
- Johnson, X., Brcich, T., Dun, E. A., Goussot, M., Haurigné, K., Beveridge, C. A., et al. (2006). Branching genes are conserved across species. Genes controlling a novel signal in pea are coregulated by other long-distance signals. *Plant Physiol.* 142, 1014–1026. doi: 10.1104/pp.106.087676
- Kebrom, T., Chandler, P., Swain, S., King, R., Richards, R., and Spielmeier, W. (2012). Inhibition of tiller bud outgrowth in the tin mutant of wheat is associated with precocious internode development. *Plant Physiol.* 160, 308–318. doi: 10.1104/pp.112.197954
- Kebrom, T. H. (2017). A growing stem inhibits bud outgrowth—the overlooked theory of apical dominance. *Front. Plant Sci.* 8:1874. doi: 10.3389/fpls.2017.01874
- Kebrom, T. H., Brutnell, T. P., and Finlayson, S. A. (2010). Suppression of sorghum axillary bud outgrowth by shade, phyB and defoliation signalling pathways. *Plant Cell Environ.* 33, 48–58. doi: 10.1111/j.1365-3040.2009.02050.x
- Kebrom, T. H., Burson, B. L., and Finlayson, S. A. (2006). Phytochrome B represses *Teosinte Branched1* expression and induces sorghum axillary bud outgrowth in response to light signals. *Plant Physiol.* 140, 1109–1117. doi: 10.1104/pp.105.074856
- Kebrom, T. H., and Mullet, J. E. (2015). Photosynthetic leaf area modulates tiller bud outgrowth in sorghum. *Plant Cell Environ.* 38, 1471–1478. doi: 10.1111/pce.12500
- Kerr, S. C., and Beveridge, C. A. (2017). IPA1: a direct target of SL signaling. *Cell Res.* 27, 1191–1192. doi: 10.1038/cr.2017.114
- Kieffer, M., Master, V., Waites, R., and Davies, B. (2011). TCP14 and TCP15 affect internode length and leaf shape in *Arabidopsis*. *Plant J.* 68, 147–158. doi: 10.1111/j.1365-313X.2011.04674.x
- Kohlen, W., Charnikhova, T., Liu, Q., Bours, R., Domagalska, M. A., Beguerie, S., et al. (2011). Strigolactones are transported through the xylem and play a key role in shoot architectural response to phosphate deficiency in nonarbuscular mycorrhizal host *Arabidopsis*. *Plant Physiol.* 155, 974–987. doi: 10.1104/pp.110.164640
- Kong, X., Zhang, M., and Ding, Z. (2014). D53: the missing link in strigolactone signaling. *Mol. Plant* 7, 761–763. doi: 10.1093/mp/ssu016
- Kosugi, S., and Ohashi, Y. (1997). PCF1 and PCF2 specifically bind to cis elements in the rice proliferating cell nuclear antigen gene. *Plant Cell* 9, 1607–1619. doi: 10.1105/tpc.9.9.1607
- Kosugi, S., and Ohashi, Y. (2002). DNA binding and dimerization specificity and potential targets for the TCP protein family. *Plant J.* 30, 337–348. doi: 10.1046/j.1365-313X.2002.01294.x
- Lantzouni, O., Klermund, C., and Schwechheimer, C. (2017). Largely additive effects of gibberellin and strigolactone on gene expression in *Arabidopsis thaliana* seedlings. *Plant J.* 92, 924–938. doi: 10.1111/tpj.13729
- Le Moigne, M. A., Guérin, V., Furet, P. M., Billard, V., Lebrec, A., Spíchal, L., et al. (2018). Asparagine and sugars are both required to sustain secondary axis elongation after bud outgrowth in *Rosa hybrid*. *J. Plant Physiol.* 222, 17–27. doi: 10.1016/j.jplph.2017.12.013
- Leduc, N., Roman, H., Barbier, F., Péron, T., Huché-Théliér, L., Lothier, J., et al. (2014). Light signaling in bud outgrowth and branching in plants. *Plants* 3, 223–250. doi: 10.3390/plants3020223
- Lemerle, D., Verbeek, B., Cousens, R. D., and Coombes, N. E. (1996). The potential for selecting wheat varieties strongly competitive against weeds. *Weed Res.* 36, 505–513. doi: 10.1111/j.1365-3180.1996.tb01679.x
- Leyser, O. (2008). Strigolactones and shoot branching: a new trick for a young dog. *Dev. Cell* 15, 337–338. doi: 10.1016/j.devcel.2008.08.008
- Leyser, O. (2009). The control of shoot branching: an example of plant information processing. *Plant Cell Environ.* 32, 694–703. doi: 10.1111/j.1365-3040.2009.01930.x
- Li, C. J., and Bangerth, F. (1999). Autoinhibition of indoleacetic acid transport in the shoots of two-branched pea (*Pisum sativum*) plants and its relationship to correlative dominance. *Physiol. Plant.* 106, 415–420. doi: 10.1034/j.1399-3054.1999.106409.x

- Li, S. (2015). The *Arabidopsis thaliana* TCP transcription factors: a broadening horizon beyond development. *Plant Signal. Behav.* 10:e1044192. doi: 10.1080/15592324.2015.1044192
- Li, S., and Zachgo, S. (2013). TCP3 interacts with R2R3-MYB proteins, promotes flavonoid biosynthesis and negatively regulates the auxin response in *Arabidopsis thaliana*. *Plant J.* 76, 901–913. doi: 10.1111/tpj.12348
- Li, X., Xia, K., Liang, Z., Chen, K., Gao, C., and Zhang, M. (2016). MicroRNA393 is involved in nitrogen-promoted rice tillering through regulation of auxin signal transduction in axillary buds. *Sci. Rep.* 6:32158. doi: 10.1038/srep32158
- Lillier, C. B., Neuhaus, R., Von Korff, M., Koornneef, M., and Van Esse, W. (2015). Mutations in barley row type genes have pleiotropic effects on shoot branching. *PLoS One* 10:e0140246. doi: 10.1371/journal.pone.0140246
- Li-Marchetti, C., Le Bras, C., Chastellier, A., Relion, D., Morel, P., Sakr, S., et al. (2017). 3D phenotyping and QTL analysis of a complex character: rose bush architecture. *Tree Genet. Genomes* 13:112. doi: 10.1007/s11295-017-1194-0
- Li-Marchetti, C., Le Bras, C., Relion, D., Citerne, S., Huché-Théliér, L., Sakr, S., et al. (2015). Genotypic differences in architectural and physiological responses to water restriction in rose bush. *Front. Plant Sci.* 6:355. doi: 10.3389/fpls.2015.00355
- Lin, H., Wang, R., Qian, Q., Yan, M., Meng, X., Fu, Z., et al. (2009). DWARF27, an iron-containing protein required for the biosynthesis of strigolactones, regulates rice tiller bud outgrowth. *Plant Cell* 21, 1512–1525. doi: 10.1105/tpc.109.065987
- Liu, J., Cheng, X., Liu, P., and Sun, J. (2017). miR156-targeted SBP-Box transcription factors interact with DWARF53 to regulate *TEOSINTE BRANCHED1* and *BARREN STALK1* expression in bread wheat. *Plant Physiol.* 174, 1931–1948. doi: 10.1104/pp.17.00445
- Lo, S. F., Yang, S. Y., Chen, K. T., Hsing, Y. I., Zeevaert, J. A., Chen, L. J., et al. (2008). A novel class of gibberellin 2-oxidases control semidwarfism, tillering, and root development in rice. *Plant Cell* 20, 2603–2618. doi: 10.1105/tpc.108.060913
- López-Ráez, J. A., Charnikhova, T., Gómez-Roldán, V., Matusova, R., Kohlen, W., De Vos, R., et al. (2008). Tomato strigolactones are derived from carotenoids and their biosynthesis is promoted by phosphate starvation. *New Phytol.* 178, 863–874. doi: 10.1111/j.1469-8137.2008.02406.x
- Lortie, C. J., and Aarssen, L. W. (1997). Apical dominance as an adaptation in *Verbascum thapsus*: effects of water and nutrients on branching. *Int. J. Plant Sci.* 158, 461–464. doi: 10.1086/297456
- Lu, Z., Yu, H., Xiong, G., Wang, J., Jiao, Y., Liu, G., et al. (2013). Genome-wide binding analysis of the transcription activator ideal plant architecture1 reveals a complex network regulating rice plant architecture. *Plant Cell* 25, 3743–3759. doi: 10.1105/tpc.113.113639
- Luo, D., Carpenter, R., Vincent, C., Copsey, L., and Coen, E. (1996). Origin of floral asymmetry in *Antirrhinum*. *Nature* 383, 794–799. doi: 10.1038/383794a0
- Luo, L., Pan, S., Liu, X., Wang, H., and Xu, G. (2017). Nitrogen deficiency inhibits cell division-determined elongation, but not initiation, of rice tiller buds. *Isr. J. Plant Sci.* 64, 32–40. doi: 10.1080/07929978.2016.1275367
- Lupas, A., Van Dyke, M., and Stock, J. (1991). Predicting coiled coils from protein sequences. *Science* 252, 1162–1164. doi: 10.1126/science.252.5009.1162
- Malaguti, D., Millard, P., Wendler, R., Hepburn, A., and Tagliavini, M. (2001). Translocation of amino acids in the xylem of apple (*Malus domestica* Borkh.) trees in spring as a consequence of both N remobilization and root uptake. *J. Exp. Bot.* 52, 1665–1671. doi: 10.1093/jxb/52.361.1665
- Martín-Fontecha, E. S., Taracón, C., and Cubas, P. (2018). To grow or not to grow, a power-saving program induced in dormant buds. *Curr. Opin. Plant Biol.* 41, 102–109. doi: 10.1016/j.pbi.2017.10.001
- Martín-Trillo, M., and Cubas, P. (2010). TCP genes: a family snapshot ten years later. *Trends Plant Sci.* 15, 31–39. doi: 10.1016/j.tplants.2009.11.003
- Martín-Trillo, M., Grandío, E. G., Serra, F., Marcel, F., Rodríguez-Buey, M. L., Schmitz, G., et al. (2011). Role of tomato *BRANCHED1*-like genes in the control of shoot branching. *Plant J.* 67, 701–714. doi: 10.1111/j.1365-313X.2011.04629.x
- Mason, M. G., Ross, J. J., Babst, B. A., Wienclaw, B. N., and Beveridge, C. A. (2014). Sugar demand, not auxin, is the initial regulator of apical dominance. *Proc. Natl. Acad. Sci. U.S.A.* 111, 6092–6097. doi: 10.1073/pnas.1322045111
- Maurel, K., Sakr, S., Gerbe, F., Guillot, A., Bonhomme, M., Rageau, R., et al. (2004). Sorbitol uptake is regulated by glucose through the hexokinase pathway in vegetative peach-tree buds. *J. Exp. Bot.* 55, 879–888. doi: 10.1093/jxb/erh087
- McSteen, P. (2009). Hormonal regulation of branching in grasses. *Plant Physiol.* 149, 46–55. doi: 10.1104/pp.108.129056
- Médiène, S., Pagès, L., Jordan, M. O., Le Bot, J., and Adamowicz, S. (2002). Influence of nitrogen availability on shoot development in young peach trees [*Prunus persica* (L.) Batsch]. *Trees* 16, 547–554. doi: 10.1007/s00468-002-0204-4
- Millard, P., Wendler, R., Hepburn, A., and Smith, A. (1998). Variations in the amino acid composition of xylem sap of *Betula pendula* Roth. trees due to remobilization of stored N in the spring. *Plant Cell Environ.* 21, 715–722. doi: 10.1046/j.1365-3040.1998.00313.x
- Minakuchi, K., Kameoka, H., Yasuno, N., Umehara, M., Luo, L., Kobayashi, K., et al. (2010). *FINE CULM1 (FC1)* works downstream of strigolactones to inhibit the outgrowth of axillary buds in rice. *Plant Cell Physiol.* 51, 1127–1135. doi: 10.1093/pcp/pcq083
- Miura, K., Ikeda, M., Matsubara, A., Song, X. J., Ito, M., Asano, K., et al. (2010). OsSPL14 promotes panicle branching and higher grain productivity in rice. *Nat. Genet.* 42, 545–549. doi: 10.1038/ng.592
- Miyawaki, K., Matsumoto-Kitano, M., and Kakimoto, T. (2004). Expression of cytokinin biosynthetic isopentenyltransferase genes in *Arabidopsis*: tissue specificity and regulation by auxin, cytokinin, and nitrate. *Plant J.* 37, 128–138. doi: 10.1046/j.1365-313X.2003.01945.x
- Morris, D. A. (1977). Transport of exogenous auxin in two-branched dwarf pea seedlings (*Pisum sativum* L.). *Planta* 136, 91–96. doi: 10.1007/BF00387930
- Morris, S. E., Cox, M. C., Ross, J. J., Krisantini, S., and Beveridge, C. A. (2005). Auxin dynamics after decapitation are not correlated with the initial growth of axillary buds. *Plant Physiol.* 138, 1665–1672. doi: 10.1104/pp.104.058743
- Muhr, M., Paulat, M., Awwanah, M., Brinkkötter, M., and Teichmann, T. (2018). CRISPR/Cas9-mediated knockout of *Populus BRANCHED1* and *BRANCHED2* orthologs reveals a major function in bud outgrowth control. *Tree Physiol.* 38, 1588–1597. doi: 10.1093/treephys/tpy088
- Muhr, M., Prüfer, N., Paulat, M., and Teichmann, T. (2016). Knockdown of strigolactone biosynthesis genes in *Populus* affects *BRANCHED1* expression and shoot architecture. *New Phytol.* 212, 613–626. doi: 10.1111/nph.14076
- Müller, D., and Leyser, O. (2011). Auxin, cytokinin and the control of shoot branching. *Ann. Bot.* 107, 1203–1212. doi: 10.1093/aob/mcr069
- Nath, U., Crawford, B. C., Carpenter, R., and Coen, E. (2003). Genetic control of surface curvature. *Science* 299, 1404–1407. doi: 10.1126/science.1079354
- Navaud, O., Dabos, P., Carnus, E., Tremousaygue, D., and Hervé, C. (2007). TCP transcription factors predate the emergence of land plants. *J. Mol. Evol.* 65, 23–33. doi: 10.1007/s00239-006-0174-z
- Ni, J., Gao, C., Chen, M. S., Pan, B. Z., Ye, K., and Xu, Z. F. (2015). Gibberellin promotes shoot branching in the perennial woody plant *Jatropha curcas*. *Plant Cell Physiol.* 56, 1655–1666. doi: 10.1093/pcp/pcv089
- Nicolas, M., and Cubas, P. (2016). TCP factors: new kids on the signaling block. *Curr. Opin. Plant Biol.* 33, 33–41. doi: 10.1016/j.pbi.2016.05.006
- Nicolas, M., Rodríguez-Buey, M. L., Franco-Zorrilla, J. M., and Cubas, P. (2015). A recently evolved alternative splice site in the *BRANCHED1a* gene controls potato plant architecture. *Curr. Biol.* 25, 1799–1809. doi: 10.1016/j.cub.2015.05.053
- Niwa, M., Daimon, Y., Kurotani, K. I., Higo, A., Pruneda-Paz, J. L., Breton, G., et al. (2013). *BRANCHED1* interacts with *FLOWERING LOCUS T* to repress the floral transition of the axillary meristems in *Arabidopsis*. *Plant Cell* 25, 1228–1242. doi: 10.1105/tpc.112.109090
- Nordström, A., Tarkowski, P., Tarkowska, D., Norbaek, R., Åstot, C., Dolezal, K., et al. (2004). Auxin regulation of cytokinin biosynthesis in *Arabidopsis thaliana*: a factor of potential importance for auxin-cytokinin-regulated development. *Proc. Natl. Acad. Sci. U.S.A.* 101, 8039–8044. doi: 10.1073/pnas.0402504101
- Ohashi, M., Ishiyama, K., Kusano, M., Fukushima, A., Kojima, S., Hanada, A., et al. (2015). Lack of cytosolic glutamine synthetase1; 2 in vascular tissues of axillary buds causes severe reduction in their outgrowth and disorder of metabolic balance in rice seedlings. *Plant J.* 81, 347–356. doi: 10.1111/tpj.12731
- Ortiz-Ramírez, C., Hernandez-Coronado, M., Thamm, A., Catarino, B., Wang, M., Dolan, L., et al. (2016). A transcriptome atlas of *Physcomitrella patens* provides insights into the evolution and development of land plants. *Mol. Plant* 9, 205–220. doi: 10.1016/j.molp.2015.12.002



- Pal, P. K., Prasad, R., and Pathania, V. (2013). Effect of decapitation and nutrient applications on shoot branching, yield, and accumulation of secondary metabolites in leaves of *Stevia rebaudiana* Bertoni. *J. Plant Physiol.* 170, 1526–1535. doi: 10.1016/j.jplph.2013.06.017
- Palatnik, J. F., Allen, E., Wu, X., Schommer, C., Schwab, R., Carrington, J. C., et al. (2003). Control of leaf morphogenesis by microRNAs. *Nature* 425, 257–263. doi: 10.1038/nature01958
- Pei, Z. M., Ghassemian, M., Kwak, C. M., McCourt, P., and Schroeder, J. I. (1998). Role of farnesyltransferase in ABA regulation of guard cell anion channels and plant water loss. *Science* 282, 287–290. doi: 10.1126/science.282.5387.287
- Petrásek, J., and Friml, J. (2009). Auxin transport routes in plant development. *Development* 136, 2675–2688. doi: 10.1242/dev.030353
- Poza-Carrión, C., Aguilar-Martínez, J. A., and Cubas, P. (2007). Role of TCP gene *BRANCHED1* in the control of shoot branching in *Arabidopsis*. *Plant Signal. Behav.* 2, 551–552. doi: 10.4161/psb.2.6.4811
- Price, J., Laxmi, A., Martin, S. K. S., and Jang, J. C. (2004). Global transcription profiling reveals multiple sugar signal transduction mechanisms in *Arabidopsis*. *Plant Cell* 16, 2128–2150. doi: 10.1105/tpc.104.022616
- Proietti, P., and Tombesi, A. (1996). Effects of gibberellic acid, asparagine and glutamine on flower bud induction in olive. *J. Horticult. Sci.* 71, 383–388. doi: 10.1080/14620316.1996.11515418
- Qin, L. J., Guo, X. Z., Feng, X. Z., Weng, L., Yan, J., Hu, X. H., et al. (2004). Cloning of *LjCYC1* gene and nuclear localization of LjCYC1 protein in *Lotus japonicus*. *J. Plant Physiol. Mol. Biol.* 30, 523–532.
- Rabot, A., Henry, C., Baaziz, K. B., Mortreau, E., Azri, W., Lother, J., et al. (2012). Insight into the role of sugars in bud burst under light in the rose. *Plant Cell Physiol.* 53, 1068–1082. doi: 10.1093/pcp/pcs051
- Rameau, C., Bertheloot, J., Leduc, N., Andrieu, B., Foucher, F., Sakr, S., et al. (2015). Multiple pathways regulate shoot branching. *Front. Plant Sci.* 5:741. doi: 10.3389/fpls.2014.00741
- Ramsay, L., Comadran, J., Druka, A., Marshall, D. F., Thomas, W. T., Macaulay, M., et al. (2011). *INTERMEDIUM-C*, a modifier of lateral spikelet fertility in barley, is an ortholog of the maize domestication gene *TEOSINTE BRANCHED 1*. *Nat. Genet.* 43, 169–172. doi: 10.1038/ng.745
- Revel, S. M., Bart, J. J., Luo, Z., Oplaat, C., Susan, E. L., Mark, W. W., et al. (2015). Environmental control of branching in petunia. *Plant Physiol.* 168, 735–751. doi: 10.1104/pp.15.00486
- Richards, R. A. (2000). Selectable traits to increase crop photosynthesis and yield of grain crops. *J. Exp. Bot.* 51, 447–458. doi: 10.1093/jexbot/51.suppl\_1.447
- Rodriguez, D., Andrade, F. H., and Goudriaan, J. (1999). Effects of phosphorus nutrition on tiller emergence in wheat. *Plant Soil* 209, 283–295. doi: 10.1023/A:1004690404870
- Roman, H., Girault, T., Barbier, F., Péron, T., Brouard, N., Pencik, A., et al. (2016). Cytokinins are initial targets of light in the control of bud outgrowth. *Plant Physiol.* 172, 489–509. doi: 10.1104/pp.16.00530
- Ruyter-Spira, C., Kohlen, W., Charnikhova, T., van Zeijl, A., van Bezouwen, L., de Ruijter, N., et al. (2011). Physiological effects of the synthetic strigolactone analog GR24 on root system architecture in *Arabidopsis*: another below-ground role for strigolactones? *Plant Physiol.* 155, 721–734. doi: 10.1104/pp.110.166645
- Sachs, T., and Thimann, K. V. (1967). The role of auxins and cytokinins in the release of buds from dominance. *Am. J. Bot.* 54, 136–144. doi: 10.1002/j.1537-2197.1967.tb06901.x
- Sakakibara, H., Suzuki, M., Takei, K., Deji, A., Taniguchi, M., and Sugiyama, T. (1998). A response-regulator homologue possibly involved in nitrogen signal transduction mediated by cytokinin in maize. *Plant J.* 14, 337–344. doi: 10.1046/j.1365-313X.1998.00134.x
- Sakr, S., Wang, M., Dédaldéchamp, F., Perez-Garcia, M. D., Ogé, L., Hamama, L., et al. (2018). The sugar-signaling hub: overview of regulators and interaction with the hormonal and metabolic network. *Int. J. Mol. Sci.* 19:E2506. doi: 10.3390/ijms19092506
- Sarvepalli, K., and Nath, U. (2018). CIN-TCP transcription factors: transiting cell proliferation in plants. *IUBMB Life* 70, 718–731. doi: 10.1002/iub.1874
- Schommer, C., Palatnik, J. F., Aggarwal, P., Chételat, A., Cubas, P., Farmer, E. E., et al. (2008). Control of jasmonate biosynthesis and senescence by miR319 targets. *PLoS Biol.* 6:e230. doi: 10.1371/journal.pbio.0060230
- Seale, M., Bennett, T., and Leyser, O. (2017). BRC1 expression regulates bud activation potential but is not necessary or sufficient for bud growth inhibition in *Arabidopsis*. *Development* 144, 1661–1673. doi: 10.1242/dev.145649
- Seo, M., and Koshiba, T. (2002). Complex regulation of ABA biosynthesis in plants. *Trends Plant Sci.* 7, 41–48. doi: 10.1016/S1360-1385(01)02187-2
- Shimizu, S., and Mori, H. (1998). Analysis of cycles of dormancy and growth in pea axillary buds based on mRNA accumulation patterns of cell cycle-related genes. *Plant Cell Physiol.* 39, 255–262. doi: 10.1093/oxfordjournals.pcp.a029365
- Shinohara, N., Taylor, C., and Leyser, O. (2013). Strigolactone can promote or inhibit shoot branching by triggering rapid depletion of the auxin efflux protein PIN1 from the plasma membrane. *PLoS Biol.* 11:e1001474. doi: 10.1371/journal.pbio.1001474
- Simon, S., Morel, K., Durand, E., Brevalle, G., Girard, T., and Lauri, P. -É. (2011). Aphids at crossroads: when branch architecture alters aphid infestation patterns in the apple tree. *Trees* 26, 273–282. doi: 10.1007/s00468-011-0629-8
- Song, X., Lu, Z., Yu, H., Shao, G., Xiong, J., Meng, X., et al. (2017). IPA1 functions as a downstream transcription factor repressed by D53 in strigolactone signaling in rice. *Cell Res.* 27, 1128–1141. doi: 10.1038/cr.2017.102
- Sorefan, K., Booker, J., Haurogné, K., Goussot, M., Bainbridge, K., Foo, E., et al. (2003). *MAX4* and *RMS1* are orthologous dioxygenase-like genes that regulate shoot branching in *Arabidopsis* and pea. *Genes Dev.* 17, 1469–1474. doi: 10.1101/gad.256603
- Soundappan, I., Bennett, T., Morffy, N., Liang, Y., Stanga, J. P., Abbas, A., et al. (2015). *SMAX1-LIKE/D53* family members enable distinct *MAX2*-dependent responses to strigolactones and karrikins in *Arabidopsis*. *Plant Cell* 27, 3143–3159. doi: 10.1105/tpc.15.00562
- Stafstrom, J. P., and Sussex, I. M. (1988). Patterns of protein synthesis in dormant and growing vegetative buds of pea. *Planta* 176, 497–505. doi: 10.1007/BF00397656
- Stirnberg, P., Chatfield, S. P., and Leyser, H. O. (1999). *AXR1* acts after lateral bud formation to inhibit lateral bud growth in *Arabidopsis*. *Plant Physiol.* 121, 839–847. doi: 10.1104/pp.121.3.839
- Studer, A., Zhao, Q., Ross-Ibarra, J., and Doebley, J. (2011). Identification of a functional transposon insertion in the maize domestication gene *tb1*. *Nat. Genet.* 43, 1160–1163. doi: 10.1038/ng.942
- Su, H., Abernathy, S. D., White, R. H., and Finlayson, S. A. (2011). Photosynthetic photon flux density and phytochrome B interact to regulate branching in *Arabidopsis*. *Plant Cell Environ.* 34, 1986–1998. doi: 10.1111/j.1365-3040.2011.02393.x
- Suzuki, T., Sakurai, K., Ueguchi, C., and Mizuno, T. (2001). Two types of putative nuclear factors that physically interact with histidine-containing phosphotransfer (Hpt) domains, signaling mediators in his-to-asp phosphorelay, in *Arabidopsis thaliana*. *Plant Cell Physiol.* 42, 37–45. doi: 10.1093/pcp/pce011
- Ta, T. C., MacDowall, F. D. H., and Faris, M. A. (1987). Utilization of carbon from shoot photosynthesis and nodule CO<sub>2</sub> fixation in the fixation and assimilation of nitrogen by alfalfa root nodules. *Can. J. Bot.* 65, 2537–2541. doi: 10.1139/b87-344
- Takeda, T., Suwa, Y., Suzuki, M., Kitano, H., Ueguchi-Tanaka, M., Ashikari, M., et al. (2003). The *OsTB1* gene negatively regulates lateral branching in rice. *Plant J.* 33, 513–520. doi: 10.1046/j.1365-313X.2003.01648.x
- Takei, K., Sakakibara, H., Taniguchi, M., and Sugiyama, T. (2001). Nitrogen-dependent accumulation of cytokinins in root and the translocation to leaf: implication of cytokinin species that induces gene expression of maize response regulator. *Plant Cell Physiol.* 42, 85–93. doi: 10.1093/pcp/pce009
- Takei, K., Takahashi, T., Sugiyama, T., Yamaya, T., and Sakakibara, H. (2002). Multiple routes communicating nitrogen availability from roots to shoots: a signal transduction pathway mediated by cytokinin. *J. Exp. Bot.* 53, 971–977. doi: 10.1093/jexbot/53.370.971
- Tanaka, M., Takei, K., Kojima, M., Sakakibara, H., and Mori, H. (2006). Auxin controls local cytokinin biosynthesis in the nodal stem in apical dominance. *Plant J.* 45, 1028–1036. doi: 10.1111/j.1365-313X.2006.02656.x
- Tarancón, C., González-Grandío, E., Oliveros, J. C., Nicolas, M., and Cubas, P. (2017). A conserved carbon starvation response underlies bud dormancy in woody and herbaceous species. *Front. Plant Sci.* 8:788. doi: 10.3389/fpls.2017.00788
- Teichmann, T., and Muhr, M. (2015). Shaping plant architecture. *Front. Plant Sci.* 6:233. doi: 10.3389/fpls.2015.00233



- Thitithanakul, S. (2012). *Effect of Nitrogen Supply Before Bud Break on Early Development of the Young Hybrid Poplar*. Doctoral dissertation, Université Blaise Pascal, Clermont-Ferrand.
- Thitithanakul, S., Pétel, G., Chalot, M., and Beaujard, F. (2012). Supplying nitrate before bud break induces pronounced changes in nitrogen nutrition and growth of young poplars. *Funct. Biol.* 39, 795–803. doi: 10.1071/FP12129
- Umehara, M., Hanada, A., Yoshida, S., Akiyama, K., Arite, T., Takeda-Kamiya, N., et al. (2008). Inhibition of shoot branching by new terpenoid plant hormones. *Nature* 455, 195–200. doi: 10.1038/nature07272
- Waldie, T., and Leyser, O. (2018). Cytokinin targets auxin transport to promote shoot branching. *Plant Physiol.* 177, 803–818. doi: 10.1104/pp.17.01691
- Wang, L., Wang, B., Jiang, L., Liu, X., Li, X., Lu, Z., et al. (2015). Strigolactone signaling in Arabidopsis regulates shoot development by targeting D53-like SMXL repressor proteins for ubiquitination and degradation. *Plant Cell* 27, 3128–3142. doi: 10.1105/tpc.15.00605
- Wang, P., Zhao, Y., Li, Z., Hsu, C. C., Liu, X., Fu, L., et al. (2018). Reciprocal regulation of the TOR kinase and ABA receptor balances plant growth and stress response. *Mol. Cell* 69, 100–112. doi: 10.1016/j.molcel.2017.12.002
- Wang, R. L., Stec, A., Hey, J., Lukens, L., and Doebley, J. (1999). The limits of selection during maize domestication. *Nature* 398, 236–239. doi: 10.1038/18435
- Wang, Y., and Jiao, Y. (2018). Axillary meristem initiation—a way to branch out. *Curr. Opin. Plant Biol.* 41, 61–66. doi: 10.1016/j.pbi.2017.09.001
- Wang, Y., Sun, S., Zhu, W., Jia, K., Yang, H., and Wang, X. (2013). Strigolactone/MAX2-induced degradation of brassinosteroid transcriptional effector BES1 regulates shoot branching. *Dev. Cell* 27, 681–688. doi: 10.1016/j.devcel.2013.11.010
- Waters, M. T., Gutjahr, C., Bennett, T., and Nelson, D. C. (2017). Strigolactone signaling and evolution. *Annu. Rev. Plant Biol.* 68, 291–322. doi: 10.1146/annurev-arplant-042916-040925
- Wickson, M., and Thimann, K. V. (1958). The antagonism of auxin and kinetin in apical dominance. *Physiol. Plant.* 11, 62–74. doi: 10.1111/j.1399-3054.1958.tb08426.x
- Xu, J., Zha, M., Li, Y., Ding, Y., Chen, L., Ding, C., et al. (2015). The interaction between nitrogen availability and auxin, cytokinin, and strigolactone in the control of shoot branching in rice (*Oryza sativa* L.). *Plant Cell Rep.* 34, 1647–1662. doi: 10.1007/s00299-015-1815-8
- Yamada, Y., Furusawa, S., Nagasaka, S., Shimomura, K., Yamaguchi, S., and Umehara, M. (2014). Strigolactone signaling regulates rice leaf senescence in response to a phosphate deficiency. *Planta* 240, 399–408. doi: 10.1007/s00425-014-2096-0
- Yang, Y., Nicolas, M., Zhang, J., Yu, H., Guo, D., Yuan, R., et al. (2018). The TIE1 transcriptional repressor controls shoot branching by directly repressing *BRANCHED1* in Arabidopsis. *PLoS Genet.* 14:e1007296. doi: 10.1371/journal.pgen.1007296
- Yao, C., and Scott, A. F. (2015). Abscisic acid is a general negative regulator of Arabidopsis axillary bud growth. *Plant Physiol.* 169, 611–626. doi: 10.1104/pp.15.00682
- Yoneyama, K., Yoneyama, K., Takeuchi, Y., and Sekimoto, H. (2007). Phosphorus deficiency in red clover promotes exudation of orobanchol, the signal for mycorrhizal symbionts and germination stimulant for root parasites. *Planta* 225, 1031–1038. doi: 10.1007/s00425-006-0410-1
- Yoshida, T., Fujita, Y., Sayama, H., Kidokoro, S., Maruyama, K., Mizoi, J., et al. (2010). AREB1, AREB2, and ABF3 are master transcription factors that cooperatively regulate ABRE-dependent ABA signaling involved in drought stress tolerance and require ABA for full activation. *Plant J.* 61, 672–685. doi: 10.1111/j.1365-313X.2009.04092.x
- Zhang, Z. B., Yang, G., Arana, F., Chen, Z., Li, Y., and Xia, H. J. (2007). Arabidopsis inositol polyphosphate 6-/3-kinase (*AtIpk2β*) is involved in axillary shoot branching via auxin signaling. *Plant Physiol.* 144, 942–951. doi: 10.1104/pp.106.092163
- Zhao, D. L., Atlin, G. N., Bastiaans, L., and Spiertz, J. H. J. (2006). Developing selection protocols for weed competitiveness in aerobic rice. *Field Crops Res.* 97, 272–285. doi: 10.1016/j.fcr.2005.10.008
- Zhou, F., Lin, Q., Zhu, L., Ren, Y., Zhou, K., Shabek, N., et al. (2013). D14-SCFD3-dependent degradation of D53 regulates strigolactone signalling. *Nature* 504, 406–410. doi: 10.1038/nature12878
- Zhou, L., Zhang, J., Yan, J., and Song, R. (2011). Two transposable element insertions are causative mutations for the major domestication gene *teosinte branched 1* in modern maize. *Cell Res.* 21, 1267–1270. doi: 10.1038/cr.2011.104
- Zou, J., Zhang, S., Zhang, W., Li, G., Chen, Z., Zhai, W., et al. (2006). The rice *HIGH-TILLERING DWARF1* encoding an ortholog of Arabidopsis MAX3 is required for negative regulation of the outgrowth of axillary buds. *Plant J.* 48, 687–698. doi: 10.1111/j.1365-313X.2006.02916.x

**Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2019 Wang, Le Moigne, Bertheloot, Crespel, Perez-Garcia, Ogé, Demotes-Mainard, Hamama, Davière and Sakr. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.