



**HAL**  
open science

## Dynamic stability of the actin ecosystem

Julie Plastino, Laurent Blanchoin

► **To cite this version:**

Julie Plastino, Laurent Blanchoin. Dynamic stability of the actin ecosystem. *Journal of Cell Science*, 2019, 132 (4), pp.jcs219832. 10.1242/jcs.219832 . hal-02048154

**HAL Id: hal-02048154**

**<https://hal.science/hal-02048154>**

Submitted on 22 Oct 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## REVIEW

## SPECIAL ISSUE: RECONSTITUTING CELL BIOLOGY

# Dynamic stability of the actin ecosystem

Julie Plastino<sup>1,2,\*</sup> and Laurent Blanchoin<sup>3,4,\*</sup>

## ABSTRACT

In cells, actin filaments continuously assemble and disassemble while maintaining an apparently constant network structure. This suggests a perfect balance between dynamic processes. Such behavior, operating far out of equilibrium by the hydrolysis of ATP, is called a dynamic steady state. **This** dynamic steady state confers a high degree of plasticity to cytoskeleton networks that allows them to adapt and optimize their architecture in response to external changes on short time-scales, thus permitting cells to adjust to their environment. In this Review, we summarize what is known about the cellular actin steady state, and what gaps remain in our understanding of this fundamental dynamic process that balances the different forms of actin organization in a cell. We focus on the minimal steps to achieve a steady state, discuss the potential feedback mechanisms at play to balance this steady state and conclude with an outlook on what is needed to fully understand its molecular nature.

**KEY WORDS:** Actin, Cytoskeleton, Steady state

## Introduction

A key property of most living systems is their ability to move and/or change shape according to environmental cues. This is instrumental for the cell or the tissue to carry out its biological program, including cell processes **such as** division and motility, and developmental processes like morphogenesis. Understanding the dynamic steady state of actin is a major challenge in cell and developmental biology because actin is a key player and driving force in the construction of the complex and dynamic scaffolding that makes up the internal architecture of eukaryotic cells (Blanchoin et al., 2014; Chhabra and Higgs, 2007). Whereas many molecules that are involved in building the actin cytoskeleton are known, the basic rules that control the coordinated dynamics of structures that exist at the same time in the cell, such as branched networks, parallel bundles and antiparallel contractile bundles, are still poorly understood. The dynamic steady state of actin has four facets: nucleation (formation of actin dimers or trimers), elongation and capping (controlled polymer growth), disassembly (breakdown of actin structures) and recycling (replenishment of the pool of actin monomers that are charged with ATP) (Fig. 1). Here, **we consider** each facet in detail **and** highlight the potential existence of feedback mechanisms that could balance the dynamics of cellular actin, and the need for reconstitution experiments to fully dissect the balance of the steady state of actin.

<sup>1</sup>Institut Curie, PSL Research University, CNRS, 75005 Paris, France. <sup>2</sup>Sorbonne Université, 75005 Paris, France. <sup>3</sup>CytomorphaLab, Biosciences & Biotechnology Institute of Grenoble, Laboratoire de Physiologie Cellulaire & Végétale, Université Grenoble-Alpes/CEA/CNRS/INRA, 38054 Grenoble, France. <sup>4</sup>CytomorphaLab, Hôpital Saint Louis, Institut Universitaire d'Hématologie, UMRS1160, INSERM/AP-HP/Université Paris Diderot, 75010 Paris, France.

\*Authors for correspondence (julie.plastino@curie.fr, laurent.blanchoin@cea.fr)

 L.B., 0000-0001-8146-9254


## The pool of actin monomers

The elementary building block for actin assembly is the actin monomer. The concentration of actin monomers is quite variable in different living systems, ranging from potentially as low as 0.01  $\mu\text{M}$  in the yeast *Saccharomyces cerevisiae* to 300  $\mu\text{M}$  in unactivated platelets (Karpova et al., 1995; Pollard et al., 2000). Since the rate of actin assembly is directly proportional to the concentration of monomers (Pollard et al., 2000), this variability means that the dynamic steady state of actin is not the same in different cell types; actin filaments could potentially grow orders of magnitude faster in platelets than in yeast.

In the cell, most of the pool of polymerizable actin is bound to profilin (Fig. 1 and Kaiser et al., 1999). However, some actin monomers in a given cell type might not be polymerizable as they can be sequestered by proteins such as thymosin  $\beta$ 4 (T $\beta$ 4) (Fig. 1 and Pantaloni and Carlier, 1993). In this context, determining the exact concentration of polymerizable actin at a given time is challenging (Raz-Ben Aroush et al., 2017) as the balance between sequestered and polymerizable actin during dynamic actin assembly is not well characterized (Skruber et al., 2018). In addition, T $\beta$ 4 has been proposed to play an active role in preventing monomer incorporation into branched networks and in targeting cytosolic actin monomers to formin-mediated assembly at the leading edge of cells (Vitriol et al., 2015). This polymerizable pool of actin monomers needs further characterization at the cellular level, but also at the subcellular level – at sites of active actin assembly – where the pool of polymerizable actin can become depleted (Boujemaa-Paterski et al., 2017). In addition, different actin networks compete for actin monomers, and this is crucial for determining network density and size (Burke et al., 2014; Suarez and Kovar, 2016). To understand the landscape of the monomer pool in a complex, actively polymerizing, network as **found** in keratocyte fragments (Raz-Ben Aroush et al., 2017), it will be necessary to use methods such as fluorescence recovery after photobleaching (FRAP) or photoactivation and/or photoconversion experiments coupled with mathematical modeling to assess actin monomer dynamics in different *in vivo* contexts (Skruber et al., 2018) and/or to use cell-size confinement to generate reconstituted systems with well-defined but limited sources of actin monomers. Given the importance of the local polymerizable actin concentration for determining actin dynamics, evaluating the pool of polymerizable actin – and potentially its gradients – in different cellular contexts is one of the key challenges of the coming years.

## Nucleation

Profilin prevents the spontaneous association of actin monomers into actin dimers and trimers, which are necessary intermediates prior to the formation of actin filaments (Dominguez, 2009; Sept and McCammon, 2001). These nucleation steps, which are thermodynamically unfavorable, are accelerated by actin nucleators (Fig. 1). Three main classes of actin nucleators have been characterized: the Arp2/3 complex, the formin family and the tandem monomer-binding protein family, including the proteins



**Fig. 1. The dynamic steady state of actin.** Minimal steps needed to reach a dynamic steady state of different actin architectures are illustrated. Sequestered and polymerizable actin represent the pool of actin monomers. Nucleation is the formation of actin dimers or trimers. Elongation and capping modulate controlled growth of the different forms of actin organization (branched networks that are generated by the Arp2/3 complex or bundles generated by formins). Disassembly and depolymerization results in the breakdown of actin structures to monomer subunits. Recycling renews the pool of actin monomers that are charged with ATP. The different nucleotide states of actin, barbed and pointed ends and different proteins or complexes are represented by the indicated symbols.

Spire and Cobl, which are proposed to cause nucleation by tethering three or more actin monomers together (Campellone and Welch, 2010). The different mechanisms of action and the cellular localization of these nucleators influence their physiological properties and abilities to build specific actin structures. The Arp2/3 complex and formins are also influenced by external signals as their activity is controlled either directly or indirectly by membrane-bound Rho GTPases (Lawson and Ridley, 2018; Ridley, 2015). In addition, profilin has been shown to favor formin-mediated assembly over Arp2/3-based nucleation, thus modulating homeostasis between different networks (Suarez et al., 2015; Rotty et al., 2015).

**Elongation and capping**

Pointed ends do not grow in profilin-actin since profilin masks the barbed face of the actin monomer, preventing addition to pointed ends (Pollard et al., 2000). Therefore, in cellular conditions of profilin-actin, networks that are generated by the Arp2/3 complex grow with a rate that depends on the association rate constant for

monomer addition at the barbed ends (which is typically 10 per μM per second, (Pollard, 1986) and the concentration of polymerizable actin monomers. Growth is terminated by a lack of monomers or by capping proteins (Fig. 1). Therefore, the balance between rates of growth and capping needs to be well-adjusted for the formation of defined actin networks (Akin and Mullins, 2008; Blanchoin et al., 2000a; Kawska et al., 2012). This balance is illustrated by a comparison of the dynamics of two similar structures found in different cells, where the actin monomer concentrations are very different. For example, in yeast, actin patches and cell lamellipodia are both generated by the Arp2/3 complex and consist of highly branched and intertwined actin networks (Young et al., 2004; Svitkina and Borisy, 1999). Based on the concentration of available monomers, the growth of a lamellipodium should be orders of magnitude faster than a patch, but this is not the case because capping proteins regulate growth in lamellipodia (Moseley and Goode, 2006). Indeed, capping proteins are in fact necessary for correct lamellipodia formation in motile cells with a high concentration of monomers (Iwasa and Mullins, 2007). In

contrast, in *S. cerevisiae*, where monomer concentration is very low, capping protein can be removed and actin patches still show a qualitatively normal organization (Young et al., 2004). Growth of actin filaments that is mediated by formin is even more complex because different formins produce different association rate constants at filament barbed ends (Chesarone and Goode, 2009). In addition, formin and capping proteins function antagonistically at the barbed ends of actin filaments to control their length through the formation of a ‘decision complex’, where capping protein and formin are simultaneously bound to a paused barbed end. Depending on how the complex decomposes, growth will resume or the filament will be permanently capped, lending an extra layer of control to the dynamics of the barbed end (Fig. 1, Bombardier et al., 2015; Shekhar et al., 2015). Finally, formins are known to cooperate with the Arp2/3 complex for the elongation of protrusive networks (Block et al., 2012; Kage et al., 2017). Taken together, elongation and capping, in the context of the amount of polymerizable monomeric actin and the nature of the nucleating agent, are what defines the dynamics of actin network growth.

### Disassembly

Actin disassembly takes place in two steps: fragmentation of actin networks into small filaments, and depolymerization into monomers of the small actin fragments that were generated in this process (Fig. 1, Blanchoin et al., 2014). Like nucleation or elongation, disassembly depends on the nature of the network (Gressin et al., 2015). A branched network disassembles mainly through debranching that is mediated by the actin depolymerizing factor (ADF)/cofilin family proteins [comprising ADF (also known as destrin), cofilin-1 (CFL1) and cofilin-2 (CFL2)] or glia maturation factor-like protein (GMF) (Blanchoin et al., 2000b; Chan et al., 2009; Gandhi et al., 2010; Gressin et al., 2015). For parallel or antiparallel networks of bundled filaments, the severing of filaments by ADF/cofilin on its own is not sufficient to dismantle the structure entirely (Gressin et al., 2015), but enough to maintain a steady state length, at least for parallel bundles that are initiated by formin (Michelot et al., 2007). Actin-interacting protein 1 (Aip1, also known as WDR1) is a necessary cofactor that synergizes with ADF/cofilin and coronin proteins to disassemble actin bundles (Nadkarni and Briher, 2014; Gressin et al., 2015; Jansen et al., 2015). For antiparallel contractile actin networks, the contribution of the myosin motor protein during disassembly is unclear, but actin filament buckling that is produced by myosin contraction has been shown to lead to filament breakage (Murrell and Gardel, 2012), and buckling might also favor severing by ADF/cofilin. To be fully efficient, ADF/cofilin must work in concert with capping proteins (Suarez et al., 2011). Indeed, the presence of ATP or ADP-Pi-loaded subunits near a growing barbed end prevents this region of the filament from being decorated by ADF/cofilin (Suarez et al., 2011; Wioland et al., 2017). Barbed-end capping terminates the growth of filaments within a structure, and thus favors decoration of filaments by ADF/cofilin (Suarez et al., 2011). Therefore, the disassembly of actin networks is intricately linked to the growth and capping balance of the actin steady state, as discussed above.

Until recently, a few puzzling questions remained concerning the disassembly step. First, why do fragments that are generated by ADF/cofilin not elongate rapidly until capped, thus reversing the disassembly effect of ADF/cofilin? Second, how does rapid disassembly from capped fragments occur? The rate constant of depolymerization at pointed ends is only 0.27 per second, and depolymerization would be much more efficient if it occurred at the barbed end with a rate constant of 7.2 per second (Pollard, 1986).

Recently, these two questions have been elegantly addressed by Wioland and co-workers, who showed that decoration with ADF/cofilin – upon nearing the barbed end – dissociates capping protein from that end (Wioland et al., 2017). Even more striking, barbed ends of filaments that are saturated with ADF/cofilin do not grow because ADF/cofilin prevents monomer addition (Wioland et al., 2017). In other words, ADF/cofilin alters barbed end dynamics by binding to the sides of the filament and changing its structure (Tanaka et al., 2018), thus preventing capping protein binding and monomer addition, while still allowing subunit dissociation (Fig. 1). Other proteins, such as twinfilin and Srv2/profilin and cyclase-associated protein (CAP), which accelerate depolymerization at the barbed and pointed ends, can also participate in actin disassembly (Johnston et al., 2015). Their collaborative effort depolymerizes a filament of one micrometer length in less than a minute (Johnston et al., 2015). How such depolymerization occurs on ADF/cofilin-decorated filaments, or on small fragments that are generated by both ADF/cofilin and Aip1, needs to be investigated. Srv2/CAP has also been shown to enhance severing by ADF/cofilin (Chaudhry et al., 2013). It is still unclear exactly how small fragments depolymerize into single subunits. Recent advances in time-resolved electron microscopy, combined with static and dynamic light scattering might help in the capture and identification of these entities, which fall below the diffraction limit (Frank, 2017; Lopez et al., 2016).

### Recycling

An assembly-competent actin oligomer pool has previously been proposed (Okreglak and Drubin, 2010; Smith et al., 2013); however, for most reassembly processes, actin filaments must be broken down into their individual monomers. Actin monomers are bound to ADP when they dissociate from a filament (Blanchoin and Pollard, 1999). These subunits therefore need to be reloaded with ATP to reintegrate into the pool of sequestered or polymerizable actin (Fig. 1). As ADF/cofilin bound to an actin subunit blocks nucleotide exchange (the rate of nucleotide dissociation of ADF/cofilin-bound ADP-actin is 0.006 per second (Blanchoin and Pollard, 1998), profilin or Srv2/CAP act as nucleotide exchanging factors: they dissociate ADF/cofilin from ADP-actin subunits and load subunits with ATP (Blanchoin and Pollard, 1998; Chaudhry et al., 2010; Gurel et al., 2015; Kotila et al., 2018). This replenishes the pool of polymerizable actin (Fig. 1). In the presence of high concentrations of T $\beta$ 4, as in platelets, the situation is more complex, because thymosins also block nucleotide exchange (Goldschmidt-Clermont et al., 1992; Xue et al., 2014). However, T $\beta$ 4 has a 100-fold higher affinity for ATP-actin compared with ADP-actin monomers (Jean et al., 1994), so nucleotide exchange probably occurs before thymosin binds monomers. This might occur through formation of a transient ternary complex between T $\beta$ 4, actin monomer and profilin, or other nucleotide-exchanging factors (Yarmola et al., 2001). Overall, a complex choreography of actin-binding proteins controls the recycling of ADP-actin monomers to the polymerizable or sequestered ATP form.

### Feedback

The huge variability in the structure of different forms of actin organization, their growth rates and lifetimes beg the question as to how the perfect match between assembly and disassembly rates and maintenance of the pool of actin monomers is ensured in these different contexts. This must be controlled by as yet unidentified feedback mechanisms. Is it a structural feedback, where network size, structure and filament density affect actin dynamics, a mechanical feedback, where tension and pressure regulate dynamics, or even a

AQ4

AQ3

373 biochemical feedback, where polymerization depletes factors, thus  
 374 limiting assembly or disassembly? It is likely that it is a combination  
 375 of these different types of feedback. Structural feedback has been  
 376 observed recently in a reconstituted lamellipodium, where network  
 377 size and filament density have been shown to control network growth  
 378 rate (Boujemaa-Paterski et al., 2017). Force-dependent feedback  
 379 controls both the growth of branched networks generated by the  
 380 Arp2/3 complex (Bieling et al., 2016; Mueller et al., 2017; Plastino  
 381 and Blanchoin, 2017) and formin-mediated actin filament assembly  
 382 (Courtemanche et al., 2013; Jégou et al., 2013; Zimmermann et al.,  
 383 2017). Biochemical control is seen in the competition for actin  
 384 monomers between formin-based and Arp2/3-based actin networks  
 385 (Burke et al., 2014) or in local monomer depletion at sites of active  
 386 assembly that negatively impacts growth rate (Boujemaa-Paterski  
 387 et al., 2017). The identification and mechanistic understanding of the  
 388 different feedback loops that control cellular actin dynamics will  
 389 require a huge effort from both top-down and bottom-up approaches,  
 390 bridging the gap between **investigations** at molecular, cellular and  
 391 **tissue levels**.

### 393 Conclusions and perspectives

394 What are the limitations to achieving a complete understanding of  
 395 the dynamic steady state of actin networks? *In vivo*, the biggest  
 396 limitation is the observation of individual actin filaments whose  
 397 average lengths are 10 to 100 nm, which is below the diffraction  
 398 limit of light microscopy (Anderson et al., 2017). The development  
 399 of new super-resolution approaches and new fluorescent markers,  
 400 combined with electron microscopy, might help to fill this gap (Gao  
 401 et al., 2018; Skruber et al., 2018). However, a true understanding  
 402 will require visualization of the actin cytoskeleton in its native state,  
 403 imaging the coordinated dynamics of different subcellular actin  
 404 organizations. A step in this direction is the *in vitro* reconstitution of  
 405 a complete dynamic system, where branched networks, parallel  
 406 bundles and contractile antiparallel structures maintain a  
 407 coordinated steady state regime in a cell-sized environment that  
 408 mimics the limited supply of biochemical components in a real cell  
 409 (Burke et al., 2014). Growing different actin organizations has been  
 410 partially achieved using micropatterning approaches (Reymann  
 411 et al., 2010), but never with a combination of different nucleation  
 412 machineries. Growth must be initiated in the presence of both the  
 413 disassembly machinery and the proteins necessary to recycle actin  
 414 subunits back to the pool of polymerizable actin. Ideally, this  
 415 reconstituted system would allow for the modulation in real time of  
 416 the different actin organizations, **such as** changing the pattern of  
 417 nucleation to evaluate how the system responds and adapts to this  
 418 new configuration. One of the biggest challenges is determining the  
 419 operating concentrations for the different components of a complex  
 420 mixture incorporating nucleation, turnover and actin recycling  
 421 machinery. Parallelizing the experiments by means of microfluidics  
 422 will probably be necessary. The field is technically ready to tackle  
 423 this challenge, both *in vitro* and *in vivo*, but it will be necessary to  
 424 join forces, as the task is too complex for a single laboratory.

### 426 Acknowledgements

427 We would like to thank Manuel Théry for intensive discussions on the actin dynamic  
 428 steady state and Agnieszka Kawska for Fig. 1.

### 429 Competing interests

430 **AQ5** The authors declare no competing or financial interests.

### 432 Funding

433 This work was supported by a European Research Council grant to L.B. (Advanced  
 434 grant AAA, 741773). J.P. acknowledges financial support from Fondation ARC pour

la Recherche sur le Cancer (PJA 20151203487) and from the IDEX Université de  
 Recherche Paris Sciences et Lettres (ANR-10-IDEX-0001-01 PSL).

### 437 References

- 438 **Akin, O. and Mullins, R. D.** (2008). Capping protein increases the rate of actin-  
 439 based motility by promoting filament nucleation by the Arp2/3 complex. *Cell* **133**,  
 440 841-851.
- 441 **Anderson, K. L., Page, C., Swift, M. F., Suraneni, P., Janssen, M. E. W., Pollard,  
 442 T. D., Li, R., Volkman, N. and Hanein, D.** (2017). Nano-scale actin-network  
 443 characterization of fibroblast cells lacking functional Arp2/3 complex. *J. Struct.  
 444 Biol.* **197**, 312-321.
- 444 **Bieling, P., Li, T. D., Weichsel, J., McGorty, R., Jreij, P., Huang, B., Fletcher, D. A.  
 445 and Mullins, R. D.** (2016). Force feedback controls motor activity and mechanical  
 446 properties of self-assembling branched actin networks. *Cell* **164**, 115-127.
- 446 **Blanchoin, L. and Pollard, T. D.** (1998). Interaction of actin monomers with  
 447 *Acanthamoeba* actophorin (ADF/cofilin) and profilin. *J. Biol. Chem.* **273**,  
 448 25106-25111.
- 449 **Blanchoin, L. and Pollard, T. D.** (1999). Mechanism of interaction of  
 450 *Acanthamoeba* actophorin (ADF/cofilin) with actin filaments. *J. Biol. Chem.* **274**,  
 451 15538-15546.
- 451 **Blanchoin, L., Amann, K. J., Higgs, H. N., Marchand, J.-B., Kaiser, D. A. and  
 452 Pollard, T. D.** (2000a). Direct observation of dendritic actin filament networks  
 453 nucleated by Arp2/3 complex and WASP/Scar proteins. *Nature* **404**, 1007-1111.
- 453 **Blanchoin, L., Pollard, T. D. and Mullins, R. D.** (2000b). Interaction of ADF/cofilin,  
 454 Arp2/3 complex, capping protein and profilin in remodeling of branched actin  
 455 filament networks. *Curr. Biol.* **10**, 1273-1282.
- 455 **Blanchoin, L., Boujemaa-Paterski, R., Sykes, C. and Plastino, J.** (2014). Actin  
 456 dynamics, architecture, and mechanics in cell motility. *Physiol. Rev.* **94**, 235-263.
- 457 **Block, J., Breitsprecher, D., Kühn, S., Winterhoff, M., Kage, F., Geffers, R.,  
 458 Duwe, P., Rohn, J. L., Baum, B., Brakebusch, C. et al.** (2012). FMNL2 drives  
 459 actin-based protrusion and migration downstream of Cdc42. *Curr. Biol.* **22**,  
 460 1005-1012.
- 460 **Bombardier, J. P., Eskin, J. A., Jaiswal, R., Correa, I. R., Jr, Xu, M.-Q., Goode,  
 461 B. L. and Gelles, J.** (2015). Single-molecule visualization of a formin-capping  
 462 protein 'decision complex' at the actin filament barbed end. *Nat. Commun.*  
 463 **6**, 8707.
- 463 **Boujemaa-Paterski, R., Suarez, C., Klar, T., Zhu, J., Guérin, C., Mogilner, A.,  
 464 Théry, M. and Blanchoin, L.** (2017). Network heterogeneity regulates steering in  
 465 actin-based motility. *Nat. Commun.* **8**, 655.
- 465 **Burke, T. A., Christensen, J. R., Barone, E., Suarez, C., Sirotkin, V. and Kovar,  
 466 D. R.** (2014). Homeostatic actin cytoskeleton networks are regulated by assembly  
 467 factor competition for monomers. *Curr. Biol.* **24**, 579-585.
- 467 **Campellone, K. G. and Welch, M. D.** (2010). A nucleator arms race: cellular control  
 468 of actin assembly. *Nat. Rev. Mol. Cell Biol.* **11**, 237-251.
- 468 **Chan, C., Beltzner, C. C. and Pollard, T. D.** (2009). Cofilin dissociates Arp2/3  
 469 complex and branches from actin filaments. *Curr. Biol.* **19**, 537-545.
- 469 **Chaudhry, F., Little, K., Talarico, L., Quintero-Monzon, O. and Goode, B. L.**  
 470 (2010). A central role for the WH2 domain of Srv2/CAP in recharging actin  
 471 monomers to drive actin turnover in vitro and in vivo. *Cytoskeleton (Hoboken)* **67**,  
 472 120-133.
- 472 **Chaudhry, F., Breitsprecher, D., Little, K., Sharov, G., Sokolova, O. and Goode,  
 473 B. L.** (2013). Srv2/cyclase-associated protein forms hexameric shirikens that  
 474 directly catalyze actin filament severing by cofilin. *Mol. Biol. Cell* **24**, 31-41.
- 474 **Chesarone, M. A. and Goode, B. L.** (2009). Actin nucleation and elongation factors:  
 475 mechanisms and interplay. *Curr. Opin. Cell Biol.* **21**, 28-37.
- 475 **Chhabra, E. S. and Higgs, H. N.** (2007). The many faces of actin: matching  
 476 assembly factors with cellular structures. *Nat. Cell Biol.* **9**, 1110-1121.
- 476 **Courtemanche, N., Lee, J. Y., Pollard, T. D. and Greene, E. C.** (2013). Tension  
 477 modulates actin filament polymerization mediated by formin and profilin. *Proc.  
 478 Natl. Acad. Sci. USA* **110**, 9752-9757.
- 478 **Dominguez, R.** (2009). Actin filament nucleation and elongation factors—structure-  
 479 function relationships. *Crit. Rev. Biochem. Mol. Biol.* **44**, 351-366.
- 479 **Frank, J.** (2017). Time-resolved cryo-electron microscopy: recent progress.  
 480 *J. Struct. Biol.* **200**, 303-306.
- 480 **Gandhi, M., Smith, B. A., Bovellan, M., Paavilainen, V., Daugherty-Clarke, K.,  
 481 Gelles, J., Lappalainen, P. and Goode, B. L.** (2010). GMF is a cofilin homolog  
 482 that binds Arp2/3 complex to stimulate filament debranching and inhibit actin  
 483 nucleation. *Curr. Biol.* **20**, 861-867.
- 483 **Gao, M., Maraspin, R., Beutel, O., Zehtabian, A., Eickholt, B., Honigsmann, A.  
 484 and Ewers, H.** (2018). Expansion stimulated emission depletion microscopy  
 485 (ExSTED). *ACS Nano* **12**, 4178-4185.
- 485 **Goldschmidt-Clermont, P. J., Furman, M. I., Wachsstock, D., Safer, D.,  
 486 Nachmias, V. T. and Pollard, T. D.** (1992). The control of actin nucleotide  
 487 exchange by thymosin $\beta$ 4 and profilin. A potential regulatory mechanism for actin  
 488 polymerization in cells. *Mol. Biol. Cell* **3**, 1015-1024.
- 488 **Gressin, L., Guillotin, A., Guérin, C., Blanchoin, L. and Michelot, A.** (2015).  
 489 Architecture dependence of actin filament network disassembly. *Curr. Biol.* **25**,  
 490 1437-1447.

497 Gurel, P. S., A. M., Guo, B., Shu, R., Mierke, D. F. and Higgs, H. N. (2015). Assembly and turnover of short actin filaments by the formin INF2 and profilin. *J Biol. Chem.* **290**, 22494-22506.

498

499 Iwasa, J. H. and Mullins, R. D. (2007). Spatial and temporal relationships between actin-filament nucleation, capping, and disassembly. *Curr. Biol.* **17**, 395-406.

500

501 Jansen, S., Collins, A., Chin, S. M., Ydenberg, C. A., Gelles, J. and Goode, B. L. (2015). Single-molecule imaging of a three-component ordered actin disassembly mechanism. *Nat. Commun.* **6**, 7202.

502

503 Jean, C., Rieger, K., Blanchoin, L., Carlier, M.-F., Lenfant, M. and Pantaloni, D. (1994). Interaction of G-actin with thymosin beta 4 and its variants thymosin beta 9 and thymosin beta met9. *J. Muscle Res. Cell Motil.* **15**, 278-286.

504

505 Jégou, A., Carlier, M.-F. and Romet-Lemonne, G. (2013). Formin mDia1 senses and generates mechanical forces on actin filaments. *Nat. Commun.* **4**, 1883.

506

507 Johnston, A. B., Collins, A. and Goode, B. L. (2015). High-speed depolymerization at actin filament ends jointly catalysed by Twinfilin and Srv2/CAP. *Nat. Cell Biol.* **17**, 1504-1511.

508

509 Kage, F., Winterhoff, M., Dimchev, V., Mueller, J., Thalheim, T., Freise, A., Bruhmann, S., Kollasser, J., Block, J., Dimchev, G. et al. (2017). FMNL formins boost lamellipodial force generation. *Nat. Commun.* **8**, 14832.

510

511 Kaiser, D. A., Vinson, V. K., Murphy, D. B. and Pollard, T. D. (1999). Profilin is predominantly associated with monomeric actin in *Acanthamoeba*. *J. Cell Sci.* **112**, 3779-3790.

512

513 Karpova, T. S., Tatchell, K. and Cooper, J. A. (1995). Actin filaments in yeast are unstable in the absence of capping protein or fimbrin. *J. Cell Biol.* **131**, 1483-1493.

514

515 Kawska, A., Carvalho, K., Manzi, J., Boujemaa-Paterski, R., Blanchoin, L., Martiel, J.-L. and Sykes, C. (2012). How actin network dynamics control the onset of actin-based motility. *Proc. Natl. Acad. Sci. USA* **109**, 14440-14445.

516

517 Kotila, T., Kogan, K., Enkavi, G., Guo, S., Vattulainen, I., Goode, B. L. and Lappalainen, P. (2018). Structural basis of actin monomer re-charging by cyclase-associated protein. *Nat. Commun.* **9**, 1892.

518

519 Lawson, C. D. and Ridley, A. J. (2018). Rho GTPase signaling complexes in cell migration and invasion. *J. Cell Biol.* **217**, 447-457.

520

521 Lopez, C. G., Saldanha, O., Huber, K. and Köster, S. (2016). Lateral association and elongation of vimentin intermediate filament proteins: a time-resolved light-scattering study. *Proc. Natl. Acad. Sci. USA* **113**, 11152-11157.

522

523 Michelot, A., Berro, J., Guérin, C., Boujemaa-Paterski, R., Staiger, C. J., Martiel, J.-L. and Blanchoin, L. (2007). Actin-filament stochastic dynamics mediated by ADF/cofilin. *Curr. Biol.* **17**, 825-833.

524

525 Moseley, J. B. and Goode, B. L. (2006). The yeast actin cytoskeleton: from cellular function to biochemical mechanism. *Microbiol. Mol. Biol. Rev.* **70**, 605-645.

526

527 Mueller, J., Szep, G., Nemethova, M., de Vries, I., Lieber, A. D., Winkler, C., Kruse, K., Small, J. V., Schmeiser, C., Keren, K. et al. (2017). Load adaptation of lamellipodial actin networks. *Cell* **171**, 188-200.e16.

528

529 Murrell, M. P. and Gardel, M. L. (2012). F-actin buckling coordinates contractility and severing in a biomimetic actomyosin cortex. *Proc. Natl. Acad. Sci. USA* **109**, 20820-20825.

530

531 Nadkarni, A. V. and Briehner, W. M. (2014). Aip1 destabilizes cofilin-saturated actin filaments by severing and accelerating monomer dissociation from ends. *Curr. Biol.* **24**, 2749-2757.

532

533 Okreglak, V. and Drubin, D. G. (2010). Loss of Aip1 reveals a role in maintaining the actin monomer pool and an in vivo oligomer assembly pathway. *J. Cell Biol.* **188**, 769-777.

534

535 Pantaloni, D. and Carlier, M.-F. (1993). How profilin promotes actin filament assembly in the presence of thymosinβ4. *Cell* **75**, 1007-1014.

536

537 Plastino, J. and Blanchoin, L. (2017). Adaptive actin networks. *Dev. Cell* **42**, 565-566.

538

539 Pollard, T. D. (1986). Rate constants for the reactions of ATP- and ADP-actin with the ends of actin filaments. *J. Cell Biol.* **103**, 2747-2754.

540

541

542

543

544

545

546

547

548

549

550

551

552

553

554

555

556

557

558

559 Pollard, T. D., Blanchoin, L. and Mullins, R. D. (2000). Molecular mechanisms controlling actin filament dynamics in nonmuscle cells. *Annu. Rev. Biophys.* **29**, 545-576.

560

561 Raz-Ben Aroush, D., Ofer, N., Abu-Shah, E., Allard, J., Krichevsky, O., Mogilner, A. and Keren, K. (2017). Actin turnover in lamellipodial fragments. *Curr. Biol.* **27**, 2963-2973 e14.

562

563 Reymann, A.-C., Martiel, J.-L., Cambier, T., Blanchoin, L., Boujemaa-Paterski, R. and Théry, M. (2010). Nucleation geometry governs ordered actin networks structures. *Nat. Mat.* **9**, 827-832.

564

565 Ridley, A. J. (2015). Rho GTPase signalling in cell migration. *Curr. Opin. Cell Biol.* **36**, 103-112.

566

567 Rotty, J. D., Wu, C., Haynes, E. M., Suarez, C., Winkelman, J. D., Johnson, H. E., Haugh, J. M., Kovar, D. R. and Bear, J. E. (2015). Profilin-1 serves as a gatekeeper for actin assembly by Arp2/3-dependent and -independent pathways. *Dev. Cell* **32**, 54-67.

568

569 Sept, D. and McCammon, J. A. (2001). Thermodynamics and kinetics of actin filament nucleation. *Biophys. J.* **81**, 667-674.

570

571 Shekhar, S., Kerleau, M., Kühn, S., Pernier, J., Romet-Lemonne, G., Jégou, A. and Carlier, M.-F. (2015). Formin and capping protein together embrace the actin filament in a menage a trois. *Nat. Commun.* **6**, 8730.

572

573 Skrubber, K., Read, T. A. and Vitriol, E. A. (2018). Reconsidering an active role for G- actin in cytoskeletal regulation. *J. Cell Sci.* **131**.

574

575 Smith, M. B., Kiuchi, T., Watanabe, N. and Vavylonis, D. (2013). Distributed actin turnover in the lamellipodium and FRAP kinetics. *Biophys. J.* **104**, 247-257.

576

577 Suarez, C. and Kovar, D. R. (2016). Internetwork competition for monomers governs actin cytoskeleton organization. *Nat. Rev. Mol. Cell Biol.* **17**, 799-810.

578

579 Suarez, C., Roland, J., Boujemaa-Paterski, R., Kang, H., McCullough, B. R., Reymann, A.-C., Guérin, C., Martiel, J.-L., De la Cruz, E. M. and Blanchoin, L. (2011). Cofilin tunes the nucleotide state of actin filaments and severs at bare and decorated segment boundaries. *Curr. Biol.* **21**, 862-868.

580

581 Suarez, C., Carroll, R. T., Burke, T. A., Christensen, J. R., Bestul, A. J., Sees, J. A., James, M. L., Sirotkin, V. and Kovar, D. R. (2015). Profilin regulates F-actin network homeostasis by favoring formin over Arp2/3 complex. *Dev. Cell* **32**, 43-53.

582

583 Svitkina, T. M. and Borisy, G. C. (1999). Arp2/3 complex and actin depolymerizing factor/cofilin in dendritic organization and treadmilling of actin filament array in lamellipodia. *J. Cell Biol.* **145**, 1009-1026.

584

585 Tanaka, K., Takeda, S., Mitsuoka, K., Oda, T., Kimura-Sakiyama, C., Maeda, Y. and Narita, A. (2018). Structural basis for cofilin binding and actin filament disassembly. *Nat. Commun.* **9**, 1860.

586

587 Vitriol, E. A., McMillen, L. M., Kapustina, M., Gomez, S. M., Vavylonis, D. and Zheng, J. Q. (2015). Two functionally distinct sources of actin monomers supply the leading edge of lamellipodia. *Cell Rep.* **11**, 433-445.

588

589 Wioland, H., Guichard, B., Senju, Y., Myram, S., Lappalainen, P., Jegou, A. and Romet-Lemonne, G. (2017). ADF/Cofilin accelerates actin dynamics by severing filaments and promoting their depolymerization at both ends. *Curr. Biol.* **27**, 1956-1967 e7.

590

591 Xue, B., Leyrat, C., Grimes, J. M. and Robinson, R. C. (2014). Structural basis of thymosin-beta4/profilin exchange leading to actin filament polymerization. *Proc. Natl. Acad. Sci. USA* **111**, E4596-E4605.

592

593 Yarmola, E. G., Parikh, S. and Bubb, M. R. (2001). Formation and implications of a ternary complex of profilin, thymosin beta 4, and actin. *J. Biol. Chem.* **276**, 45555-45563.

594

595 Young, M. E., Cooper, J. A. and Bridgman, P. C. (2004). Yeast actin patches are networks of branched actin filaments. *J. Cell Biol.* **166**, 629-635.

596

597 Zimmermann, D., Homa, K. E., Hocky, G. M., Pollard, L. W., De La Cruz, E. M., Voth, G. A., Trybus, K. M. and Kovar, D. R. (2017). Mechanoregulated inhibition of formin facilitates contractile actomyosin ring assembly. *Nat. Commun.* **8**, 7.

598

599

600

601

602

603

604

605

606

607

608

609

610

611

612

613

614

615

616

617

618

619

620

AQ6

**AQ8 Summary:**

**Funding details**

S.No.	Funder name	Funder ID	Grant ID
1	European Research Council	<a href="http://dx.doi.org/10.13039/100010663">http://dx.doi.org/10.13039/100010663</a>	Advanced grant AAA 741773
2	Fondation ARC pour la Recherche sur le Cancer	<a href="http://dx.doi.org/10.13039/501100004097">http://dx.doi.org/10.13039/501100004097</a>	PJA 20151203487
3	Université de Recherche Paris Sciences et Lettres		ANR-10-IDEX-0001-01 PSL