



HAL
open science

Hardy-Sobolev inequalities with singularities on non smooth boundary. Part 2: Influence of the global geometry in small dimensions

Hussein Cheikh Ali

► **To cite this version:**

Hussein Cheikh Ali. Hardy-Sobolev inequalities with singularities on non smooth boundary. Part 2: Influence of the global geometry in small dimensions. *Journal of Differential Equations*, 2021, 270, pp.185-216. 10.1016/j.jde.2020.07.043 . hal-02048901v2

HAL Id: hal-02048901

<https://hal.science/hal-02048901v2>

Submitted on 5 Oct 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**HARDY-SOBOLEV INEQUALITIES WITH SINGULARITIES ON
NON SMOOTH BOUNDARY. PART 2: INFLUENCE OF THE
GLOBAL GEOMETRY IN SMALL DIMENSIONS.**

HUSSEIN CHEIKH ALI

ABSTRACT. We consider Hardy-Sobolev nonlinear equations on domains with singularities. We introduced this problem in Cheikh-Ali [4]. Under a local geometric hypothesis, namely that the generalized mean curvature is negative (see (7) below), we proved the existence of extremals for the relevant Hardy-Sobolev inequality for large dimensions. In the present paper, we tackle the question of small dimensions that was left open. We introduce a "mass", that is a global quantity, the positivity of which ensures the existence of extremals in small dimensions. As a byproduct, we prove the existence of solutions to a perturbation of the initial equation via the Mountain-Pass Lemma.

CONTENTS

| | | |
|----|--|----|
| 1. | Introduction | 1 |
| 2. | Definition of the generalized curvature and the mass | 4 |
| 3. | Some background results | 5 |
| 4. | Test-functions estimates for the mass: proof of Theorem 1.2 | 6 |
| 5. | Examples of mass | 14 |
| 6. | Proof of Theorem 1.3: functional background for the perturbed equation | 15 |
| 7. | Proof of Theorem 1.3: Test-Functions estimates | 19 |
| | References | 24 |

1. INTRODUCTION

Let Ω be a bounded domain of \mathbb{R}^n , $n \geq 3$. We fix $s \in [0, 2]$ and $\gamma \in \mathbb{R}$. It follows from the classical Caffarelli-Kohn-Nirenberg inequalities [3] that if $\gamma < \frac{(n-2)^2}{4}$, there exists $K > 0$ such that

$$(1) \quad \left(\int_{\Omega} \frac{|u|^{2^*(s)}}{|x|^s} dx \right)^{\frac{2}{2^*(s)}} \leq K \int_{\Omega} \left(|\nabla u|^2 - \gamma \frac{u^2}{|x|^2} \right) dx,$$

for all $u \in D^{1,2}(\Omega)$, where $2^*(s) := \frac{2(n-s)}{n-2}$ and $D^{1,2}(\Omega)$ is the completion of $C_c^\infty(\Omega)$ with respect to the norm $u \mapsto \|\nabla u\|_2$. We define the Hardy constant by

$$\gamma_H(\Omega) := \inf \left\{ \frac{\int_{\Omega} |\nabla u|^2 dx}{\int_{\Omega} \frac{u^2}{|x|^2} dx}; u \in D^{1,2}(\Omega) \setminus \{0\} \right\} > 0.$$

The classical Hardy inequality reads $\gamma_H(\mathbb{R}^n) = \frac{(n-2)^2}{4}$ and therefore, we have that $\gamma_H(\Omega) \geq \frac{(n-2)^2}{4}$. We refer to [4] for discussions and properties of the Hardy constant. As one checks, for any $\gamma < \gamma_H(\Omega)$, there exists $K = K(\Omega, \gamma, s) > 0$ such that (1) holds for all $u \in D^{1,2}(\Omega)$. For $a \in L^\infty(\Omega)$, we define

$$\mu_{\gamma,s,a}(\Omega) = \inf_{u \in D^{1,2}(\Omega) \setminus \{0\}} J_{\gamma,s,a}^\Omega(u),$$

where

$$J_{\gamma,s,a}^\Omega(u) := \frac{\int_\Omega \left(|\nabla u|^2 - \left(\frac{\gamma}{|x|^2} + a(x) \right) u^2 \right) dx}{\left(\int_\Omega \frac{|u|^{2^*(s)}}{|x|^s} dx \right)^{\frac{2}{2^*(s)}}},$$

so that

$$(2) \quad \mu_{\gamma,s,a}(\Omega) \left(\int_\Omega \frac{|u|^{2^*(s)}}{|x|^s} dx \right)^{\frac{2}{2^*(s)}} \leq \int_\Omega \left(|\nabla u|^2 - \left(\frac{\gamma}{|x|^2} + a(x) \right) u^2 \right) dx,$$

for all $u \in D^{1,2}(\Omega)$. As in [4], we address the question of the existence of extremals for (2), more precisely

Q: Does there exist $u \in D^{1,2}(\Omega) \setminus \{0\}$ for which equality holds in (2)?

When $0 \in \Omega$, there are no extremals for $\mu_{\gamma,s,0}(\Omega)$ (see [7]). From now on, we assume that $0 \in \partial\Omega$. When Ω is a smooth domain, criteria for existence are in Ghoussoub-Robert [8]: in particular, there is a dichotomy between large dimension (where the criterion is local) and the small dimensions (where the criterion is global). In [4], we studied the case of domains that are modeled on cones:

Definition 1. We fix $1 \leq k \leq n$. Let Ω be a domain of \mathbb{R}^n . We say that $x_0 \in \partial\Omega$ is a singularity of type $(k, n-k)$ if there exist U, V open subsets of \mathbb{R}^n such that $0 \in U$, $x_0 \in V$ and there exists a diffeomorphism $\phi \in C^\infty(U, V)$ such that $\phi(0) = x_0$ and

$$\phi(U \cap (\mathbb{R}_+^k \times \mathbb{R}^{n-k})) = \phi(U) \cap \Omega \text{ and } \phi(U \cap \partial(\mathbb{R}_+^k \times \mathbb{R}^{n-k})) = \phi(U) \cap \partial\Omega,$$

with the additional hypothesis that the differential at 0, namely $d\phi_0$, is an isometry.

In the sequel, we write $\mathbb{R}^{k+,n-k} := \mathbb{R}_+^k \times \mathbb{R}^{n-k}$. We have that (see [4])

$$\gamma_H(\mathbb{R}^{k+,n-k}) = \frac{(n-2+2k)^2}{4}.$$

We have proved the following:

Theorem 1.1 (Cheikh-Ali [4]). *Let Ω be a bounded domain in \mathbb{R}^n , $n \geq 3$, such that $0 \in \partial\Omega$ is a singularity of type $(k, n-k)$ for some $k \in \{1, \dots, n\}$. We fix $0 \leq s < 2$ and $0 \leq \gamma < \gamma_H(\Omega)$. In addition, suppose that either $\{s > 0\}$ or $\{s = 0, n \geq 4 \text{ and } \gamma > 0\}$. We assume that*

$$(3) \quad \gamma \leq \gamma_H(\mathbb{R}^{k+,n-k}) - \frac{1}{4} \text{ that is } n \geq n_{\gamma,k} := \sqrt{4\gamma+1} + 2 - 2k.$$

Then there are extremals for $\mu_{\gamma,s,0}(\Omega)$ if

$$GH_{\gamma,s}(\Omega) < 0$$

where $GH_{\gamma,s}(\Omega)$ is the **generalized mean curvature** defined below in (7).

The assumption

$$(4) \quad \{s > 0\} \text{ or } \{s = 0, n \geq 4 \text{ and } \gamma > 0\}$$

will be reminiscent in the statements below. Its sole utility is to ensure the existence of extremals for $\mu_{\gamma,s,0}(\mathbb{R}^{k+,n-k})$ (see Ghoussoub-Robert [8]).

This result is for large dimension $n \geq n_{\gamma,k}$ (see (3)). In the present article, we tackle the case of the remaining small dimensions. The argument based on local geometry performed for the proof of Theorem 1.1 is not working here. Here, the global geometry has an impact: in order to obtain extremals, we must introduce a "mass" in the spirit of Schoen [14] and Schoen-Yau [15]. Concerning low dimension phenomena, we refer to the pioneer work of Brezis-Nirenberg [2], Jannelli [13] and the more recent reference Ghoussoub-Robert [7] for further discussions. Our main theorem is the following:

Theorem 1.2. *Let Ω be a bounded domain in \mathbb{R}^n , $n \geq 3$, such that $0 \in \partial\Omega$ is a singularity of type $(k, n-k)$ for some $k \in \{1, \dots, n\}$. We fix $0 \leq s < 2$, $\gamma < \gamma_H(\Omega)$ and $a \in C^{0,\theta}(\Omega)$ ($\theta \in (0, 1)$). We assume that condition (4) holds and that*

$$\gamma > \gamma_H(\mathbb{R}^{k+,n-k}) - \frac{1}{4} \text{ that is } n < n_{\gamma,k}.$$

We assume that the operator $-\Delta - (\gamma|x|^{-2} + a(x))$ is coercive and has a mass $m_{\gamma,a}(\Omega)$ (see Definition 3), and that $m_{\gamma,a}(\Omega) > 0$. Then there are extremals for $\mu_{\gamma,s,a}(\Omega)$. In particular, there exists $u \in C^{2,\theta}(\Omega) \cap D^{1,2}(\Omega)$ such that

$$(5) \quad \begin{cases} -\Delta u - \left(\frac{\gamma}{|x|^2} + a(x)\right) u = \frac{u^{2^*(s)-1}}{|x|^s} & \text{in } \Omega, \\ u > 0 & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega. \end{cases}$$

In the second part of this paper, we consider the perturbative Hardy-Schrödinger equation. Given $a, h \in C^{0,\theta}(\Omega)$ for some $\theta \in (0, 1)$ and $q \in (1, 2^* - 1)$ where $2^* = 2^*(0)$, we investigate the existence of solutions $u \in C^2(\Omega) \cap D^{1,2}(\Omega)$ to

$$(6) \quad \begin{cases} -\Delta u - \left(\frac{\gamma}{|x|^2} + a(x)\right) u = \frac{u^{2^*(s)-1}}{|x|^s} + h(x)u^q & \text{in } \Omega, \\ u > 0 & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega. \end{cases}$$

We refer to Brezis-Nirenberg [2] ($\gamma = 0$ and $s = 0$ on a smooth domain Ω), Ghoussoub-Yuan [10] ($\gamma = 0$, $s > 0$ and $0 \in \Omega$), Ghoussoub-Kang [9] and Jaber [12] ($\gamma = 0$, $s > 0$ and $0 \in \partial\Omega$). In the Riemannian context with no boundary, still for $\gamma = 0$, we refer to Djadli [5] when $s = 0$, and to Jaber [11] for $s > 0$ and $h \equiv 0$.

The case $a, h \equiv 0$ was tackled in [4] for $n \geq n_{\gamma,k}$ for nonsmooth domains. We prove the following:

Theorem 1.3. *Let Ω be a bounded domain in \mathbb{R}^n , $n \geq 3$, such that $0 \in \partial\Omega$ is a singularity of type $(k, n-k)$ for some $k \in \{1, \dots, n\}$. Let $a, h \in C^{0,\theta}(\Omega)$ ($\theta \in (0, 1)$) be such that $-\Delta - (\gamma|x|^{-2} + a)$ is coercive and $h \geq 0$. Consider $s \in [0, 2)$ and $\gamma < \gamma_H(\mathbb{R}^{k+,n-k})$. We assume that condition (4) holds and we fix $q \in (1, 2^* - 1)$, $2^* = 2^*(0)$. Then, there exists a positive Mountain-Pass solution $u \in D^{1,2}(\Omega)$ to the perturbative Hardy-Schrödinger equation (6) under one of the following conditions:*

- $0 \leq \gamma < \gamma_H(\mathbb{R}^{k+,n-k}) - \frac{1}{4}$, and

$$\begin{cases} GH_{\gamma,s}(\Omega) < 0 & \text{if } q+1 < \frac{2n-2}{n-2}, \\ c_1 GH_{\gamma,s}(\Omega) - c_2 h(0) < 0 & \text{if } q+1 = \frac{2n-2}{n-2}, \\ h(0) > 0 & \text{if } q+1 > \frac{2n-2}{n-2}, \end{cases}$$
- $0 \leq \gamma = \gamma_H(\mathbb{R}^{k+,n-k}) - \frac{1}{4}$, and

$$\begin{cases} GH_{\gamma,s}(\Omega) < 0 & \text{if } q+1 \leq \frac{2n-2}{n-2}, \\ h(0) > 0 & \text{if } q+1 > \frac{2n-2}{n-2}, \end{cases}$$
- $\gamma > \gamma_H(\mathbb{R}^{k+,n-k}) - \frac{1}{4}$, and

$$\begin{cases} m_{\gamma,a}(\Omega) > 0 & \text{if } q+1 < \frac{2n-2(\alpha_+-\alpha_-)}{n-2}, \\ c_3 m_{\gamma,a}(\Omega) + c_2 h(0) > 0 & \text{if } q+1 = \frac{2n-2(\alpha_+-\alpha_-)}{n-2}, \\ h(0) > 0 & \text{if } q+1 > \frac{2n-2(\alpha_+-\alpha_-)}{n-2}, \end{cases}$$

where $GH_{\gamma,s}(\Omega)$ is the generalized curvature (see (7)), $m_{\gamma,a}(\Omega)$ is the mass of Ω at 0 (see Definition 3), $\alpha_+ - \alpha_- = 2\sqrt{\gamma_H(\mathbb{R}^{k+,n-k}) - \gamma}$ (see (8) below) and $c_1, c_2, c_3 > 0$ are defined in (70).

This result shows how the subcritical nonlinearity has an impact on the existence of solutions. When the subcritical nonlinearity is close to being linear, only the geometry of Ω commands the existence. Conversely, when it is close to being critical, the subcritical nonlinearity commands the existence, whatever the geometry is.

Notation: In the sequel, C denotes a positive constant, the value of which may change from one page to another and even from one line to the next.

This paper is organized as follows. Section 2 is devoted to the definitions of the generalized curvature and the singular interior mass. In Section 3, we introduce some preliminary results that will be of use in the sequel. In Section 4, we prove Theorem 1.2, which is a existence of extremals for $\mu_{\gamma,s,a}(\Omega)$ is ensured for small dimensions when the mass $m_{\gamma,a}(\Omega)$ is positive. In Section 6, we prove Theorem 1.3, which is a general existence result for a Mountain-Pass solution for equation (6). In Section 7, we make test-function estimates in order to obtain a sufficient condition of existence for (6).

2. DEFINITION OF THE GENERALIZED CURVATURE AND THE MASS

Generalized curvature.

Definition 2. Let Ω be a domain in \mathbb{R}^n with $n \geq 3$ such that $0 \in \partial\Omega$ is a singularity of type $(k, n-k)$. We define

$$\Omega_i := \phi(U \cap \{x_i > 0\}) \text{ for all } i = 1, \dots, k,$$

where (ϕ, U) is a chart as in Definition 1. We have that:

- For all $i = 1, \dots, k$, Ω_i is smooth around $0 \in \partial\Omega_i$.
- Up to permutation, the Ω_i 's are locally independent of the chart ϕ .
- The Ω_i 's define locally Ω : there exists $\delta > 0$ such that

$$\Omega \cap B_\delta(0) = \bigcap_{i=1}^k \Omega_i \cap B_\delta(0).$$

We set $\Sigma := \cap_{i=1}^k \partial\Omega_i$ where $k \in \{1, \dots, n\}$. The vector \vec{H}_0^Σ denotes the mean-curvature vector at 0 of the $(n-k)$ -submanifold Σ . For any $m = 1, \dots, k$, $II_0^{\partial\Omega_m}$ denotes the second fundamental form at 0 of the oriented $(n-1)$ -submanifold $\partial\Omega_m$. The **generalized mean curvature** of Ω is defined by:

$$(7) \quad GH_{\gamma,s}(\Omega) := c_{\gamma,s}^1 \sum_{m=1}^k \langle \vec{H}_0^\Sigma, \vec{\nu}_m \rangle + c_{\gamma,s}^2 \sum_{i,m=1, i \neq m}^k II_0^{\partial\Omega_m}(\vec{\nu}_i, \vec{\nu}_i) \\ + c_{\gamma,s}^3 \sum_{p,q,m=1, |\{p,q,m\}|=3}^k II_0^{\partial\Omega_m}(\vec{\nu}_p, \vec{\nu}_q)$$

where for any $m = 1, \dots, k$, $\vec{\nu}_m$ is the outward normal vector at 0 of $\partial\Omega_m$ and $c_{\gamma,s}^1, c_{\gamma,s}^2, c_{\gamma,s}^3$ are positive explicit constants. We refer to [4] for details on this curvature.

The mass. Let $\alpha \in \mathbb{R}$ be a real number and fix $\gamma < \gamma_H(\mathbb{R}^{k+,n-k})$. Then

$$\left(-\Delta - \frac{\gamma}{|x|^2}\right) S_\alpha = 0 \Leftrightarrow \alpha \in \{\alpha_-, \alpha_+\},$$

where:

$$(8) \quad S_\alpha := |x|^{-\alpha-k} \prod_{i=1}^k x_i \text{ and } \alpha_\pm = \alpha_\pm(\gamma, n, k) := \frac{n-2}{2} \pm \sqrt{\gamma_H(\mathbb{R}^{k+,n-k}) - \gamma}.$$

The functions $S_{\alpha_-}, S_{\alpha_+}$ are prototypes of solution to (5) vanishing on $\partial\mathbb{R}^{k+,n-k}$.

Definition 3. Let Ω be a bounded domain in \mathbb{R}^n , $n \geq 3$, such that $0 \in \partial\Omega$ is a singularity of type $(k, n-k)$ for some $k \in \{1, \dots, n\}$. We fix $\gamma < \gamma_H(\Omega)$ and $a \in C^{0,\theta}(\Omega)$ ($\theta \in (0, 1)$). We say that a coercive operator $-\Delta - (\gamma|x|^{-2} + a)$ has a mass if there exists $G \in C^2(\Omega) \cap D_{loc,0}^{1,2}(\Omega)$ such that

$$(9) \quad \begin{cases} -\Delta G - \left(\frac{\gamma}{|x|^2} + a(x)\right) G = 0 & \text{in } \Omega, \\ G > 0 & \text{in } \Omega, \\ G = 0 & \text{on } \partial\Omega \setminus \{0\}, \end{cases}$$

and there exists $c \in \mathbb{R}$ such that

$$(10) \quad G(x) = \prod_{i=1}^k d(x, \partial\Omega_i) (|x|^{-\alpha_+ - k} + c|x|^{-\alpha_- - k} + o(|x|^{-\alpha_- - k})) \text{ as } x \rightarrow 0,$$

where α_\pm is defined in (8). We can therefore define the quantity $m_{\gamma,a}(\Omega) := c$ as the boundary mass of the operator $-\Delta - (\gamma|x|^{-2} + a)$. The function G is unique, so that the definition of the mass makes sense.

Examples of domains with positive or negative mass are in Section 5 below.

3. SOME BACKGROUND RESULTS

We start with the following result that is reminiscent for critical elliptic problems:

Theorem 3.1. [see Cheikh-Ali [4]] Let Ω be a bounded domain in \mathbb{R}^n , $n \geq 3$, such that $0 \in \partial\Omega$ is a singularity of type $(k, n-k)$ for some $k \in \{1, \dots, n\}$. Assume that $\gamma < \gamma_H(\mathbb{R}^{k+,n-k})$, $0 \leq s \leq 2$, and $\mu_{\gamma,s,a}(\Omega) < \mu_{\gamma,s,0}(\mathbb{R}^{k+,n-k})$. Then there are extremals for $\mu_{\gamma,s,a}(\Omega)$.

Indeed, Theorem 3.1 was proved in [4] when $a \equiv 0$. The proof extends to the general case with no effort. Recall now an optimal regularity theorem.

Theorem 3.2. [See Felli-Ferrero [6] and [4]] Let Ω be a bounded domain in \mathbb{R}^n , $n \geq 3$, such that $0 \in \partial\Omega$ is a singularity of type $(k, n-k)$ for some $k \in \{1, \dots, n\}$. We fix $\gamma < \gamma_H(\mathbb{R}^{k+, n-k})$. Let $f : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$ be a Caratheodory function such that

$$(11) \quad |f(x, v)| \leq C|v| \left(1 + \frac{|v|^{2^*(s)-2}}{|x|^s} \right) \text{ for all } x \in \Omega, v \in \mathbb{R}.$$

Let $u \in D^{1,2}(\Omega)_{loc,0}$, be a weak solution to

$$-\Delta u - \frac{\gamma + O(|x|^\tau)}{|x|^2} u = f(x, u) \text{ in } D^{1,2}(\Omega)_{loc,0}$$

for some $\tau > 0$. Then there exists $K \in \mathbb{R}$ such that

$$\lambda^{\alpha_-} u(\lambda\phi(x)) \rightarrow K|x|^{-\alpha_-} \frac{\prod_{i=1}^k x_i}{|x|^k} \text{ in } B_1(0) \cap \mathbb{R}^{k+, n-k},$$

uniformly in C^1 as $\lambda \rightarrow 0$, where ϕ is a chart as in Definition 1.

In section 4, we will need the following lemma:

Lemma 3.1. [See [4]] Assume that $u \in D^{1,2}(\mathbb{R}^{k+, n-k})_{loc,0}$ is a weak solution of

$$\begin{cases} -\Delta u - \frac{\gamma + O(|x|^\tau)}{|x|^2} u = 0 & \text{in } D^{1,2}(\mathbb{R}^{k+, n-k})_{loc,0}, \\ u = 0 & \text{on } B_{2\delta}(0) \cap \partial\mathbb{R}^{k+, n-k}, \end{cases}$$

for some $\tau > 0$ and $\alpha \in \{\alpha_-, \alpha_+\}$. Assume there exists $c > 0$ such that

$$|u(x)| \leq c|x|^{-\alpha} \text{ for } x \rightarrow 0, x \in \mathbb{R}^{k+, n-k}.$$

- Then, there exists $c_1 > 0$ such that

$$|\nabla u(x)| \leq c_1|x|^{-\alpha-1} \text{ as } x \rightarrow 0, x \in \mathbb{R}^{k+, n-k}.$$

- If $\lim_{x \rightarrow 0} |x|^\alpha u(x) = 0$, then $\lim_{x \rightarrow 0} |x|^{\alpha+1} |\nabla u(x)| = 0$.

4. TEST-FUNCTIONS ESTIMATES FOR THE MASS: PROOF OF THEOREM 1.2

Let $U \in D^{1,2}(\mathbb{R}^{k+, n-k})$ be a positive extremal for $\mu_{\gamma, s, 0}(\mathbb{R}^{k+, n-k})$. Then

$$J_{\gamma, s, 0}^{\mathbb{R}^{k+, n-k}}(U) = \frac{\int_{\mathbb{R}^{k+, n-k}} (|\nabla U|^2 - \gamma|x|^{-2}U^2) dx}{\left(\int_{\mathbb{R}^{k+, n-k}} |x|^{-s}|U|^{2^*(s)} dx \right)^{\frac{2}{2^*(s)}}} = \mu_{\gamma, s, 0}(\mathbb{R}^{k+, n-k}).$$

Therefore, there exists $\xi > 0$ such that

$$(12) \quad \begin{cases} -\Delta U - \gamma|x|^{-2}U = \xi|x|^{-s}U^{2^*(s)-1} & \text{in } \mathbb{R}^{k+, n-k}, \\ U > 0 & \text{in } \mathbb{R}^{k+, n-k}, \\ U = 0 & \text{on } \partial\mathbb{R}^{k+, n-k}. \end{cases}$$

For $r > 0$, we define

$$(13) \quad B_r := B_r(0) \text{ and } B_{r,+} := B_r(0) \cap \mathbb{R}^{k+, n-k}.$$

Therefore, with $\delta > 0$ small, the chart ϕ of Definition 1 yields

$$\phi(B_{3\delta} \cap \mathbb{R}^{k+, n-k}) = \phi(B_{3\delta}) \cap \Omega \text{ and } \phi(B_{3\delta} \cap \partial\mathbb{R}^{k+, n-k}) = \phi(B_{3\delta}) \cap \partial\Omega.$$

We fix $\eta \in C_c^\infty(\mathbb{R}^n)$ such that

$$(14) \quad \eta(x) = \begin{cases} 1 & \text{for } x \in B_\delta, \\ 0 & \text{for } x \notin B_{2\delta}. \end{cases}$$

Define also for convenience,

$$(15) \quad p(x) := \prod_{i=1}^k d(x, \partial\Omega_i) \text{ for all } x \in \Omega \text{ and } v(x) := \prod_{i=1}^k x_i \text{ for all } x \in \mathbb{R}^{k+, n-k}.$$

Equation (10) allows us to define $\Theta : \Omega \rightarrow \mathbb{R}$ such that

$$G(x) = (\eta v |x|^{-\alpha_+ - k}) \circ \phi^{-1}(x) + \Theta(x) \text{ for any } x \in \Omega,$$

where ϕ is as in Definition 1. We then get that $\Theta \in D^{1,2}(\Omega)$ and

$$(16) \quad \Theta(x) = m_{\gamma, a}(\Omega) p(x) |x|^{-\alpha_- - k} + o(p(x) |x|^{-\alpha_- - k}) \text{ as } x \rightarrow 0.$$

Note that

$$(17) \quad \left\{ \gamma > \gamma_H(\mathbb{R}_+^k \times \mathbb{R}^{n-k}) - \frac{1}{4} \right\} \Leftrightarrow \{ \alpha_+ - \alpha_- < 1 \} \Leftrightarrow \{ n < n_{\gamma, k} \}.$$

Since U satisfies (12), Theorem 3.2 yields $K_1 > 0$ such that

$$(18) \quad \lim_{\lambda \rightarrow 0^+} \lambda^{\alpha_-} U(\lambda x) = K_1 v(x) |x|^{-\alpha_- - k} \text{ in } B_1(0) \cap \mathbb{R}^{k+, n-k}.$$

The regularity applied to the Kelvin transform $x \mapsto \bar{U}(x) := |x|^{2-n} U(\frac{x}{|x|^2})$ yields

$$(19) \quad \lim_{\lambda \rightarrow +\infty} \lambda^{\alpha_+} U(\lambda x) = K_2 v(x) |x|^{-\alpha_+ - k} \text{ in } B_1(0) \cap \mathbb{R}^{k+, n-k},$$

for some $K_2 > 0$. Up to multiplying U by a positive constant, we assume that $K_2 = 1$. Equation (18), the Kelvin transform and Lemma 3.1 yield

$$(20) \quad |U(x)| \leq C |x|^{-\alpha_+} \text{ and } |\nabla U(x)| \leq C |x|^{-1-\alpha_+} \text{ for any } x \in \mathbb{R}^{k+, n-k}.$$

For $\epsilon > 0$, we define

$$(21) \quad U_\epsilon(x) := \epsilon^{-\frac{n-2}{2}} U(\epsilon^{-1}x) \text{ for all } x \in \mathbb{R}^{k+, n-k}$$

and

$$(22) \quad u_\epsilon(x) := (\eta U_\epsilon) \circ \phi^{-1}(x) \text{ for } x \in \Omega \text{ and } \tilde{u}_\epsilon := u_\epsilon + \epsilon^{\frac{\alpha_+ - \alpha_-}{2}} \Theta.$$

The main result of this paper is the following:

Proposition 4.1. *Let Ω be a bounded domain in \mathbb{R}^n , $n \geq 3$ such that $0 \in \partial\Omega$ is a singularity of type $(k, n-k)$ for some $k \in \{1, \dots, n\}$. We fix $0 \leq s < 2$, $\gamma < \gamma_H(\Omega)$ and $a \in C^{0, \theta}(\Omega)$ ($\theta \in (0, 1)$). Assume that there exists a positive extremal U for $\mu_{\gamma, s, 0}(\mathbb{R}^{k+, n-k})$. We assume that*

$$\gamma > \gamma_H(\mathbb{R}^{k+, n-k}) - \frac{1}{4} \text{ that is } n < n_{\gamma, k},$$

and that the operator $-\Delta - (\gamma|x|^{-2} + a(x))$ is coercive with the mass $m_{\gamma, a}(\Omega)$. We let $(\tilde{u}_\epsilon)_\epsilon \in D^{1,2}(\Omega)$ be as in (22). Then

$$J_{\gamma, s, a}^\Omega(\tilde{u}_\epsilon) = \mu_{\gamma, s, 0}(\mathbb{R}^{k+, n-k}) \left(1 - \zeta_{\gamma, s}^0 m_{\gamma, a}(\Omega) \epsilon^{\alpha_+ - \alpha_-} + o(\epsilon^{\alpha_+ - \alpha_-}) \right) \text{ as } \epsilon \rightarrow 0,$$

where

$$(23) \quad \zeta_{\gamma, s}^0 := (\alpha_+ - \alpha_-) C_{k, n} \left(\xi \int_{\mathbb{R}^{k+, n-k}} \frac{U^{2^*(s)}}{|x|^s} dx \right)^{-1} > 0,$$

where $C_{k,n}$ is defined in (25).

Remark: as noted in the introduction, the existence of extremals for $\mu_{\gamma,s,0}(\mathbb{R}^{k+,n-k})$ is a consequence of (4).

As one checks, Theorem 1.2 is a direct consequence of the combination of Proposition 4.1 and Theorem 3.1.

This section is devoted to the proof of Proposition 4.1.

Proof of Proposition 4.1: From the definitions of \tilde{u}_ϵ and G , and the uniform C^1 -convergence in (19), it follows that

$$(24) \quad \lim_{\epsilon \rightarrow 0} \frac{\tilde{u}_\epsilon}{\epsilon^{\frac{\alpha_+ - \alpha_-}{2}}} = G \text{ in } C_{loc}^1(\Omega) \cap D_{loc,0}^{1,2}(\Omega).$$

Define the constant

$$(25) \quad C_{k,n} := \int_{S^{n-1} \cap \mathbb{R}^{k+,n-k}} \left(\prod_{i=1}^k x_i \right)^2 d\sigma.$$

In the sequel, ϑ_ρ^ϵ will denote a quantity such

$$\lim_{\rho \rightarrow 0} \lim_{\epsilon \rightarrow 0} \vartheta_\rho^\epsilon = 0.$$

For convenience, we define

$$N_{\gamma,a}(w) := |\nabla w|^2 - (\gamma|x|^{-2} + a)w^2.$$

Step 4.1. For any $\rho > 0$, we claim that

$$\int_{\Omega \setminus \phi(B_{\rho,+})} N_{\gamma,a}(\tilde{u}_\epsilon) dx = \epsilon^{\alpha_+ - \alpha_-} (\alpha_+ C_{k,n} \rho^{n-2\alpha_+ - 2} + m_{\gamma,a}(\Omega)(n-2)C_{k,n} + \vartheta_\rho^\epsilon),$$

as $\epsilon \rightarrow 0$ where the constant $C_{k,n}$ is defined in (25).

Proof of Step 4.1: Indeed, it follows from (24) that

$$\lim_{\epsilon \rightarrow 0} \epsilon^{-(\alpha_+ - \alpha_-)} \int_{\Omega \setminus \phi(B_{\rho,+})} N_{\gamma,a}(\tilde{u}_\epsilon) dx = \int_{\Omega \setminus \phi(B_{\rho,+})} N_{\gamma,a}(G) dx.$$

Since G satisfies (9) and vanishes on $\partial\Omega \setminus \{0\}$, integrating by parts yields

$$(26) \quad \begin{aligned} \int_{\Omega \setminus \phi(B_{\rho,+})} N_{\gamma,a}(G) dx &= \int_{\Omega \setminus \phi(B_{\rho,+})} G(-\Delta G - (\gamma|x|^{-2} + a(x))G) dx \\ &\quad - \int_{\phi(\partial(B_{\rho,+}))} G \partial_\nu G d\sigma \\ &= - \int_{(\partial B_\rho(0)) \cap \mathbb{R}^{k+,n-k}} (G \circ \phi) \partial_{\phi_* \nu} (G \circ \phi) d(\phi^* \sigma), \end{aligned}$$

where $\nu(x)$ is the outer normal vector of $B_\rho(0)$ at $x \in \partial B_\rho(0)$. We will now find the value of $(G \circ \phi) \partial_{\phi_* \nu} (G \circ \phi)$. Using (15) and the definition of G , we have that

$$(27) \quad (G \circ \phi)(x) = v(x)|x|^{-\alpha_+ - k} + m_{\gamma,a}(\Omega)v(x)|x|^{-\alpha_- - k} + o(v(x)|x|^{-\alpha_- - k}) \text{ as } x \rightarrow 0.$$

From Θ and the uniform convergence in C^1 of G , we have for all $l = 1, \dots, n$ that

$$(28) \quad \partial_l(\Theta \circ \phi) = \partial_l(m_{\gamma,a}(\Omega)v|x|^{-\alpha_- - k}) + o(|x|^{-\alpha_- - 1}) \text{ as } x \rightarrow 0.$$

Moreover, it follows from the definition of G that

$$\begin{aligned}\partial_l(G \circ \phi) &= \partial_l v (|x|^{-\alpha_+ - k} + m_{\gamma,a}(\Omega)|x|^{-\alpha_- - k}) \\ &\quad - x_l v ((\alpha_+ + k)|x|^{-\alpha_+ - k - 2} + (\alpha_- + k)m_{\gamma,a}(\Omega)|x|^{-\alpha_- - k - 2}) + o(|x|^{-\alpha_- - 1}).\end{aligned}$$

Since

$$\phi_* \nu(x) = \frac{x}{|x|} + O(|x|) \text{ as } x \rightarrow 0 \text{ and } \alpha_+ < \alpha_- + 1,$$

we obtain as $x \rightarrow 0$ that,

$$(29) \quad \partial_{\phi_* \nu}(G \circ \phi) = -v (\alpha_+ |x|^{-\alpha_+ - k - 1} + m_{\gamma,a}(\Omega)\alpha_- |x|^{-\alpha_- - k - 1}) + o(|x|^{-\alpha_- - 1}).$$

We combine the equations (27), (29) and since $\alpha_+ + \alpha_- = n - 2$, $-2\alpha_- - 1 > 1 - n$, $\alpha_+ - \alpha_- < 1$, we get

$$-(G \circ \phi)\partial_{\phi_* \nu}(G \circ \phi) = v^2 (\alpha_+ |x|^{-2\alpha_+ - 2k - 1} + m_{\gamma,a}(\Omega)(n - 2)|x|^{-n+1-2k}) + o(|x|^{1-n}).$$

Moreover, using again the definition of v ,

$$-\int_{\partial B_{\rho,+}} (G \circ \phi)\partial_{\phi_* \nu}(G \circ \phi)d(\phi^* \sigma) = \alpha_+ C_{k,n} \rho^{n-2\alpha_+ - 2} + m_{\gamma,a}(\Omega)(n - 2)C_{k,n} + \vartheta_\rho,$$

where $\lim_{\rho \rightarrow 0} \vartheta_\rho = 0$ and $C_{k,n}$ is defined in (25). Plugging the last equation in (26) yields Step 4.1. \square

Step 4.2. We claim that, as $\epsilon \rightarrow 0$,

$$\int_{\Omega} N_{\gamma,a}(\tilde{u}_\epsilon) dx = \xi \int_{\mathbb{R}^{k_+, n-k}} |x|^{-s} U^{2^*(s)} dx + m_{\gamma,a}(\Omega)(n-2)C_{k,n} \epsilon^{\alpha_+ - \alpha_-} + o(\epsilon^{\alpha_+ - \alpha_-}).$$

Proof of Step 4.2: The definition (22) of \tilde{u}_ϵ rewrites

$$(30) \quad \tilde{u}_\epsilon \circ \phi(x) = U_\epsilon(x) + \epsilon^{\frac{\alpha_+ - \alpha_-}{2}} \Theta \circ \phi(x) \text{ for all } x \in B_{\delta,+}.$$

Fix $\rho \in]0, \delta[$ that we will eventually let go to 0. We define

$$I_{\epsilon,\rho} := \int_{\phi(B_{\rho,+})} (|\nabla \tilde{u}_\epsilon|^2 - (\gamma|x|^{-2} + a) \tilde{u}_\epsilon^2) dx.$$

Let $\phi^* Eucl$ be the pullback of the Euclidean metric. With (30), we get

$$\begin{aligned}I_{\epsilon,\rho} &= \int_{B_{\rho,+}} \left(|\nabla(\tilde{u}_\epsilon \circ \phi)|_{\phi^* Eucl}^2 - \left(\frac{\gamma}{|\phi(x)|^2} + a \circ \phi \right) (\tilde{u}_\epsilon \circ \phi)^2 \right) |Jac(\phi)| dx \\ &= \int_{B_{\rho,+}} \left(|\nabla U_\epsilon|_{\phi^* Eucl}^2 - \left(\frac{\gamma}{|\phi(x)|^2} + a \circ \phi \right) U_\epsilon^2 \right) |Jac(\phi)| dx \\ &\quad + 2\epsilon^{\frac{\alpha_+ - \alpha_-}{2}} \int_{B_{\rho,+}} \left(\langle \nabla U_\epsilon, \nabla(\Theta \circ \phi) \rangle_{\phi^* Eucl} - \left(\frac{\gamma}{|\phi(x)|^2} + a \circ \phi \right) (\Theta \circ \phi) U_\epsilon \right) |Jac(\phi)| dx \\ &\quad + \epsilon^{\alpha_+ - \alpha_-} \int_{B_{\rho,+}} \left(|\nabla(\Theta \circ \phi)|_{\phi^* Eucl}^2 - \left(\frac{\gamma}{|\phi(x)|^2} + a \circ \phi \right) (\Theta \circ \phi)^2 \right) |Jac(\phi)| dx.\end{aligned}$$

Since $d\phi_0 = Id_{\mathbb{R}^n}$, $\phi^* Eucl = Eucl + O(|x|)$ and $\Theta \in D^{1,2}(\Omega)$, we get that

$$\begin{aligned}
I_{\epsilon,\rho} &= \int_{B_{\rho,+}} \left(|\nabla U_\epsilon|_{Eucl}^2 - \left(\frac{\gamma}{|x|^2} + a \circ \phi \right) U_\epsilon^2 \right) dx \\
&+ O \left(\int_{B_{\rho,+}} |x| (|\nabla U_\epsilon|_{Eucl}^2 + |x|^{-2} U_\epsilon^2) dx \right) \\
&+ 2\epsilon^{\frac{\alpha_+ - \alpha_-}{2}} \int_{B_{\rho,+}} \left(\langle \nabla U_\epsilon, \nabla(\Theta \circ \phi) \rangle_{Eucl} - \left(\frac{\gamma}{|x|^2} + a \circ \phi \right) (\Theta \circ \phi) U_\epsilon \right) dx \\
&+ O \left(\epsilon^{\frac{\alpha_+ - \alpha_-}{2}} \int_{B_{\rho,+}} |x| (|\nabla U_\epsilon| \cdot |\nabla(\Theta \circ \phi)| + |x|^{-2} (\Theta \circ \phi) U_\epsilon) dx \right) + \epsilon^{\alpha_+ - \alpha_-} \vartheta_\rho^\epsilon
\end{aligned}$$

as $\epsilon \rightarrow 0$. The explicit expression (21) of U_ϵ , (20) and $n > 2\alpha_+$ yield

$$\begin{aligned}
\int_{B_{\rho,+}} U_\epsilon^2 dx &= O \left(\epsilon^{\alpha_+ - \alpha_-} \int_0^\rho r^{n-2\alpha_+ - 1} dr \right) \\
(31) \qquad \qquad &= \epsilon^{\alpha_+ - \alpha_-} \vartheta_\rho^\epsilon.
\end{aligned}$$

The definition of Θ and $\alpha_+ + \alpha_- = n - 2$ give

$$\begin{aligned}
\epsilon^{\frac{\alpha_+ - \alpha_-}{2}} \int_{B_{\rho,+}} a \circ \phi (\Theta \circ \phi) U_\epsilon dx &= O \left(\epsilon^{\alpha_+ - \alpha_-} \int_0^\rho r dr \right) \\
(32) \qquad \qquad \qquad &= \epsilon^{\alpha_+ - \alpha_-} \vartheta_\rho^\epsilon.
\end{aligned}$$

We combine the equations (18), (20), (28), (31) and (32),

$$\begin{aligned}
I_{\epsilon,\rho} &= \int_{B_{\rho,+}} \left(|\nabla U_\epsilon|_{Eucl}^2 - \frac{\gamma}{|x|^2} U_\epsilon^2 \right) dx \\
&+ 2\epsilon^{\frac{\alpha_+ - \alpha_-}{2}} \int_{B_{\rho,+}} \left(\langle \nabla U_\epsilon, \nabla(\Theta \circ \phi) \rangle_{Eucl} - \frac{\gamma}{|x|^2} (\Theta \circ \phi) U_\epsilon \right) dx + \epsilon^{\alpha_+ - \alpha_-} \vartheta_\rho^\epsilon \text{ as } \epsilon \rightarrow 0.
\end{aligned}$$

Integrating by parts and since both U_ϵ and $\Theta \circ \phi$ vanish on $\partial\mathbb{R}^{k_+,n-k_-} \setminus \{0\}$, we get as $\epsilon \rightarrow 0$ that

$$\begin{aligned}
I_{\epsilon,\rho} &= \int_{B_{\rho,+}} U_\epsilon (-\Delta U_\epsilon - \gamma|x|^{-2} U_\epsilon) dx + \int_{\mathbb{R}^{k_+,n-k_-} \cap \partial B_\rho(0)} U_\epsilon \partial_\nu U_\epsilon d\sigma \\
(33) \qquad &+ 2\epsilon^{\frac{\alpha_+ - \alpha_-}{2}} \left(\int_{B_{\rho,+}} (\Theta \circ \phi) (-\Delta U_\epsilon - \gamma|x|^{-2} U_\epsilon) dx \right. \\
&\left. + \int_{\mathbb{R}^{k_+,n-k_-} \cap \partial B_\rho(0)} (\Theta \circ \phi) \partial_\nu U_\epsilon d\sigma \right) + \epsilon^{\alpha_+ - \alpha_-} \vartheta_\rho^\epsilon.
\end{aligned}$$

We claim as $\epsilon \rightarrow 0$ that

$$(34) \quad \int_{\mathbb{R}^{k_+,n-k_-} \cap \partial B_\rho(0)} (\Theta \circ \phi) \partial_\nu U_\epsilon d\sigma = -\alpha_+ \epsilon^{\frac{\alpha_+ - \alpha_-}{2}} m_{\gamma,a}(\Omega) C_{k,n} + o(\epsilon^{\frac{\alpha_+ - \alpha_-}{2}}),$$

and

$$(35) \quad \int_{\mathbb{R}^{k_+,n-k_-} \cap \partial B_\rho(0)} U_\epsilon \partial_\nu U_\epsilon d\sigma = -\alpha_+ C_{k,n} \epsilon^{\alpha_+ - \alpha_-} \rho^{n-2\alpha_+ - 2} + o(\epsilon^{\alpha_+ - \alpha_-} \rho^{n-2-2\alpha_+}).$$

We prove the claim. It follows from the uniform C^1 -convergence in (19) that we have for all $l = 1, \dots, n$

$$(36) \quad \lim_{\lambda \rightarrow +\infty} \lambda^{\alpha_+} \partial_l U(\lambda x) = |x|^{-\alpha_+ - k} \left(\delta_{l \leq k} \prod_{j=1; j \neq l}^k x_j - (\alpha_+ + k) \frac{v(x)x_l}{|x|^2} \right),$$

where $\delta_{l \leq k}$ is such that $\delta_{l \leq k} = 1$ if $l \leq k$, and $\delta_{l \leq k} = 0$ otherwise, and v is defined in (15). The definition of U_ϵ and (20) yield

$$\partial_l U_\epsilon = \epsilon^{\frac{\alpha_+ - \alpha_-}{2}} \left(|x|^{-\alpha_+ - k} \left(\delta_{l \leq k} \prod_{j=1; j \neq l}^k x_j - (\alpha_+ + k) \frac{x_l}{|x|^2} v \right) + o(|x|^{-\alpha_+ - 1}) \right).$$

Since $\nu(x) = |x|^{-1}x$ is the outer normal vector of $B_\rho(0)$, we then get

$$(37) \quad \partial_\nu U_\epsilon = \epsilon^{\frac{\alpha_+ - \alpha_-}{2}} (-\alpha_+ v |x|^{-\alpha_+ - k - 1} + o(|x|^{-\alpha_+ - 1})),$$

as $\epsilon \rightarrow 0$ uniformly on compact subsets of $\overline{\mathbb{R}^{k_+, n-k}} \setminus \{0\}$. From the definition of Θ and $\alpha_+ + \alpha_- = n - 2$, and (17), we obtain as $\epsilon \rightarrow 0$ that

$$(\Theta \circ \phi) \partial_\nu U_\epsilon = \epsilon^{\frac{\alpha_+ - \alpha_-}{2}} (-\alpha_+ m_{\gamma, a}(\Omega) v^2 |x|^{-n+1-2k} + o(|x|^{1-n})).$$

Therefore, we get (34). The definition of U_ϵ and the equations (19) and (37) yield

$$U_\epsilon \partial_\nu U_\epsilon = -\alpha_+ \epsilon^{\alpha_+ - \alpha_-} v^2 |x|^{-2\alpha_+ - 2k - 1} + o(\epsilon^{\alpha_+ - \alpha_-} |x|^{-2\alpha_+ - 1}),$$

as $\epsilon \rightarrow 0$ uniformly locally in $\overline{\mathbb{R}^{k_+, n-k}} \setminus \{0\}$. This yields (35) and proves the claim.

We combine equations (33), (34) and (35) to get

$$\begin{aligned} I_{\epsilon, \rho} &= \int_{B_{\rho, +}} U_\epsilon (-\Delta U_\epsilon - \gamma |x|^{-2} U_\epsilon) dx - \alpha_+ C_{k, n} \epsilon^{\alpha_+ - \alpha_-} \rho^{n-2\alpha_+ - 2} \\ &+ 2\epsilon^{\frac{\alpha_+ - \alpha_-}{2}} \int_{B_{\rho, +}} (\Theta \circ \phi) (-\Delta U_\epsilon - \gamma |x|^{-2} U_\epsilon) dx \\ &- 2\alpha_+ \epsilon^{\alpha_+ - \alpha_-} m_{\gamma, a}(\Omega) C_{k, n} + \epsilon^{\alpha_+ - \alpha_-} \vartheta_\rho^\epsilon. \end{aligned}$$

Since U satisfies equation (12), with the definition (21) of U_ϵ , we get

$$-\Delta U_\epsilon - \gamma |x|^{-2} U_\epsilon = \xi |x|^{-s} U_\epsilon^{2^*(s)-1}.$$

Therefore, we get as $\epsilon \rightarrow 0$ that

$$(38) \quad \begin{aligned} I_{\epsilon, \rho} &= \xi \int_{B_{\rho, +}} \frac{U_\epsilon^{2^*(s)}}{|x|^s} dx - \alpha_+ C_{k, n} \epsilon^{\alpha_+ - \alpha_-} \rho^{n-2\alpha_+ - 2} \\ &+ 2\epsilon^{\frac{\alpha_+ - \alpha_-}{2}} \xi \int_{B_{\rho, +}} (\Theta \circ \phi) \frac{U_\epsilon^{2^*(s)-1}}{|x|^s} dx \\ &- 2\alpha_+ \epsilon^{\alpha_+ - \alpha_-} m_{\gamma, a}(\Omega) C_{k, n} + \epsilon^{\alpha_+ - \alpha_-} \vartheta_\rho^\epsilon. \end{aligned}$$

It follow from the definition (21) of U_ϵ and the first estimate in (20) that

$$\left| \xi \int_{\overline{\mathbb{R}^{k_+, n-k}} \setminus (B_{\rho, +})} \frac{U_\epsilon^{2^*(s)}}{|x|^s} dx \right| \leq C \epsilon^{\frac{2^*(s)}{2}(\alpha_+ - \alpha_-)}.$$

Therefore, with $2^*(s) > 2$, we get

$$(39) \quad \xi \int_{B_{\rho,+}} \frac{U_\epsilon^{2^*(s)}}{|x|^s} dx = \xi \int_{\mathbb{R}^{k_+,n-k}} \frac{U^{2^*(s)}}{|x|^s} dx + o(\epsilon^{\alpha_+ - \alpha_-}) \text{ as } \epsilon \rightarrow 0.$$

The definition (15), (21) and the first control in (20) yield

$$\begin{aligned} \int_{\mathbb{R}^{k_+,n-k} \setminus (B_{\rho,+})} v \frac{U_\epsilon^{2^*(s)-1}}{|x|^{s+\alpha_-+k}} dx &= O\left(\epsilon^{\frac{\alpha_+ - \alpha_-}{2}} \int_{\epsilon^{-1}\rho}^{+\infty} r^{(1-\frac{2^*(s)}{2})(\alpha_+ - \alpha_-) - 1} dr\right) \\ &= \epsilon^{\frac{2^*(s)-1}{2}(\alpha_+ - \alpha_-)} \vartheta_\rho^\epsilon. \end{aligned}$$

Therefore, with the definition of Θ , we get as $\epsilon \rightarrow 0$ that

$$(40) \quad \begin{aligned} \xi \int_{B_{\rho,+}} \frac{U_\epsilon^{2^*(s)-1}}{|x|^s} \Theta \circ \phi dx &= \xi m_{\gamma,a}(\Omega) \int_{B_{\rho,+}} v \frac{U_\epsilon^{2^*(s)-1}}{|x|^{s+\alpha_-+k}} dx \\ &+ o\left(\int_{B_{\rho,+}} v \frac{U_\epsilon^{2^*(s)-1}}{|x|^{s+\alpha_-+k}} dx\right) \\ &= \epsilon^{\frac{\alpha_+ - \alpha_-}{2}} \left(m_{\gamma,a}(\Omega) \xi \int_{\mathbb{R}^{k_+,n-k}} v \frac{U^{2^*(s)-1}}{|x|^{s+\alpha_-+k}} dx + \vartheta_\epsilon^\rho\right). \end{aligned}$$

Since $(-\Delta - \gamma|x|^{-2})(v|x|^{-\alpha_- - k}) = 0$ and U vanishes on $\partial\mathbb{R}^{k_+,n-k} \setminus \{0\}$, integrating by parts, we get that

$$(41) \quad \begin{aligned} \xi \int_{\mathbb{R}^{k_+,n-k}} v \frac{U^{2^*(s)-1}}{|x|^{s+\alpha_-+k}} dx &= \lim_{R \rightarrow +\infty} \int_{B_{R,+}} v|x|^{-\alpha_- - k} (-\Delta U - \gamma|x|^{-2}U) dx \\ &= \lim_{R \rightarrow +\infty} \left[\int_{B_{R,+}} U (-\Delta - \gamma|x|^{-2})(v|x|^{-\alpha_- - k}) dx \right. \\ &\quad \left. - \int_{\mathbb{R}^{k_+,n-k} \cap \partial B_R} \partial_\nu U v|x|^{-\alpha_- - k} d\sigma \right]. \end{aligned}$$

Arguing as for (37), it follows from (36) that, as $R \rightarrow +\infty$

$$\partial_\nu U = -\alpha_+ v|x|^{-\alpha_+ - k - 1} + o(|x|^{-\alpha_+ - 1}) \text{ uniformly for } x \in \partial B_R(0) \cap \mathbb{R}^{k_+,n-k}.$$

Moreover, since $\alpha_+ + \alpha_- = n - 2$ we get

$$\partial_\nu U v|x|^{-\alpha_- - k} = -\alpha_+ v^2|x|^{-(n+2k-1)} + o(|x|^{1-n}).$$

This latest equation yields

$$\lim_{R \rightarrow +\infty} \int_{\mathbb{R}^{k_+,n-k} \cap \partial B_R(0)} \partial_\nu U v|x|^{-\alpha_- - k} d\sigma = -\alpha_+ C_{k,n}.$$

Then, by (41)

$$(42) \quad \xi \int_{\mathbb{R}^{k_+,n-k}} v \frac{U^{2^*(s)-1}}{|x|^{s+\alpha_-+k}} dx = \alpha_+ C_{k,n}.$$

Combining (40) and (42), we get

$$(43) \quad \xi \int_{B_{\rho,+}} \frac{U_\epsilon^{2^*(s)-1}}{|x|^s} \Theta \circ \phi dx = \epsilon^{\frac{\alpha_+ - \alpha_-}{2}} (\alpha_+ m_{\gamma,a}(\Omega) C_{k,n} + \vartheta_\epsilon^\rho) \text{ as } \epsilon \rightarrow 0.$$

Next, the equations (38), (39) and (43) yield

$$I_{\epsilon, \rho} = \xi \int_{\mathbb{R}^{k_+, n-k}} \frac{U_{\epsilon}^{2^*(s)}}{|x|^s} dx - \alpha_+ C_{k,n} \epsilon^{\alpha_+ - \alpha_-} \rho^{n-2\alpha_+ - 2} + o(\epsilon^{\alpha_+ - \alpha_-}).$$

On the other hand, using Step 4.1 the definition of $I_{\epsilon, \rho}$ and the last equation, we get Step 4.2. \square

Step 4.3. *We claim as $\epsilon \rightarrow 0$ that,*

$$\int_{\Omega} \frac{\tilde{u}_{\epsilon}^{2^*(s)}}{|x|^s} dx = \int_{\mathbb{R}^{k_+, n-k}} \frac{U_{\epsilon}^{2^*(s)}}{|x|^s} dx + 2^*(s) \alpha_+ m_{\gamma, a}(\Omega) \xi^{-1} C_{k,n} \epsilon^{\alpha_+ - \alpha_-} + o(\epsilon^{\alpha_+ - \alpha_-}).$$

Proof of Step 4.3: We fix $\rho > 0$. The definitions of \tilde{u}_{ϵ} and Θ , and $2^*(s) > 2$ yield

$$(44) \quad \int_{\phi(B_{2\delta, +} \setminus B_{\rho, +})} \frac{\tilde{u}_{\epsilon}^{2^*(s)}}{|x|^s} dx = o(\epsilon^{\alpha_+ - \alpha_-}),$$

as $\epsilon \rightarrow 0$. Equations (16), (18), (30) and (44) yield

$$\int_{\Omega} \frac{\tilde{u}_{\epsilon}^{2^*(s)}}{|x|^s} dx = \int_{B_{\rho, +}} \frac{|U_{\epsilon} + \epsilon^{\frac{\alpha_+ - \alpha_-}{2}} (\Theta \circ \phi)|^{2^*(s)}}{|x|^s} |(1 + O(|x|))| dx + o(\epsilon^{\alpha_+ - \alpha_-}),$$

as $\epsilon \rightarrow 0$, and

$$\begin{aligned} \int_{\Omega} \frac{\tilde{u}_{\epsilon}^{2^*(s)}}{|x|^s} dx &= \int_{B_{\rho, +}} \left(\frac{U_{\epsilon}^{2^*(s)}}{|x|^s} + 2^*(s) \epsilon^{\frac{\alpha_+ - \alpha_-}{2}} \frac{U_{\epsilon}^{2^*(s)-1}}{|x|^s} (\Theta \circ \phi) \right) dx \\ &+ \int_{B_{\rho, +}} O \left(\epsilon^{\alpha_+ - \alpha_-} \frac{U_{\epsilon}^{2^*(s)-2}}{|x|^s} (\Theta \circ \phi)^2 + \epsilon^{\frac{2^*(s)}{2}(\alpha_+ - \alpha_-)} |\Theta \circ \phi|^{2^*(s)} \right) dx + o(\epsilon^{\alpha_+ - \alpha_-}). \end{aligned}$$

Using the asymptotics (16) and (18) of Θ and U , we get that

$$(45) \quad \begin{aligned} \int_{B_{\rho, +}} \frac{U_{\epsilon}^{2^*(s)-2}}{|x|^s} (\epsilon^{\frac{\alpha_+ - \alpha_-}{2}} (\Theta \circ \phi))^2 dx &= O \left(\epsilon^{2(\alpha_+ - \alpha_-)} \int_0^{\epsilon^{-1}\rho} r^{\frac{2^*(s)}{2}(\alpha_+ - \alpha_-) - 1} dr \right) \\ &= \epsilon^{\alpha_+ - \alpha_-} \vartheta_{\epsilon}^{\rho}, \end{aligned}$$

and, from the definition of Θ and the control (16), we get that

$$(46) \quad \begin{aligned} \int_{B_{\rho, +}} (\epsilon^{\frac{\alpha_+ - \alpha_-}{2}} \Theta \circ \phi)^{2^*(s)} |x|^{-s} dx &= O \left(\epsilon^{(\alpha_+ - \alpha_-) \frac{2^*(s)}{2}} \int_0^{\rho} r^{\frac{2^*(s)}{2}(\alpha_+ - \alpha_-) - 1} dr \right) \\ &= \epsilon^{\alpha_+ - \alpha_-} \vartheta_{\epsilon}^{\rho}. \end{aligned}$$

The equations (45) and (46) yield as $\epsilon \rightarrow 0$ that

$$(47) \quad \int_{\Omega} \frac{\tilde{u}_{\epsilon}^{2^*(s)}}{|x|^s} dx = \int_{B_{\rho, +}} \left(\frac{U_{\epsilon}^{2^*(s)}}{|x|^s} + 2^*(s) \epsilon^{\frac{\alpha_+ - \alpha_-}{2}} \frac{U_{\epsilon}^{2^*(s)-1}}{|x|^s} (\Theta \circ \phi) \right) dx + \epsilon^{\alpha_+ - \alpha_-} \vartheta_{\epsilon}^{\rho}.$$

Therefore, for all $\xi > 0$ the equations (39), (43) and (47) yield the result. \square

Step 4.4. *We are now in a position to prove Proposition 4.1.*

Proof of Step 4.4: By Step 4.3, we have that

$$\begin{aligned}
\left(\int_{\Omega} \frac{\tilde{u}_{\epsilon}^{2^*(s)}}{|x|^s} dx \right)^{\frac{2}{2^*(s)}} &= \left(\int_{\mathbb{R}^{k_+, n-k}} \frac{U_{\epsilon}^{2^*(s)}}{|x|^s} dx \right)^{\frac{2}{2^*(s)}} \\
(48) \quad &+ 2\alpha_+ m_{\gamma, a}(\Omega) \xi^{-1} C_{k, n} \epsilon^{\alpha_+ - \alpha_-} \left(\int_{\mathbb{R}^{k_+, n-k}} \frac{U_{\epsilon}^{2^*(s)}}{|x|^s} dx \right)^{\frac{2}{2^*(s)} - 1} \\
&+ o(\epsilon^{\alpha_+ - \alpha_-}).
\end{aligned}$$

We go back to the definition of $J_{\gamma, a, s}^{\Omega}$. Step 4.3, Equation (48) and (12) yield

$$J_{\gamma, s, a}^{\Omega}(\tilde{u}_{\epsilon}) = J_{\gamma, s, 0}^{\mathbb{R}^{k_+, n-k}}(U) (1 - m_{\gamma, a}(\Omega) \zeta_{\gamma, s}^0 \epsilon^{\alpha_+ - \alpha_-} + o(\epsilon^{\alpha_+ - \alpha_-})),$$

as $\epsilon \rightarrow 0$, where $\zeta_{\gamma, s}^0$ is defined in (23). This ends the proof of Proposition 4.1. \square

Combining Proposition 4.1 and Theorem 3.1 yields Theorem 1.2 .

5. EXAMPLES OF MASS

In this section, we discuss the existence and the sign of the mass. An example of existence of mass is as follows:

Proposition 5.1. *Let Ω be a bounded domain in \mathbb{R}^n , $n \geq 3$ such that $0 \in \partial\Omega$ is a singularity of type $(k, n-k)$ for some $k \in \{1, \dots, n\}$. We assume that $\gamma > \gamma_H(\mathbb{R}^{k_+, n-k}) - 1/4$ and that*

$$(49) \quad \Omega \cap B_{\delta}(0) = \mathbb{R}^{k_+, n-k} \cap B_{\delta}(0) \text{ for some } \delta > 0.$$

We assume that $\gamma_H(\mathbb{R}^{k_+, n-k}) - \frac{1}{4} < \gamma < \gamma_H(\Omega)$, that $a \in C^{0, \theta}(\Omega)$ vanishes around 0 and that $-\Delta - (\gamma|x|^{-2} + a(x))$ is coercive. Then the mass is defined.

Proof of Proposition 5.1. We fix η as in (14). For $a \in C^{0, \theta}(\Omega)$ that vanishes around 0, define on Ω the function

$$g := \left(-\Delta - \frac{\gamma}{|x|^2} - a(x) \right) (\eta S_{\alpha_+}),$$

where S_{α_+} is defined in (8) such that $-\Delta S_{\alpha_+} - \gamma|x|^{-2} S_{\alpha_+} = 0$ on $\mathbb{R}^{k_+, n-k}$. Note that this definition makes sense when the support of η is small enough due to (49) and a vanishes around 0. In particular $g(x) = 0$ around 0. Therefore, we have that $g \in L^{\frac{2n}{n+2}}(\Omega) = (L^{2^*}(\Omega))' \subset (D^{1,2}(\Omega))'$. Since the operator $-\Delta - (\gamma|x|^{-2} + a)$ is coercive, there exists $w \in D^{1,2}(\Omega)$ such that

$$\begin{cases} \left(-\Delta - \frac{\gamma}{|x|^2} - a(x) \right) w = g & \text{in } \Omega, \\ w = 0 & \text{on } \partial\Omega. \end{cases}$$

Since g vanishes around 0, Theorem 3.2 yields the existence of $K \in \mathbb{R}$ such that

$$w(x) = K \frac{v(x)}{|x|^{\alpha_- + k}} + o\left(\frac{v(x)}{|x|^{\alpha_- + k}} \right) \text{ as } x \rightarrow 0,$$

where v is as in (15). For all $x \in \overline{\Omega} \setminus \{0\}$, we define the function $G_0 := \eta S_{\alpha_+} - w$. The definition of w yields

$$\begin{cases} \left(-\Delta - \frac{\gamma}{|x|^2} - a(x) \right) G_0 = 0 & \text{in } \Omega, \\ G_0 = 0 & \text{on } \partial\Omega \setminus \{0\}. \end{cases}$$

For $\delta_0 > 0$ small enough, the definitions of S_{α_+} , w and $\alpha_- < \alpha_+$ yield

$$G_0(x) = v(x)|x|^{-\alpha_+-k} (1 + o(1)) \text{ in } \mathbb{R}^{k_+,n-k} \cap B_{\delta_0},$$

with $o(1) \rightarrow 0$ as $x \rightarrow 0$. Therefore, $G_0 > 0$ in $\mathbb{R}^{k_+,n-k} \cap B_{\delta_0}$. Then coercivity and the comparison principle yield $G_0 > 0$ in Ω . Moreover, we have that

$$G_0(x) = v(x) (|x|^{-\alpha_+-k} - K|x|^{-\alpha_--k} + o(|x|^{-\alpha_--k})),$$

as $x \rightarrow 0$. Then the mass at 0 of $-\Delta - (\gamma|x|^{-2} + a(x))$ is defined and $m_{\gamma,a}(\Omega) = -K$. This proves Proposition 5.1.

We now discuss briefly examples of negative and positive mass. Here, the reference is Section 9 of Ghoussoub-Robert [8]. We still assume (49) and that $\gamma > \gamma_H(\mathbb{R}^{k_+,n-k}) - 1/4$, so that the mass $m_{\gamma,0}(\Omega)$ is defined. When $\Omega \subset \mathbb{R}^{k_+,n-k}$, due to the comparison principle, we get that $G_0 < S_{\alpha_+}$, and $m_{\gamma,0}(\Omega) < 0$. Arguing as in [8], we are able to define the mass of a domain $\tilde{\Omega} \supset \mathbb{R}^{k_+,n-k}$, for which $m_{\gamma,0}(\tilde{\Omega}) > 0$: then, defining $\tilde{\Omega}_R := \tilde{\Omega} \cap B_R(0)$, we get that $\lim_{R \rightarrow +\infty} m_{\gamma,0}(\tilde{\Omega}_R) = m_{\gamma,0}(\tilde{\Omega}) > 0$. So for $R > 0$ large, we get examples of bounded domains with a singularity of type $(k, n - k)$ at 0 and with positive mass.

6. PROOF OF THEOREM 1.3: FUNCTIONAL BACKGROUND FOR THE PERTURBED EQUATION

In this section, we proceed as in Jaber [11]. For any function $G \in C^1(E, \mathbb{R})$ where $(E, \|\cdot\|)$ is a Banach space, we say that $(u_m)_{m \in \mathbb{N}} \in E$ is a Palais-Smale sequence of G if there exists $\beta \in \mathbb{R}$ such that

$$G(u_m) \rightarrow \beta \text{ and } G'(u_m) \rightarrow 0 \text{ in } E' \text{ as } m \rightarrow +\infty.$$

Here, we say that the Palais-Smale sequence is at level β . The main tool is the Mountain-Pass Lemma of Ambrosetti-Rabinowitz [1]:

Theorem 6.1 (Mountain-Pass Lemma [1]). *Consider $G \in C^1(E, \mathbb{R})$ where $(E, \|\cdot\|)$ is a Banach space. We assume that $G(0) = 0$ and that*

- *There exists $\lambda, r > 0$ such that $G(u) \geq \lambda$ for all $u \in E$ such that $\|u\| = r$,*
- *There exists u_0 in E such that $\limsup_{t \rightarrow +\infty} G(tu_0) < 0$.*

We consider $t_0 > 0$ sufficiently large such that $\|t_0u_0\| > r$ and $G(t_0u_0) < 0$, and

$$\beta = \inf_{c \in \Gamma} \sup_{t \in [0,1]} G(c(t)),$$

where

$$\Gamma = \{c : [0, 1] \rightarrow E \text{ s.t. } c(0) = 0, c(1) = t_0u_0\}.$$

Then, there exists a Palais-Smale sequence at level β for G . Moreover, we have that $\beta \leq \sup_{t \geq 0} G(tu_0)$.

Weak solutions to (6) are to the nonzero critical points of the functional

$$E_q(u) := \frac{1}{2} \int_{\Omega} \left(|\nabla u|^2 - \left(\frac{\gamma}{|x|^2} + a \right) u^2 \right) dx - \int_{\Omega} \frac{u_+^{2^*(s)}}{2^*(s)|x|^s} dx - \int_{\Omega} \frac{hu_+^{q+1}}{q+1} dx,$$

for any $u \in D^{1,2}(\Omega)$ and where $u_+ = \max\{u, 0\}$. In the sequel, we assume that the operator $-\Delta - \left(\frac{\gamma}{|x|^2} + a(x) \right)$ is coercive, so that there exists $c_0 > 0$ such that

$$(50) \quad \int_{\Omega} \left(|\nabla w|^2 - \left(\frac{\gamma}{|x|^2} + a \right) w^2 \right) dx \geq c_0 \int_{\Omega} |\nabla w|^2 dx \text{ for all } w \in D^{1,2}(\Omega).$$

Proposition 6.1. *Let Ω be a bounded domain in \mathbb{R}^n , $n \geq 3$ such that $0 \in \partial\Omega$ is a singularity of type $(k, n-k)$ for some $k \in \{1, \dots, n\}$. We fix $a, h \in C^{0,\theta}(\Omega)$, $\theta \in (0, 1)$. We assume that $h \geq 0$ and that (50) holds. Fix $u_0 \in D^{1,2}(\Omega)$ such that $u_0 \geq 0$, $u_0 \not\equiv 0$, and $q \in (1, 2^* - 1)$. Then there exists a sequence $(u_m)_{m \in \mathbb{N}} \in D^{1,2}(\Omega)$ that is a Palais-Smale sequence for E_q at level β such that $0 < \beta \leq \sup_{t \geq 0} E_q(tu_0)$.*

Proof of Proposition 6.1: Clearly $E_q \in C^1(D^{1,2}(\Omega))$. Note that $E_q(0) = 0$. It follows from (50) and the Sobolev and Hardy-Sobolev embeddings that there exist $c_0, c_1, c_2 > 0$ such that

$$(51) \quad E_q(u) \geq c_0 \|u\|^2 - c_1 \|u\|^{2^*(s)} - c_2 \|u\|^{q+1} \text{ for all } u \in D^{1,2}(\Omega).$$

Define $f(r) = r^2 [c_0 - c_1 r^{2^*(s)-2} - c_2 r^{q-1}] := r^2 g(r)$ and since $2^*(s), q+1 > 2$ we have $g(r) \rightarrow c_0$ as $r \rightarrow 0$. Then there exists $r_0 > 0$ such that $r < r_0$, we have $g(r) > \frac{c_0}{2}$. Therefore, for all $u \in D^{1,2}(\Omega)$ such that $\|u\| = \frac{r_0}{2}$ and by (51), we have $E_q(u) \geq \frac{c_0 r_0^2}{8} := \lambda$. We fix $u_0 \in D^{1,2}(\Omega)$, $u_0 \not\equiv 0$. We have that

$$\begin{aligned} E_q(tu_0) &= \frac{t^2}{2} \int_{\Omega} \left(|\nabla u_0|^2 - \left(\frac{\gamma}{|x|^2} + a \right) u_0^2 \right) dx \\ &\quad - \frac{t^{2^*(s)}}{2^*(s)} \int_{\Omega} \frac{|u_0|^{2^*(s)}}{|x|^s} dx - \frac{t^{q+1}}{q+1} \int_{\Omega} h |u_0|^{q+1} dx \\ &:= \frac{t^2}{2} R_1 - \frac{t^{2^*(s)}}{2^*(s)} R_2 - \frac{t^{q+1}}{q+1} R_3 \leq t^{2^*(s)} \left(\frac{t^{2-2^*(s)}}{2} R_1 - R_2 \right), \end{aligned}$$

where $R_1, R_2 > 0$ and $R_3 \geq 0$. Since $2^*(s) > 2$, we have $E_q(tu_0) \rightarrow -\infty$ as $t \rightarrow +\infty$. Then $\limsup_{t \rightarrow +\infty} E_q(tu_0) < 0$. We consider $t_0 > 0$ large such that $\|t_0 u_0\| > r$ and $E_q(t_0 u_0) < 0$. For $t \in [0, 1]$, we have $E_q(c(t)) \geq \lambda$ and then there exists

$$\beta := \inf_{c \in \Gamma} \sup E_q(c(t)) \geq \lambda > 0.$$

Proposition 6.1 then follows from Theorem 6.1. \square

Proposition 6.2. *Let Ω be a bounded domain in \mathbb{R}^n , $n \geq 3$ such that $0 \in \partial\Omega$ is a singularity of type $(k, n-k)$ for some $k \in \{1, \dots, n\}$. We fix $a, h \in C^{0,\theta}(\Omega)$, $\theta \in (0, 1)$. We assume that $h \geq 0$ and that (50) holds. We fix $\gamma < \gamma_H(\mathbb{R}^{k_+, n-k})$ and $\beta \in \mathbb{R}$ such that*

$$(52) \quad \beta < \frac{2-s}{2(n-s)} \mu_{\gamma, s, 0}(\mathbb{R}^{k_+, n-k})^{\frac{n-s}{2-s}}.$$

Then, for any Palais-Smale sequence $(u_m)_{m \in \mathbb{N}} \in D^{1,2}(\Omega)$ for E_q at level β , there exists $u \in D^{1,2}(\Omega)$ such that $E_q(u) = \beta$ and we have that (u_m) converges strongly in $D^{1,2}(\Omega)$ as $m \rightarrow +\infty$ up to a subsequence. Moreover, we have that $E'_q(u) = 0$.

Proof of Proposition 6.2: Let $(u_m)_{m \in \mathbb{N}} \in D^{1,2}(\Omega)$ be a Palais-Smale sequence for E_q such that

$$E_q(u_m) \rightarrow \beta \text{ and } E'_q(u_m) \rightarrow 0 \text{ in } D^{1,2}(\Omega)'.$$

Step 6.1. *We claim that $(u_m)_m$ is bounded in $D^{1,2}(\Omega)$.*

Proof of Step 6.1: The coercivity (50) and the definition of E_q yield

$$(53) \quad \|u_m\|^2 \leq 2c_0^{-1} \left(E_q(u_m) + \frac{1}{2^*(s)} \int_{\Omega} \frac{(u_m)_+^{2^*(s)}}{|x|^s} dx + \frac{1}{q+1} \int_{\Omega} h (u_m)_+^{q+1} dx \right).$$

Since $E'_q(u_m) \rightarrow 0$ in $D^{1,2}(\Omega)'$, we observe that

$$\int_{\Omega} \left(|\nabla u_m|^2 - \left(\frac{\gamma}{|x|^2} + a \right) u_m^2 \right) dx = \int_{\Omega} \frac{(u_m)_+^{2^*(s)}}{|x|^s} dx + \int_{\Omega} h(u_m)_+^{q+1} dx + o(\|u_m\|).$$

The definition of the energy E_q and the last equation yield

$$2E_q(u_m) = \left(1 - \frac{2}{2^*(s)} \right) \int_{\Omega} \frac{(u_m)_+^{2^*(s)}}{|x|^s} dx + \left(1 - \frac{2}{q+1} \right) \int_{\Omega} h(u_m)_+^{q+1} dx + o(\|u_m\|).$$

Moreover, since $E_q(u_m) \rightarrow \beta$ as $m \rightarrow +\infty$, $h \geq 0$ and $q+1 > 2$, we obtain that

$$\begin{aligned} \left(1 - \frac{2}{2^*(s)} \right) \int_{\Omega} \frac{(u_m)_+^{2^*(s)}}{|x|^s} dx &= 2E_q(u_m) - \left(1 - \frac{2}{q+1} \right) \int_{\Omega} h(u_m)_+^{q+1} dx + o(\|u_m\|) \\ &\leq 2\beta + o(\|u_m\|), \end{aligned}$$

therefore,

$$(55) \quad \left(1 - \frac{2}{2^*(s)} \right) \int_{\Omega} \frac{(u_m)_+^{2^*(s)}}{|x|^s} dx = O(1) + o(\|u_m\|).$$

Similarly, we have that

$$(56) \quad \left(1 - \frac{2}{q+1} \right) \int_{\Omega} h(u_m)_+^{q+1} dx = O(1) + o(\|u_m\|).$$

Relations (53) and (54) give

$$(57) \quad \|u_m\|^2 \leq c_0^{-1} \left(\int_{\Omega} \frac{(u_m)_+^{2^*(s)}}{|x|^s} dx + \int_{\Omega} h(u_m)_+^{q+1} dx \right) + o(\|u_m\|).$$

The equations (55), (56) and (57) yield,

$$\|u_m\|^2 = O(1) + o(\|u_m\|),$$

as $m \rightarrow +\infty$. This proves Step 6.1. \square

Therefore, up to a subsequence, there exists $u \in D^{1,2}(\Omega)$ such that

$$(58) \quad \begin{cases} u_m \rightharpoonup u & \text{weakly in } D^{1,2}(\Omega), \\ u_m \rightarrow u & \text{strongly in } L^p(\Omega) \text{ for all } 1 < p < 2^*. \end{cases}$$

Moreover, we have $E'_q(u) = 0$.

Step 6.2. We claim that, as $m \rightarrow +\infty$

$$(59) \quad \int_{\Omega} \left(|\nabla(u_m - u)|^2 - \gamma \frac{(u_m - u)^2}{|x|^2} \right) dx = \int_{\Omega} \frac{(u_m - u)_+^{2^*(s)}}{|x|^s} dx + o(1),$$

and,

$$(60) \quad \frac{2-s}{2(n-s)} \int_{\Omega} \left(|\nabla(u_m - u)|^2 - \gamma \frac{(u_m - u)^2}{|x|^2} \right) dx \leq \beta + o(1).$$

Proof of Step 6.2: We have that

$$\begin{aligned} \langle E'_q(u_m), \varphi \rangle &= \int_{\Omega} \left((\nabla u_m, \nabla \varphi) - \left(\frac{\gamma}{|x|^2} + a \right) u_m \varphi \right) dx \\ &\quad - \int_{\Omega} \frac{(u_m)_+^{2^*(s)-1}}{|x|^s} \varphi dx - \int_{\Omega} h(u_m)_+^q \varphi dx, \end{aligned}$$

for all $\varphi \in D^{1,2}(\Omega)$. We observe that

$$\begin{aligned}
(61) \quad o(1) &= \langle E'_q(u_m) - E'_q(u), u_m - u \rangle \\
&= \int_{\Omega} \left(|\nabla(u_m - u)|^2 - \left(\frac{\gamma}{|x|^2} + a \right) (u_m - u)^2 \right) dx \\
&\quad - \int_{\Omega} \left((u_m)_+^{2^*(s)-1} - u_+^{2^*(s)-1} \right) \frac{(u_m - u)}{|x|^s} dx \\
&\quad - \int_{\Omega} h \left((u_m)_+^q - u_+^q \right) (u_m - u) dx.
\end{aligned}$$

Since $u_m \rightharpoonup u$ weakly in $D^{1,2}(\Omega)$ as $m \rightarrow \infty$, integration theory yields

$$(62) \quad \lim_{m \rightarrow +\infty} \int_{\Omega} \frac{(u_m)_+^{2^*(s)-1}}{|x|^s} u dx = \int_{\Omega} \frac{u_+^{2^*(s)}}{|x|^s} dx = \lim_{m \rightarrow +\infty} \int_{\Omega} \frac{u_+^{2^*(s)-1}}{|x|^s} u_m dx.$$

Equation (58) yields

$$(63) \quad \int_{\Omega} h(u_m - u) \left((u_m)_+^q - u_+^q \right) dx = \int_{\Omega} h(u_m - u)^{q+1} dx + o(1) = o(1),$$

as $m \rightarrow +\infty$. Combining (61), (62) and (63), we get as $m \rightarrow +\infty$ that

$$(64) \quad \int_{\Omega} \left(|\nabla(u_m - u)|^2 - \left(\frac{\gamma}{|x|^2} + a \right) (u_m - u)^2 \right) dx = \int_{\Omega} \left((u_m)_+^{2^*(s)} - u_+^{2^*(s)} \right) \frac{dx}{|x|^s} + o(1).$$

Since $2^*(s) > 1$, we get that

$$\left| (u_m)_+^{2^*(s)} - u_+^{2^*(s)} - (u_m - u)_+^{2^*(s)} \right| \leq C \left(|u_m - u|^{2^*(s)-1} |u| + |u|^{2^*(s)-1} |u_m - u| \right),$$

for some $C > 0$ independent of m . Therefore, with (58), we have that

$$(65) \quad \int_{\Omega} \left((u_m)_+^{2^*(s)} - (u_m - u)_+^{2^*(s)} \right) \frac{dx}{|x|^s} = \int_{\Omega} \frac{u_+^{2^*(s)}}{|x|^s} dx + o(1).$$

Since $u_m \rightarrow u$ strongly in $L^2(\Omega)$ as $m \rightarrow +\infty$ and by (64), (65), we obtain (59).

With (58) we have that

$$\begin{aligned}
E_q(u_m) - E_q(u) &= \frac{1}{2} \int_{\Omega} \left(|\nabla(u_m - u)|^2 - \gamma \frac{(u_m - u)^2}{|x|^2} \right) dx \\
&\quad - \frac{1}{2^*(s)} \int_{\Omega} \left((u_m)_+^{2^*(s)} - u_+^{2^*(s)} \right) \frac{dx}{|x|^s} + o(1).
\end{aligned}$$

With (59), we get

$$E_q(u_m) - E_q(u) = \left(\frac{1}{2} - \frac{1}{2^*(s)} \right) \int_{\Omega} \left(|\nabla(u_m - u)|^2 - \gamma \frac{(u_m - u)^2}{|x|^2} \right) dx + o(1).$$

Since u is a solution to (6) then $E_q(u) \geq 0$. Moreover $E_q(u_m) \rightarrow \beta$ as $m \rightarrow +\infty$. Then we then get (60). This proves Step 6.2. \square

Step 6.3. *We claim that*

$$(66) \quad \lim_{m \rightarrow +\infty} u_m = u \text{ in } D^{1,2}(\Omega).$$

Proof of Step 6.3: Since $\gamma < \gamma_H(\mathbb{R}^{k+,n-k})$ for all $k \in \{1, \dots, n\}$ and by the Proposition 2.1 in Cheikh-Ali [4], then for all $\epsilon > 0$ there exists $c_\epsilon > 0$ such that for all $v \in D^{1,2}(\Omega)$,

$$\left(\int_{\Omega} \frac{|v|^{2^*(s)}}{|x|^s} dx \right)^{\frac{2}{2^*(s)}} \leq (\mu_{\gamma,s,0}(\mathbb{R}^{k+,n-k})^{-1} + \epsilon) \int_{\Omega} \left(|\nabla v|^2 - \frac{\gamma}{|x|^2} v^2 \right) dx + c_\epsilon \int_{\Omega} v^2 dx.$$

Take $\theta_m = u_m - u$. Since (u_m) converges strongly to u in $L^2(\Omega)$, taking $v = \theta_m$ yields

$$(67) \quad \left(\int_{\Omega} \frac{(\theta_m)_+^{2^*(s)}}{|x|^s} dx \right)^{\frac{2}{2^*(s)}} \leq (\mu_{\gamma,s,0}(\mathbb{R}^{k+,n-k})^{-1} + \epsilon) \int_{\Omega} \left(|\nabla \theta_m|^2 - \frac{\gamma}{|x|^2} \theta_m^2 \right) dx + o(1).$$

We write $N(\theta_m) := \int_{\Omega} \left(|\nabla \theta_m|^2 - \frac{\gamma}{|x|^2} \theta_m^2 \right) dx$. By (59) and (67), we get that

$$N(\theta_m)^{\frac{2}{2^*(s)}} \left(1 - (\mu_{\gamma,s,0}(\mathbb{R}^{k+,n-k})^{-1} + \epsilon) N(\theta_m)^{1 - \frac{2}{2^*(s)}} \right) \leq o(1).$$

With (60) and the last inequation, we get that, as $m \rightarrow \infty$,

$$(68) \quad N(\theta_m)^{\frac{2}{2^*(s)}} \left(1 - (\mu_{\gamma,s,0}(\mathbb{R}^{k+,n-k})^{-1} + \epsilon) \left(\frac{2(n-s)\beta}{2-s} \right)^{\frac{2^*(s)-2}{2^*(s)}} + o(1) \right) \leq o(1).$$

With the assumption (52) and (68), taking $\epsilon > 0$ small enough, we get that $N(\theta_m) \rightarrow 0$ as $m \rightarrow +\infty$ and by coercivity, we obtain (66). \square

With Step 6.3 and since $E_q(u_m) \rightarrow \beta$ as $m \rightarrow +\infty$, we get that $E_q(u) = \beta$. This ends the proof of Proposition 6.2. \square

Theorem 6.2. *Let Ω be a bounded domain in \mathbb{R}^n , $n \geq 3$, such that $0 \in \partial\Omega$ is a singularity of type $(k, n-k)$ for some $k \in \{1, \dots, n\}$. We fix $\gamma < \gamma_H(\mathbb{R}^{k+,n-k})$, $a \in C^{0,\theta}(\Omega)$ such that $-\Delta - (\gamma|x|^{-2} + a(x))$ is coercive, and $h \in C^{0,\theta}(\Omega)$ such that $h \geq 0$ and let $0 \leq s < 2$ and $1 < q < 2^* - 1$. Assume that there exists $u_0 \in D^{1,2}(\Omega)$, $u_0 \not\equiv 0$, such that*

$$(69) \quad \sup_{t \geq 0} E_q(tu_0) < \frac{2-s}{2(n-s)} \mu_{\gamma,s,0}(\mathbb{R}^{k+,n-k})^{\frac{n-s}{2-s}},$$

then equation (6) has a non-vanishing solution in $D^{1,2}(\Omega)$ of Mountain-Pass type.

Proof of Theorem 6.2: By Proposition 6.1, there exists a Palais-Smale sequence $(u_m)_{m \in \mathbb{N}} \in D^{1,2}(\Omega)$ for E_q at level $\beta > 0$ such that $\beta \leq \sup_{t \geq 0} E_q(tu_0)$. It then follows from Proposition 6.2 that, up to a subsequence, (u_m) converges strongly to u in $D^{1,2}(\Omega)$. Then $E_q(u) = \beta > 0$, so $u \not\equiv 0$, and $E'_q(u) = 0$. Coercivity and $E'_q(u)[u_-] = 0$ yield $u \geq 0$. Regularity theory and Hopf's principle yield $u \in C^{2,\theta}(\Omega)$ and $u > 0$. Then u is a solution of (6). This proves Theorem 6.2. \square

7. PROOF OF THEOREM 1.3: TEST-FUNCTIONS ESTIMATES

The main result of this section is the following:

Proposition 7.1. *We fix $\gamma < \gamma_H(\mathbb{R}^{k+,n-k})$ and $0 \leq s < 2$. We assume that there are extremals for $\mu_{\gamma,s,0}(\mathbb{R}^{k+,n-k})$ and we let U as in (12) be such an extremal. We let $(u_\epsilon)_\epsilon$ and $(\tilde{u}_\epsilon)_\epsilon$ be as in (22). Then,*

(a) For $0 \leq \gamma < \gamma_H(\mathbb{R}^{k_+, n-k}) - \frac{1}{4}$, we have that

$$\sup_{t \geq 0} E_q(tu_\epsilon) := \beta_0 + \begin{cases} c_1 GH_{\gamma, s}(\Omega) \epsilon + o(\epsilon) & \text{if } q+1 < \frac{2n-2}{n-2}, \\ (c_1 GH_{\gamma, s}(\Omega) - c_2 h(0)) \epsilon + o(\epsilon) & \text{if } q+1 = \frac{2n-2}{n-2}, \\ -c_2 h(0) \epsilon^{n - \frac{(q+1)(n-2)}{2}} + o(\epsilon^{n - \frac{(q+1)(n-2)}{2}}) & \text{if } q+1 > \frac{2n-2}{n-2}. \end{cases}$$

(b) For $0 \leq \gamma = \gamma_H(\mathbb{R}^{k_+, n-k}) - \frac{1}{4}$, we have that

$$\sup_{t \geq 0} E_q(tu_\epsilon) := \beta_0 + \begin{cases} c_1 GH_{\gamma, s}(\Omega) \epsilon \ln\left(\frac{1}{\epsilon}\right) + o(\epsilon \ln\left(\frac{1}{\epsilon}\right)) & \text{if } q+1 \leq \frac{2n-2}{n-2}, \\ -c_2 h(0) \epsilon^{n - \frac{(q+1)(n-2)}{2}} + o(\epsilon^{n - \frac{(q+1)(n-2)}{2}}) & \text{if } q+1 > \frac{2n-2}{n-2}. \end{cases}$$

(c) For $\gamma > \gamma_H(\mathbb{R}^{k_+, n-k}) - \frac{1}{4}$, we have that

$$\sup_{t \geq 0} E_q(t\tilde{u}_\epsilon) := \beta_0 + \begin{cases} -c_3 m_\gamma(\Omega) \epsilon^{\alpha_+ - \alpha_-} + o(\epsilon^{\alpha_+ - \alpha_-}) & \text{if } q+1 < \frac{2n-2(\alpha_+ - \alpha_-)}{n-2}, \\ -(c_3 m_\gamma(\Omega) + c_2 h(0)) \epsilon^{\alpha_+ - \alpha_-} + o(\epsilon^{\alpha_+ - \alpha_-}) & \text{if } q+1 = \frac{2n-2(\alpha_+ - \alpha_-)}{n-2}, \\ -c_2 h(0) \epsilon^{n - \frac{(q+1)(n-2)}{2}} + o(\epsilon^{n - \frac{(q+1)(n-2)}{2}}) & \text{if } q+1 > \frac{2n-2(\alpha_+ - \alpha_-)}{n-2}, \end{cases}$$

where $\beta_0 = \frac{2-s}{2(n-s)} \mu_{\gamma, s, 0}(\mathbb{R}^{k_+, n-k})^{\frac{n-s}{2-s}}$,

$$(70) \quad \begin{cases} c_1 = \frac{\mu_{\gamma, s, 0}(\mathbb{R}^{k_+, n-k})^{\frac{2^*(s)}{2^*(s)-2}}}{2} \left(\xi \int_{\mathbb{R}^{k_+, n-k}} \frac{U^{2^*(s)}}{|x|^s} dx \right)^{-1}, \\ c_2 = \frac{\xi^{\frac{q+1}{2^*(s)-2}}}{q+1} \int_{\mathbb{R}^{k_+, n-k}} U^{q+1} dx, \\ c_3 = \mu_{\gamma, s, 0}(\mathbb{R}^{k_+, n-k})^{\frac{2^*(s)}{2^*(s)-2}} \frac{\alpha_+ - \alpha_-}{2} \frac{\int_{S^{n-1} \cap \mathbb{R}^{k_+, n-k}} \left(\prod_{i=1}^k x_i \right)^2 d\sigma}{\xi \int_{\mathbb{R}^{k_+, n-k}} \frac{U^{2^*(s)}}{|x|^s} dx}. \end{cases}$$

Theorem 6.2 and Proposition 7.1 yield Theorem 1.3.

Proof of Proposition 7.1: We define the test-function sequence $(Z_\epsilon)_{\epsilon > 0}$ by

$$Z_\epsilon(x) := \begin{cases} u_\epsilon & \text{if } \gamma \leq \gamma_H(\mathbb{R}^{k_+, n-k}) - \frac{1}{4}, \\ \tilde{u}_\epsilon & \text{if } \gamma > \gamma_H(\mathbb{R}^{k_+, n-k}) - \frac{1}{4}, \end{cases}$$

where u_ϵ and \tilde{u}_ϵ are as in the definition (22). We have:

$$E_q(tZ_\epsilon) = \frac{t^2}{2} R_\epsilon - \frac{t^{2^*(s)}}{2^*(s)} B_\epsilon - \frac{t^{q+1}}{q+1} C_{h, \epsilon},$$

when $\epsilon \rightarrow 0$ where:

$$R_\epsilon := \int_{\Omega} \left(|\nabla Z_\epsilon|^2 - \left(\frac{\gamma}{|x|^2} + a(x) \right) Z_\epsilon^2 \right) dx$$

$$B_\epsilon := \int_{\Omega} \frac{Z_\epsilon^{2^*(s)}}{|x|^s} dx \text{ and } C_{h, \epsilon} := \int_{\Omega} h Z_\epsilon^{q+1} dx.$$

Step 7.1. We fix $f \in C^{0, \theta}(\Omega)$, $\theta \in (0, 1)$, and $p \in [1, 2^*)$. We claim that

$$\int_{\Omega} f |Z_\epsilon|^{p+1} dx = \begin{cases} f(0) \epsilon^{n - \frac{n-2}{2}(p+1)} \int_{\mathbb{R}^{k_+, n-k}} U^{p+1} dx + o\left(\epsilon^{n - \frac{n-2}{2}(p+1)}\right) & \text{if } n < (p+1)\alpha_+, \\ O\left(\epsilon^{\frac{p+1}{2}(\alpha_+ - \alpha_-)} \ln\left(\frac{1}{\epsilon}\right)\right) & \text{if } n = (p+1)\alpha_+, \\ O\left(\epsilon^{\frac{p+1}{2}(\alpha_+ - \alpha_-)}\right) & \text{if } n > (p+1)\alpha_+. \end{cases}$$

Moreover, we have that

$$(71) \quad \int_{\Omega} f |Z_\epsilon|^{p+1} dx \rightarrow 0 \text{ as } \epsilon \rightarrow 0.$$

Proof of Step 7.1: Note that it follows from (20) that

$$(72) \quad 0 < U_\epsilon(x) \leq C\epsilon^{\frac{\alpha_+ - \alpha_-}{2}} |x|^{-\alpha_+} \text{ for all } x \in \mathbb{R}^{k_+, n-k} \text{ and } \epsilon > 0.$$

We first prove Step 7.1 for u_ϵ , postponing the case of \tilde{u}_ϵ , and then Z_ϵ , to the end of the proof. We distinguish three cases:

Case 1: We assume that $n > (p+1)\alpha_+$. It follows from (72) that

$$\left| \int_{\Omega} f |u_\epsilon|^{p+1} dx \right| \leq C\epsilon^{\frac{p+1}{2}(\alpha_+ - \alpha_-)} \int_{\Omega} |x|^{-(p+1)\alpha_+} dx \leq C\epsilon^{\frac{p+1}{2}(\alpha_+ - \alpha_-)}$$

as $\epsilon \rightarrow 0$. This proves Step 7.1 for u_ϵ when $n > (p+1)\alpha_+$.

Case 2: We assume that $n = (p+1)\alpha_+$. With (72), we get that

$$\begin{aligned} \left| \int_{\Omega} f |u_\epsilon|^{p+1} dx \right| &\leq C\epsilon^{n - \frac{n-2}{2}(p+1)} + C \int_{B_{\delta, +}} |u_\epsilon|^{p+1} dx \\ &\leq C\epsilon^{n - \frac{n-2}{2}(p+1)} + C\epsilon^{n - \frac{n-2}{2}(p+1)} \int_{B_{\delta\epsilon^{-1}, +}} U^{p+1} dx \\ &\leq C\epsilon^{n - \frac{n-2}{2}(p+1)} + C\epsilon^{n - \frac{n-2}{2}(p+1)} \int_1^{\epsilon^{-1}\delta} r^{-1} dr \\ &\leq C\epsilon^{\frac{p+1}{2}(\alpha_+ - \alpha_-)} \ln\left(\frac{1}{\epsilon}\right) \end{aligned}$$

Case 3: We assume that $n < (p+1)\alpha_+$. For $\rho > 0$ small enough, it follows from (72) that

$$\int_{\Omega \setminus \phi(B_{\rho, +})} f |u_\epsilon|^{p+1} dx = O\left(\epsilon^{\frac{p+1}{2}(\alpha_+ - \alpha_-)}\right) \text{ as } \epsilon \rightarrow 0.$$

Independently, since $f \in C^{0, \theta}(\Omega)$, we have that

$$(73) \quad \begin{aligned} \int_{\phi(B_{\rho, +})} f |u_\epsilon|^{p+1} dx &= \int_{B_{\rho, +}} f \circ \phi \cdot U_\epsilon^{p+1} |\text{Jac } \phi| dx \\ &= \epsilon^{n - \frac{n-2}{2}(p+1)} f(0) \int_{B_{\rho\epsilon^{-1}, +}} U^{p+1} dx + O\left(\int_{B_{\rho, +}} |x|^\theta |U_\epsilon|^{p+1} dx\right) \end{aligned}$$

Since $n < (p+1)\alpha_+$, it follows from (20) that $U \in L^{p+1}(\mathbb{R}^{k_+, n-k})$ and that

$$(74) \quad \begin{aligned} \int_{B_{\rho\epsilon^{-1}, +}} U^{p+1} dx &= \int_{\mathbb{R}^{k_+, n-k}} U^{p+1} dx + O\left(\int_{\mathbb{R}^{k_+, n-k} \setminus B_{\rho\epsilon^{-1}, +}} U^{p+1} dx\right) \\ &= \int_{\mathbb{R}^{k_+, n-k}} U^{p+1} dx + O\left(\int_{\epsilon^{-1}\rho}^{\infty} r^{n-(p+1)\alpha_+ - 1} dr\right) \\ &= \int_{\mathbb{R}^{k_+, n-k}} U^{p+1} dx + O\left(\epsilon^{(p+1)\alpha_+ - n}\right) \end{aligned}$$

We claim that

$$(75) \quad \int_{B_{\rho, +}} |x|^\theta |U_\epsilon|^{p+1} dx = o\left(\epsilon^{n - \frac{n-2}{2}(p+1)}\right) \text{ as } \epsilon \rightarrow 0.$$

Indeed, when $\theta + n > (p+1)\alpha_+$, we argue as in Case 1. When $\theta + n = (p+1)\alpha_+$, we argue as in Case 2. When $\theta + n < (p+1)\alpha_+$, we make a change of variable

$y = \epsilon^{-1}x$ and we argue as in (74). This yields (75). Putting (74) and (75) in (73) yields Step 7.1 for u_ϵ in Case 2.

We now prove Step 7.1. When $\gamma \leq \gamma_H(\mathbb{R}^{k+,n-k}) - \frac{1}{4}$, $Z_\epsilon = u_\epsilon$, and we are done. When $\gamma > \gamma_H(\mathbb{R}^{k+,n-k}) - \frac{1}{4}$, $Z_\epsilon = \tilde{u}_\epsilon$. With the definition (22), we get that

$$(76) \quad \int_{\Omega} f|\tilde{u}_\epsilon|^{p+1} dx = \int_{\Omega} f \left| u_\epsilon + \epsilon^{\frac{\alpha_+ - \alpha_-}{2}} \Theta \right|^{p+1} dx \\ = \int_{\Omega} f|u_\epsilon|^{p+1} dx + O\left(\epsilon^{\frac{\alpha_+ - \alpha_-}{2}} \int_{\Omega} |u_\epsilon|^p |\Theta| dx\right) + O\left(\epsilon^{\frac{p+1}{2}(\alpha_+ - \alpha_-)} \int_{\Omega} |\Theta|^{p+1} dx\right)$$

Since $\Theta \in D^{1,2}(\Omega)$ and $p+1 < 2^*$, we get that $\Theta \in L^{p+1}(\Omega)$. It follows from (16) that $|\Theta(x)| \leq C|x|^{-\alpha_-}$ for all $x \in \Omega$. Arguing as in Cases 1, 2, 3 above, we get that the second term in the right-hand-side of (76) is dominated by $\int_{\Omega} |u_\epsilon|^{p+1} dx$. Then Step 7.1 for $\gamma > \gamma_H(\mathbb{R}^{k+,n-k}) - 1/4$ follows. \square

By Cheikh-Ali [4] and Step 7.1 for the case $\gamma \leq \gamma_H(\mathbb{R}^{k+,n-k}) - 1/4$ and Steps 4.2 and 4.3 for the case $\gamma > \gamma_H(\mathbb{R}^{k+,n-k}) - 1/4$, we get that, as $\epsilon \rightarrow 0$,

$$(77) \quad R_\epsilon \rightarrow R_0 := \xi \int_{\mathbb{R}^{k+,n-k}} \frac{U^{2^*(s)}}{|x|^s} dx \text{ and } B_\epsilon \rightarrow B_0 := \int_{\mathbb{R}^{k+,n-k}} \frac{U^{2^*(s)}}{|x|^s} dx.$$

Step 7.2. We claim that for all $\epsilon > 0$, then there exists a unique t_ϵ such that

$$(78) \quad \sup_{t \geq 0} E_q(tZ_\epsilon) = E_q(t_\epsilon Z_\epsilon).$$

Moreover, t_ϵ verifies

$$(79) \quad t_\epsilon = S_\epsilon [1 - C_0 C_{h,\epsilon} + o(C_{h,\epsilon})],$$

where $S_\epsilon := (R_\epsilon B_\epsilon^{-1})^{\frac{1}{2^*(s)-2}}$, $C_0 > 0$ and $t_\epsilon \rightarrow t_0$ as $\epsilon \rightarrow 0$.

Proof of Step 7.2: We have that $\partial_t E_q(tZ_\epsilon) = 0$ iff $t = 0$ or $g_\epsilon(t) = R_\epsilon$ where $g_\epsilon(t) := B_\epsilon t^{2^*(s)-2} + C_{h,\epsilon} t^{q-1}$. Since $B_\epsilon, C_{h,\epsilon} \geq 0$ and g_ϵ is a strictly increasing map i.e $g_\epsilon(t) - R_\epsilon$ also, and since $R_\epsilon > 0$ we have $g_\epsilon(0) - R_\epsilon < 0$ then, there exists $t_\epsilon > 0$ unique verifying $g_\epsilon(t_\epsilon) = R_\epsilon$ such that (78) holds. Since $g_\epsilon(t_\epsilon) = R_\epsilon$, we get

$$t_\epsilon \leq S_\epsilon := (R_\epsilon B_\epsilon^{-1})^{\frac{1}{2^*(s)-2}}.$$

We are using (77), (71) and (12) to get that $S_\epsilon \rightarrow (R_0 B_0^{-1})^{\frac{1}{2^*(s)-2}} = \xi^{\frac{1}{2^*(s)-2}}$ as $\epsilon \rightarrow 0$. Therefore, t_ϵ is bounded and there exists t_0 such that $t_\epsilon \rightarrow t_0$ up to extraction. Since $g_\epsilon(t_\epsilon) = R_\epsilon$ and $C_{h,\epsilon} \rightarrow 0$ as $\epsilon \rightarrow 0$, we obtain that

$$t_\epsilon = [R_\epsilon B_\epsilon^{-1} - C_{h,\epsilon} B_\epsilon^{-1} t_\epsilon^{q-1}]^{\frac{1}{2^*(s)-2}} \\ = S_\epsilon [1 - C_{h,\epsilon} R_\epsilon^{-1} t_\epsilon^{q-1}]^{\frac{1}{2^*(s)-2}} = S_\epsilon [1 - C_0 C_{h,\epsilon} + o(C_{h,\epsilon})],$$

where $C_0 := \frac{R_0^{-1} t_0^{q-1}}{2^*(s)-2}$ and $t_0 = \xi^{\frac{1}{2^*(s)-2}}$. This yields (79) and Step 7.2. \square

Step 7.3. We claim that, as $\epsilon \rightarrow 0$,

$$E_q(t_\epsilon Z_\epsilon) = \frac{2-s}{2(n-s)} (J_{\gamma,s,a}^\Omega(Z_\epsilon))^{\frac{2^*(s)}{2^*(s)-2}} - \frac{\xi^{\frac{q+1}{2^*(s)-2}}}{q+1} C_{h,\epsilon} + o(C_{h,\epsilon}).$$

Proof of Step 7.3: The expression (79) of Step 7.2 and (71) yield

$$\begin{aligned}
E_q(t_\epsilon Z_\epsilon) &= \frac{t_\epsilon^2}{2} R_\epsilon - \frac{t_\epsilon^{2^*(s)}}{2^*(s)} B_\epsilon - \frac{t_\epsilon^{q+1}}{q+1} C_{h,\epsilon} \\
&= \frac{S_\epsilon^2 [1 - C_0 C_{h,\epsilon} + o(C_{h,\epsilon})]^2}{2} R_\epsilon - \frac{S_\epsilon^{2^*(s)} [1 - C_0 C_{h,\epsilon} + o(C_{h,\epsilon})]^{2^*(s)}}{2^*(s)} B_\epsilon \\
&\quad - \frac{S_\epsilon^{q+1} [1 - C_0 C_{h,\epsilon} + o(C_{h,\epsilon})]^{q+1}}{q+1} C_{h,\epsilon} \\
&= \frac{S_\epsilon^2 [1 - 2C_0 C_{h,\epsilon} + o(C_{h,\epsilon})]}{2} R_\epsilon - \frac{S_\epsilon^{2^*(s)} [1 - C_0 2^*(s) C_{h,\epsilon} + o(C_{h,\epsilon})]}{2^*(s)} B_\epsilon \\
&\quad - \frac{S_\epsilon^{q+1} [1 - (q+1)C_0 C_{h,\epsilon} + o(C_{h,\epsilon})]}{q+1} C_{h,\epsilon},
\end{aligned}$$

then,

$$\begin{aligned}
E_q(t_\epsilon Z_\epsilon) &= \frac{S_\epsilon^2}{2} R_\epsilon - \frac{S_\epsilon^{2^*(s)}}{2^*(s)} B_\epsilon - \frac{S_\epsilon^{q+1}}{q+1} C_{h,\epsilon} \\
&\quad - C_0 C_{h,\epsilon} \left[S_\epsilon^2 R_\epsilon - S_\epsilon^{2^*(s)} B_\epsilon - S_\epsilon^{q+1} C_{h,\epsilon} \right] + o(C_{h,\epsilon}).
\end{aligned}$$

Since $S_\epsilon := (R_\epsilon B_\epsilon^{-1})^{\frac{1}{2^*(s)-2}}$ and $C_{h,\epsilon} \rightarrow 0$ as $\epsilon \rightarrow 0$, this yields Step 7.3. \square

Proof of Proposition 7.1 when $0 \leq \gamma \leq \gamma_H(\mathbb{R}^{k+,n-k}) - \frac{1}{4}$. In this case, we recall that $Z_\epsilon(x) = u_\epsilon(x)$. Note that

$$\left\{ \gamma < (=) \gamma_H(\mathbb{R}^{k+,n-k}) - \frac{1}{4} \right\} \Leftrightarrow \{ \alpha_+ - \alpha_- > (=) 1 \}.$$

It was proved in Proposition 5.1 in Cheikh-Ali [4] that

- For $\gamma < \gamma_H(\mathbb{R}^{k+,n-k}) - \frac{1}{4}$, we have that

$$(80) \quad J_{\gamma,s,0}^\Omega(u_\epsilon) = \mu_{\gamma,s,0}(\mathbb{R}^{k+,n-k}) (1 + \kappa GH_{\gamma,s}(\Omega)\epsilon + o(\epsilon)).$$

- For $\gamma = \gamma_H(\mathbb{R}^{k+,n-k}) - \frac{1}{4}$, we have that

$$(81) \quad J_{\gamma,s,0}^\Omega(u_\epsilon) = \mu_{\gamma,s,0}(\mathbb{R}^{k+,n-k}) \left(1 + \kappa GH_{\gamma,s}(\Omega)\epsilon \ln\left(\frac{1}{\epsilon}\right) + o\left(\epsilon \ln\left(\frac{1}{\epsilon}\right)\right) \right),$$

where $\kappa := \left(\xi \int_{\mathbb{R}^{k+,n-k}} \frac{U^{2^*(s)}}{|x|^s} dx \right)^{-1}$ and $GH_{\gamma,s}(\Omega)$ is defined in (7). It follows from Step 7.1 that $\int_\Omega u_\epsilon^2 dx = o(\epsilon)$ if $\alpha_+ - \alpha_- > 1$, and $O(\epsilon)$ if $\alpha_+ - \alpha_- = 1$. Therefore (80) and (81) hold unchanged with the potential a .

Case 1: We assume that $n < (q+1)\alpha_+$. It follows from Step 7.1 that

$$C_{h,\epsilon} = \int_\Omega h |u_\epsilon|^{q+1} dx = h(0) \epsilon^{n - \frac{n-2}{2}(q+1)} \int_{\mathbb{R}^{k+,n-k}} U^{q+1} dx + o\left(\epsilon^{n - \frac{n-2}{2}(q+1)}\right)$$

as $\epsilon \rightarrow 0$. Then, when $n < (q+1)\alpha_+$, Case (a) of Proposition 7.1 follows by combining Step 7.3, (80), (81), the estimate of $C_{h,\epsilon}$ and studying the relative positions of $n - \frac{n-2}{2}(q+1)$ and 1.

Case 2: We assume that $n \geq (q+1)\alpha_+$. Since $\alpha_+ - \alpha_- \geq 1$ and $q > 1$, we then get that

$$n - \frac{n-2}{2}(q+1) - 1 = (n - (q+1)\alpha_+) + \frac{q+1}{2}(\alpha_+ - \alpha_-) - 1 > 0.$$

Then, for $n \geq (q+1)\alpha_+$, Cases (a) and (b) of Proposition 7.1 follows by the same arguments as in Case 1.

This proves Cases (a) and (b) of Proposition 7.1. \square

Proof of Proposition 7.1 when $\gamma > \gamma_H(\mathbb{R}^{k_+, n-k}) - \frac{1}{4}$. Proposition 4.1 yields

$$(82) \quad J_{\gamma, s, a}^\Omega(\tilde{u}_\epsilon) = \mu_{\gamma, s, 0}(\mathbb{R}^{k_+, n-k}) \left(1 - \zeta_{\gamma, s}^0 m_{\gamma, a}(\Omega) \epsilon^{\alpha_+ - \alpha_-} + o(\epsilon^{\alpha_+ - \alpha_-})\right),$$

as $\epsilon \rightarrow 0$. Here, we compare $n - \frac{n-2}{2}(q+1)$ and $\alpha_+ - \alpha_-$. Note that

$$n - \frac{n-2}{2}(q+1) - (\alpha_+ - \alpha_-) = n - (q+1)\alpha_+ + \frac{q-1}{2}(\alpha_+ - \alpha_-).$$

Therefore, since $q > 1$, when $n \geq (q+1)\alpha_+$, we have that $n - \frac{n-2}{2}(q+1) > \alpha_+ - \alpha_-$. As for the case $\gamma \leq \gamma_H(\mathbb{R}^{k_+, n-k}) - \frac{1}{4}$, we get Case (b) of Proposition 7.1 by studying the relative positions of $n - \frac{n-2}{2}(q+1)$ and $\alpha_+ - \alpha_-$ and using Step 7.1 and (82).

This proves Case (c) of Proposition 7.1. \square

All these cases prove Proposition 7.1. As already mentioned, Theorem 6.2 and Proposition 7.1 yield Theorem 1.3.

REFERENCES

- [1] A. Ambrosetti and P.H. Rabinowitz, *Dual variational methods in critical point theory and applications*, J. Functional Analysis **14** (1973), 349–381.
- [2] H. Brezis and L. Nirenberg, *Positive solutions of nonlinear elliptic equations involving critical Sobolev exponents*, Comm. Pure Appl. Math. **36** (1983), 437–477.
- [3] L. Caffarelli, R. Kohn, and L. Nirenberg, *First order interpolation inequalities with weights*, Compositio Math. **53** (1984), no. 3, 259–275.
- [4] H. Cheikh Ali, *Hardy–Sobolev inequalities with singularities on non smooth boundary: Hardy constant and extremals. Part I: Influence of local geometry*, Nonlinear Anal. **182** (2019), 316–349.
- [5] Z. Djadli, *Nonlinear elliptic equations with critical Sobolev exponent on compact Riemannian manifolds*, Calc. Var. Partial Differential Equations **8** (1999), no. 4, 293–326.
- [6] V. Felli and A. Ferrero, *Almgreen-type monotonicity methods for the classification of behaviour at corners of solutions to semilinear elliptic equations*, Proc. Roy. Soc. Edinburgh Sect. A **143** (2013), no. 5, 957–1019.
- [7] N. Ghoussoub and F. Robert, *Sobolev inequalities for the Hardy–Schrödinger operator: extremals and critical dimensions*, Bull. Math. Sci. **6** (2016), no. 1, 89–144.
- [8] ———, *Hardy-singular boundary mass and Sobolev-critical variational problems*, Anal. PDE **10** (2017), no. 5, 1017–1079.
- [9] N. Ghoussoub and X. S. Kang, *Hardy–Sobolev critical elliptic equations with boundary singularities*, Ann. Inst. H. Poincaré Anal. Non Linéaire **21** (2004), 767–793.
- [10] N. Ghoussoub and C. Yuan, *Multiple solutions for quasi-linear PDEs involving the critical Sobolev and Hardy exponents*, Trans. Amer. Math. Soc. **352** (2000), 5703–5743.
- [11] H. Jaber, *Mountain pass solutions for perturbed Hardy–Sobolev equations on compact manifolds*, Analysis (Berlin) **36** (2016), no. 4, 287–296.
- [12] ———, *Influence of Mean Curvature on Mountain-Pass Solutions for Hardy–Sobolev Equations*, Rocky Mountain J. Math. **49** (2019), no. 2, 505–519.
- [13] E. Jannelli, *The role played by space dimension in elliptic critical problems*, J. Differential Equations **156** (1999), no. 2, 407–426.
- [14] R. Schoen, *Conformal deformation of a Riemannian metric to constant scalar curvature*, J. Differential Geom. **20** (1984), no. 2, 479–495.

- [15] R. Schoen and S.-T. Yau, *Conformally flat manifolds, Kleinian groups and scalar curvature*, Invent. Math. **92** (1988), no. 1, 47–71.

DÉPARTEMENT DE MATHÉMATIQUE, UNIVERSITÉ LIBRE DE BRUXELLES, CP 214, BOULEVARD DU TRIOMPHE, B-1050 BRUXELLES, BELGIUM; INSTITUT ÉLIE CARTAN, UNIVERSITÉ DE LORRAINE, BP 70239, F-54506 VANDŒUVRE-LÈS-NANCY, FRANCE

Email address: Hussein.cheikh-ali@ulb.ac.be; Hussein.cheikh-ali@univ-lorraine.fr