



HAL
open science

Perspective-12-Quadric: An analytical solution to the camera pose estimation problem from conic - quadric correspondences

Vincent Gaudillère, Gilles Simon, Marie-Odile Berger

► **To cite this version:**

Vincent Gaudillère, Gilles Simon, Marie-Odile Berger. Perspective-12-Quadric: An analytical solution to the camera pose estimation problem from conic - quadric correspondences. 2019. hal-02054882v3

HAL Id: hal-02054882

<https://hal.science/hal-02054882v3>

Preprint submitted on 5 Mar 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Perspective-12-Quadric: An analytical solution to the camera pose estimation problem from conic - quadric correspondences.

Vincent Gaudillière, Gilles Simon, Marie-Odile Berger

Inria Nancy Grand - Est / Loria, Nancy, France
{firstname}.{lastname}@inria.fr

Goal: To estimate the pose of a perspective camera given a set of n 3D quadrics in the world and their corresponding 2D projections (conics) in the image.

1 Problem statement.

- **2D:** Homogeneous quadratic form of a conic equation:

$$u^\top C_i u = 0,$$

where $u \in \mathbb{R}^3$ is the homogeneous vector of a generic 2D point belonging to the conic defined by the symmetric matrix $C_i \in \mathbb{R}^{3 \times 3}$. The conic has 5 degrees of freedom given by the 6 elements of the lower triangular part of the symmetric matrix C_i , except one for the scale. The dual conic is defined by $C_i^* = \text{adj}(C_i)$.

- **3D:** Homogeneous quadratic form of a quadric equation:

$$x^\top Q_i x = 0,$$

where $x \in \mathbb{R}^4$ is the homogeneous vector of a generic 3D point belonging to the quadric defined by the symmetric matrix $Q_i \in \mathbb{R}^{4 \times 4}$. The quadric has 9 degrees of freedom given by the 10 elements of the lower triangular part of the symmetric matrix Q_i , except one for the scale. The dual quadric is defined by $Q_i^* = \text{adj}(Q_i)$.

- **Projection equation:** Let's consider the projection matrix

$$P = K[R|t] = \begin{bmatrix} p_{11} & p_{12} & p_{13} & p_{14} \\ p_{21} & p_{22} & p_{23} & p_{24} \\ p_{31} & p_{32} & p_{33} & p_{34} \end{bmatrix}$$

The projection equation [1] is:

$$\beta_i C_i^* = P Q_i^* P^\top \tag{1}$$

2 Solution.

In order to recover P in closed form from the set of conic-quadratic pairs, we have to re-arrange Eq. (1) into a linear system [2, 3]. Let us define the operator $vec()$ that serialises all the elements of a generic matrix, and the operator $vech()$ that serialises the elements of the lower triangular part of a symmetric matrix. Then let us define $v_i^* = vech(Q_i^*)$ and $c_i^* = vech(C_i^*)$. Then let us arrange the product of the elements of P and P^\top in a single matrix $G \in \mathbb{R}^{6 \times 10}$ as follows:

$$G = D(P \otimes P)E$$

where \otimes is the Kronecker product, and matrices $D \in \mathbb{R}^{6 \times 9}$ and $E \in \mathbb{R}^{16 \times 10}$, are two matrices such that $vech(X) = Dvec(X)$ and $vec(Y) = Evech(Y)$ respectively, where $X \in \mathbb{R}^{9 \times 9}$ and $Y \in \mathbb{R}^{16 \times 16}$ are two symmetric matrices.

$$D = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}, E = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

$$G = \begin{bmatrix} p_{11}^2 & 2p_{11}p_{12} & p_{12}^2 & 2p_{11}p_{13} & 2p_{12}p_{13} & p_{13}^2 & 2p_{11}p_{14} & 2p_{12}p_{14} & 2p_{13}p_{14} & p_{14}^2 \\ p_{11}p_{21} & p_{11}p_{22} + p_{12}p_{21} & p_{12}p_{22} & p_{11}p_{23} + p_{13}p_{21} & p_{12}p_{23} + p_{13}p_{22} & p_{13}p_{23} & p_{11}p_{24} + p_{14}p_{21} & p_{12}p_{24} + p_{14}p_{22} & p_{13}p_{24} + p_{14}p_{23} & p_{14}p_{24} \\ p_{21}^2 & 2p_{21}p_{22} & p_{22}^2 & 2p_{21}p_{23} & 2p_{22}p_{23} & p_{23}^2 & 2p_{21}p_{24} & 2p_{22}p_{24} & 2p_{23}p_{24} & p_{24}^2 \\ p_{11}p_{31} & p_{11}p_{32} + p_{12}p_{31} & p_{12}p_{32} & p_{11}p_{33} + p_{13}p_{31} & p_{12}p_{33} + p_{13}p_{32} & p_{13}p_{33} & p_{11}p_{34} + p_{14}p_{31} & p_{12}p_{34} + p_{14}p_{32} & p_{13}p_{34} + p_{14}p_{33} & p_{14}p_{34} \\ p_{21}p_{31} & p_{21}p_{32} + p_{22}p_{31} & p_{22}p_{32} & p_{21}p_{33} + p_{23}p_{31} & p_{22}p_{33} + p_{23}p_{32} & p_{23}p_{33} & p_{21}p_{34} + p_{24}p_{31} & p_{22}p_{34} + p_{24}p_{32} & p_{23}p_{34} + p_{24}p_{33} & p_{24}p_{34} \\ p_{31}^2 & 2p_{31}p_{32} & p_{32}^2 & 2p_{31}p_{33} & 2p_{32}p_{33} & p_{33}^2 & 2p_{31}p_{34} & 2p_{32}p_{34} & 2p_{33}p_{34} & p_{34}^2 \end{bmatrix}$$

Given G we can rewrite Eq. (1) as a linear equation:

$$\beta_i c_i^* = G v_i^* \quad (2)$$

Eq. (2) can thus be solved using DLT algorithm, *e.g.* can be written:

$$0 = Bg,$$

where $g = \text{vec}(G) = [g_{11} \cdots g_{610}]^\top \in \mathbb{R}^{60}$, and $B \in \mathbb{R}^{15n \times 60}$ is the concatenation of matrices $b_i \in \mathbb{R}^{15 \times 60}$:

$$b_i = \begin{bmatrix} c_{2i} \mathbf{v}_i^\top & -c_{1i} \mathbf{v}_i^\top & \mathbf{0}_{10}^\top & \mathbf{0}_{10}^\top & \mathbf{0}_{10}^\top & \mathbf{0}_{10}^\top \\ \mathbf{0}_{10}^\top & c_{3i} \mathbf{v}_i^\top & -c_{2i} \mathbf{v}_i^\top & \mathbf{0}_{10}^\top & \mathbf{0}_{10}^\top & \mathbf{0}_{10}^\top \\ \mathbf{0}_{10}^\top & \mathbf{0}_{10}^\top & c_{4i} \mathbf{v}_i^\top & -c_{3i} \mathbf{v}_i^\top & \mathbf{0}_{10}^\top & \mathbf{0}_{10}^\top \\ \mathbf{0}_{10}^\top & \mathbf{0}_{10}^\top & \mathbf{0}_{10}^\top & c_{5i} \mathbf{v}_i^\top & -c_{4i} \mathbf{v}_i^\top & \mathbf{0}_{10}^\top \\ \mathbf{0}_{10}^\top & \mathbf{0}_{10}^\top & \mathbf{0}_{10}^\top & \mathbf{0}_{10}^\top & c_{6i} \mathbf{v}_i^\top & -c_{5i} \mathbf{v}_i^\top \\ -c_{3i} \mathbf{v}_i^\top & \mathbf{0}_{10}^\top & c_{1i} \mathbf{v}_i^\top & \mathbf{0}_{10}^\top & \mathbf{0}_{10}^\top & \mathbf{0}_{10}^\top \\ \mathbf{0}_{10}^\top & -c_{4i} \mathbf{v}_i^\top & \mathbf{0}_{10}^\top & c_{2i} \mathbf{v}_i^\top & \mathbf{0}_{10}^\top & \mathbf{0}_{10}^\top \\ \mathbf{0}_{10}^\top & \mathbf{0}_{10}^\top & -c_{5i} \mathbf{v}_i^\top & \mathbf{0}_{10}^\top & c_{3i} \mathbf{v}_i^\top & \mathbf{0}_{10}^\top \\ \mathbf{0}_{10}^\top & \mathbf{0}_{10}^\top & \mathbf{0}_{10}^\top & -c_{6i} \mathbf{v}_i^\top & \mathbf{0}_{10}^\top & c_{4i} \mathbf{v}_i^\top \\ c_{4i} \mathbf{v}_i^\top & \mathbf{0}_{10}^\top & \mathbf{0}_{10}^\top & -c_{1i} \mathbf{v}_i^\top & \mathbf{0}_{10}^\top & \mathbf{0}_{10}^\top \\ \mathbf{0}_{10}^\top & c_{5i} \mathbf{v}_i^\top & \mathbf{0}_{10}^\top & \mathbf{0}_{10}^\top & -c_{2i} \mathbf{v}_i^\top & \mathbf{0}_{10}^\top \\ \mathbf{0}_{10}^\top & \mathbf{0}_{10}^\top & c_{6i} \mathbf{v}_i^\top & \mathbf{0}_{10}^\top & \mathbf{0}_{10}^\top & -c_{4i} \mathbf{v}_i^\top \\ -c_{5i} \mathbf{v}_i^\top & \mathbf{0}_{10}^\top & \mathbf{0}_{10}^\top & \mathbf{0}_{10}^\top & c_{1i} \mathbf{v}_i^\top & \mathbf{0}_{10}^\top \\ \mathbf{0}_{10}^\top & -c_{6i} \mathbf{v}_i^\top & \mathbf{0}_{10}^\top & \mathbf{0}_{10}^\top & \mathbf{0}_{10}^\top & c_{2i} \mathbf{v}_i^\top \\ c_{6i} \mathbf{v}_i^\top & \mathbf{0}_{10}^\top & \mathbf{0}_{10}^\top & \mathbf{0}_{10}^\top & \mathbf{0}_{10}^\top & -c_{1i} \mathbf{v}_i^\top \end{bmatrix},$$

with:

$$\begin{aligned} \mathbf{c}_i^\top &= [c_{1i} \ c_{2i} \ c_{3i} \ c_{4i} \ c_{5i} \ c_{6i}], \\ \mathbf{v}_i^\top &= [v_{1i} \ v_{2i} \ v_{3i} \ v_{4i} \ v_{5i} \ v_{6i} \ v_{7i} \ v_{8i} \ v_{9i} \ v_{10i}], \\ \mathbf{0}_{10}^\top &= [0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0] \end{aligned}$$

Each pair of conic-quadric provides 15 homogeneous linear equations in the unknown elements of G . However, only 5 are linearly independent. Thus 12 pairs of 3D-2D correspondences are required to solve the system. A total least squares solution can then be found by choosing g as a right singular vector corresponding to the smallest singular value of B . Finally, P can be inferred from G .

References

- [1] R.I. Hartley and A. Zisserman, *Multiple View Geometry in Computer Vision*, Cambridge University Press, second edition, 2004.
- [2] M. Crocco and C. Rubino and A. Del Bue, *Structure from Motion with Objects*, CVPR, 2016.
- [3] C. Rubino and M. Crocco and A. Del Bue, *3D Object Localisation from Multi-View Image Detections*, TPAMI, 2018.