



HAL
open science

Design, synthesis and evaluation of 2-aryl benzoxazoles as promising hit for the A2A receptor

Romain Duroux, Nicolas Renault, Joana Esteves, Laurence Agouridas, David
Blum, Luisa Lopes, Patricia Melnyk, Saïd Yous

► **To cite this version:**

Romain Duroux, Nicolas Renault, Joana Esteves, Laurence Agouridas, David Blum, et al.. Design, synthesis and evaluation of 2-aryl benzoxazoles as promising hit for the A2A receptor. *Journal of Enzyme Inhibition and Medicinal Chemistry*, 2017, 32 (1), pp.850-864. 10.1080/14756366.2017.1334648 . hal-02060508

HAL Id: hal-02060508

<https://hal.science/hal-02060508>

Submitted on 20 Dec 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

RESEARCH PAPER



Design, synthesis and evaluation of 2-aryl benzoxazoles as promising hit for the A_{2A} receptor

Romain Duroux^a, Nicolas Renault^b, Joana Esteves Cuelho^c, Laurence Agouridas^a, David Blum^a, Luisa V. Lopes^c, Patricia Melnyk^a and Saïd You^a

^aINSERM, CHU Lille, UMR-S 1172 – JPArc – Centre de Recherche Jean-Pierre AUBERT Neurosciences et Cancer, Université de Lille, Lille, France; ^bINSERM, CHU Lille, U995 – LIRIC – Lille Inflammation Research International Center, Université de Lille, Lille, France; ^cInstituto de Medecina Molecular, Lisbon, Portugal

ABSTRACT

The development of adenosine A_{2A} receptor antagonists has received much interest in recent years for the treatment of neurodegenerative diseases. Based on docking studies, a new series of 2-arylbenzoxazoles has been identified as potential A_{2A}R antagonists. Structure-affinity relationship was investigated in position 2, 5 and 6 of the benzoxazole heterocycle leading to compounds with a micromolar affinity towards the A_{2A} receptor. Compound **F1**, with an affinity of 1 μm, presented good absorption, distribution, metabolism and excretion properties with an excellent aqueous solubility (184 μm) without being cytotoxic at 100 μm. This compound, along with low-molecular weight compound **D1** (K_i = 10 μm), can be easily modulated and thus considered as relevant starting points for further hit-to-lead optimisation.

ARTICLE HISTORY

Received 24 March 2017
Revised 10 May 2017
Accepted 18 May 2017

KEYWORDS

Benzoxazole; A_{2A} receptor; solubility; neurodegenerative disease

Introduction

Alzheimer's (AD) and Parkinson's diseases (PD), currently the most important neurodegenerative pathologies, are characterised by a progressive neuronal death¹. Current therapies are restricted to symptomatic interventions and do not prevent progressive neuronal loss. Therefore, novel therapeutic solutions are needed and one of the promising strategies consists of targeting the adenosine A_{2A} receptor.

Epidemiological studies have shown that people consuming regularly over a lifetime caffeine-based beverages are substantially less likely to develop PD and AD^{2,3}. Caffeine exerts its biological effects primarily by antagonising adenosine receptors (GPCRs). The adenosine A_{2A} receptor subtype, one of the four adenosine receptors, was shown in multiple studies to be responsible for the neuroprotective effects of caffeine in experimental models of AD and PD^{4–6}. Besides, many A_{2A} antagonists have been synthesised over the past few years and some showed promising results in managing cognitive dysfunction in both diseases⁷. For example, antagonists (Figure 1) such as Istradefylline (KW-6002), Preladenant (SCH 4208⁸) and Tozadenant (SYN 115)⁹ have been investigated clinically in PD with promising results, especially Istradefylline which has been approved in Japan as an adjunct to L-DOPA therapy^{10,11}. Less research work has been undertaken regarding AD but it is now well established that A_{2A} antagonists lead to the improvement of spatial memory accompanied by the decrease of Aβ amyloid burden, Tau hyperphosphorylation and neurotoxicity^{6,12}.

Therefore, developing A_{2A} antagonists constitutes a promising therapeutic strategy for the treatment of both AD and PD. However, although many antagonists have been developed so far,



constant drawbacks remain such as high toxicity and poor solubility^{7,8,13,14}. These important limitations have obstructed the development of drugs targeting this receptor. Therefore, one of the main challenges regarding A_{2A} antagonist development is to improve solubility and lower toxicity while keeping good affinity at A_{2A} receptor. Selectivity parameters are now more debated since studies have highlighted the therapeutic potential of dual A₁/A_{2A} antagonists¹⁵ as well as a non-selective ligand proven by caffeine, for neurodegenerative disease.


With the aim of developing novel A_{2A} antagonists with good water solubility, we designed a series of benzoxazoles bearing a protonable amine function (Figure 2(A)). We first focused on a diffuse hydrophobic zone located at the top of the active site that is generally occupied by an aryl group in well-known antagonists (Figure 2(B)). Because of the presence of an acidic cluster (Glu169, Asp170) in this pocket, a tertiary amine which can allow for an ionic interaction could be a good alternative to an aryl group (Figure 2(C)). The present work describes the medicinal chemistry approach leading to a series of 2-arylbenzoxazoles which best compounds display micromolar affinity towards the A_{2A} receptor and good water solubility.

Methods

Chemistry

All reagents and solvents were purchased and used without further purification. Reactions were monitored by TLC performed on MachereyNagel Alugram[®] Sil 60/UV254 sheets (thickness 0.2 mm). Some purification of products was carried out by column

CONTACT Saïd You  said.yous@univ-lille2.fr  Jean-Pierre Aubert Research Center, UMR-S1172, Faculté de Pharmacie (Université Lille), 3 rue du Pr Laguesse, BP 83, 59006 Lille Cedex, France

 Supplemental data for this article can be accessed [here](#).

© 2017 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.




Figure 1. Selective adenosine A_{2A} receptor antagonists.




Figure 2. Molecular modelling-guided design. (A) Representation of various modulations around the benzoxazole scaffold. (B) Predicted binding mode of ZM-241385 in the apoA2AR-T4E pocket (dark) compared with the X-ray binding mode (gray). (C) Putative binding mode of compound F1 in the apoA2AR-T4E pocket.

chromatography using MachereyNagel silica gel (230e400 mesh). Melting points were determined on a BÜCHI B-540 apparatus and are uncorrected. NMR spectra were recorded on a Bruker Avance 300 spectrometer operating at 300 MHz (^1H) or 75 MHz (^{13}C). Chemical shifts are in parts per million (ppm) and were referenced to the residual proton peaks in deuterated solvents. Mass spectra were recorded with an LCMS (Waters Alliance Micromass ZQ 2000). LCMS analysis was performed using a Waters XBridge C18 column (5 μm particle size column, dimensions 50 \times 4.6 mm). A gradient starting from 98% H_2O /formate buffer 5 mm (pH 3.8) and reaching 100% CH_3CN /formate buffer 5 mm (pH 3.8) within 4 min at a flow rate of 2 ml/min was used followed by a return to the starting conditions within 1 min. FT-IR spectra were recorded on a Thermo Nicolet Avatar 320 FT-IR spectrometer. The purity of final compounds was verified by high-pressure liquid chromatography (HPLC) columns: C4 Interchrom UPTISPHERE. Analytical HPLC was performed on a Shimadzu LC-2010AHT system equipped with a UV detector set at 254 and 215 nm. Compounds were dissolved in 50 ml acetonitrile and 950 ml buffer A and injected into the system. The following eluent systems were used: buffer A (H_2O /TFA, 100:0.1) and buffer B (CH_3CN / H_2O /TFA, 80:20:0.1). HPLC retention times (HPLC tR) were obtained at a flow rate of 0.2 ml/min for 35 min using the following conditions: a gradient run from 100% of buffer A over 1 min, then to 100% of buffer B over the next 30 min.

General procedure for the synthesis of amide (2a–2d)

To a solution of acid (17.8 mmol) in DCM (100 ml) at 0 °C were added thionyl chloride (71.4 mmol) and 5 drops of DMF. This solution was stirred for 3 h at reflux, cooled to room temperature and concentrated *in vacuo*. The residue was diluted in 50 ml of EtOAc and added to a solution of aminophenol (16.6 mmol) and Et_3N (35.6 mmol) in 35 ml of EtOAc at 0 °C. The reaction mixture was stirred at room temperature overnight, hydrolysed with water and extracted twice with EtOAc. An acid–base workup with saturated NaHCO_3 and 1 M HCl solution was performed and the organic layer was concentrated *in vacuo*. The solid was then suspended in a mixture of EtOH/ H_2O (250 ml/10 ml) and NaOH was added (54 mmol). The reaction mixture was stirred at reflux for 4 h, cooled to room temperature, acidified slowly with 6 M HCl up to acidic pH. Resulting solid was filtered, washed with water and recrystallised in an appropriate solvent.

N-(2-Hydroxy-5-methylphenyl)furan-2-carboxamide (2a): The title compound was prepared from 2-furoic acid and 4-methyl-2-aminophenol to afford **2a** as a beige solid (80%): mp 186 °C (acetonitrile). ^1H NMR (300 MHz, $[\text{D}_6]\text{DMSO}$): 9.76 (br s, 1H, OH), 9.09 (br s, 1H, NH), 7.92 (m, 1H, H_5), 7.72 (m, 1H, H_6), 7.28 (dd, 1H, H_4 , $J = 2.2$ Hz and $J = 8.4$ Hz), 6.79 (m, 2H, H_3 and H_3), 6.70 (dd, 1H, H_4 , $J = 1.7$ Hz and $J = 3.5$ Hz), 2.21 (s, 3H, CH_3). IR (ν , cm^{-1}): 3380 (NH), 2750–3100 (OH), 1640 (C=O). LC-MS (ESI) m/z found: 218 $[\text{M} + \text{H}]^+$.

N-(2-Hydroxy-5-methylphenyl)-3,4-dimethoxybenzamide (2b): The title compound was prepared from 3,4-dimethoxybenzoic acid and 4-methyl-2-aminophenol to afford **2b** as a white solid (82%): mp 164 °C (acetonitrile). ^1H NMR (300 MHz, CDCl_3): 9.47 (br s, 1H, OH), 9.41 (br s, 1H, NH), 7.61 (dd, 1H, H_4 , $J = 2.0$ Hz and $J = 8.4$ Hz), 7.55 (d, 1H, H_6 , $J = 2.0$ Hz), 7.42 (m, 1H, H_2), 7.07 (d, 1H, H_3 , $J = 8.4$ Hz), 6.83 (dd, 1H, H_6 , $J = 1.7$ Hz and $J = 8.3$ Hz), 6.79 (d, 1H, H_5 , $J = 8.3$ Hz), 3.84 (s, 3H, OMe), 3.82 (s, 3H, OMe), 2.22 (s, 3H, CH_3). IR (ν , cm^{-1}): 3368 (NH), 3182–2854 (OH), 1653 (C=O). LC-MS (ESI) m/z found: 288 $[\text{M} + \text{H}]^+$.

N-(2-Hydroxy-4-methylphenyl)furan-2-carboxamide (2c): The title compound was prepared from furoic acid and 5-methyl-2-aminophenol to afford **2c** as a brown solid (72%): mp 170 °C (acetonitrile).

^1H NMR (300 MHz, CDCl_3): 8.85 (br s, 1H, OH), 8.34 (br s, 1H, NH), 7.50 (m, 1H, H_5), 7.28 (m, 1H, H_3), 7.10 (d, 1H, H_6 , $J = 8.1$ Hz), 6.87 (m, 1H, H_3), 6.72 (m, 1H, H_5), 6.58 (dd, 1H, H_4 , $J = 1.7$ Hz and $J = 3.5$ Hz), 2.30 (s, 3H, CH_3). IR (ν , cm^{-1}): 3388 (NH), 3182–2854 (OH), 1649 (C=O).

N-(2-Hydroxy-5-nitro-phenyl) furan-2-carboxamide (2d): The title compound was prepared from furoic acid and 4-nitro-2-aminophenol to afford **2d** as a yellow solid (84%): mp 165 °C (methanol). ^1H NMR (300 MHz, CDCl_3): 9.77 (br s, 1H, OH), 9.13 (br s, 1H, NH), 8.79 (d, 1H, H_6 , $J = 1.9$ Hz), 8.25 (dd, 1H, H_4 , $J = 1.9$ Hz and $J = 8.6$ Hz), 8.07 (m, 1H, H_5), 7.81 (d, 1H, H_3 , $J = 3.5$ Hz), 7.32 (d, 1H, H_3 , $J = 8.6$ Hz), 6.78 (dd, 1H, H_4 , $J = 1.7$ Hz and $J = 3.5$ Hz). IR (ν , cm^{-1}): 3376 (NH), 3000–2500 (OH), 1640 (C=O).

General procedure for the synthesis of benzoxazole (3a–3d)

A suspension of amide **2** (2.21 mmol) and TsOH (5.53 mmol) was refluxed in toluene (150 ml) equipped with a Dean–Stark apparatus until complete dissolution for 17 h. The solution was then cooled to room temperature, hydrolysed with water and basified with a 6 M solution of NaOH up to basic pH (10–12). The organic layer was separated, dried over K_2CO_3 and concentrated *in vacuo*. Solid was suspended from the appropriate solvent and filtered.

2-(Furan-2-yl)-5-methyl-1,3-benzoxazole (3a): The title compound was prepared from amide **2a** to afford **3a** as a beige solid (82%): mp 64 °C (petroleum ether). ^1H NMR (300 MHz, CDCl_3): 7.67 (m, 1H, H_5), 7.54 (m, 1H, H_3), 7.44 (d, 1H, H_7 , $J = 8.3$ Hz), 7.25 (m, 1H, H_4), 7.17 (m, 1H, H_6), 6.70 (dd, 1H, H_4 , $J = 1.7$ Hz and $J = 3.5$ Hz), 2.49 (s, 3H, CH_3). LC-MS (ESI) m/z found: 200 $[\text{M} + \text{H}]^+$.

2-(3,4-Dimethoxyphenyl)-5-methyl-1,3-benzoxazole (3b): The title compound was prepared from amide **2b** to afford **3b** as a beige solid (80%): mp 136 °C (diethyl ether). ^1H NMR (300 MHz, CDCl_3): 7.83 (dd, 1H, H_6 , $J = 1.9$ Hz and $J = 8.4$ Hz), 7.75 (d, 1H, H_2 , $J = 1.9$ Hz), 7.53 (m, 1H, H_6), 7.43 (d, 1H, H_5 , $J = 8.4$ Hz), 7.13 (m, 1H, H_4), 6.98 (d, 1H, H_7 , $J = 8.4$ Hz), 4.02 (s, 3H, OMe), 3.97 (s, 3H, OMe), 2.48 (s, 3H, CH_3). LC-MS (ESI) m/z found: 270 $[\text{M} + \text{H}]^+$.

2-(Furan-2-yl)-6-methyl-1,3-benzoxazole (3c): The title compound was prepared from amide **2c** to afford **3c** as a beige solid (68%): mp 54 °C (diethyl ether). ^1H NMR (300 MHz, CDCl_3): 8.04 (m, 1H, H_5), 7.61 (d, 1H, H_4 , $J = 8.1$ Hz), 7.40 (d, 1H, H_3 , $J = 3.5$ Hz), 7.38 (m, 1H, H_7), 7.27–7.18 (m, 1H, H_5), 6.79 (dd, 1H, H_4 , $J = 1.7$ Hz and $J = 3.5$ Hz), 2.44 (s, 3H, CH_3). LC-MS (ESI) m/z found: 200 $[\text{M} + \text{H}]^+$.

2-(Furan-2-yl)-5-nitro-1,3-benzoxazole (3d): The title compound was prepared from amide **2b** to afford **3d** as a yellow solid (82%): mp 182 °C (diethyl ether). ^1H NMR (300 MHz, CDCl_3): 8.62 (d, 1H, H_4 , $J = 2.1$ Hz), 8.34 (dd, 1H, H_6 , $J = 2.1$ Hz and $J = 8.4$ Hz), 7.75 (d, 1H, H_7 , $J = 8.4$ Hz), 7.68 (m, 1H, H_5), 7.39 (d, 1H, H_3 , $J = 3.5$ Hz), 6.68 (dd, 1H, H_4 , $J = 1.7$ Hz and $J = 3.5$ Hz). IR (ν , cm^{-1}): 1519 (Ar-NO_2). LC-MS (ESI) m/z found: 231 $[\text{M} + \text{H}]^+$.

Methyl 2-[2-(3,4-dimethoxyphenyl)-1,3-benzoxazol-5-yl] acetate (3e): To a suspension of methyl 2-(3-amino-4-hydroxyphenyl) acetate (1.6 g, 8.83 mmol) in T_3P (solution in EtOAc) (4.2 g, 13.3 mmol), were added 3,4-dimethoxybenzoic acid (1.61 g, 8.83 mmol) and DIPEA (1.46 ml, 8.83 mmol). The reaction mixture was heated overnight at 120 °C, cooled to room temperature, suspended in water and extracted three times with EtOAc. Combined organic layers were washed 1 M NaOH solution, dried over MgSO_4 and concentrated *in vacuo*. Solid was recrystallised from EtOH to afford compound **3e** as a beige solid (1.59 g, 55%): mp 110 °C. ^1H NMR (300 MHz, CDCl_3): 7.82 (dd, 1H, H_6 , $J = 8.4$ Hz and $J = 1.8$ Hz), 7.74 (d, 1H, H_4 , $J = 1.8$ Hz), 7.64 (m, 1H, H_2), 7.49 (d, 1H, H_7 , $J = 8.4$ Hz), 7.24 (dd, 1H, H_6 , $J = 8.4$ Hz and $J = 1.8$ Hz), 6.98 (d, 1H, H_5 , $J = 8.4$ Hz), 4.01 (s, 3H, OMe), 3.95 (s, 3H, OMe), 3.75 (s, 2H, CH_2),

3.70 (s, 3H, CO₂Me). IR (ν , cm⁻¹): 1730 (ester C=O). LC-MS (ESI) *m/z* found: 328 [M + H]⁺.

General procedure for the synthesis of compound (4a–4c)

To a solution of compound (**3a–3c**) (20.1 mmol) in CCl₄ (150 ml) was added *N*-bromosuccinimide (NBS) (24.1 mmol) and benzoyl peroxide (1.41 mmol) and the reaction mixture was refluxed under a halogen lamp (230 W). After 3 h and 30 min stirring, the mixture was cooled to room temperature and the succinimide was filtered off. Then, the solution was concentrated *in vacuo*, and the solid was suspended in diethyl ether and filtered.

5-(Bromomethyl)-2-(furan-2-yl)-1,3-benzoxazole (**4a**): The title compound was prepared from benzoxazole **3a** to afford **4a** as a beige solid (70%): mp 130 °C. ¹H NMR (300 MHz, CDCl₃): 7.77 (m, 1H, H₅), 7.67 (m, 1H, H₃), 7.54 (d, 1H, H₇, *J* = 8.4 Hz), 7.42 (dd, 1H, H₆, *J* = 1.8 Hz and *J* = 8.4 Hz), 7.30 (m, 1H, H₄), 6.64 (dd, 1H, H₄, *J* = 1.7 Hz and *J* = 3.5 Hz), 4.64 (s, 2H, CH₂). LC-MS (ESI) *m/z* found: 278 [M + H]⁺, 280 [M + H]⁺.

6-(Bromomethyl)-2-(furan-2-yl)-1,3-benzoxazole (**4b**): The title compound was prepared from benzoxazole **3b** to afford **4b** as a brown solid (65%): mp 122 °C. ¹H NMR (300 MHz, CDCl₃): 7.72–7.68 (m, 2H, H₅ and H₄), 7.60 (m, 1H, H₇), 7.40 (dd, 1H, H₅, *J* = 8.2 Hz and *J* = 1.6 Hz), 7.30 (dd, 1H, H₃, *J* = 0.7 Hz and *J* = 3.5 Hz), 6.63 (dd, 1H, H₄, *J* = 1.7 Hz and *J* = 3.5 Hz), 4.64 (s, 2H, CH₂). IR (ν , cm⁻¹): 750 (C-Br).

5-(Bromomethyl)-2-(3,4-dimethoxyphenyl)-1,3-benzoxazole (**4c**): The title compound was prepared from benzoxazole **3c** to afford **4c** as a beige solid (75%): mp 140 °C. ¹H NMR (300 MHz, CDCl₃): 7.86 (dd, 1H, H₆, *J* = 1.9 Hz and *J* = 8.4 Hz), 7.76 (m, 2H, H₄ and H₂), 7.53 (d, 1H, H₇ or H₅, *J* = 8.5 Hz), 7.39 (dd, 1H, H₆, *J* = 1.7 Hz and *J* = 8.4 Hz), 7.00 (d, 1H, H₇ or H₅, *J* = 8.4 Hz), 4.65 (s, 2H, CH₂), 4.02 (s, 3H, OMe), 3.99 (s, 3H, OMe). IR (ν , cm⁻¹): 750 (C-Br).

General procedure for the synthesis of compound (A1–A3 and A5–A8)

To a solution of amine (2.05 mmol) in acetone (15 ml) were added compound (**4a–4c**) (1.87 mmol) and Et₃N (2.05 mmol). The reaction mixture was refluxed for 1 h, cooled to room temperature and concentrated *in vacuo*. Solid was suspended in water and extracted three times with EtOAc. Combined organic layers were dried over MgSO₄ and concentrated *in vacuo* to give a solid which was then purified.

2-(Furan-2-yl)-5-(piperidin-1-ylmethyl)-1,3-benzoxazole hydrochloride (**A1**): The title compound was prepared from compound **4a** and piperidine. Solid was suspended in diethyl ether with HCl gas, concentrated *in vacuo* and recrystallised from acetonitrile to afford **A1** as a white solid (74%): mp >300 °C. ¹H NMR (300 MHz, [D₆]DMSO): 12.43 (br s, 1H, NH⁺), 8.02 (m, 1H, H₆), 7.75 (m, 1H, H₅), 7.70 (m, 1H, H₄), 7.66 (d, 1H, H₇, *J* = 8.1 Hz), 7.32 (d, 1H, H₃, *J* = 3.5 Hz), 6.65 (dd, 1H, H₄, *J* = 1.7 Hz and *J* = 3.5 Hz), 4.26 (d, 2H, CH₂, *J* = 5.0 Hz), 3.47 (m, 2H, H_{piperidine}), 2.61 (m, 2H, H_{piperidine}), 2.33 (m, 2H, H_{piperidine}), 1.92–1.80 (m, 3H, H_{piperidine}), 1.35 (m, 1H, H_{piperidine}). ¹³C NMR (75 MHz, [D₆]DMSO): 156.5 (C), 151.0 (C), 146.3 (CH), 142.1 (C), 142.0 (C), 129.1 (CH), 125.1 (C), 122.9 (CH), 115.3 (CH), 112.5 (CH), 111.8 (CH), 60.9 (CH₂), 52.3 (2 CH₂), 22.5 (2 CH₂), 22.1 (CH₂). LC-MS (ESI) *m/z* found: 283 [M + H]⁺. HPLC: C₄ column: t_R = 16.2 min, purity 98%.

2-(Furan-2-yl)-5-[(4-phenylpiperazin-1-yl)methyl]-1,3-benzoxazole hydrochloride (**A2**): The title compound was prepared from compound **4a** and phenylpiperazine. Solid was suspended in diethyl ether with HCl gas, concentrated *in vacuo* and recrystallised from ethanol to afford **A2** as a white solid (65%): mp

>300 °C. ¹H NMR (300 MHz, [D₆]DMSO): 13.12 (br s, 1H, NH⁺), 8.04 (m, 1H, H₅), 7.83 (m, 1H, H₆), 7.68 (m, 2H), 7.33 (d, 1H, H₃, *J* = 3.5 Hz), 7.30–7.25 (m, 2H), 6.98–6.89 (m, 3H), 6.66 (dd, 1H, H₄, *J* = 1.7 Hz and *J* = 3.5 Hz), 4.37 (s, 2H, CH₂), 3.75–3.52 (m, 6H, H_{piperazine}), 3.06 (m, 2H, H_{piperazine}). ¹³C NMR (75 MHz, [D₆]DMSO): 156.0 (C), 150.6 (C), 150.0 (C), 148.0 (CH), 141.9 (C), 141.7 (C), 129.6 (2 CH), 129.4 (CH), 127.5 (C), 123.5 (CH), 120.4 (CH), 116.4 (2 CH), 116.2 (CH), 113.4 (CH), 111.6 (CH), 58.7 (CH₂), 50.6 (2 CH₂), 45.7 (2 CH₂). LC-MS (ESI) *m/z* Found: 360 [M + H]⁺. HPLC: C₄ column: t_R = 15.5 min, purity >99%. tert-Butyl-4-[[2-(Furan-2-yl)-1,3-benzoxazol-5-yl]methyl]piperazine-1-carboxylate (**A3**): The title compound was prepared from compound **4a** and boc-piperazine. Solid was recrystallised from methanol to afford **A3** as a white solid (83%): mp 128 °C (methanol). ¹H NMR (300 MHz, CDCl₃): 7.69 (m, 1H, H₅), 7.67 (m, 1H, H₄), 7.49 (d, 1H, H₇, *J* = 8.3 Hz), 7.34 (dd, 1H, H₆, *J* = 1.5 Hz and *J* = 8.3 Hz), 7.26 (m, 1H, H₃), 6.62 (dd, 1H, H₄, *J* = 1.7 Hz and *J* = 3.5 Hz), 3.63 (s, 2H, CH₂), 3.44 (t, 4H, H_{piperazine}, *J* = 4.9 Hz), 2.42 (t, 4H, H_{piperazine}, *J* = 3.5 Hz), 1.45 (s, 9H, (3 CH₃)). NMR ¹³C (CDCl₃, δ ppm): 163.0 (C), 155.6 (C), 154.0 (C), 149.8 (C), 147.7 (C), 141.9 (C), 141.8 (CH), 132.2 (CH), 127.9 (C), 121.5 (CH), 115.8 (CH), 113.3 (CH), 111.2 (CH), 79.6 (2 CH₂), 60.8 (CH₂), 51.9 (2 CH₂), 26.5 (3 CH₃). IR (ν , cm⁻¹): 1679 (C=O). LC-MS (ESI) *m/z* found: 328 [M-tBu + H]⁺, 284 [M + H]⁺. HPLC: C₄ column: t_R = 14.7 min, purity >99%.

4-[[2-(Furan-2-yl)-1,3-benzoxazol-5-yl]methyl]piperazin-1-ylphenol (**A5**): The title compound was prepared from compound **4a** and 4-piperazinephenol. Solid was purified by flash chromatography using DCM/EtOAc (9/1) to afford **A5** as a white solid (83%): mp 212 °C. ¹H NMR (300 MHz, [D₆]DMSO): 8.79 (br s, 1H, OH), 8.07 (m, 1H, H₅), 7.71 (m, 2H, H₇ and H₄), 7.46 (d, 1H, H₃, *J* = 3.5 Hz), 7.41 (dd, 1H, H₆, *J* = 1.2 Hz and *J* = 8.4 Hz), 6.82 (dd, 1H, H₄, *J* = 1.8 Hz and *J* = 3.5 Hz), 6.78 (d, 2H, H_{phenyl}, *J* = 8.7 Hz), 6.63 (d, 2H, H_{phenyl}, *J* = 8.7 Hz), 3.64 (s, 2H, CH₂), 2.96 (m, 4H, H_{piperazine}), 2.55 (m, 4H, H_{piperazine}). ¹³C NMR (75 MHz, [D₆]DMSO): 155.4 (C), 151.3 (C), 149.2 (C), 147.6 (CH), 144.7 (C), 142.1 (C), 141.7 (C), 135.9 (C), 126.9 (CH), 120.2 (CH), 118.2 (2 CH), 115.9 (2 CH), 115.5 (CH), 113.3 (CH), 110.8 (CH), 62.2 (CH₂), 53.2 (2 CH₂), 50.5 (2 CH₂). IR (ν , cm⁻¹): 3195–2780 (OH). LC-MS (ESI) *m/z* found: 376 [M + H]⁺. HPLC: C₄ column: t_R = 10.2 min, purity 99% C₁₈ column: t_R = 14.3 min, purity >99%.

2-(Furan-2-yl)-5-[(4-[(2-methoxyethoxy)phenyl]piperazin-1-yl)methyl]-1,3-benzoxazole (**A6**): The title compound was prepared from compound **4a** and 1-[4-(2-methoxyethoxy)phenyl]piperazine. Solid was recrystallised from ethanol to afford **A6** as a white solid (33%): mp 123 °C. RMN ¹H (300 MHz, [D₆]DMSO): 8.08 (m, 1H, H₅), 7.72 (m, 2H, H₇ and H₄), 7.46 (dd, 1H, H₃, *J* = 0.7 Hz and *J* = 3.5 Hz), 7.40 (dd, 1H, H₆, *J* = 1.4 Hz and *J* = 8.4 Hz), 6.88–6.79 (m, 5H, H₄, H_{phenyl}), 3.99 (m, 2H, OCH₂), 3.64 (s, 2H, CH₂), 3.61 (m, 2H, CH₂O), 3.29 (s, 3H, OCH₃), 3.01 (m, 4H, H_{piperazine}), 2.54 (m, 4H, H_{piperazine}). RMN ¹³C (75 MHz, [D₆]DMSO): 155.4 (C), 152.5 (C), 149.2 (CH), 147.5 (C), 145.9 (C), 142.1 (C), 141.7 (C), 135.9 (C), 126.1 (C), 119.9 (CH), 118.1 (2 CH), 115.4 (2 CH), 114.1 (CH), 112.3 (CH), 110.2 (CH), 71.2 (CH₂), 67.7 (CH₂), 60.8 (CH₂), 53.4 (2 CH₂), 50.6 (2 CH₂), 33.7 (CH₃). LC-MS (ESI) *m/z* found: 434 [M + H]⁺. HPLC: C₄ column: t_R = 16.1 min, purity >99%.

2-(Furan-2-yl)-6-(piperidin-1-ylmethyl)-1,3-benzoxazole hydrochloride (**A7**): The title compound was prepared from compound **4c** and piperidine. Solid was suspended in diethyl ether with HCl gas, concentrated *in vacuo* and recrystallised from acetonitrile to afford **A7** (276 mg, 68%): mp >300 °C. ¹H NMR (300 MHz, [D₆]DMSO): 11.08 (br s, 1H, NH⁺), 8.16–8.10 (m, 2H, H₇ and H₅), 7.82 (d, 1H, H₄, *J* = 8.1 Hz), 7.63 (m, 1H, H₅), 7.51 (d, 1H, H₃, *J* = 3.2 Hz), 6.83 (m, 1H, H₄), 4.38 (m, 2H, CH₂), 3.28 (m, 2H, H_{piperidine}), 2.83 (m, 2H, H_{piperidine}), 1.76–1.66 (m, 5H, H_{piperidine}),

1.35 (m, 1H, $H_{\text{piperidine}}$). ^{13}C NMR (75 MHz, $[\text{D}_6]\text{DMSO}$): 156.1 (C), 149.9 (C), 148.0 (CH), 142.4 (C), 141.7 (C), 129.0 (C), 127.8 (CH), 120.1 (CH), 116.3 (CH), 114.4 (CH), 113.4 (CH), 59.1 (CH_2), 52.0 (2 CH_2), 22.6 (2 CH_2), 21.9 (CH_2). LC-MS (ESI) m/z found: 283 $[\text{M} + \text{H}]^+$. HPLC: C_4 column: $t_{\text{R}} = 15.7$ min, purity 96%.

2-(3,4-Dimethoxyphenyl)-5-(piperidin-1-ylmethyl)-1,3-benzoxazole hydrochloride (**A8**): The title compound was prepared from compound **4b** and piperidine. Solid was suspended in diethyl ether with HCl gas, concentrated *in vacuo* and recrystallised from acetonitrile to afford **A8** as a white solid (200 mg, 60%): mp $>300^\circ\text{C}$, ^1H NMR (300 MHz, $[\text{D}_6]\text{DMSO}$): 12.31 (br s, 1H, NH^+), 7.90 (dd, 1H, H_6 , $J = 1.6$ Hz and $J = 8.4$ Hz), 7.82 (dd, 1H, H_6 , $J = 1.9$ Hz and $J = 8.4$ Hz), 7.78 (d, 1H, H_2 , $J = 1.6$ Hz), 7.71 (d, 1H, H_4 , $J = 1.9$ Hz), 7.62 (d, 1H, H_5 , $J = 8.4$ Hz), 6.97 (d, 1H, H_7 , $J = 8.4$ Hz), 4.27 (d, 2H, CH_2 , $J = 5.1$ Hz), 3.99 (s, 3H, OMe), 3.96 (s, 3H, OMe), 3.45 (m, 2H, $H_{\text{piperidine}}$), 2.64 (m, 2H, $H_{\text{piperidine}}$), 2.34 (m, 2H, $H_{\text{piperidine}}$), 1.86 (m, 3H, $H_{\text{piperidine}}$), 1.35 (m, 1H, $H_{\text{piperidine}}$). ^{13}C NMR (75 MHz, $[\text{D}_6]\text{DMSO}$): δ 164.4 (C), 152.4 (C), 151.5 (C), 149.3 (C), 142.6 (C), 128.4 (CH), 124.7 (C), 122.5 (CH), 121.5 (C), 119.0 (CH), 111.5 (CH), 111.1 (CH), 110.1 (CH), 60.9 (CH_2), 56.1 (CH_3), 56.1 (CH_3), 52.6 (2 CH_2), 22.5 (2 CH_2), 22.0 (CH_2). LC-MS (ESI) m/z found: 353 $[\text{M} + \text{H}]^+$. HPLC: C_4 column: $t_{\text{R}} = 14.3$ min, purity 99%.

Synthesis of 2-(Furan-2-yl)-5-(piperazin-1-ylmethyl)-1,3-benzoxazole (**A4**): A solution of compound (**A3**) (100 mg, 0.35 mmol) diluted in 5 ml of DCM with TFA (2 ml, 26 mmol) was stirred for 3 h at room temperature, hydrolysed with water and basified with 1 M solution of NaOH up to basic pH and extracted three times with DCM. Combined organic layers were dried over MgSO_4 and concentrated *in vacuo*. Solid was suspended in diethyl ether with a drop of ethanol and filtered to afford compound (**A4**) as a beige solid (60%): mp 192°C . ^1H NMR (300 MHz, $[\text{D}_6]\text{DMSO}$): 8.08 (m, 1H, H_5), 7.74–7.71 (m, 2H, H_4 and H_7), 7.46 (dd, 1H, H_3 , $J = 0.7$ Hz and $J = 3.5$ Hz), 7.38 (dd, 1H, H_6 , $J = 1.6$ Hz and $J = 8.3$ Hz), 6.83 (dd, 1H, H_4 , $J = 1.7$ Hz and $J = 3.5$ Hz), 3.65 (s, 2H, CH_2), 3.45–3.39 (m, 4H, $H_{\text{piperazine}}$), 2.43 (m, 4H, $H_{\text{piperazine}}$). ^{13}C NMR (75 MHz, $[\text{D}_6]\text{DMSO}$): 155.4 (C), 149.3 (C), 147.6 (CH), 142.0 (C), 141.7 (C), 135.2 (C), 127.0 (CH), 120.3 (CH), 115.6 (CH), 113.3 (CH), 110.9 (CH), 61.9 (CH_2), 50.6 (2 CH_2), 44.1 (2 CH_2). LC-MS (ESI) m/z found: 284 $[\text{M} + \text{H}]^+$. HPLC: C_4 column: $t_{\text{R}} = 15.8$ min, purity 99%.

2-[2-(Furan-2-yl)-1,3-benzoxazol-5-yl]acetonitrile (**5**): To a solution of 5-(bromomethyl)-2-(3,4-dimethoxyphenyl)-1,3-benzoxazole (**4a**) (1.23 g, 3.53 mmol) in a mixture of EtOH (56 ml) and H_2O (15 ml) was added KCN (1.15 g, 17.7 mmol). After one night stirring at reflux, the reaction mixture was cooled to room temperature and concentrated *in vacuo*. Solid was suspended in water and extracted three times with EtOAc. Combined organic layer were dried over MgSO_4 and concentrated *in vacuo*. Solid was then recrystallised in methanol to afford (**5**) as a beige solid (582 mg, 56%): mp 140°C . ^1H NMR (300 MHz, CDCl_3): 7.71–7.70 (m, 2H, H_5 and H_3), 7.58 (d, 1H, H_7 , $J = 8.3$ Hz), 7.36–7.31 (m, 2H, H_4 and H_6), 6.64 (dd, 1H, H_4 , $J = 1.7$ Hz and $J = 3.5$ Hz), 3.90 (s, 2H, CH_2). IR (ν , cm^{-1}): 2240 (CN). LC-MS (ESI) m/z found: 225 $[\text{M} + \text{H}]^+$.

2-[2-(Furan-2-yl)-1,3-benzoxazol-5-yl] acetic acid (**6**): A solution of compound (**5**) (1.1 g, 4.91 mmol) in a mixture of AcOH (5.5 ml), H_2SO_4 (5.5 ml) and H_2O (5.5 ml) was refluxed for 17 h. After cooling to room temperature, cold water was added and the solution was extracted with EtOAc. The organic layer was extracted twice with a saturated NaHCO_3 solution, and then the aqueous layer was acidified with 1 M HCl solution and extracted twice with EtOAc. Combined organic layer were dried over MgSO_4 and concentrated *under vacuum*. Solid was suspended in diethyl ether and filtered to afford compound (**6**) as a white solid (702 mg, 60%): mp 203°C . ^1H NMR (300 MHz, CDCl_3): 12.38 (br s, 1H, OH), 8.07 (m, 1H, H_5), 7.68 (d, 1H, H_7 , $J = 8.4$ Hz), 7.66 (m, 1H, H_3), 7.45 (m, 1H, H_4), 7.32

(dd, 1H, H_6 , $J = 1.8$ Hz and $J = 8.4$ Hz), 6.80 (dd, 1H, H_4 , $J = 1.7$ Hz and $J = 3.5$ Hz), 3.72 (s, 2H, CH_2). IR (ν , cm^{-1}): 1710 (acid C=O). LC-MS (ESI) m/z found: 244 $[\text{M} + \text{H}]^+$.

General procedure for the synthesis of amide (7a–7d)

To a solution of acid (**6**) (1.5 mmol) in toluene (4 ml) was added at 0°C SOCl_2 (5.97 mmol). The mixture was refluxed during 1 h, cooled to room temperature and concentrated *in vacuo*. The oil was then diluted with EtOAc (26 ml) and added dropwise to a solution of amine (1.6 mmol) and Et_3N (2.25 mmol) in EtOAc (30 ml) while being stirred and cooled in an ice bath. After 1 h and 30 min stirring at room temperature, the mixture was hydrolysed with water, extracted twice with EtOAc and combined organic layers were washed with a saturated NaHCO_3 solution, brine, dried over MgSO_4 and concentrated *in vacuo* to give a solid which was suspended in diethyl ether and filtered.

2-[2-(Furan-2-yl)-1,3-benzoxazol-5-yl]-1-(piperidin-1-yl)ethan-1-one (**7a**): The title compound was prepared from compound **6** and piperidine to afford **7a** as a white solid (76%): mp 112°C . ^1H NMR (300 MHz, CDCl_3): 8.16 (d, 1H, H_5 , $J = 1.7$ Hz), 7.71 (m, 2H, H_4 and H_7), 7.68 (m, 1H, H_6), 7.34 (d, 1H, H_3 , $J = 3.5$ Hz), 6.81 (dd, 1H, H_4 , $J = 1.7$ Hz and $J = 3.5$ Hz), 3.78 (s, 2H, CH_2), 3.39 (m, 4H, $H_{\text{piperidine}}$), 1.68 (m, 4H, $H_{\text{piperidine}}$), 1.47 (m, 2H, $H_{\text{piperidine}}$). IR (ν , cm^{-1}): 1640 (C=O).

2-[2-(Furan-2-yl)-1,3-benzoxazol-5-yl]-1-(4-phenylpiperazin-1-yl)ethan-1-one (**7b**): The title compound was prepared from compound **6** and phenylpiperazine to afford **7b** as a white solid (78%): mp 128°C . ^1H NMR (300 MHz, CDCl_3): 7.68 (m, 1H, H_5), 7.63 (m, 1H, H_3), 7.52 (d, 1H, H_7 , $J = 8.4$ Hz), 7.32–7.23 (m, 4H, H_{phenyl}), 6.89–6.87 (m, 3H, H_4 , H_6 and H_{phenyl}), 6.63 (dd, 1H, H_4 , $J = 1.7$ Hz and $J = 3.5$ Hz), 3.91 (s, 2H, CH_2), 3.83 (t, 2H, $H_{\text{piperazine}}$, $J = 4.6$ Hz), 3.66 (t, 2H, $H_{\text{piperazine}}$, $J = 4.6$ Hz), 3.16 (t, 2H, $H_{\text{piperazine}}$, $J = 4.6$ Hz), 3.02 (t, 2H, $H_{\text{piperazine}}$, $J = 4.6$ Hz). IR (ν , cm^{-1}): 1640 (C=O).

2-[2-(Furan-2-yl)-1,3-benzoxazole-5-yl]-1-(morpholine-4-yl)ethan-1-one (**7c**): The title compound was prepared from compound **6** and morpholine to afford **7c** as a white solid (57%): mp 118°C . ^1H NMR (300 MHz, CDCl_3): 7.67 (m, 1H, H_5), 7.59 (m, 1H, H_3), 7.52 (d, 1H, H_7 , $J = 8.4$ Hz), 7.27 (m, 2H, H_4 and H_6), 6.63 (dd, 1H, H_4 , $J = 1.7$ Hz and $J = 3.5$ Hz), 3.85 (s, 2H, CH_2), 3.66 (m, 4H, $H_{\text{morpholine}}$), 3.50 (m, 4H, $H_{\text{morpholine}}$). IR (ν , cm^{-1}): 1630 (C=O).

2-[2-(Furan-2-yl)-1,3-benzoxazol-5-yl]-1-[4-[4-(2-methoxyethoxy)phenyl]piperazin-1-yl]ethan-1-one (**7d**): The title compound was prepared from compound **6** and 1-[4-(2-methoxyethoxy)phenyl]piperazine to afford **7d** as a beige solid (73%): mp 114°C . ^1H NMR (300 MHz, $[\text{D}_6]\text{DMSO}$): 7.67 (m, 1H, H_5), 7.62 (m, 1H, H_3), 7.51 (d, 1H, H_7 , $J = 8.4$ Hz), 7.29 (m, 2H, H_6 and H_4), 6.84 (m, 4H, H_{phenyl} and H_{phenyl}), 6.63 (dd, 1H, H_4 , $J = 1.7$ Hz and $J = 3.5$ Hz), 4.05 (t, 2H, CH_2O , $J = 4.6$ Hz), 3.89 (s, 2H, CH_2), 3.81 (t, 2H, $H_{\text{piperazine}}$, $J = 4.7$ Hz), 3.71 (t, 2H, CH_2O , $J = 4.6$ Hz), 3.63 (t, 2H, $H_{\text{piperazine}}$, $J = 4.7$ Hz), 3.43 (s, 3H, OCH_3), 3.02 (t, 2H, $H_{\text{piperazine}}$, $J = 4.7$ Hz), 2.88 (t, 2H, $H_{\text{piperazine}}$, $J = 4.7$ Hz). IR (ν , cm^{-1}): 1630 (C=O).

General procedure for the synthesis of compound (B1–B4)

To a solution of amide (1.06 mmol) (**7a–7d**) in THF (5 ml) was added LiAlH_4 (2.65 mmol). After 1 h stirring at room temperature, water (0.1 ml), 1 M NaOH solution (0.1 ml) and water (0.3 ml) were added successively to get a white mineral solid which was filtered off and washed with EtOAc (50 ml). Organic layer was then washed with water, brine solution, dried over MgSO_4 and concentrated *in vacuo* to afford a solid which was then purified.

2-(Furan-2-yl)-5-[2-(piperidine-1-yl) ethyl]-1,3-benzoxazole hydrochloride (**B1**): The title compound was prepared from amide **7a**.

Solid was suspended in diethyl ether with HCl gas, concentrated *in vacuo* and recrystallised from acetonitrile to afford **B1** as a white solid (10%): mp >300 °C. ¹H NMR (300 MHz, [D₆]DMSO): 10.53 (br s, 1H, NH⁺), 8.0 (m, 1H, H₅), 7.74 (d, 1H, H₇, *J* = 8.4 Hz), 7.69 (m, 1H, H₄), 7.46 (d, 1H, H₃, *J* = 3.5 Hz), 7.33 (dd, 1H, H₆, *J* = 3.4 and *J* = 8.4 Hz), 6.82 (dd, 1H, H₄, *J* = 1.7 Hz and *J* = 3.5 Hz), 3.52–3.48 (m, 2H, CH₂CH₂), 3.33–3.19 (m, 4H, H_{piperidine}), 2.93–2.90 (m, 2H, CH₂CH₂), 1.80–1.70 (m, 5H, H_{piperidine}), 1.43–1.39 (m, 1H, H_{piperidine}). ¹³C NMR (75 MHz, [D₆]DMSO): 155.5 (C), 149.0 (C), 147.7 (CH), 142.0 (C), 141.9 (C), 134.9 (C), 126.8 (CH), 120.1 (CH), 115.7 (CH), 113.3 (CH), 111.3 (CH), 57.2 (CH₂), 52.5 (2 CH₂), 29.7 (CH₂), 22.8 (2 CH₂), 21.9 (CH₂). LC-MS (ESI) *m/z* found: 297 [M + H]⁺. HPLC: C₄ column: t_R = 16.3 min, purity >99%.

2-(Furan-2-yl)-5-[2-(4-phenylpiperazin-1-yl) ethyl]-1,3-benzoxazole (**B2**): The title compound was prepared from amide **7b**. Solid was recrystallised from ethanol to afford **B2** as a white solid (12%): mp 160 °C. ¹H NMR (300 MHz, [D₆]DMSO): 8.06 (m, 1H, H₅), 7.65 (m, 1H, H₄), 7.56 (1H, d, H₇, *J* = 8.2 Hz), 7.43 (d, 1H, H₃, *J* = 3.5 Hz), 7.31 (dd, 1H, H₆, *J* = 1.3 Hz and *J* = 8.2 Hz), 7.22–7.17 (m, 2H, H_{phenyl}), 6.92 (d, 2H, H_{phenyl}, *J* = 7.9 Hz), 6.80 (dd, 1H, H₄, *J* = 1.7 Hz and *J* = 3.5 Hz), 6.78–6.73 (m, 1H, H_{phenyl}), 3.12 (m, 4H, H_{piperazine}), 2.88 (t, 2H, CH₂CH₂, *J* = 7.3 Hz), 2.64–2.57 (m, 6H, CH₂CH₂ and H_{piperazine}). ¹³C NMR (75 MHz, [D₆]DMSO): 155.2 (C), 151.5 (C), 148.5 (C), 147.5 (CH), 141.7 (C), 142.1 (C), 138.2 (C), 129.4 (CH), 126.9 (CH), 120.0 (CH), 119.2 (CH), 115.8 (CH), 115.4 (CH), 113.2 (CH), 110.8 (CH), 60.4 (CH₂), 53.1 (2 CH₂), 48.7 (2 CH₂), 33.0 (CH₂). LC-MS (ESI) *m/z* found: 374 [M + H]⁺. HPLC: C₄ column: t_R = 15.8 min, purity >99%.

2-(Furan-2-yl)-5-[2-(morpholin-4-yl) ethyl]-1,3-benzoxazole hydrochloride (**B3**): The title compound was prepared from amide **7c**. Solid was suspended in diethyl ether with HCl gas, concentrated *in vacuo* and recrystallised from acetonitrile to afford **B3** as a white solid (80%): mp >300 °C. ¹H NMR (300 MHz, [D₆]DMSO): 11.36 (br s, 1H, NH⁺), 8.06 (m, 1H, H₅), 7.73–7.67 (m, 2H, H₇ and H₄), 7.43 (m, 1H, H₃), 7.31 (m, 1H, H₆), 7.80 (m, 1H, H₄), 3.92–3.82 (m, 4H, CH₂CH₂), 3.31–3.18 (m, 8H, H_{morpholine}). LC-MS (ESI) *m/z* found: 299 [M + H]⁺. HPLC: C₄ column: t_R = 14.8 min, purity 99%.

2-(Furan-2-yl)-5-[2-[4-(2-methoxyethoxy)phenyl]piperazin-1-yl]ethyl-1,3-benzoxazole (**B4**): The title compound was prepared from amide **7d**. Solid was suspended recrystallised from acetonitrile to afford **B4** as a beige solid (73%): mp >300 °C. ¹H NMR (300 MHz, CDCl₃): 7.67 (m, 1H, H₅), 7.62 (m, 1H, H₃), 7.48 (d, 1H, H₇, *J* = 8.3 Hz), 7.27 (m, 2H, H₆ and H₄), 7.22 (m, 1H, H_{phenyl}), 6.90 (m, 4H, H_{phenyl}), 6.63 (dd, 1H, H₄, *J* = 1.7 Hz and *J* = 3.5 Hz), 4.09 (t, 2H, OCH₂, *J* = 4.6 Hz), 3.73 (t, 2H, MeOCH₂, *J* = 4.6 Hz), 3.45 (s, 3H, OMe), 3.15 (m, 4H, H_{piperazine}), 2.98 (m, 2H, CH₂CH₂), 2.73 (m, 6H, H_{piperazine} and CH₂CH₂). ¹³C NMR (75 MHz, CDCl₃): 155.5 (C), 152.9 (C), 148.8 (C), 145.9 (C), 145.7 (CH), 142.7 (C), 141.9 (C), 137.3 (C), 126.1 (CH), 119.9 (CH), 118.1 (2 CH), 115.4 (2 CH), 114.1 (CH), 112.3 (CH), 110.2 (CH), 71.2 (CH₂), 67.7 (CH₂), 60.8 (CH₂), 59.2 (CH₂), 53.4 (2 CH₂), 50.6 (2 CH₂), 33.7 (CH₃). LC-MS (ESI) *m/z* found: 448 [M + H]⁺. HPLC: C₄ column: t_R = 15.2 min, purity 99%.

General procedure for the synthesis of acid (8a–8b)

A solution of dimethyl malonate (9.06 mmol) in acetone (33 ml) with K₂CO₃ (13.6 mmol) was stirred for 20 min at room temperature. Then compound (**4a**, **4c**) (4.53 mmol) was added and the mixture was stirred at reflux for 2 h, cooled to room temperature and concentrated *in vacuo*. Solid was suspended in water and extracted twice with EtOAc. Combined organic layer were dried over MgSO₄ and then concentrated *in vacuo*. The solid was suspended in H₂O (7 ml) and NaOH (18.1 mmol) was added. The mixture was stirred overnight at 40 °C and then washed with EtOAc.

The aqueous layer was acidified with 6 M HCl solution up to acid pH (1–3) and the formed solid was filtered. Crude was heated in DMF (5 ml) at 80 °C for 3 h, cooled to room temperature, hydrolysed with water and acidified with 1 M HCl solution and extracted three times with EtOAc. Combined organic layers were washed with brine, dried over MgSO₄ and concentrated *in vacuo*. Solid was suspended in diethyl ether and filtered.

3-[2-(Furan-2-yl)-1,3-benzoxazol-5-yl] propanoic acid (**8a**): The title compound was prepared from compound **4a** to afford **8a** as a white solid (43%): mp 207 °C. ¹H NMR (300 MHz, CDCl₃): 12.19 (br s, 1H, OH), 8.05 (m, 1H, H₅), 7.66–7.62 (m, 2H, H₆ and H₄), 7.42 (d, 1H, H₃, *J* = 3.5 Hz), 7.28 (d, 1H, H₇, *J* = 8.2 Hz), 6.79 (m, 1H, H₄), 2.94 (t, 2H, CH₂CH₂, *J* = 7.5 Hz), 2.59 (t, 2H, CH₂CH₂, *J* = 7.5 Hz). IR (ν, cm⁻¹): 1686 (C = O). LC-MS (ESI) *m/z* found: 258 [M + H]⁺.

3-[2-(3,4-Dimethoxyphenyl)-1,3-benzoxazol-5-yl] propanoic acid (**8b**): The title compound was prepared from compound **4c** to afford **8b** as a white solid (47%): mp 182 °C. ¹H NMR (300 MHz, CDCl₃): 12.14 (br s, 1H, OH), 7.77 (dd, 1H, H₆, *J* = 2.0 Hz and *J* = 8.4 Hz), 7.67–7.60 (m, 3H, H₄, H₂ and H₅), 7.25 (dd, 1H, H₆, *J* = 1.6 Hz and *J* = 8.3 Hz), 7.17 (d, 1H, H₇, *J* = 8.3 Hz), 3.88 (s, 3H, OMe), 3.86 (s, 3H, OMe), 2.94 (t, 2H, CH₂CH₂, *J* = 7.7 Hz), 2.59 (t, 2H, CH₂CH₂, *J* = 7.7 Hz). IR (ν, cm⁻¹): 1710 (C = O).

Synthesis of compounds (9a–9c)

Same procedure as described for compounds (7a–7d) has been used.

3-(2-(Furan-2-yl)-1,3-benzoxazol-5-yl)-1-(4-phenylpiperazin-1-yl)propan-1-one (**9a**): The title compound was prepared from compound **8a** and phenylpiperazine to afford **9a** as a white solid (55%): mp 164 °C. ¹H NMR (300 MHz, CDCl₃): 7.66 (m, 1H, H₅), 7.58 (m, 1H, H₃), 7.48 (d, 1H, H₇, *J* = 8.3 Hz), 7.29–7.23 (m, 4H, H_{phenyl}), 6.91–6.87 (m, 3H, H₄, H₆ and H_{phenyl}), 6.62 (dd, 1H, H₄, *J* = 1.7 Hz and *J* = 3.5 Hz), 3.57 (m, 2H, H_{piperazine}, *J* = 5.7 Hz), 3.55 (m, 2H, H_{piperazine}), 3.17–3.11 (m, 4H, CH₂CH₂ and H_{piperazine}), 3.04 (m, 2H, H_{piperazine}), 2.73 (t, 2H, CH₂CH₂, *J* = 7.7 Hz). IR (ν, cm⁻¹): 1658 (C = O).

3-[2-(3,4-Dimethoxyphenyl)-1,3-benzoxazole-5-yl]-1-(piperidin-1-yl)propan-1-one (**9b**): The title compound was prepared from compound **8b** and piperidine to afford **9b** as a white solid (66%): mp 130 °C. ¹H NMR (300 MHz, CDCl₃): 7.85 (dd, 1H, H₆, *J* = 8.4 Hz and *J* = 1.9 Hz), 7.75 (d, 1H, H₂, *J* = 1.9 Hz), 7.56 (m, 1H, H₄), 7.58 (d, 1H, H₇, *J* = 8.4 Hz), 7.21 (dd, 1H, H₆, *J* = 1.5 Hz and *J* = 8.1 Hz), 7.17 (d, 1H, H₇, *J* = 8.4 Hz), 4.02 (s, 3H, OMe), 3.97 (s, 3H, OMe), 3.57 (t, 1H, CH₂CH₂, *J* = 5.6 Hz), 3.35 (t, 2H, H_{piperidine}, *J* = 5.6 Hz), 3.10 (t, 2H, CH₂CH₂, *J* = 7.7 Hz), 2.67 (t, 2H, CH₂CH₂, *J* = 7.7 Hz), 1.68–1.47 (m, 6H, H_{piperidine}). IR (ν, cm⁻¹): 1625 (C = O).

3-[2-(3,4-Dimethoxyphenyl)-1,3-benzoxazol-5-yl]-1-(4-phenylpiperazin-1-yl)propan-1-one (**9c**): The title compound was prepared from compound **8b** and phenylpiperazine to afford **9c** as a white solid (54%): mp 160 °C. ¹H NMR (300 MHz, CDCl₃): 7.85 (dd, 1H, H₆, *J* = 2.0 Hz and *J* = 8.5 Hz), 7.75 (d, 1H, H₂, *J* = 2.0 Hz), 7.60 (d, 1H, H₄, *J* = 1.2 Hz), 7.47 (d, 1H, H₇, *J* = 8.3 Hz), 7.28–7.20 (m, 3H, H₆ and H_{phenyl}), 6.99 (d, 1H, H₅, *J* = 8.5 Hz), 6.87–6.82 (m, 3H, H_{phenyl}), 4.02 (s, 3H, OMe), 3.98 (s, 3H, OMe), 3.79 (m, 2H, H_{piperazine}), 3.55 (m, 2H, H_{piperazine}), 3.16–3.11 (m, 4H, CH₂CH₂ and H_{piperazine}), 3.03 (m, 2H, H_{piperazine}), 2.74 (t, 2H, CH₂CH₂, *J* = 7.4 Hz). IR (ν, cm⁻¹): 1635 (C = O).

Synthesis of compounds (C1–C3)

Same procedure as described for compounds (**B1–B4**) has been used.

2-(Furan-2-yl)-5-(3-(4-phenylpiperazin-1-yl)propyl)-1,3-benzoxazole hydrochloride (**C1**): The title compound was prepared from

compound **9a**. Solid was suspended in diethyl ether with HCl gas, concentrated *in vacuo* and recrystallised from acetonitrile to afford **C1** as a white solid (12%): mp >300 °C. ¹H NMR (300 MHz, CDCl₃): 11.00 (br s, 1H, NH⁺), 8.07 (m, 1H, H₅), 7.71–7.67 (m, 2H, H₄ and H₃), 7.44 (m, 1H, H₆), 7.32–7.24 (m, 3H, H₇ and H_{phenyl}), 6.98 (m, 2H, H_{phenyl}), 6.81 (m, 2H, H_{phenyl} and H₄), 3.76 (m, 2H), 3.57 (m, 2H), 3.12 (m, 6H), 2.80 (m, 2H, CH₂CH₂CH₂), 2.14 (m, 2H, CH₂CH₂CH₂). ¹³C NMR (75 MHz, [D₆]DMSO): 155.3 (C), 150.1 (C), 148.7 (C), 147.6 (CH), 142.0 (C), 141.9 (C), 138.3 (C), 129.6 (2 CH), 126.9 (CH), 120.4 (CH), 119.6 (CH), 116.4 (2 CH), 115.5 (CH), 113.3 (CH), 111.0 (CH), 55.5 (CH₂), 51.1 (2 CH₂), 45.9 (2 CH₂), 32.4 (CH₂), 25.6 (CH₂). LC-MS (ESI) *m/z* found: 388 [M + H]⁺. HPLC: C₄ column: t_R = 15.8 min, purity 98%.

2-(3,4-Dimethoxyphenyl)-5-[3-(piperidin-1-yl)propyl]-1,3-benzoxazole hydrochloride (**C2**): The title compound was prepared from compound **9b**. Solid was suspended in diethyl ether with HCl gas, concentrated *in vacuo* and recrystallised from ethyl acetate to afford **C2** as a white solid (72%): mp 214 °C. ¹H NMR (300 MHz, [D₆]DMSO): 10.20 (br s, 1H, NH⁺), 7.77 (dd, 1H, H₆, *J* = 2.0 Hz and *J* = 8.4 Hz), 7.68 (d, 1H, H₅, *J* = 8.4 Hz), 7.66–7.63 (m, 2H, H₂ and H₄), 7.21 (dd, 1H, H₆, *J* = 1.5 Hz and *J* = 8.3 Hz), 7.17 (d, 1H, H₇, *J* = 8.3 Hz), 3.88 (s, 3H, OMe), 3.86 (s, 3H, OMe), 3.38 (m, 2H, CH₂CH₂CH₂), 2.99 (m, 2H, CH₂CH₂CH₂), 2.76 (m, 2H, CH₂CH₂CH₂), 2.08 (m, 2H, H_{piperidine}), 1.75–1.65 (m, 5H, H_{piperidine}), 1.35 (m, 1H, H_{piperidine}). ¹³C NMR (75 MHz, [D₆]DMSO): 163.2 (C), 152.5 (C), 149.5 (C), 149.3 (C), 142.4 (C), 137.8 (C), 125.9 (C), 121.3 (CH), 119.2 (CH), 119.2 (CH), 112.4 (CH), 110.9 (CH), 110.2 (CH), 56.2 (CH₃), 56.1 (CH₃), 55.9 (CH₂), 52.4 (2 CH₂), 32.5 (CH₂), 25.6 (CH₂), 22.9 (2 CH₂), 21.9 (CH₂). LC-MS (ESI) *m/z* found: 381 [M + H]⁺. HPLC: C₄ column: t_R = 17.3 min, purity 98%.

2-(3,4-Dimethoxyphenyl)-5-[3-(4-phenylpiperazin-1-yl)propyl]-1,3-benzoxazole hydrochloride (**C3**): The title compound was prepared from compound **9c**. Solid was suspended in diethyl ether with HCl gas, concentrated *in vacuo* and recrystallised from ethanol to afford **C3** as a white solid (73%): mp 194 °C. ¹H NMR (300 MHz, [D₆]DMSO): 10.81 (br s, 1H, NH⁺), 7.77 (dd, 1H, H₆, *J* = 1.8 Hz and *J* = 8.4 Hz), 7.70–7.65 (m, 3H, H₄, H₆ and H₇), 7.29–7.22 (m, 3H, H₂ and H_{phenyl}), 7.17 (d, 1H, H₅, *J* = 8.4 Hz), 6.98 (m, 2H, H_{phenyl}), 6.87–6.82 (m, 1H, H_{phenyl}), 3.88 (s, 3H, OMe), 3.86 (s, 3H, OMe), 3.77 (m, 2H, CH₂CH₂CH₂), 3.56 (m, 2H, H_{piperazine}), 3.11 (m, 6H, H_{piperazine}), 2.79 (t, 2H, CH₂CH₂CH₂, *J* = 7.4 Hz), 2.13 (m, 2H, CH₂CH₂CH₂). ¹³C NMR (75 MHz, [D₆]DMSO): 163.2 (C), 152.5 (C), 150.1 (C), 149.5 (C), 149.3 (C), 142.4 (C), 137.8 (C), 129.6 (2 CH), 125.9 (CH), 121.3 (CH), 120.4 (C), 119.3 (CH), 119.2 (CH), 116.4 (2 CH), 112.4 (CH), 110.9 (CH), 110.2 (CH), 56.2 (CH₂), 56.1 (CH₃), 55.5 (CH₃), 51.1 (2 CH₂), 45.9 (2 CH₂), 32.5 (CH₂), 25.7 (CH₂). LC-MS (ESI) *m/z* found: 458 [M + H]⁺. HPLC: C₄ column: t_R = 16.6 min, purity >99%.

2-(2-(3,4-dimethoxyphenyl)-1,3-benzoxazol-5-yl)ethanol (10). To a solution of compound (**3e**) (1 g, 3.06 mmol) in THF (10 ml) was added LiAlH₄ (340 mg, 9.2 mmol). After 1 h stirring at room temperature, water (0.34 ml), 1 M NaOH solution (0.34 ml) and water (1.02 ml) were added successively until get a white solid which was filtered off and washed with EtOAc (60 ml). Organic layer was washed with water, dried over MgSO₄ and concentrated *in vacuo*. Solid was recrystallised in acetonitrile to afford compound (**10**) as a beige solid (603 mg, 86%): mp 210 °C. ¹H NMR (300 MHz, CDCl₃): 9.13 (br s, 1H, OH) 7.86 (dd, 1H, H₆, *J* = 1.8 Hz and *J* = 8.4 Hz), 7.76 (d, 1H, H₂, *J* = 1.8 Hz), 7.61 (m, 1H, H₄), 7.50 (d, 1H, H₇, *J* = 8.4 Hz), 7.24 (dd, 1H, H₆, *J* = 1.2 Hz and *J* = 8.4 Hz), 7.00 (d, 1H, H₅, *J* = 8.4 Hz), 4.03 (s, 3H, OMe), 3.98 (s, 3H, OMe), 3.92 (m, 2H,

CH₂CH₂), 3.64 (t, 2H, CH₂CH₂, *J* = 6.6 Hz). IR (ν, cm⁻¹): 3400 (OH). LC-MS (ESI) *m/z* found: 300 [M + H]⁺.

2-(2-(3,4-Dimethoxyphenyl)-1,3-benzoxazol-5-yl)ethyl methanesulfonate (11). To a solution of compound (**10**) (800 mg, 2.37 mmol) in DCM (20 ml) with Et₃N (0.49 ml, 3.55 mmol) at 0 °C, was added dropwise mesyl chloride (0.28 ml, 3.55 mmol). After 4 h stirring at room temperature, mixture was hydrolysed with water and extracted twice with DCM. Combined organic layers were dried over MgSO₄ and concentrated *in vacuo*. Solid was suspended in diethyl ether and filtered to afford compound (**11**) as a beige solid (895 mg, 100%): mp 112 °C. ¹H NMR (300 MHz, CDCl₃): 7.85 (dd, 1H, H₆, *J* = 8.4 Hz and *J* = 2.0 Hz), 7.76 (d, 1H, H₄, *J* = 2.0 Hz), 7.61 (d, 1H, H₄, *J* = 1.6 Hz), 7.51 (d, 1H, H₇, *J* = 8.3 Hz), 7.21 (dd, 1H, H₆, *J* = 8.3 Hz and *J* = 1.6 Hz), 7.00 (d, 1H, H₅, *J* = 8.4 Hz), 4.47 (t, 2H, CH₂CH₂, *J* = 6.8 Hz), 4.02 (s, 3H, OMe), 3.97 (s, 3H, OMe), 3.18 (t, 2H, CH₂CH₂, *J* = 6.9 Hz), 2.88 (s, 3H, CH₃). LC-MS (ESI) *m/z* found: 378 [M + H]⁺.

General procedure for the synthesis of compound (B5–B6)

To a solution of compound **11** (0.79 mmol) in DMF (8 ml), were added K₂CO₃ (1.46 mmol) and amine (1.03 mmol). After overnight stirring at 80 °C, the reaction mixture was cooled, hydrolysed with water and extracted three times with EtOAc. Combined organic layers were dried over MgSO₄ and concentrated *in vacuo* to afford a solid which was then purified.

2-(3,4-Dimethoxyphenyl)-5-(2-(piperidin-1-yl)ethyl)-1,3-benzoxazole hydrochloride (**B5**): The title compound was prepared from compound **11** and piperidine. Solid was suspended in diethyl ether with HCl gas, concentrated *in vacuo* and recrystallised from ethanol to afford **B5** as a white solid (25%): mp 260 °C. ¹H NMR (300 MHz, [D₆]DMSO): 10.63 (br m, 1H, NH⁺), 7.79 (dd, 1H, H₆, *J* = 8.4 Hz and *J* = 2.0 Hz), 7.73 (d, 1H, H₇, *J* = 8.3 Hz), 7.68 (d, 1H, H₂, *J* = 1.3 Hz), 7.66 (d, 1H, H₄, *J* = 2.0 Hz), 7.31 (dd, 1H, H₆, *J* = 8.4 Hz and *J* = 1.7 Hz), 7.18 (d, 1H, H₅, *J* = 8.6 Hz), 3.89 (s, 3H, OMe), 3.87 (s, 3H, OMe), 3.50 (m, 2H, CH₂CH₂), 3.27–3.20 (m, 4H, CH₂CH₂ and H_{piperidine}), 2.92 (m, 2H, H_{piperidine}), 1.82–1.70 (m, 5H, H_{piperidine}), 1.42 (m, 1H, H_{piperidine}). ¹³C NMR (75 MHz, [D₆]DMSO): 163.4 (C), 152.5 (C), 149.5 (C), 149.3 (C), 142.4 (C), 137.8 (C), 125.9 (C), 121.3 (CH), 119.2 (CH), 119.2 (CH), 112.4 (CH), 110.9 (CH), 110.2 (CH), 57.3 (CH₂), 56.2 (CH₃), 56.1 (CH₃), 52.4 (2 CH₂), 29.7 (CH₂), 22.8 (2 CH₂), 21.9 (CH₂). LC-MS (ESI) *m/z* found: 367 [M + H]⁺. HPLC: C₄ column: t_R = 16.1 min, purity 99%.

2-(3,4-Dimethoxyphenyl)-5-[2-(4-phenylpiperazin-1-yl)ethyl]-1,3-benzoxazole (**B6**): The title compound was prepared from compound **11** and phenylpiperazine. Solid was recrystallised from ethanol to afford **B6** as a white solid (28%): mp 140 °C. ¹H NMR (300 MHz, CDCl₃): 7.85 (dd, 1H, H₆, *J* = 2.0 Hz and *J* = 8.4 Hz), 7.76 (d, 1H, H₂, *J* = 2.0 Hz), 7.61 (m, 1H, H₄), 7.48 (d, 1H, H₅, *J* = 8.4 Hz), 7.31–7.26 (m, 2H, H₇ and H_{phenyl}), 7.20 (dd, 1H, H₆, *J* = 1.5 Hz and *J* = 8.3 Hz), 7.01–6.85 (m, 4H, H_{phenyl}), 4.03 (s, 3H, OMe), 3.98 (s, 3H, OMe), 3.27–3.24 (t, 4H, H_{piperazine}, *J* = 4.8 Hz), 3.01–2.96 (m, 2H, CH₂CH₂), 2.75–2.71 (m, 6H, CH₂CH₂ and H_{piperazine}). ¹³C NMR (300 MHz, CDCl₃): 163.4 (C), 152.0 (C), 151.3 (C), 149.4 (C), 149.2 (C), 142.5 (C), 136.9 (C), 129.1 (2 CH), 125.5 (CH), 121.1 (CH), 119.9 (C), 119.7 (CH), 119.4 (CH), 116.1 (2 CH), 111.0 (CH), 110.0 (2 CH), 60.9 (CH₂), 56.1 (CH₃), 56.0 (CH₃), 53.3 (2 CH₂), 49.2 (2 CH₂), 33.7 (CH₂). LC-MS (ESI) *m/z* found: 444 [M + H]⁺. HPLC: C₄ column: t_R = 15.7 min, purity 99%.

2-(Furan-2-yl)-1,3-benzoxazol-5-amine (D1). To a solution of 2-(furan-2-yl)-5-nitro-1,3-benzoxazole (3.07 g, 13.3 mmol) (**3d**) in

EtOH (130 ml) were added Pd/C (10%, 100 mg) and hydrazine monohydrate (0.78 ml, 16 mmol). The mixture was heated at 70 °C for 3 h, cooled to room temperature, the Pd/C was filtered off and the filtrate was concentrated *in vacuo*. Crude was suspended in water and extracted three times with EtOAc. Combined organic layer were washed with brine, dried over MgSO₄ and concentrated *in vacuo*. Solid was recrystallised from hexane to afford compound **(D1)** as grey solid (2.16 g, 81%): mp 112 °C. ¹H NMR (300 MHz, [D₆]DMSO): 8.01 (m, 1H, H₅), 7.40 (d, 1H, H₇, *J* = 8.7 Hz), 7.34 (d, 1H, H₃, *J* = 3.5 Hz), 6.84 (d, 1H, H₄, *J* = 2.1 Hz), 6.77 (dd, 1H, H₄, *J* = 1.7 Hz and *J* = 3.5 Hz), 6.66 (dd, 1H, H₆, *J* = 2.1 Hz and *J* = 8.7 Hz), 5.16 (br s, 2H, NH₂). ¹³C NMR (75 MHz, [D₆]DMSO): 154.9 (C), 147.3 (C), 147.0 (CH), 142.6 (C), 142.5 (C), 142.4 (C), 114.4 (CH), 113.6 (CH), 113.1 (CH), 110.9 (CH), 103.0 (CH). IR (ν , cm⁻¹): 3424 (NH₂). LC-MS (ESI) *m/z* found: 201 [M + H]⁺. HPLC: C₄ column: t_R = 14.2 min, purity 98%.

N-(4-(Benzyloxy)phenethyl)-2-(furan-2-yl)-1,3-benzoxazol-5-amine

(12). To a solution of compound **D1** (800 mg, 4 mmol) in DMF (15 ml) were added 1-(benzyloxy)-4-(2-bromoethyl)benzene (1.4 g, 4.8 mmol)¹⁶ and K₂CO₃ (828 mg, 5.99 mmol). The reaction mixture was stirred at 80 °C overnight, cooled to room temperature, hydrolysed with water, acidified with 1 M HCl solution, and extracted with EtOAc three times. Combined organic layers were washed with brine, dried over MgSO₄ and concentrated *in vacuo* to give a yellow oil. Purification by flash chromatography was realised with EtOAc/Cyclohexane as solvent (1/9 up to 3/7) to give the product as a yellow oil which was suspended in diethyl ether and filtered to afford compound **12** as a pale brown solid (430 mg, 26%): mp 184 °C. ¹H NMR (75 MHz, [D₆]DMSO, *J* Hz): 8.10 (m, 1H, H₅), 7.75 (m, 1H), 7.58 (m, 1H), 7.46–7.26 (m, 7H), 7.21 (d, 2H, H_{phenyl}, *J* = 8.7 Hz), 6.96 (d, 2H, H_{phenyl}, *J* = 8.7 Hz), 6.83 (dd, 1H, H₄, *J* = 1.7 Hz and *J* = 3.5 Hz), 5.06 (s, 2H, CH₂), 3.52 (t, 2H, CH₂CH₂, *J* = 7.5 Hz), 2.99 (t, 2H, CH₂CH₂, *J* = 7.5 Hz). LC-MS (ESI) *m/z* found: 411 [M + H]⁺.

4-(2-[(2-(Furan-2-yl)-1,3-benzoxazol-5-yl)amino]ethyl)phenol (E2). A solution of compound **12** (740 mg, 1.66 mmol) in MeOH (15 ml) with Pd/C (50 mg) under H₂ atmosphere was stirred at 25 °C for overnight. Pd/C was filtered off and the filtrate was concentrated *in vacuo*. Solid was recrystallised in CH₃CN to afford compound **E2** as a yellow crystal (104 mg, 18%): mp 174 °C. ¹H NMR (300 MHz, [D₆]DMSO): 9.16 (br s, 1H, OH), 8.01 (m, 1H, H₅), 7.43 (d, 1H, H₇, *J* = 8.7 Hz), 7.33 (d, 1H, H₃, *J* = 3.5 Hz), 7.07 (d, 2H, H_{phenyl}, *J* = 8.4 Hz), 6.80 (m, 1H, H₄), 6.77 (dd, 1H, H₄, *J* = 1.7 Hz and *J* = 3.5 Hz), 6.69 (d, 2H, H_{phenyl}, *J* = 8.4 Hz), 6.71 (m, 1H, H₆), 5.73 (br t, 1H, *J* = 5.6 Hz, NH), 3.24–3.17 (m, 2H, *J* = 6.9 Hz, CH₂CH₂), 2.75 (t, 2H, *J* = 7.6 Hz, CH₂CH₂). ¹³C NMR (75 MHz, [D₆]DMSO): 156.1 (C), 154.9 (C), 147.7 (C), 147.0 (CH), 142.8 (C), 142.5 (C), 142.2 (C); 130.3 (C), 130.0 (2 CH), 115.5 (2 CH), 114.4 (CH), 113.1 (CH), 112.8 (CH), 111.1 (CH), 100.2 (CH); 46.1 (CH₂), 34.4 (CH₂). IR (ν , cm⁻¹): 3340 (OH). LC-MS (ESI) *m/z* found: 321 [M + H]⁺. HPLC: C₄ column: t_R = 14.8 min, purity 97%.

2-(Furan-2-yl)-N-(2-(piperidin-1-yl)ethyl)-1,3-benzoxazol-5-amine

hydrochloride (E1). To a solution of compound **D1** (800 mg, 4 mmol) in DMF (16 ml) were added K₂CO₃ (1.7 g, 12 mmol) and *N*-2-chloroethyl piperidine hydrochloride (1471 mg, 7.99 mmol). The reaction mixture was stirred at 70 °C overnight, cooled to room temperature, hydrolysed with water and extracted three times with EtOAc. Combined organic layer were dried over MgSO₄ and concentrated *in vacuo*. Purification by flash chromatography was performed with DCM/EtOH/NH₃ (90/10/1) as a solvent.

The obtained yellow oil was suspended in diethyl ether with HCl gas to formed a solid which was filtered to afford compound **E1** as a beige solid (11 mg, 8%): mp 169 °C. ¹H NMR (300 MHz, [D₆]DMSO): 10.69 (br m, 1H, NH⁺), 8.04 (m, 1H, H₅), 7.55 (d, 1H, H₇, *J* = 8.8 Hz), 7.38 (m, 1H, H₃), 7.09 (m, 1H, H₄), 6.88 (dd, 1H, H₆, *J* = 1.9 Hz and *J* = 8.8 Hz), 6.79 (dd, 1H, H₄, *J* = 1.7 Hz and *J* = 3.5 Hz), 5.37 (br m, 1H, NH), 3.57 (t, 2H, CH₂CH₂, *J* = 6.3 Hz), 3.49–3.46 (m, 2H, H_{piperidine}), 3.24 (t, 2H, CH₂CH₂, *J* = 6.2 Hz), 2.90 (m, 2H, H_{piperidine}), 1.79 (m, 5H, H_{piperidine}), 1.39 (m, 1H, H_{piperidine}). ¹³C NMR (75 MHz, [D₆]DMSO): 155.3 (C), 147.2 (CH), 145.0 (C), 143.6 (C), 142.7 (C), 124.3 (C), 114.9 (CH), 114.1 (CH), 113.2 (CH), 111.5 (CH), 102.6 (CH), 54.5 (CH₂), 52.7 (2 CH₂), 39.2 (CH₂), 22.8 (2 CH₂), 21.8 (CH₂). LC-MS (ESI) *m/z* found: 312 [M + H]⁺. HPLC: C₄ column: t_R = 15.9 min, purity 99%.

2-Bromo-N-(2-(furan-2-yl)-1,3-benzoxazol-5-yl)acetamide (13): To a solution of compound **D1** (210 mg, 1.05 mmol) and Et₃N (0.18 ml, 1.26 mmol) in DCM (10 ml) at 0 °C was added dropwise a solution of bromoacetyl bromide (0.11 ml, 1.26 mmol) diluted in DCM (5 ml). The reaction mixture was stirred at room temperature for 2 h, hydrolysed with water and extracted twice with DCM. Combined organic layers were washed with brine, dried over MgSO₄ and concentrated *in vacuo*. The solid was suspended in diethyl ether and filtered to afford compound **13** as a white solid (249 mg, 74%): mp 203 °C. ¹H NMR (300 MHz, CDCl₃): 8.27 (br s, 1H, NH), 7.96 (m, 1H, H₅), 7.70–7.69 (m, 1H, H₄), 7.56–7.50 (m, 2H, H₇ and H₆), 7.30 (dd, 1H, H₃, *J* = 0.7 Hz and *J* = 3.5 Hz), 6.39 (dd, 1H, H₄, *J* = 1.7 Hz and *J* = 3.5 Hz), 4.08 (s, 2H, CH₂). IR (ν , cm⁻¹): 3246 (NH), 1643 (C=O).

General procedure for the synthesis of compounds (F1–F3)

To a solution of compound **13** (0.59 mmol) in acetone (5 ml) were added K₂CO₃ (0.88 mmol) and the amine (0.65 mmol). The reaction mixture was refluxed for 2 h, cooled to room temperature and concentrated *in vacuo*. The crude was suspended in water and extracted three times with EtOAc. Combined organic layer were washed with brine, dried over MgSO₄ and concentrated *in vacuo* to afford a solid which was then purified.

N-[2-(Furan-2-yl)-1,3-benzoxazol-5-yl]-2-(piperidin-1-yl)acetamide hydrochloride (F1): The title compound was prepared from compound **13** and piperidine. Solid was suspended in diethyl ether with HCl gas, concentrated *in vacuo* and recrystallised from ethanol to afford **F1** as a white solid (57%): mp >300 °C. ¹H NMR (300 MHz, [D₆]DMSO): 11.26 (br s, 1H, NH⁺), 9.99 (br s, 1H, NH), 8.17 (m, 1H, H₅), 8.08 (m, 1H, H₄), 7.76 (d, 1H, H₇, *J* = 8.8 Hz), 7.58 (dd, 1H, H₆, *J* = 1.9 Hz and *J* = 8.8 Hz), 7.46 (d, 1H, H₃, *J* = 3.5 Hz), 6.82 (dd, 1H, H₄, *J* = 1.7 Hz and *J* = 3.5 Hz), 4.18 (s, 2H, CH₂), 3.50–3.47 (m, 2H, H_{piperidine}), 3.10 (m, 2H, H_{piperidine}), 1.80–1.67 (m, 5H, H_{piperidine}), 1.40 (m, 1H, H_{piperidine}). ¹³C NMR (75 MHz, [D₆]DMSO): 163.4 (C), 156.0 (C), 147.7 (CH), 146.6 (C), 141.9 (C), 141.8 (C), 135.9 (C), 118.2 (CH), 115.8 (CH), 113.3 (CH), 111.4 (CH), 110.8 (CH), 57.6 (CH₂), 53.4 (2 CH₂), 22.7 (2 CH₂), 21.6 (CH₂). LC-MS (ESI) *m/z* found: 326 [M + H]⁺. HPLC: C₄ column: t_R = 15.1 min, purity 98%.

N-[2-(Furan-2-yl)-1,3-benzoxazol-5-yl]-2-(4-phenylpiperazin-1-yl)acetamide (F2): The title compound was prepared from compound **13** and phenylpiperazine. Solid was recrystallised from ethanol to afford **F2** as a white solid (48%): mp 174 °C. ¹H NMR (300 MHz, CDCl₃): 9.26 (br s, 1H, NH⁺), 8.02 (m, 1H, H₄), 7.68–7.67 (m, 1H, H₅), 7.56 (dd, 1H, H₆, *J* = 2.0 Hz and *J* = 8.8 Hz), 7.51 (d, 1H, H₇, *J* = 8.6 Hz), 7.3–7.29 (m, 3H, H_{phenyl} and H₃), 6.98–6.88 (m, 3H, H_{phenyl} and H_{phenyl}), 6.63 (dd, 1H, H₄, *J* = 1.7 Hz and *J* = 3.5 Hz), 3.30 (t, 4H, H_{piperazine}, *J* = 4.9 Hz), 3.26 (s, 2H, CH₂), 2.83 (t, 4H, H_{piperazine}, *J* = 4.9 Hz). ¹³C NMR (75 MHz, CDCl₃): 168.2 (C), 156.2 (C),

152.9 (C), 146.9 (C), 145.8 (CH), 142.5 (C), 142.2 (C), 134.9 (C), 129.2 (2 CH), 120.2 (CH), 117.8 (CH), 116.3 (2 CH), 114.5 (CH), 112.3 (CH), 111.2 (CH), 110.5 (CH), 62.0 (CH₂), 50.6 (2 CH₂), 49.5 (2 CH₂). LC-MS (ESI) *m/z* found: 403 [M + H]⁺. HPLC: C₄ column: *t_R* = 14.7 min, purity >99%. *tert*-Butyl-4-([2-(furan-2-yl)-1,3-benzoxazol-5-yl]carbamoyl)methylpiperazine-1-carboxylate (**F3**): The title compound was prepared from compound **13** and *tert*-butyl piperazine-1-carboxylate. Solid was recrystallised from ethanol to afford **F3** as a white solid (90%): mp 186 °C. ¹H NMR (300 MHz, CDCl₃): 9.15 (br s, 1H, NH), 7.98 (d, 1H, H₄, *J* = 2.0 Hz), 7.68–7.67 (m, 1H, H₅), 7.55 (dd, 1H, H₆, *J* = 2.0 Hz and *J* = 8.6 Hz), 7.50 (d, 1H, H₇, *J* = 8.6 Hz), 7.28 (m, 1H, H₃), 6.62 (dd, 1H, H₄, *J* = 1.7 Hz and *J* = 3.5 Hz), 3.54 (t, 4H, H_{piperazine}, *J* = 4.9 Hz), 3.19 (s, 2H, CH₂), 2.60 (t, 4H, H_{piperazine}, *J* = 4.9 Hz), 1.48 (s, 9H, 3 CH₃). ¹³C NMR (75 MHz, CDCl₃): 167.9 (C), 156.2 (C), 154.6 (C), 147.0 (C), 145.9 (CH), 142.5 (C), 142.2 (C), 134.8 (C), 117.8 (CH), 114.5 (CH), 112.3 (CH), 111.2 (CH), 110.5 (CH), 80.1 (CH₂), 62.1 (2 CH₂), 53.3 (2 CH₂), 28.4 (3 CH₃). IR (*μ*, cm⁻¹): 1684 (C=O), 1673 (C=O). LC-MS (ESI) *m/z* found: 371 [M-tBu + H]⁺, 427 [M + H]⁺. HPLC: C₄ column: *t_R* = 15.1 min, purity 99%.

N-(2-(Furan-2-yl)-1,3-benzoxazol-5-yl)-2-(piperazin-1-yl)acetamide dihydrochloride (**F4**): To a solution of (**F3**) (180 mg, 0.422 mmol) in MeOH (10 ml) was added 6 M HCl (1.41 ml, 8.44 mmol) and the reaction mixture was stirred at 40 °C for 15 h. The precipitated product was filtered and washed with diethyl ether to afford compound (**F4**) as a white solid (121 mg, 72%): mp 192 °C. ¹H NMR (300 MHz, [D₆]DMSO): 11.16 (br s, 1H, NH), 9.88 (br s, 2H, NH₂⁺), 8.16 (d, 1H, H₄, *J* = 1.8 Hz), 8.07 (m, 1H, H₅), 7.75 (d, 1H, H₇, *J* = 8.8 Hz), 7.59 (dd, 1H, H₆, *J* = 1.8 Hz and *J* = 8.8 Hz), 7.45–7.44 (m, 1H, H₃), 6.81 (dd, 1H, H₄, *J* = 1.7 Hz and *J* = 3.5 Hz), 4.22 (s, 2H, CH₂), 3.57 (s, 4H, H_{piperazine}, *J* = 4.9 Hz), 3.43 (s, 4H, H_{piperazine}, *J* = 4.9 Hz). ¹³C NMR (75 MHz, [D₆]DMSO): 163.7 (C), 155.9 (C), 147.7 (CH), 146.5 (C), 141.9 (C), 141.8 (C), 135.9 (C), 118.2 (CH), 115.7 (CH), 113.3 (CH), 111.4 (CH), 110.8 (CH), 57.6 (CH₂), 49.1 (4 CH₂). LC-MS (ESI) *m/z* found: 327 [M + H]⁺. HPLC: C₄ column: *t_R* = 14.9 min, purity 99%.

Molecular docking

Molecular docking was performed using Gold suite v5.2¹⁷ within the Hermes v1.6 GUI (CCDC[®]). Thus after adding hydrogens, water molecules were deleted and docking was performed in a 10 Å around co-crystallised ligands then ligands deleted. Early termination of three docking solutions within 1.5 Å was set up in order to highlight ligands converging towards a few binding modes.

Pharmacological assays

Displacement binding assays

Radioligands were obtained from the following source: [³H]-ZM241385 from IsoBio (10–50 Ci/ml), [3H]-DPCPX from PerkinElmer (Waltham, MA) (120 Ci/mmol), [3H]-CPX from Perkin Elmer (120 Ci/mmol), [125I]-AB-MECA from Chelatec (2200 Ci/mmol) (Saint-Herblain, France).

For A_{2A} receptor binding evaluation, the stock solution of compounds was prepared in DMSO. The final concentration of DMSO was no more than 3% for radioligand binding assay.

Competition binding curves of the A_{2A} receptor antagonist [³H]-ZM241385 by the designed A_{2A} antagonists described above, were performed as before¹⁸ in Human HEK293 A_{2A}R membranes (PerkinElmer). 0.5 ml of membranes (0.5 U of A_{2A}R) were incubated with [³H]-ZM241385 (2 nm) and increasing concentrations of the designed A_{2A}R antagonists (0–600 nm) in a final volume of 300 μl

in the presence of 1 U/ml of adenosine deaminase (Roche, Basel, Switzerland). All samples were assayed in duplicate. Non-specific binding was determined for each assay in the presence of the antagonist ZM-24135 (8.3 nm). Microplates were incubated for 1 h at room temperature and the reaction was stopped by vacuum filtration with a Skatron semi-automatic cell harvester with chilled incubation solution (pH 7.4, Tris 50 mm MgCl 10 mm) to filter mats 1.5 mm (Molecular Devices, Sunnyvale, CA). Three millilitres of scintillation cocktail (OptiPhase "HiSafe" 2, PerkinElmer) were added and radioactivity bound to the filters was determined after 12 h. Molecules inhibited binding by ≥30% at 10 μm were submitted to K_i evaluation. This percentage was calculated using Excel 2013 as a ratio of data with ligands to data without ligands. Data were analysed using Graph Pad Prism, version 5.01 (San Diego, CA). Inhibition constants (K_i) were calculated from the IC₅₀ values by non-linear regression analysis, the Cheng and Prusoff equation and K_D value of 1.0 nm were used. Displacement reference curves were performed with ZM-24135 (0–6 nm in 6%, 40% or 60% of DMSO). No difference was observed between each concentration.

Affinity towards A₁R (human recombinant CHO cells, [3H]-DPCPX (1 nm), Cerep catalogue reference 0002, as described by Townsend-Nicholson¹⁹), A_{2B}R (human recombinant HEK-293 cells, [3H]-CPX (5 nm), Cerep catalogue reference 0005, as described by Stehle²⁰) and A₃R (human recombinant HEK-293 cells, [125I]-AB-MECA (0.15 nm), Cerep catalogue reference 0006, as described by Salvatore²¹) was determined by CEREP laboratories. K_D values used were: 1.7 nm for [3H]-DPCPX, 65 nm for [3H]-CPX and 0.15 nm for [125I]-AB-MECA. For these three receptors, the stock solution of compounds was prepared in DMSO. The final concentration of DMSO was no more than 1%. Data were analysed using SigmaPlot[®] version 4.0 for Windows[®] (© 1997 by SPSS Inc., Chicago, IL).

Cell culture and cytotoxicity assay

The human neuroblastoma cell line (SY5Y) was cultured in Dulbecco's modified Eagle medium (DMEM) (Gibco, Waltham, MA) supplemented with 2 mm L-glutamine, 100 mg/ml streptomycin, 100 IU/ml penicillin, 1 mm non-essential amino acids and 10% (v/v) heat-inactivated foetal bovine serum (Sigma-Aldrich, Saint-Louis, MO), and grown at 37 °C in a humidified incubator with 5% CO₂. Cells were seeded at 2000 cells per well onto 96-well plates in DMEM medium. Cells were starved for 24 h to obtain synchronous cultures and were then incubated in a culture medium that contained various concentrations of test compounds, each dissolved in less than 0.1% DMSO. After 72 h of incubation, cell growth was estimated by the colorimetric MTT (thiazolyl blue tetrazolium bromide) assay.

Absorption, distribution, metabolism and excretion (ADME) assessment

Aqueous solubility (in phosphate-buffered saline, PBS, pH 7.4; incubation room temperature for 24 h as described by Lipinski²² Eurofins Cerep SA catalogue reference G235), partition coefficient (log D, n-octanol/PBS, pH 7.4, room temperature for 60 min as described by Sangster²³; Eurofins Cerep SA catalogue reference 0417), human plasma protein binding evaluated at 10 μm concentration for 4 h at 37 °C as described by Banker²⁴ (Eurofins Cerep SA catalogue reference 2194), A-B and B-A permeability coefficient evaluated at 10 μm for 40 min as described by Hidalgo²⁵ (Papp, Caco-2 cells, pH 6.5/7.4; Eurofins Cerep SA catalogue reference G228), metabolic stability in human liver microsomes evaluated at

0.1 μM concentration for 0, 15, 30, 45, 60 min at 37 °C as described by Obach²⁶ (Eurofins Cerep SA catalogue reference 0607) were determined in standard assays by Eurofins Cerep SA, France www.cerep.fr).

Results and discussion

Structure-based insights

Our design was guided by molecular modelling studies which took into consideration the two structures of adenosine A_{2A} receptors co-crystallised with high-affinity A_{2A} antagonists ZM-241385 and T4E (1,2,4-triazines) in respective 3EML^{27,28} and 3UZC²⁹ PDB entries. Although several molecular dynamics studies have shown the importance of water molecules for ligand recognition^{30,31}, these molecules represent a bias in molecular docking with many options like their rigid pre-docking displacement, a tolerance for moving them or simply delete them. Since T4E-bound structures show that the key water molecules in ZM-241385-bound crystal structures could be displaced by aryl substituents, we decided to avoid this experimental bias in order to benefit with the whole cavity.

In contrast with ZM-241385 bound structure, the T4E-bound one was identified as the most suitable target to predict correct docking poses of both T4E (no data show) and ZM-241385 within a 2.0 Å structural deviation in comparison with experimental co-crystallised poses (Figure 2(B)). This is due to the greater volume of T4E-bound pocket that allows to accommodate bulky di-aryl substituted triazine as well as linear ZM-241385 ligands. Consequently, apoA2AR-T4E pocket appears to be more relevant to accommodate a wide diversity of chemical structures and was therefore used to perform our docking calculations.

A set of benzoxazole-based molecules have been docked, using Gold suite v5.2 within the Hermes v1.6 GUI (CCDC[®]), into the apoA2AR-T4E pocket and the ones that satisfied interactions with essential amino acids³² Phe168, Glu169, Trp246 and Asn253 were selected. ZM-241385 was shown as an example (Figure 2(B)) rather than the triazine because it is structurally closer to our benzoxazole ligands (Figure 2(C)). As illustrated in Figure 2(B), the central heterocycle of ZM-241385 interacts through a hydrogen bond with Asn253 and π -stacking with Phe168. Benzoxazole ring seems to recapitulate these interactions and could, therefore, constitute a good alternative as a central core. The furan, found in many A_{2A} antagonists, was selected to interact with Trp246 by aromatic interaction and thus made the antagonist character of our ligands.

Indeed, interaction with this key amino acid is well known to lock the A_{2A} receptor³³ and more generally class-A GPCR³⁴ in their inactive conformation. The 3,4 dimethoxyphenyl, found on Istradefylline, was also used to create this interaction. Furthermore, different nature and size of linkers was used to bring selected amine function. Indeed, these basic functions in designed ligands (Figure 2(C)) occupy the same pocket as the phenol part of the ZM-241385 and allow not only to interact with Glu169 through an ionic interaction but also to improve solubility.


Chemistry

Synthetic routes used to prepare benzoxazole derivatives are depicted in Schemes 1–3. The first set of benzoxazoles (**3a–d**) was prepared from commercially available aminophenols using two different synthetic routes (Scheme 1). The first one led to molecules **3a–d**, in two steps. Indeed, aminophenol derivatives **1a–e** reacted with appropriate acyl chlorides to give an equimolar mixture of mono and diacyl compounds which, when treated under basic conditions, gave amides **2a–d**. Subsequent cyclisation under acidic conditions afforded benzoxazoles **3a–d** in good yield³⁵. The second synthetic route allowed to transform aminophenol **1e** into compound **3e** in one step using T3P^{®36}.


A radical bromination of compounds **3a–c** was performed to generate the key brominated intermediates **4a–c** (Scheme 2) which allowed the introduction of the tertiary amine at different distances from the central heterocycle. Molecules **A1–8** displaying a one-methylene linker were obtained by reacting **4a–c** with various amines. Compound **A4** was synthesised from **A3** deprotection using TFA. Based on binding results (see Table 1), we focused the further synthetic effort on C-5 substituted compounds. Treatment of **4a** with potassium cyanide followed by an acidic hydrolysis gave carboxylic acid **6** which was subjected to amidification and reduction to afford target compounds **B1–4** (two-methylene linker).

To get molecules **C1–3** (three-methylene linker), the same procedure was used starting from carboxylic acids **8a–b**, obtained by malonic substitution performed on compounds **4a–b** followed by a basic hydrolysis and then a decarboxylation reaction by heating in DMF.


To obtain molecules **B5–6** (Scheme 3), ester function of compound **3e** was first reduced to alcohol **10** using LiAlH₄. The latter was then activated by the action of mesyl chloride to afford molecule **11**. The classical nucleophilic substitution was then performed to give compounds **B5–6**.



Scheme 1. Reagents and conditions: (a) i) ArCO₂H, SOCl₂, DMF, DCM, ii) Et₃N, EtOAc, aminophenol (**1a–d**); iii) NaOH, EtOH/H₂O, then 6 M HCl, 60–80% over 2 steps; (b) APTS, toluene, reflux, 70–80%, (c) T3P[®] (50% in EtOAc), DIPEA, 3,4-dimethoxybenzoic acid, 35%.



Scheme 2. Reagents and conditions: (a) NBS, benzoyl peroxide, CCl₄, reflux/hν (230 W), 60–75%; (b) R₂R₁NH, Et₃N, acetone, 50–85%; (c) TFA, DCM, 60%; (d) KCN, EtOH/H₂O, 50–60%; (e) H₂O/H₂SO₄/AcOH, reflux, 60–70%; (f) *i*) SOCl₂, toluene; *ii*) secondary amine, Et₃N, EtOAc, 50–60%; (g) LiAlH₄, THF, 60–75%; (h) *i*) dimethyl malonate, K₂CO₃, acetone; *ii*) NaOH, H₂O then 6 M HCl; *iii*) DMF, reflux, 40–43%.



Scheme 3. Reagents and conditions: (a) LiAlH₄, THF, 86%; (b) MsCl, Et₃N, DCM, quant.; (c) secondary amine, K₂CO₃, DMF, 60 °C, 25–28%.

To get target molecules with an amide (**F1–4**) or ethylamine linker (**E1–2**), 5-nitro benzoxazole derivative (**3d**) was reduced with hydrazine hydrate in the presence of Pd/C to afford compound **D1** (Scheme 4). Nucleophilic substitution gave compounds **E1** and **12**. The latter was deprotected under H₂ atmosphere afforded **E2**. To get molecules **F1–3**, treatment of **D1** with bromoacetyl bromide gave compound **13** that was substituted with various amines. Compound **F4** was obtained from **F3** deprotection using 6 M HCl in MeOH.

Structure-affinity relationship and early ADME studies































Affinities of benzoxazole derivatives for the human adenosine receptor were determined by a competitive radioligand displacement assay using [³H]-ZM241385 as radioligand¹⁸. All compounds were first screened at 10 μM concentration, and K_i values were

determined for those exhibiting a specific displacement superior to 35%.

The first set of derivatives **A1–8**, allowed drawing some early SARs (Table 1). First, comparing molecules **A1–6**, piperidine emerged as the preferred amine since a dramatic decrease in affinity was observed with all other selected ones. Surprisingly, the phenylpiperazine group featured in many A_{2A} antagonists was not tolerated in our series.

Regarding the position of the protonable amine function, compounds with a chain at the C-5 position appeared to exhibit a higher affinity than a compound with a chain at the C-6 position (molecules **A1** versus **A7**). Therefore, the protonable amine was placed in the C-5 position for following molecules. Concerning the aryl in position 2 of the benzoxazole, replacing the furan of **A1** by a 3,4-dimethoxyphenyl (**A8**) totally abolished affinity. This result is in agreement with literature since the preference of the adenosine A_{2A} receptor for furan substituent is well documented even if the 3,4-dimethoxyphenyl is found on the Istradefylline molecule^{7,13}.

Table 1. A_{2A} receptor affinity and cytotoxicity data of compounds A1–8, B1–6 and C1–3.

					
Cpd.	Ar		Ki ± SEM (μM) ^a / % inhib ^b	CC ₅₀ ^c / % inhib ^d	
ZM - 241385			0.0013 ± 0.002		
n = 1	A1		19 ± 3.1	1%	
	A2		10%	33%	
	A3		15%	30%	
	A4			15%	0%
	A5			11%	27%
	A6			14%	37%
	A7		-	5%	0%
A8			18%	0%	
n = 2	B1		7 ± 1.4	1%	
	B2		3%	0%	
	B3			11 ± 0.25	11%
	B4			10%	-
	B5			10%	40%
	B6			5%	19%
n = 3	C1			28%	-
	C2			10%	73 μM
	C3			11%	

^aDisplacement of specific [³H]-ZM 241385 binding to hA_{2A} receptors stably expressed in HEK293 cells. ^bDisplacement percentage of specific [³H]-ZM 241385 at 10 μM. ^cCompound concentration causing 50% of SY5Y cell death after 24 h treatment. ^dPercentage of dead SY5Y cells after 24 h treatment at 100 μM.




Table 2. A_{2A} receptor affinity and cytotoxicity data of compounds D1, E1–2 and F1–4.

Cpd.	R	D1, E1–E2		Cpd.	R	F1–F4	
		Ki (μM) ^a / % inhib ^b	CC ₅₀ ^c / % inhib ^d			Ki (μM) ^a / % inhib ^b	CC ₅₀ ^c / % inhib ^d
D1	H	10 ± 0.5	4%	F1		1 ± 0.08	1%
E1		11 ± 0.02	3%	F2		12%	1%
E2		15%	55 μM	F3		25%	12%
				F4		8%	0%

^aDisplacement of specific [³H]-ZM 241385 (2 nM) binding to hA_{2A} receptors stably expressed in HEK293 cells. ^bDisplacement percentage of specific [³H]-ZM 241385 at 10 μM. ^cCompound concentration causing 50% of SY5Y cell death after 24 h treatment. ^dPercentage of dead SY5Y cells after 24 h treatment at 100 μM.

Table 3. Preliminary ADME studies of F1.

Cpd.	Solubility (μM) PBS, pH 7.4 ^a	PPB (%) ^b	Permeability Caco-2- (10 ⁻⁶ cm/s) ^c		LogD _{7.4} ^d	HLM ^e t _{1/2} (min)	Cl _{int} ^f (μl/min/mg)
			A/B	B/A			
F1	184	83	50	24	2.33	11	630

^aEvaluated after 24 h stirring. ^bPPB = plasma protein binding. Compound was tested at 10 μM concentration. ^cPermeability = Compound was tested at 10 μM concentration at pH 6.5/7.4. ^dDetermined between a mixture PBS_{7.4}/octanol. ^eHLM = human liver microsomes. ^fCl_{int} = Compound was tested at 0.1 μM concentration.

Throughout Table 1, a similar tendency was observed for the other analogues suggesting that the piperidine chain at C-5 and the furan at C-2 of the benzoxazole ring are the best moieties for A_{2A} receptor affinity.

Concerning the linker, the comparison between **B1** and **A1** revealed that increasing the length to two carbons is beneficial for affinity. Moreover, as can be seen from affinity data of **E1** (11 μm) and **F1** (1 μm) (Table 2), a three-atom linker is also well tolerated. When comparing the latter two compounds, rigidifying the linker by incorporating an amide in place of an amine allows an increase in affinity. Besides, **F1** was found to be the most active compound in this series with a K_i value of 1 μm . As can be seen from Figure 2(C) the three-atom linker seems necessary to allow the interaction between the piperidine and Glu169. This ligand also exhibited a slight selectivity (see Table 4 of the Supplementary Material) versus A_1 receptor (2.5-fold). Concerning the two others adenosine receptors, preliminary studies showed a probably good interaction with hA_{2B} (73% inhibition at 10 μm) but a highly selectivity over adenosine A_3 receptor (11% inhibition at 10 μm). These informations do not constitute a brake for the development of **F1** as a potential drug candidate at this "hit" identification stage.

Interestingly, replacing the protonable amine of **E1** by the aminoethyl phenol chain present in ZM-241385 (**E2**) led to a drastic drop in affinity, highlighting the importance of the tertiary amine in this series.

Finally, compound **D1** ($K_i = 10 \mu\text{m}$), despite the lack of the protonable amine, could be considered a promising hit for future A_{2A} antagonist development. Indeed, given its low molecular weight ($\text{MW} = 200 \text{ g}\cdot\text{mol}^{-1}$), it offers multiple possibilities for further modifications.

The most interesting compounds of this series, **F1** and **D1**, were subjected to preliminary pharmacokinetics studies (Table 3). As expected, molecule **F1** displaying a protonable amine exhibits a higher solubility (184 μm) than molecule **D1** (28 μm) in PBS solution at pH 7.4. When compared to reference A_{2A} antagonists, these two hits show a higher solubility^{37–40}. Indeed, Istradefylline (KW-6002) and Preladenant (SCH 4208⁸), the two reference antagonists exhibit solubility values of 1.5 μm and 20 nm, respectively.

Moreover, **F1** displayed a good partition coefficient ($\log D_{7.4}$) of 2.33 which is within the same range as reference A_{2A} antagonists currently in clinical studies⁴¹. These results confirm the importance of having a tertiary amine which allows a sharp increase in solubility while keeping a good $\log D_{7.4}$ value. At 10 μm concentration, a correct value of permeability coefficient (Caco-2 cells, pH 6.5/7.4) is observed ($50 \times 10^{-6} \text{ cm/s}$). The efflux ratio ($B/A_{A/B}$) of 0.5 also suggested that **F1** was probably distributed through P-glycoprotein (P-gp), an important transporter protein found in cell throughout the body. A good human plasma protein binding (mean of 83%) was also observed compared to reference A_{2A} antagonist which expressed high PPB around 98%. Nevertheless, this compound also exhibited a high clearance and thus a short half time in human liver microsome at 0.1 μm .

Finally, no toxicity was observed for active compounds at 100 μm when tested on neuroblastoma cell lines (SY5Y, Tables 1,2).

Conclusions

Reported results showed a set of benzoxazole derivatives, diversely substituted at the C-2 and C-5 positions, as new "hits" molecules for adenosine A_{2A} receptor. Among the synthesised compounds, those featured by a furan at the C-2 position combined with a piperidine and an amide-based linker at C-5 resulted in the most

active compound (**F1**) toward the $hA_{2A}R$ ($K_i = 1 \mu\text{m}$). Furthermore, the latter presented good preliminary ADME properties with a very interesting solubility (184 μm) as well as good $\log D_{7.4}$ (2.33) without cytotoxicity at 100 μm . Thus, both **F1** and **D1**, which can be easily modulated in position 7, appear to be promising starting points for further optimisation.

Acknowledgements

We express our thanks to the 300MHz NMR facilities funded by the Region Nord-Pas de Calais (France), the Ministère de la Jeunesse, de l'Éducation Nationale et de la Recherche (MJENR) and the Fonds Européens de Développement Regional (FEDER). This work was supported by Lille 2 University, ANR "Adoratau", Comue Univ Lille Nord de France. Romain Duroux is the recipient of a fellowship from Lille 2 University.

Disclosure statement

No potential conflict of interest was reported by the authors.

References

- Fargo K, Bleiler L. Alzheimer's disease facts and figures. *Alzheimers Dement* 2014;10:47–92.
- Panza F, Solfrizzi V, Barulli MR, et al. Coffee, tea, and caffeine consumption and prevention of late-life cognitive decline and dementia: a systematic review. *J Nutr Health Aging* 2015;19:313–28.
- Solfrizzi V, Panza F, Imbimbo BP, et al. Coffee consumption habits and the risk of mild cognitive impairment: the Italian longitudinal study on aging. *J Alzheimers Dis* 2015;47:889–99.
- Flaten V, Laurent C, Coelho JE, et al. From epidemiology to pathophysiology: what about caffeine in Alzheimer's disease? *Biochem Soc Trans* 2014;42:587–92.
- Xu K, Di Luca DG, Orrú M, et al. Neuroprotection by caffeine in the MPTP model of Parkinson's disease and its dependence on adenosine A_{2A} receptors. *Neuroscience* 2016;322:129–37.
- Dall'igna OP, Fett P, Gomes MW, et al. Caffeine and adenosine A_{2A} receptor antagonists prevent beta-amyloid (25–35)-induced cognitive deficits in mice. *Exp Neurol* 2007;203:241–5.
- Preti D, Baraldi PG, Moorman AR, et al. History and perspectives of A_{2A} adenosine receptor antagonists as potential therapeutic agents. *Med Res Rev* 2015;35:790–848.
- Shook BC, Jackson PF. Adenosine A_{2A} receptor antagonists and Parkinson's disease. *ACS Chem Neurosci* 2011;2:555–67.
- Hauser RA, Olanow CW, Kieburtz KD, et al. Tozadenant (SYN115) in patients with Parkinson's disease who have motor fluctuations on levodopa: a phase 2b, double-blind, randomised trial. *Lancet Neurol* 2014;13:767–76.
- Pinna A. Adenosine A_{2A} receptor antagonists in Parkinson's disease: progress in clinical trials from the newly approved istradefylline to drugs in early development and those already discontinued. *CNS Drugs* 2014;28:455–74.
- Kondo T, Mizuno Y, Japanese Istradefylline Study Group. A long-term study of istradefylline safety and efficacy in patients with Parkinson disease. *Clin Neuropharmacol* 2015;38:41–6.

12. Laurent C, Burnouf S, Ferry B, et al. A_{2A} adenosine receptor deletion is protective in a mouse model of Tauopathy. *Mol Psychiatry* 2016;21:97–107.
13. de Lera Ruiz M, Lim YH, Zheng J. Adenosine A_{2A} receptor as a drug discovery target. *J Med Chem* 2014;57:3623–50.
14. Müller CE, Ferré S. Blocking striatal adenosine A_{2A} receptors: a new strategy for basal ganglia disorders. *Recent Pat CNS Drug Discov* 2007;2:1–21.
15. Robinson SJ, Petzer JP, Terre'Blanche G, et al. 2-Aminopyrimidines as dual adenosine A_1/A_{2A} antagonists. *Eur J Med Chem* 2015;104:177–88.
16. Tokuyama H, Okano K, Fujiwara H, et al. Total synthesis of dictyodendrins A-E. *Chem Asian J* 2011;6:560–72.
17. Jones G, Willett P, Glen RC. Molecular recognition of receptor sites using a genetic algorithm with a description of desolvation. *J Mol Biol* 1995;245:43–53.
18. Lopes LV, Cunha RA, Ribeiro JA. Cross talk between A_1 and A_{2A} adenosine receptors in the hippocampus and cortex of young adult and old rats. *J Neurophysiol* 1999;82:3196–203.
19. Townsend-Nicholson A, Schofield PR. A threonine residue in the seventh transmembrane domain of the human A_1 adenosine receptor mediates specific agonist binding. *J Biol Chem* 1994;269:2373–6.
20. Stehle JH, Rivkees SC, Lee JJ, et al. Molecular cloning and expression of the cDNA for a novel A_2 -adenosine receptor subtype. *Mol Endocrinol* 1992;6:384–93.
21. Salvatore CA, Jacobson MA, Taylor HE, et al. Molecular cloning and characterization of the human A_3 adenosine receptor. *Proc Natl Acad Sci U S A* 1993;90:10365–9.
22. Lipinski CA, Lombardo F, Dominy BW, et al. Experimental and computational approaches to estimate solubility and permeability in drug discovery and development settings. *Adv Drug Deliv Rev* 1997;46:3–26.
23. Sangster J. Wiley series in solution chemistry. Volume 2. Chichester: John Wiley and Sons; 1997.
24. Banker MJ, Clark TH, Williams JA. Development and validation of a 96-well equilibrium dialysis apparatus for measuring plasma protein binding. *J Pharm Sci* 2003;92:967.
25. Hidalgo IJ, Raub TJ, Borchardt RT. Characterization of the human colon carcinoma cell line (Caco-2) as a model system for intestinal epithelial permeability. *Gastroenterology* 1989;96:736–49.
26. Obach RS, Baxter JG, Liston TE, et al. The prediction of human pharmacokinetic parameters from preclinical and *in vitro* metabolism data. *J Pharmacol Exp Ther* 1997;283:46–58.
27. Jaakola VP, Griffith MT, Hanson MA, et al. The 2.6 angstrom crystal structure of a human A_{2A} adenosine receptor bound to an antagonist. *Science* 2008;322:1211–17.
28. Liu W, Chun E, Thompson AA, et al. Structural basis for allosteric regulation of GPCRs by sodium ions. *Science* 2012;337:232–6.
29. Congreve M, Andrews SP, Doré AS, et al. Discovery of 1,2,4-triazine derivatives as adenosine A_{2A} antagonists using structure based drug design. *J Med Chem* 2012;55:1898–903.
30. Sabbadin D, Ciancetta A, Moro S. Perturbation of fluid dynamics properties of water molecules during G protein-coupled receptor-ligand recognition: the human A_{2A} adenosine receptor as a key study. *J Chem Inf Model* 2014;54:2846–55.
31. Lenselink EB, Beuming T, Sherman W, et al. Selecting an optimal number of binding site waters to improve virtual screening enrichments against the adenosine A_{2A} receptor. *J Chem Inf Model* 2014;54:1737–46.
32. Keränen H, Gutiérrez-de-Terán H, Åqvist J. Structural and energetic effects of A_{2A} adenosine receptor mutations on agonist and antagonist binding. *PLoS One* 2014;9:e108492.
33. Yuan S, Hu Z, Filipek S, et al. W246(6.48) opens a gate for a continuous intrinsic water pathway during activation of the adenosine A_{2A} receptor. *Angew Chem Int Ed Engl* 2015;54:556–9.
34. Venkatakrisnan AJ, Deupi X, Lebon G, et al. Molecular signatures of G-protein-coupled receptors. *Nature* 2013;494:185–94.
35. Wisastra R, Ghizzoni M, Boltjes A, et al. Anacardic acid derived salicylates are inhibitors or activators of lipoxigenases. *Bioorg Med Chem* 2012;20:5027–32.
36. Wen X, Bakali JE, Deprez-Poulain R, et al. Efficient propylphosphonic anhydride (@T3P) mediated synthesis of benzothiazoles, benzoxazoles and benzimidazoles. *Tetrahedron Lett* 2012;53:2440–3.
37. Evaluation and Licensing Division, Pharmaceutical and Food Safety Bureau Ministry of Health. Labour and Welfare [Online] 2015. Available from: <https://www.pmda.go.jp/files/000153870.pdf>
38. Müller CE. Prodrug approaches for enhancing the bioavailability of drugs with low solubility. *Chem Biodivers* 2009;6:2071–83.
39. Slee DH, Zhang X, Moorjani M, et al. Identification of novel, water-soluble, 2-amino-N-pyrimidin-4-yl acetamides as A_{2A} receptor antagonists with *in vivo* efficacy. *J Med Chem* 2008;51:400–6.
40. Mikkelsen GK, Langgård M, Schroder TJ, et al. Synthesis and SAR studies of analogues of 4-(3,3-dimethyl-butylamino)-3,5-difluoro-N-thiazol-2-yl-benzamide (Lu AA41063) as adenosine A_{2A} receptor ligands with improved aqueous solubility. *Bioorg Med Chem Lett* 2015;25:1212–16.
41. Zheng J, Yang Z, Li X, et al. Optimization of 6-heterocyclic-2-(1H-pyrazol-1-yl)-N-(pyridin-2-yl)pyrimidin-4-amine as potent adenosine A_{2A} receptor antagonists for the treatment of Parkinson's disease. *ACS Chem Neurosci* 2014;5:674–82.