



Mixed quantum skew Howe duality and link invariants of type A

Hoel Queffelec, Antonio Sartori

► To cite this version:

Hoel Queffelec, Antonio Sartori. Mixed quantum skew Howe duality and link invariants of type A. Journal of Pure and Applied Algebra, 2019, 223 (7), pp.2733-2779. <10.1016/j.jpaa.2018.09.014>. <hal-02063209>

HAL Id: hal-02063209

<https://hal.science/hal-02063209v1>

Submitted on 21 Oct 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons CC BY-NC 4.0 - Attribution - Non-commercial use - International License

Mixed quantum skew Howe duality and link invariants of type A

Hoel Queffelec^a, Antonio Sartori¹

^a*Institut Montpellierain Alexander Grothendieck, CNRS, Univ. Montpellier*

Abstract

We define a ribbon category $\mathbf{Sp}(\beta)$, depending on a parameter β , which encompasses Cautis, Kamnitzer and Morrison's spider category, and describes for $\beta = m - n$ the monoidal category of representations of $U_q(\mathfrak{gl}_{m|n})$ generated by exterior powers of the vector representation and their duals. We identify this category $\mathbf{Sp}(\beta)$ with a direct limit of quotients of a dual idempotented quantum group $\mathbf{U}_q(\mathfrak{gl}_{r+s})$, proving a mixed version of skew Howe duality in which exterior powers and their duals appear at the same time. We show that the category $\mathbf{Sp}(\beta)$ gives a unified natural setting for defining the colored $\mathfrak{gl}_{m|n}$ link invariant (for $\beta = m - n$) and the colored HOMFLY-PT polynomial (for β generic).

Keywords: Webs, spider category, quantum Lie superalgebras, skew-Howe duality, HOMFLY-PT polynomial, Reshetikhin-Turaev invariants.

Contents

1	Introduction	2
2	The category of tangles	9
3	The general linear Lie superalgebra	11

Email addresses: hoel.queffelec@umontpellier.fr (Hoel Queffelec),
antoniosartorimath@gmail.com (Antonio Sartori)

Preprint submitted to Journal of Pure and Applied Algebra

August 16, 2018

4	Super skew Howe duality	19
5	The category Sp^+	25
6	Graphical calculus for duals	34
7	Mixed skew Howe duality and the idempotented quantum group	56
8	Link invariants	64

1. Introduction

In 1985 Jones [Jones \(1985\)](#) defined a remarkable new polynomial invariant of links. A few years later, Reshetikhin and Turaev [Reshetikhin and Turaev \(1990\)](#) reinterpreted the Jones polynomial and generalized it by defining a whole class of quantum link invariants. Their work, based on the use of quantum groups and built upon the introduction of ribbon categories, opened the way to a world of connections between knot theory and representation theory. In particular, they defined quantum link invariants attached to finite dimensional representations of semisimple Lie algebras, recovering the Jones polynomial for \mathfrak{sl}_2 . Although the Alexander polynomial [Alexander \(1928\)](#) was not originally covered by Reshetikhin and Turaev's process, it has been well-known to experts that it is possible to recover it using the super quantum group $U_q(\mathfrak{gl}_{1|1})$ [Kauffman and Saleur \(1991\)](#); [Viro \(2006\)](#) (see also [Sartori \(2015\)](#) for a more accessible explanation and for additional references). (An alternative and earlier construction uses the quantum group $U_q(\mathfrak{sl}_2)$ for q a root of unity [Murakami \(1992\)](#), but we will not pursue this approach.) In particular, both the Jones and the Alexander polynomials fit together inside the family of quantum link invariants associated to the Lie superalgebras $\mathfrak{gl}_{m|n}$. We shall mention another link invariant, the HOMFLY-PT polynomial [Freyd et al. \(1985\)](#); [Przytycki and Traczyk \(1987\)](#), which in a certain sense could be interpreted as a limit version for $m \rightarrow \infty$ of the \mathfrak{gl}_m link invariants, but does not fit *a priori* in the setting of quantum link invariants.

A major new step in knot theory was the idea of *categorification*. Khovanov's seminal work [Khovanov \(2000, 2002, 2006\)](#) lifting the Jones polynomial to a homology theory, initiated an effort to categorify the polynomial knot

invariants, which produced an \mathfrak{sl}_3 [Khovanov \(2004\)](#) and then an \mathfrak{sl}_m link homology [Khovanov and Rozansky \(2008\)](#) via matrix factorizations, as well as a triply graded link homology theory [Khovanov \(2007\)](#), based on Soergel bimodules, categorifying the HOMFLY-PT polynomial. In a parallel and connected way, a categorification program for quantum groups and their representations started and prospered in the last fifteen years, using tools from representation theory and geometry. Milestones were the categorifications using the BGG category \mathcal{O} , initiated in [Bernstein et al. \(1999\)](#); [Frenkel et al. \(2006\)](#) and the definition of the Khovanov-Lauda-Rouquier algebra [Khovanov and Lauda \(2009, 2011, 2010\)](#); [Rouquier \(2008\)](#). We point out here that, although a categorification of the Alexander polynomial does exist [Manolescu et al. \(2007\)](#), it comes from a completely different theory (symplectic geometry tools and in particular Heegaard-Floer homology), and a representation theoretical interpretation of it is still missing. Conjecturally, there should be a whole family of $\mathfrak{gl}_{m|n}$ link homologies which should encompass all previously mentioned categorifications (see also [Gorsky et al. \(2013\)](#)).

In a remarkable paper, Cautis, Kamnitzer and Morrison [Cautis et al. \(2014\)](#) showed how the classical skew Howe duality [Howe \(1989, 1995\)](#) for the pair $(\mathfrak{gl}_m, \mathfrak{gl}_k)$ can be used to interpret the braiding of $U_q(\mathfrak{gl}_m)$ -representations as the action of a dual quantum group $U_q(\mathfrak{gl}_k)$. In particular, they describe the combinatorics of intertwiners of tensor products of exterior powers of the natural representation \mathbb{C}_q^m of $U_q(\mathfrak{gl}_m)$ using a spider category, which is equivalent to the direct limit for $k \rightarrow \infty$ of some quotients of the idempotented versions of a dual quantum group $U_q(\mathfrak{gl}_k)$. This approach seems to give the right combinatorial setting for categorification [Lauda et al. \(2015\)](#); [Mackaay and Yonezawa \(2013\)](#); [Tubbenhauer \(2014\)](#); [Queffelec and Rose \(2016\)](#), and has been recently used to give uniqueness results for link homology theories [Mackaay and Webster \(2015\)](#).

The work of Cautis, Kamnitzer and Morrison actually really applies to braids. Indeed, in the process of assigning an intertwiner of representations to an oriented tangle (and, in particular, a knot or link), one needs to have both the standard representation \mathbb{C}_q^m and its dual at ones disposal. The fact that Cautis, Kamnitzer and Morrison's result yields \mathfrak{gl}_m link invariants (and can be used for their categorification) is based on the identification of the exterior powers

$$\bigwedge^a \mathbb{C}^m \cong \left(\bigwedge^{m-a} \mathbb{C}^m \right)^* \quad \text{as } \mathfrak{sl}_m\text{-representations.} \quad (1.1)$$

With the aim of studying $\mathfrak{gl}_{m|n}$ link invariants and possibly their categorification, this has two main limitations.

The first one concerns its extension to the super case, which is needed to define $\mathfrak{gl}_{m|n}$ link invariants (and, in particular, the Alexander polynomial). Here we denote by $\mathfrak{gl}_{m|n}$ the general Lie superalgebra and by $\mathbb{C}_q^{m|n}$ its standard representation. A super version of skew Howe duality does hold for the pair $(\mathfrak{gl}_{m|n}, \mathfrak{gl}_k)$ and can be used, in a similar way as in [Cautis et al. \(2014\)](#), to describe intertwiners of tensor products of exterior powers of $\mathbb{C}_q^{m|n}$. This approach can even be used to categorify these representations, see [Sartori \(2016\)](#). However, since in the case $n \neq 0$ no exterior power of the standard representation of $\mathfrak{gl}_{m|n}$ is isomorphic to its dual, the dual of the standard representation does not appear in skew Howe duality. Indeed, there is a qualitative distinction: the representation appearing in skew Howe duality is always semisimple, while in general mixed tensor products of the vector representation together with its dual are not.

The second limitation appears when one wants to inspect the HOMFLY-PT polynomial. In this case, roughly speaking, one would like to take a limit for $m \rightarrow \infty$. The identification (1.1) above then becomes meaningless and again the dual of the standard representation disappears from the picture.

The goal of this paper is to overcome these two limitations. Our main result is an extension of skew Howe duality to a more general picture where both exterior powers of the vector representation and of its dual appear at the same time, and where one can make sense of the limit $m \rightarrow \infty$ by taking a generic value of the parameter β .

An overview of the categories and functors that we will consider is presented in figure 1. The top row of figure 1 is the generalization of [Cautis et al. \(2014\)](#) to the super case of $\mathfrak{gl}_{m|n}$, and describes intertwiners of exterior powers of the natural representation $\mathbb{C}_q^{m|n}$. The second row is our extension to exterior powers and their duals. The third row will serve as auxiliary tool. We will describe some of the categories and functors involved in the following.

Graphical calculus for duals

Let m, n be two non-negative integers. We will always set $d = m - n$. We denote by $\text{Rep}_{m|n}^+$ (respectively, $\text{Rep}_{m|n}$) the monoidal category of $U_q(\mathfrak{gl}_{m|n})$ -representations generated by exterior powers of the natural representation




Figure 1: Overview of the categories and functors involved in our construction. (The third column makes sense only for $\beta = m - n$.)

$\mathbb{C}_q^{m|n}$ (respectively, exterior powers of $\mathbb{C}_q^{m|n}$ and their duals). The first goal of our paper is to obtain a unified graphical calculus for $\text{Rep}_{m|n}$. This extends in two directions the work of Cautis, Kamnitzer and Morrison, who described the category $\text{Rep}_{m|0}^+$ (and with the trick (1.1) actually described $\text{Rep}_{m|0}$, if restricting to \mathfrak{sl}_m).

The graphical calculus of Cautis et al. (2014) is given by a spider category (introduced in rank 2 already in Kuperberg (1996)) describing $\text{Rep}_{m|0}^+$. The extension to the super case (replacing \mathfrak{gl}_m by $\mathfrak{gl}_{m|n}$), which we will develop in all details (see section 4), is relatively easy. Similarly to Cautis et al. (2014) we can define a monoidal category Sp^+ together with a full functor $\mathcal{G}_{m|n}^+ : \text{Sp}^+ \rightarrow \text{Rep}_{m|n}^+$ (theorem 5.2). The category Sp^+ is some sort of universal tool, built in order to describe all categories $\text{Rep}_{m|n}^+$ at the same time. It is an interesting task to find all relations in $\text{Rep}_{m|n}^+$, i.e. to describe a monoidal ideal $\mathcal{I}_{m|n}$ of Sp^+ such that $\mathcal{G}_{m|n}^+$ descends to a fully faithful functor $\text{Sp}^+/\mathcal{I}_{m|n} \rightarrow \text{Rep}_{m|n}^+$. This is very easy for $n = 0$ (one just kills objects of Sp^+ labeled by integers greater than m , whence the notation $\text{Sp}_{\leq m}^+ = \text{Sp}^+/\mathcal{I}_{m|0}$) and it is possible also in general, although the relations are not so nice (see Grant (2016) for the case of $\mathfrak{gl}_{1|1}$). An interesting fact is that we can lift the braiding of all the categories $\text{Rep}_{m|n}^+$ directly to Sp^+ in a uniform way, turning it into a braided category (proposition 5.12).

Now, with the goal of studying link invariants and their categorification, we are interested in understanding the full category $\text{Rep}_{m|n}$, and not only its “positive part” $\text{Rep}_{m|n}^+$. But, as we already explained, in the case $n \neq 0$ the trick (1.1) does not work anymore. Hence we define a rigid category \mathbf{Sp} by adding duals to \mathbf{Sp}^+ , and we show that the functor $\mathcal{G}_{m|n}^+$ extends to a full functor $\mathcal{G}_{m|n}: \mathbf{Sp} \rightarrow \text{Rep}_{m|n}$ (proposition 6.2). The category \mathbf{Sp} , however, is too big: its endomorphism spaces are infinite-dimensional and the functor $\mathcal{G}_{m|n}$, hence, cannot be faithful. Moreover, there is no map from the category of oriented framed tangles to \mathbf{Sp} which could give link invariants. Hence we introduce a quotient $\mathbf{Sp}(\beta)$ of \mathbf{Sp} , depending on a parameter β which can be either a generic variable (corresponding to the second variable of the HOMFLY-PT polynomial) or an integer. Our first main result then is a graphical complete description of the category $\text{Rep}_{m|n}$, modulo the relevant ideal of relations (not described in this paper):

Main Theorem A (See theorem 6.6 and proposition 6.16). *For $\beta = d = m - n$ we get an induced functor $\mathcal{G}_{m|n}: \mathbf{Sp}(d) \rightarrow \text{Rep}_{m|n}$ which is still full. Moreover, this descends to a fully faithful functor $\mathbf{Sp}(d)/(\mathcal{I}_{m|n}) \rightarrow \text{Rep}_{m|n}$. (Here $(\mathcal{I}_{m|n})$ denotes the monoidal ideal generated by the image of $\mathcal{I}_{m|n}$ in $\mathbf{Sp}(d)$).*

Similarly as before, the ribbon structure of the representation categories $\text{Rep}_{m|n}$ lifts in a unified way to $\mathbf{Sp}(\beta)$, which hence is a ribbon category (theorem 6.13).

Mixed skew-Howe duality and the doubled Schur algebra

In Cautis et al. (2014) the main tool yielding the description of the category Rep_m^+ is skew Howe duality, which gives two commuting actions

$$U_q(\mathfrak{gl}_m) \curvearrowright \bigwedge_q^N (\mathbb{C}_q^m \otimes \mathbb{C}_q^r) \curvearrowright U_q(\mathfrak{gl}_r) \quad (1.2)$$

producing a description of $U_q(\mathfrak{gl}_m)$ intertwiners by elements of $U_q(\mathfrak{gl}_r)$, and *vice versa*. In particular, this provides an identification of the category \mathbf{Sp}^+ with the direct limit for $r \rightarrow \infty$ of a quotient of the idempotent version of $U_q(\mathfrak{gl}_r)$. Furthermore, the action of the braid group on the $U_q(\mathfrak{gl}_m)$ -modules is identified as the quantum Weyl group action generated by Lusztig’s symmetries Lusztig (1993); Kamnitzer and Tingley (2009); Cautis et al. (2010, 2014).

Surprisingly, it is possible to generalize skew Howe duality and get the dual $(\mathbb{C}_q^{m|n})^*$, together with its exterior powers, into the picture. In our setting, we proceed the other way around with respect to [Cautis et al. \(2014\)](#), and we deduce such generalization from our graphical calculus:

Main Theorem B (See theorem [7.7](#) and corollary [7.12](#)). *On the representation*

$$\bigoplus_{\substack{a_1 + \dots + a_r - a_{r+1} \\ - \dots - a_{r+s} + (m-n)s = N}} \bigwedge_q^{a_1} \mathbb{C}_q^{m|n} \otimes \dots \otimes \bigwedge_q^{a_r} \mathbb{C}_q^{m|n} \otimes \bigwedge_q^{a_{r+1}} (\mathbb{C}_q^{m|n})^* \otimes \dots \otimes \bigwedge_q^{a_{r+s}} (\mathbb{C}_q^{m|n})^* \quad (1.3)$$

there is a naturally defined $U_q(\mathfrak{gl}_{m|n})$ -equivariant action of the quantum group $U_q(\mathfrak{gl}_{r+s})$ and of its idempotent version $\dot{U}_q(\mathfrak{gl}_{r+s})$. The latter action generates the full centralizer of the $U_q(\mathfrak{gl}_{m|n})$ -action. This induces an equivalence of categories between $\mathbf{Sp}(d)$ and the direct limit of a quotient of $\dot{U}_q(\mathfrak{gl}_{r+s})$ for $r, s \rightarrow \infty$.

Such a quotient was already introduced by the two authors in [Queffelec and Sartori \(2018\)](#) with the name of *doubled Schur algebra*. For an illustrative picture see figure [2](#) and example [7.10](#) on page [62](#).

To the best of our knowledge, the dual actions on [\(1.3\)](#) are new, also in the non-quantized setting. A related duality in the non-quantized setting was analyzed in ([Cheng et al., 2004](#), section 3), where the roles of $\mathfrak{gl}_{m|n}$ and \mathfrak{gl}_{r+s} are swapped. However, the duality from [Cheng et al. \(2004\)](#) is somehow easier, since the representation involved is semisimple, while [\(1.3\)](#) is in general not semisimple. Moreover, it is not clear how to quantize the results from ([Cheng et al., 2004](#), section 3); indeed, it is not even evident how to define the dual action in the quantized setting. We point out that it should be possible to generalize our methods in order to obtain dual actions of $U_q(\mathfrak{gl}_{m|n})$ and $U_q(\mathfrak{gl}_{r+s|r'+s'})$ on a bigger space than [\(1.3\)](#) (which would contain also symmetric powers). For $q = 1$, this would include the duality of [Cheng et al. \(2004\)](#) as a special case.

Link invariants, categorification and the parameter β

Let us denote by $\mathbf{Tangles}^{\text{lab}}$ the category of oriented framed tangles whose strands are labeled by positive numbers. For all β we have a functor

$$\mathcal{Q}_\beta: \mathbf{Tangles}^{\text{lab}} \rightarrow \mathbf{Sp}(\beta),$$

which for $\beta = d$, composed with the action of $\mathbf{Sp}(d)$ on $\mathbf{Rep}_{m|n}$, gives the Reshetikhin-Turaev invariant (proposition 6.15) in the uniting framework of Murakami et al. (1998). This allows us to define inside $\mathbf{Sp}(d)$ the $\mathfrak{gl}_{m|n}$ link invariant labeled by exterior powers of the vector representation. The study of the link invariants is only briefly outlined in this paper, but the results we obtain here are central in the second part of our first paper Queffelec and Sartori (2018). The reader interested in the consequences of our approach in the study of link invariants should refer to Queffelec and Sartori (2018).

We stress that this could also be of great interest for *categorification*. Indeed, after Cautis et al. (2014), the categorification of \mathfrak{gl}_m link invariants can be interpreted as a categorification of the functor $\mathbf{Tangles}^{\ell ab} \rightarrow \mathbf{Sp}_{\leq m}^+$, i.e. as a 2-functor from the 2-category of tangles to a 2-categorical lift of $\mathbf{Sp}_{\leq m}^+$ (cf. Lauda et al. (2015); Queffelec and Rose (2016)). What such a lift should be becomes clear after $\mathbf{Sp}_{\leq m}^+$ is identified by Cautis et al. (2014) with a sum of quotients of idempotented version of quantum groups: the natural target of such 2-functor then is some version of the KLR 2-category Khovanov and Lauda (2010); Rouquier (2008).

Similarly, in our case, the categorification of $\mathfrak{gl}_{m|n}$ link invariants should be a categorification of the functor $\mathbf{Tangles}_q^{\ell ab} \rightarrow \mathbf{Sp}(d)$. We believe that the identification of $\mathbf{Sp}(d)$ with the direct limit of doubled Schur algebras can point in the right direction for lifting $\mathbf{Sp}(d)$ to a 2-category and hence constructing a categorification of $\mathfrak{gl}_{m|n}$ link invariants via categorification of such doubled Schur algebras.

Moreover, we point out that the definition of the category $\mathbf{Sp}(\beta)$ depends on a parameter β , and the action of $\mathbf{Sp}(\beta)$ on $U_q(\mathfrak{gl}_{m|n})$ representation comes when β is specialized to $d = m - n$. Nevertheless, our construction of $\mathbf{Sp}(\beta)$ makes sense also for a generic value of β . In this case, the natural functor $Q_\beta: \mathbf{Tangles}^{\ell ab} \rightarrow \mathbf{Sp}(\beta)$ can be used to define the HOMFLY-PT polynomial Freyd et al. (1985); Przytycki and Traczyk (1987). A categorification of this functor, via a similar strategy as in the case $\beta = d$ explained above, could be useful for better understanding the triply graded link homology Khovanov (2007) and its relation with \mathfrak{gl}_m and more generally $\mathfrak{gl}_{m|n}$ link homology (see also Gorsky et al. (2013) for some conjectural properties).

Structure of the paper

In section 2 we will fix some notations and recall the definition of the quantized oriented Brauer algebra. In section 3 we will collect some basic facts on the quantum enveloping superalgebra $U_q(\mathfrak{gl}_{m|n})$ and its representations. Section 4 will be devoted to the proof of the quantized super version of classical skew Howe duality. In section 5 we will recall the definition of the braided category \mathbf{Sp}^+ from Cautis et al. (2014) and we will construct its action on $\mathbf{Rep}_{m|n}^+$. In the following section 6 we will define our main object, the category $\mathbf{Sp}(\beta)$, by adding duals to \mathbf{Sp}^+ . We will prove that this category acts fully on $\mathbf{Rep}_{m|n}$ for $\beta = m - n$ and we will construct its ribbon structure. In section 7 we will give the quantum group interpretation of $\mathbf{Sp}(\beta)$, proving our main result. Finally, in section 8 we describe some applications to link invariants.

Acknowledgments

We would like to thank Christian Blanchet, Aaron Lauda, Tony Licata, Scott Morrison, David Rose, Catharina Stroppel, Daniel Tubbenhauer, Emmanuel Wagner, Weiqiang Wang and Paul Wedrich for useful discussions and comments, as well as the anonymous referee.

2. The category of tangles

The parameter β

All our diagrammatic categories will be defined over a field \mathbb{k} containing the complex numbers \mathbb{C} and two elements q and q^β which are not roots of the unity. The first main case we are interested in is $\mathbb{k} = \mathbb{C}(q, q^\beta)$, i.e. a transcendental extension of $\mathbb{C}(q)$ by a formal variable q^β . In this case we say that β is *generic*. The second main example is when β is some integer d and $\mathbb{k} = \mathbb{C}(q)$. We will sometimes write \mathbb{C}_q instead of $\mathbb{C}(q)$ for shortness.

For $x \in \mathbb{Z}\beta + \mathbb{Z}$ and $k \in \mathbb{N}$ we define

$$[x] = \frac{q^x - q^{-x}}{q - q^{-1}}, \quad (2.1)$$

$$\begin{bmatrix} x \\ k \end{bmatrix} = \frac{[x][x-1] \cdots [x-k+1]}{[k][k-1] \cdots [1]}. \quad (2.2)$$

The following identities hold for all $x, y \in \mathbb{Z}\beta + \mathbb{Z}$ and $k \in \mathbb{N}$, and will be used often through the paper (the last one will be proved in detail in Lemma 6.14):

$$[x][y+1] - [x+1][y] = [x-y], \quad (2.3)$$

$$[x][y] - [x-1][y-1] = [x+y-1], \quad (2.4)$$

$$[x+y] = q^y[x] + q^{-x}[y] = q^{-y}[x] + q^x[y], \quad (2.5)$$

$$\begin{bmatrix} x+1 \\ k \end{bmatrix} = q^{x+1-k} \begin{bmatrix} x \\ k-1 \end{bmatrix} + q^{-k} \begin{bmatrix} x \\ k \end{bmatrix} = q^k \begin{bmatrix} x \\ k \end{bmatrix} + q^{-x-1+k} \begin{bmatrix} x \\ k-1 \end{bmatrix}, \quad (2.6)$$

$$\sum_{\ell=0}^k (-q)^\ell \begin{bmatrix} x-\ell \\ k \end{bmatrix} \begin{bmatrix} k \\ \ell \end{bmatrix} = q^{-kx+k^2}. \quad (2.7)$$

Tangles

Let **Tangles** be the monoidal category of oriented framed tangles. Its objects are sequences of $\{1, 1^*\}$ and its morphisms are oriented framed tangles modulo isotopy. The category **Tangles** can be described in terms of generators and relations, see for example [Ohtsuki \(2002\)](#). Let also $\mathbf{Tangles}_q$ be the additive closure of its \mathbb{k} -linear version, with morphisms being \mathbb{k} -vector spaces

$$\mathbf{Tangles}_q(\boldsymbol{\eta}, \boldsymbol{\eta}') = \text{span}_{\mathbb{k}} \mathbf{Tangles}(\boldsymbol{\eta}, \boldsymbol{\eta}') \quad (2.8)$$

and where as objects we have formal (finite) direct sums of objects of **Tangles**.

We let $\mathbf{Tangles}^{\text{lab}}$ be the category consisting of oriented framed tangles whose connected components are labeled by positive integers. The objects of $\mathbf{Tangles}^{\text{lab}}$ are sequences of $\{a, a^* \mid a \in \mathbb{Z}_{>0}\}$. Similarly as before, we denote by $\mathbf{Tangles}_q^{\text{lab}}$ its additive \mathbb{k} -linear version. There are obvious inclusions $\mathbf{Tangles} \hookrightarrow \mathbf{Tangles}^{\text{lab}}$ and $\mathbf{Tangles}_q \hookrightarrow \mathbf{Tangles}_q^{\text{lab}}$ given by labeling all strands by 1.

We will regard all these tangle categories as monoidal categories, with monoidal unit being the empty sequence.

The quantized oriented Brauer category

We recall the definition of the quantized oriented Brauer algebra/category, following [Weimer \(2005\)](#) and [Dipper et al. \(2014\)](#) (see also [Blanchet \(2000\)](#); [Geer and Patureau-Mirand \(2008\)](#) for a related approach, as well as literature

on (HOMFLY-PT) skein modules ([Turaev \(1991\)](#), [Przytycki \(1999\)](#), [Morton \(2002\)](#) and related papers)).

Definition 2.1. *The quantized oriented Brauer category $\mathbf{Br}(\beta)$ is the quotient of $\mathbf{Tangles}_q$ modulo the following relations*

$$\begin{array}{c} \uparrow \quad \downarrow \\ \diagdown \quad \diagup \\ \uparrow \quad \downarrow \end{array} - \begin{array}{c} \uparrow \quad \downarrow \\ \diagup \quad \diagdown \\ \uparrow \quad \downarrow \end{array} = (q - q^{-1}) \left(\begin{array}{c} \uparrow \quad \downarrow \\ \diagdown \quad \diagup \\ \uparrow \quad \downarrow \end{array} \right) \left(\begin{array}{c} \uparrow \quad \downarrow \\ \diagup \quad \diagdown \\ \uparrow \quad \downarrow \end{array} \right), \quad \bigcirc = \bigcirc = [\beta], \quad (2.9a)$$

$$\begin{array}{c} \uparrow \quad \downarrow \\ \bigcirc \end{array} = \begin{array}{c} \uparrow \quad \downarrow \\ \bigcirc \end{array} = q^{-\beta} \begin{array}{c} \uparrow \quad \downarrow \\ \downarrow \end{array}, \quad \begin{array}{c} \uparrow \quad \downarrow \\ \bigcirc \end{array} = \begin{array}{c} \uparrow \quad \downarrow \\ \bigcirc \end{array} = q^{+\beta} \begin{array}{c} \uparrow \quad \downarrow \\ \uparrow \end{array}. \quad (2.9b)$$

Definition 2.2. *If $\boldsymbol{\eta} \in \{1, 1^*\}^N$ is a sequence of orientations, the quantized oriented Brauer algebra $\mathbf{Br}_{\boldsymbol{\eta}}(\beta)$ is the endomorphism algebra $\mathbf{End}_{\mathbf{Br}(\beta)}(\boldsymbol{\eta})$.*

For $r, s \geq 0$ let

$$\boldsymbol{\eta}_{r,s} = (\underbrace{1, \dots, 1}_r, \underbrace{1^*, \dots, 1^*}_s). \quad (2.10)$$

Then $\mathbf{Br}_{\boldsymbol{\eta}_{r,s}}(\beta)$ is the *walled Brauer algebra* with parameter β .

It is shown in ([Dipper et al., 2014](#), Lemma 1.4) and in [Weimer \(2005\)](#) that $\mathbf{Br}_{\boldsymbol{\eta}}(\beta)$ is free of rank $k!$, where k is the length of $\boldsymbol{\eta}$.

We recall that for $r \geq 0$ the *Hecke algebra* \mathcal{H}_r is the $\mathbb{C}(q)$ algebra on generators H_i for $i = 1, \dots, r-1$ with relations

$$H_i H_j = H_j H_i \quad \text{if } |i - j| > 2, \quad (2.11a)$$

$$H_i H_{i+1} H_i = H_{i+1} H_i H_{i+1}, \quad H_i^2 = (q^{-1} - q) H_i + 1, \quad (2.11b)$$

see [Kazhdan and Lusztig \(1979\)](#), [Soergel \(1997\)](#). It follows that if $\boldsymbol{\eta} = 1^N$ then $\mathbf{Br}_{\boldsymbol{\eta}}(\beta)$, which does not depend on β , is isomorphic to \mathcal{H}_N . We denote by \mathbf{H} the full subcategory of $\mathbf{Br}(\beta)$ monoidally generated by 1, which does not depend on β .

3. The general linear Lie superalgebra

We will denote by m and n two non-negative integers, and we will always suppose that either m or n is non-zero. We set also $d = m - n$. In the following, as usual, *super* means $\mathbb{Z}/2\mathbb{Z}$ -graded.

The Lie superalgebra $\mathfrak{gl}_{m|n}$

Let $I = I_{m|n}$ denote the set $\{1, \dots, m+n\}$. We define a parity function

$$|i| = \begin{cases} 0 & \text{if } i \leq m, \\ 1 & \text{if } i > m. \end{cases} \quad (3.1)$$

We let also $I^0 = I \setminus \{m+n\}$.

Let $\mathbb{C}^{m|n}$ be the vector superspace on homogeneous basis x_i for $i \in I_{m|n}$, with $\mathbb{Z}/2\mathbb{Z}$ -degree $|x_i| = |i|$. The *general linear Lie superalgebra* $\mathfrak{gl}_{m|n}$ is the endomorphism space of $\mathbb{C}^{m|n}$ equipped with the supercommutator $[y, z] = yz - (-1)^{|y||z|}zy$. Let $\mathfrak{h} \subset \mathfrak{gl}_{m|n}$ be the Cartan subalgebra consisting of all diagonal matrices, and let $\{h_i \mid i \in I\}$ be its standard basis. Let $\{\varepsilon_i \mid i \in I\}$ be the corresponding dual basis of \mathfrak{h}^* . We define a non-degenerate symmetric bilinear form on \mathfrak{h}^* by setting on the basis

$$(\varepsilon_i, \varepsilon_j) = (-1)^{|i|} \delta_{ij}, \quad (3.2)$$

where as usual δ_{ij} denotes the Kronecker delta. We denote by \mathbf{P} the *weight lattice* and by \mathbf{P}^* its dual: they are the free \mathbb{Z} -modules spanned by the ε_i 's and the h_i 's, respectively.

The roots of $\mathfrak{gl}_{m|n}$ are $\{\varepsilon_i - \varepsilon_j \mid i, j \in I\}$. A root is said to be *even* (respectively, *odd*) if the corresponding root vector is. We fix the standard triangular decomposition, with positive roots $\{\varepsilon_i - \varepsilon_j \mid i < j\}$ and simple roots $\{\alpha_i = \varepsilon_i - \varepsilon_{i+1} \mid i \in I^0\}$. We define the simple coroots

$$\alpha_i^\vee = (-1)^{|i|} h_i - (-1)^{|i+1|} h_{i+1} \quad \text{for all } i \in I^0. \quad (3.3)$$

We let ρ_0 denote half the sum of the even positive roots and ρ_1 denote half the sum of the odd positive roots, and we set $\rho = \rho_0 - \rho_1$. Explicitly we have

$$\begin{aligned} 2\rho = & (d-1)\varepsilon_1 + (d-3)\varepsilon_2 + \dots + (d-2m+1)\varepsilon_m \\ & + (d+2n-1)\varepsilon_{m+1} + (d+2n-3)\varepsilon_{m+2} + \dots + (d+1)\varepsilon_{m+n}. \end{aligned} \quad (3.4)$$

The quantum enveloping superalgebra

The *quantum enveloping superalgebra* $U_q = U_q(\mathfrak{gl}_{m|n})$ is defined (see for example [Scheunert \(1992\)](#); [Zhang \(1998\)](#)) to be the unital superalgebra over

$\mathbb{C}(q)$ with generators E_i, F_i, \mathbf{q}^h for $i \in I, h \in \mathbf{P}^*$ subject to the following relations:

$$\mathbf{q}^0 = 1, \quad \mathbf{q}^h \mathbf{q}^{h'} = \mathbf{q}^{h+h'}, \quad (3.5a)$$

$$\mathbf{q}^h E_i = q^{\langle h, \alpha_i \rangle} E_i \mathbf{q}^h, \quad \mathbf{q}^h F_i = q^{-\langle h, \alpha_i \rangle} F_i \mathbf{q}^h, \quad (3.5b)$$

$$(-1)^{|i|} E_i F_i - (-1)^{|i+1|} F_i E_i = \frac{K_i - K_i^{-1}}{q - q^{-1}}, \quad (3.5c)$$

$$E_m^2 = F_m^2 = 0, \quad (3.5d)$$

$$E_i E_j = E_j E_i \text{ and } F_i F_j = F_j F_i \quad \text{if } |i - j| \geq 2, \quad (3.5e)$$

$$E_i F_j = F_j E_i \quad \text{if } i \neq j, \quad (3.5f)$$

$$E_i^2 E_{i+1} - [2] E_i E_{i+1} E_i + E_{i+1} E_i^2 = 0 \quad \text{for } i, i+1 \neq m, \quad (3.5g)$$

$$E_{i+1}^2 E_i - [2] E_{i+1} E_i E_{i+1} + E_i E_{i+1}^2 = 0 \quad \text{for } i, i+1 \neq m, \quad (3.5h)$$

$$F_i^2 F_{i+1} - [2] F_i F_{i+1} F_i + F_{i+1} F_i^2 = 0 \quad \text{for } i, i+1 \neq m, \quad (3.5i)$$

$$F_{i+1}^2 F_i - [2] F_{i+1} F_i F_{i+1} + F_i F_{i+1}^2 = 0 \quad \text{for } i, i+1 \neq m, \quad (3.5j)$$

$$E_m E_{m-1} E_m E_{m+1} + E_{m-1} E_m E_{m+1} E_m + E_m E_{m+1} E_m E_{m-1} \\ + E_{m+1} E_m E_{m-1} E_m - [2] E_m E_{m-1} E_{m+1} E_m = 0, \quad (3.5k)$$

$$F_m F_{m-1} F_m F_{m+1} + F_{m-1} F_m F_{m+1} F_m + F_m F_{m+1} F_m F_{m-1} \\ + F_{m+1} F_m F_{m-1} F_m - [2] F_m F_{m-1} F_{m+1} F_m = 0, \quad (3.5l)$$

where $K_i = q^{\alpha_i^\vee}$.

We define a *comultiplication* $\Delta: U_q \rightarrow U_q \otimes U_q$, a *counit* $\mathbf{u}: U_q \rightarrow \mathbb{C}(q)$ and an *antipode* $S: U_q \rightarrow U_q$ by setting on the generators

$$\begin{aligned} \Delta(E_i) &= E_i \otimes K_i^{-1} + 1 \otimes E_i, & \Delta(F_i) &= F_i \otimes 1 + K_i \otimes F_i, \\ S(E_i) &= -E_i K_i, & S(F_i) &= -K_i^{-1} F_i, \\ \Delta(\mathbf{q}^h) &= \mathbf{q}^h \otimes \mathbf{q}^h, & S(\mathbf{q}^h) &= \mathbf{q}^{-h}, \\ \mathbf{u}(E_i) &= \mathbf{u}(F_i) = 0, & \mathbf{u}(\mathbf{q}^h) &= 1, \end{aligned} \quad (3.6)$$

and extending Δ and \mathbf{u} to superalgebra homomorphisms and S to a superalgebra anti-homomorphism. As well-known (and easy to check), the maps Δ , \mathbf{u} and S turn $U_q(\mathfrak{gl}_{m|n})$ into a Hopf superalgebra.

Representations

We let $\mathbb{C}_q = \mathbb{C}(q)$ be the trivial representation of $U_q(\mathfrak{gl}_{m|n})$. We denote by $V = \mathbb{C}_q^{m|n}$ its natural (vector) representation. It is a $\mathbb{C}(q)$ -vector superspace

of homogeneous basis x_i for $i \in I$, with grading given by $|x_i| = |i|$. (Note that by a slight abuse of notation we also denoted by x_i the basis vectors of $\mathbb{C}^{m|n}$.) The action of $U_q(\mathfrak{gl}_{m|n})$ is determined by

$$E_i x_j = \delta_{i+1,j} x_i, \quad F_i x_j = \delta_{i,j} x_{i+1}, \quad \mathbf{q}^h x_i = q^{\langle h, \varepsilon_i \rangle} x_i. \quad (3.7)$$

A weight $\lambda \in \mathbf{P}$ is said to be *dominant* if it is a dominant weight for $\mathfrak{gl}_m \oplus \mathfrak{gl}_n \subset \mathfrak{gl}_{m|n}$. Given a dominant weight $\lambda \in \mathbf{P}$, one can define the quantum Kac module $K_q(\lambda)$ with highest weight λ by parabolic induction. If λ is typical (that is, if $(\lambda + \rho, \alpha) \neq 0$ for all odd roots α), then $K_q(\lambda)$ is simple, and we denote it by $L_{q,m|n}(\lambda)$ or also simply by $L_q(\lambda)$ (see (Kwon, 2014, Section 2)). If λ is not typical, one can define $L_q(\lambda)$ as the unique simple quotient of $K_q(\lambda)$. Anyway, we will only need the typical case.


We recall that a partition of N is a sequence of positive integers $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_\ell)$ for some $\ell \geq 0$ with $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_\ell$ such that $|\lambda| = \lambda_1 + \dots + \lambda_\ell = N$. To a partition we associate a Young diagram, which we also denote by λ , as in the following picture:



$$(3.8)$$

The transposed λ^T of a partition λ is obtained by taking the symmetric Young diagram around the diagonal. In formulas, $\lambda_i^T = \#\{h \mid \lambda_h \geq i\}$.

A partition is said to be a (m, n) -hook if $\lambda_i \leq n$ for all $i > m$, or equivalently if its Young diagram fits into the union of two orthogonal strips of width m and n , as in the following picture:



$$(3.9)$$

Let $H_{m|n}$ denote the set of all (m, n) -hook partitions. Let $\lambda = (\lambda_1, \dots, \lambda_\ell) \in H_{m|n}$ and write λ as concatenation of two partitions $\lambda' = (\lambda_1, \dots, \lambda_m)$ and λ''^T . Notice that λ'' is a partition with at most n parts. Then the weight

$$\lambda = \lambda'_1 \varepsilon_1 + \dots + \lambda'_m \varepsilon_m + \lambda''_1 \varepsilon_{m+1} + \dots + \lambda''_n \varepsilon_{m+n} \quad (3.10)$$

is typical and we let $L_q(\lambda) = L_q(\lambda)$. As a convention, if λ is a partition which is not in $H_{m|n}$, then we just set $L_{q,m|n}(\lambda) = 0$.

Ribbon structure

As well-known, the \hbar -version $U_\hbar(\mathfrak{gl}_{m|n})$ of the quantum enveloping algebra of $\mathfrak{gl}_{m|n}$ can be endowed with a ribbon structure, and this allows to turn the category of finite-dimensional $U_q(\mathfrak{gl}_{m|n})$ -representations into a ribbon category. Actually, in the literature it is more common to find the ribbon structure of $U_\hbar(\mathfrak{sl}_{m|n})$, see for example [Geer \(2005\)](#), [Geer and Patureau-Mirand \(2010\)](#). For choosing a universal R -matrix for $\mathfrak{gl}_{m|n}$ one has an additional degree of freedom, due to the presence of a nontrivial center. This allows to obtain slightly nicer formulas. We explain our conventions in this subsection. As a reference for ribbon algebras and \hbar -versions of the quantum enveloping algebras, see for example [Chari and Pressley \(1994\)](#) or [Kassel \(1995\)](#).

Let R' be the universal R -matrix for $\mathfrak{sl}_{m|n}$ from [Geer and Patureau-Mirand \(2010\)](#). As R -matrix for $\mathfrak{gl}_{m|n}$ we take $R = q^{\frac{1}{2}I \otimes I} (R')^{-1}$, where $I = h_1 + \dots + h_{m+n} \in \mathbb{P}^*$ is the identity matrix. Using the ribbon structure of $U_\hbar(\mathfrak{sl}_{m|n})$ it is straightforward to check that R defines a braided structure on $U_\hbar(\mathfrak{gl}_{m|n})$. Let also v' be the ribbon element from [Geer and Patureau-Mirand \(2010\)](#). Then by setting $v = q^{\frac{1}{2}I^2} v'$ we get a ribbon element for our braided structure of $U_\hbar(\mathfrak{gl}_{m|n})$, and hence a ribbon structure of the latter.

Braiding and twist

On the simple finite-dimensional representation $L_q(\lambda)$ the twist operator acts as

$$\theta_{L_q(\lambda)}(w) = q^{-(\lambda, \lambda + 2\rho)} w, \quad (3.11)$$

see also ([Geer and Patureau-Mirand, 2010](#), Lemma 3.1).¹ As for the braiding, we will only need the explicit action on tensor powers of the vector

¹Notice that in [Geer and Patureau-Mirand \(2010\)](#) (\cdot, \cdot) denotes the bilinear form of the (dual of) the Cartan of $\mathfrak{sl}_{m|n}$, while in (3.11) we denoted by (\cdot, \cdot) the bilinear form of the (dual of) the Cartan of $\mathfrak{gl}_{m|n}$. The fact that we get apparently the same formula is due to our rescaling of the universal R -matrix, see above.

representation. According to our conventions, this is given by

$$\check{R}_{V,V}(x_a \otimes x_b) = \begin{cases} q^{-1}x_a \otimes x_a & \text{if } a = b \leq m, \\ (-1)^{|a||b|}x_b \otimes x_a & \text{if } a < b, \\ (-1)^{|a||b|}x_b \otimes x_a + (q^{-1} - q)x_a \otimes x_b & \text{if } a > b, \\ -qx_a \otimes x_a & \text{if } a = b > m, \end{cases} \quad (3.12)$$

see also [Zhang \(2009\)](#).

The map $\check{R}_{V,V}$ can be used in a straightforward way to define an action of the Hecke algebra \mathcal{H}_r (see section 2) on $V^{\otimes r}$. We recall the following result, also known as super Schur-Weyl duality:

Proposition 3.1 ([Mitsuhashi \(2006\)](#)). *The actions of \mathcal{H}_r and $U_q(\mathfrak{gl}_{m|n})$ on $V^{\otimes r}$ commute with each other and generate each other's centralizer. Moreover, we have the following decomposition as a module over $U_q(\mathfrak{gl}_{m|n}) \otimes \mathcal{H}_r$:*

$$V^{\otimes r} \cong \bigoplus_{\substack{\lambda \in H_{m|n} \\ |\lambda|=r}} L_q(\lambda) \otimes S(\lambda) \quad (3.13)$$

where $S(\lambda)$ is the simple \mathcal{H}_r -module corresponding to the partition λ .

We stress that it follows in particular that $V^{\otimes r}$ is semisimple.

Braided symmetric and exterior powers

Let W be any finite dimensional representation of $U_q(\mathfrak{gl}_{m|n})$, and let $TW = \bigoplus_{\ell \geq 0} \bigotimes^{\ell} W$ denote its tensor algebra. We recall from [Berenstein and Zwinknagl \(2008\)](#) that one can define the braided exterior power $\bigwedge_q^{\bullet} W$ of W as the quotient of TW modulo the ideal generated by the $\check{R}_{W,W}$ -eigenvectors of $\bigotimes^2 W$ with positive eigenvalues. Analogously, one can define the braided symmetric power $S_q^{\bullet} W$ of W as the quotient of TW modulo the ideal generated by the $\check{R}_{W,W}$ -eigenvectors of $\bigotimes^2 W$ with negative eigenvalues. Here, the positive and negative eigenvalues of $\check{R}_{W,W}$ are $\{q^i \mid i \in \mathbb{Z}\}$ and $\{-q^i \mid i \in \mathbb{Z}\}$, respectively.

The braided exterior and symmetric algebras are naturally \mathbb{Z} -graded (note that this is an additional grading to the $\mathbb{Z}/2\mathbb{Z}$ -grading, which in this case we

will call $\mathbb{Z}/2\mathbb{Z}$ -parity to avoid confusion). Moreover, their graded dimensions are smaller or equal to the graded dimensions of the usual exterior and symmetric algebras. If the equality holds, we say that W is *flat*.

REMARK 3.2. It is easy to see that if we forget the $\mathbb{Z}/2\mathbb{Z}$ -parity, then we have an isomorphism between $\bigwedge_q^\bullet(W\langle 1 \rangle)$ and $S_q^\bullet W$, where $\langle 1 \rangle$ denotes the shift by one in the $\mathbb{Z}/2\mathbb{Z}$ -parity. For $p \in \mathbb{Z}/2\mathbb{Z}$ and $j \in \mathbb{Z}_{\geq 0}$ let $(S_q^j W)_p$ denote the component of $S_q^\bullet W$ with $\mathbb{Z}/2\mathbb{Z}$ -parity p and \mathbb{Z} -degree j , so that $S_q^\bullet(W) = \bigoplus_{p \in \mathbb{Z}_2, j \in \mathbb{Z}_{\geq 0}} (S_q^j W)_p$. If we define $\mathbb{P}S_q^\bullet W$ to have graded components $(\mathbb{P}S_q^\bullet W)_{p,j} = (S_q^j W)_{p+j}$ then we have an honest graded isomorphism $\bigwedge_q^\bullet(W\langle 1 \rangle) \cong \mathbb{P}S_q^\bullet W$ which also preserves the $\mathbb{Z}/2\mathbb{Z}$ -parity.

In the next subsection, we will analyze further braided exterior powers of V .

Exterior powers

We define I to be the vector subspace of $\bigotimes^2 V$ spanned by

$$\begin{aligned} x_a \otimes x_a & \quad \text{for } a \leq m, \\ x_a \otimes x_b + (-1)^{|a||b|} q^{-1} x_b \otimes x_a & \quad \text{for } a < b. \end{aligned} \quad (3.14)$$

Notice that the braiding $\check{R}_{V,V}$ has minimal polynomial $(t - q^{-1})(t + q)$ and by construction I is the eigenspace with eigenvalue q^{-1} . In particular, I is a $U_q(\mathfrak{gl}_{m|n})$ -subrepresentation of $\bigotimes^2 V$.

Let $TV = \bigoplus_{\ell \geq 0} \bigotimes^\ell V$ be the tensor algebra of V . We define $\bigwedge_q^\bullet V = TV / \langle I \rangle$: it is the free algebra on generators x_a with multiplication \wedge_q modulo the relations

$$\begin{aligned} x_a \wedge_q x_a &= 0 & \text{for } a \leq m, \\ x_a \wedge_q x_b &= -(-1)^{|a||b|} q^{-1} x_b \wedge_q x_a & \text{for } a < b. \end{aligned} \quad (3.15)$$

Since TV is a $U_q(\mathfrak{gl}_{m|n})$ -representation and $\langle I \rangle$ is an invariant subspace, it follows that $\bigwedge_q^\bullet V$ is naturally a $U_q(\mathfrak{gl}_{m|n})$ -representation, and it decomposes as $\bigoplus_{i=0}^\infty \bigwedge_q^i V$. It follows from proposition 3.1, or it is easy to check explicitly, that each $\bigwedge_q^i V$ is irreducible, and

$$\bigwedge_q^i V \cong \begin{cases} L_q(\varepsilon_1 + \cdots + \varepsilon_i) & \text{if } i \leq m, \\ L_q(\varepsilon_1 + \cdots + \varepsilon_m + (i - m)\varepsilon_{m+1}) & \text{if } i > m \text{ and } n \geq 1, \\ 0 & \text{if } i > m \text{ and } n = 0. \end{cases} \quad (3.16)$$

We will often write \bigwedge_q^i instead of $\bigwedge_q^i V$ for shortness.

Definition 3.3. We define $\text{Rep}_{m|n}^+$ to be the full additive subcategory of representations of $U_q(\mathfrak{gl}_{m|n})$ monoidally generated by the $\bigwedge_q^i V$'s for $i \geq 0$, and we define $\text{Rep}_{m|n}$ to be the full additive subcategory of representations of $U_q(\mathfrak{gl}_{m|n})$ monoidally generated by the $\bigwedge_q^i V$'s for $i \geq 0$ and by their duals.

Reshetikhin-Turaev invariant

Since the category of $U_q(\mathfrak{gl}_{m|n})$ -representations is ribbon, given a tangle diagram, we can label its strands by finite dimensional $U_q(\mathfrak{gl}_{m|n})$ -representations and associate to it some equivariant map which is an invariant of the given tangle (see [Reshetikhin and Turaev \(1990\)](#); [Ohtsuki \(2002\)](#) for the details).

We will not actually need the full ribbon calculus of this category; it will be enough for our purposes to restrict to labelings of the strands by representations $\bigwedge_q^a V$ for $a \geq 1$. In particular, we will just label our strands by positive integers, and to a strand labeled by a we associate the representation $\bigwedge_q^a V$. In this way we get the Reshetikhin-Turaev functor $\mathcal{RT}_{m|n}: \text{Tangles}^{\text{lab}} \rightarrow \text{Rep}_{m|n}$.

We will sometimes use the same picture for an element of $\text{Tangles}_q^{\text{lab}}$ and for its image under the functor $\mathcal{RT}_{m|n}$. This should cause no confusion. For example, the braiding of $V \otimes V$ and its inverse will be represented by

$$\begin{array}{c} V \otimes V \\ \uparrow \quad \uparrow \\ 1 \quad 1 \\ \downarrow \quad \downarrow \\ V \otimes V \end{array} \quad \text{and} \quad \begin{array}{c} V \otimes V \\ \uparrow \quad \uparrow \\ 1 \quad 1 \\ \downarrow \quad \downarrow \\ V \otimes V \end{array} \quad (3.17)$$

respectively. We also have evaluation and coevaluation maps, which for V and V^* are graphically represented by

$$\begin{array}{c} \mathbb{C}_q \\ \uparrow \\ V \otimes V^* \end{array} \quad \begin{array}{c} \mathbb{C}_q \\ \uparrow \\ V^* \otimes V \end{array} \quad \begin{array}{c} V^* \otimes V \\ \uparrow \\ \mathbb{C}_q \end{array} \quad \begin{array}{c} V \otimes V^* \\ \uparrow \\ \mathbb{C}_q \end{array} \quad (3.18)$$

We recall the following relations in $\text{Rep}_{m|n}$, which are a consequence of (3.11)

(cf. also (Geer and Patureau-Mirand, 2010, Lemma 3.1)):

$$\begin{array}{c} 1 \\ \updownarrow \end{array} \bigcirc = q^{-d} \begin{array}{c} 1 \\ \updownarrow \end{array} \quad \text{and} \quad \begin{array}{c} 1 \\ \updownarrow \end{array} \bigcirc = q^{+d} \begin{array}{c} 1 \\ \updownarrow \end{array}, \quad (3.19)$$

and more generally

$$\begin{array}{c} a \\ \updownarrow \end{array} \bigcirc = q^{-ad+a(a-1)} \begin{array}{c} a \\ \updownarrow \end{array} \quad \text{and} \quad \begin{array}{c} a \\ \updownarrow \end{array} \bigcirc = q^{ad-a(a-1)} \begin{array}{c} a \\ \updownarrow \end{array}. \quad (3.20)$$

Moreover, we have

$$1 \bigcirc = [d], \quad (3.21)$$

where $d = m - n$ is the superdimension of $\mathbb{C}^{m|n}$. We stress that these are not equalities in Tangles_q , but only in $\text{Rep}_{m|n}$.

4. Super skew Howe duality

We consider now simultaneously two quantum superalgebras $U_q(\mathfrak{gl}_{m|n})$ and $U_q(\mathfrak{gl}_{k|l})$ for some $m, n, k, l \geq 0$. We let $V_1 = \mathbb{C}_q^{m|n}$ and $V_2 = \mathbb{C}_q^{k|l}$ be their vector representations, with bases $\{x_a \mid 1 \leq a \leq m + n\}$ and $\{y_b \mid 1 \leq b \leq k + l\}$ respectively.

We will consider the representation $V_1 \otimes V_2$ of $U_q(\mathfrak{gl}_{m|n}) \otimes U_q(\mathfrak{gl}_{k|l})$, whose standard basis we denote by $z_{ab} = x_a \otimes y_b$. Our first goal is to define exterior powers of this representation, generalizing Berenstein and Zwicknagl (2008). Notice that the tensor product algebra $U_q(\mathfrak{gl}_{m|n}) \otimes U_q(\mathfrak{gl}_{k|l})$ is the quantum enveloping superalgebra of $\mathfrak{gl}_{m|n} \oplus \mathfrak{gl}_{k|l}$. It follows by general theory that its representation category is braided, and its universal R -matrix is the tensor product of the two R -matrices of $U_q(\mathfrak{gl}_{m|n})$ and $U_q(\mathfrak{gl}_{k|l})$.

Let $\sigma_{23}: V_1 \otimes V_1 \otimes V_2 \otimes V_2 \rightarrow V_1 \otimes V_2 \otimes V_1 \otimes V_2$ be the superpermutation which swaps the two middle tensor factors. The operator $\mathcal{R} = \sigma_{23} \circ (\check{R}_{V_1, V_1} \otimes \check{R}_{V_2, V_2}) \circ \sigma_{23}^{-1}$ then is the braiding on $(V_1 \otimes V_2) \otimes (V_1 \otimes V_2)$. (Although this follows by general statements, it is straightforward to see directly that it satisfies the braid relation.)

Since the minimal polynomial of both \check{R}_{V_1, V_1} and of \check{R}_{V_2, V_2} is $(t - q^{-1})(t + q)$, the minimal polynomial of \mathcal{R} is $(t - q^{-2})(t - q^2)(t + 1)$. We set I to be the sum of the eigenspaces of \mathcal{R} with eigenvalues q^{-2} and q^2 (i.e. with *positive* eigenvalues). It is easy to see that I is spanned by

$$\begin{aligned}
& z_{ab} \otimes z_{ab} && \text{for } |a| = |b|, \\
& z_{ac} \otimes z_{bc} + (-1)^{|a||b|} q^{-1} z_{bc} \otimes z_{ac} && \text{for } a < b, |c| = 0, \\
& z_{ac} \otimes z_{ad} + (-1)^{|c||d|} q^{-1} z_{ad} \otimes z_{ac} && \text{for } |a| = 0, c < d, \\
& z_{ac} \otimes z_{bd} + (-1)^{|a||b|} q^{-1} z_{bc} \otimes z_{ad} \\
& \quad + (-1)^{|c||d|} q^{-1} z_{ad} \otimes z_{bc} + (-1)^{|a||b|+|c||d|} q^{-2} z_{bd} \otimes z_{ac} && \text{for } a < b, c < d, \\
& z_{ac} \otimes z_{ad} - (-1)^{|c||d|} q z_{ad} \otimes z_{ac} && \text{for } |a| = 1, c < d, \\
& z_{ac} \otimes z_{bc} - (-1)^{|a||b|} q z_{bc} \otimes z_{ac} && \text{for } a < b, |c| = 1, \\
& z_{ac} \otimes z_{bd} - (-1)^{|a||b|} q z_{bc} \otimes z_{ad} \\
& \quad - (-1)^{|c||d|} q z_{ad} \otimes z_{bc} + (-1)^{|a||b|+|c||d|} q^2 z_{bd} \otimes z_{ac} && \text{for } a < b, c < d.
\end{aligned} \tag{4.1}$$

We let $\bigwedge_q^\bullet(V_1 \otimes V_2) = T(V_1 \otimes V_2)/\langle I \rangle$: this is the quotient of the free algebra on the set $\{z_{ab}\}$ with multiplication \wedge_q modulo the relations

$$\begin{aligned}
& z_{ab} \wedge_q z_{ab} = 0 && \text{for } |a| = |b|, \\
& z_{ac} \wedge_q z_{bc} = -(-1)^{|a||b|} q^{-1} z_{bc} \wedge_q z_{ac} && \text{for } a < b, |c| = 0, \\
& z_{ac} \wedge_q z_{ad} = -(-1)^{|c||d|} q^{-1} z_{ad} \wedge_q z_{ac} && \text{for } |a| = 0, c < d, \\
& z_{ac} \wedge_q z_{bd} = -(-1)^{|a||b|+|c||d|} z_{bd} \wedge_q z_{ac} && \text{for } a < b, c < d, \\
& z_{ac} \wedge_q z_{bd} = -(-1)^{|a||b|+|c||d|} z_{bd} \wedge_q z_{ac} \\
& \quad + (-1)^{|a||b|} (q^{-1} - q) z_{bc} \wedge_q z_{ad} && \text{for } a > b, c < d, \\
& z_{ac} \wedge_q z_{ad} = (-1)^{|c||d|} q z_{ad} \wedge_q z_{ac} && \text{for } |a| = 1, c < d, \\
& z_{ac} \wedge_q z_{bc} = (-1)^{|a||b|} q z_{bc} \wedge_q z_{ac} && \text{for } a < b, |c| = 1.
\end{aligned} \tag{4.2}$$

The tensor algebra $T(V_1 \otimes V_2)$ is a $(U_q(\mathfrak{gl}_{m|n}) \otimes U_q(\mathfrak{gl}_{k|l}))$ -module, and the ideal $\langle I \rangle$ is by construction an invariant subspace. In particular, $\bigwedge_q^\bullet(V_1 \otimes V_2)$ is also a $(U_q(\mathfrak{gl}_{m|n}) \otimes U_q(\mathfrak{gl}_{k|l}))$ -module, and it decomposes as $\bigoplus_{N=0}^{\infty} \bigwedge_q^N(V_1 \otimes V_2)$.

We want to show that $V_1 \otimes V_2$ is flat, that is, the dimension of $\bigwedge_q^N(V_1 \otimes V_2)$ coincides with the dimension of the usual exterior power $\bigwedge^N(V_1 \otimes V_2)$.

Unfortunately, it is not easy to see this from the presentation (4.2). Hence we will give now an alternative construction of $\bigwedge_q^\bullet(V_1 \otimes V_2)$. This will come only with an action of $U_q(\mathfrak{gl}_{m|n})$, and not of $U_q(\mathfrak{gl}_{k|l})$. However, the flatness will be obvious.

Let A, B be any locally finite $U_q(\mathfrak{gl}_{m|n})$ -module algebras with multiplications μ_A and μ_B , respectively. We recall from (Lehrer et al., 2011, Section 2) that one can define a $U_q(\mathfrak{gl}_{m|n})$ -module algebra structure on $A \otimes B$ with multiplication

$$\mu_{A \otimes B} = (\mu_A \otimes \mu_B)(\text{id}_A \otimes \check{R}_{A,B}^{-1} \otimes \text{id}_B), \quad (4.3)$$

This is well-defined ((Lehrer et al., 2011, Theorem 2.3)), the construction can be iterated with a third module algebra C and is associative ((Lehrer et al., 2011, Lemma 2.5)).

We apply this construction iteratively to the algebras $S_q^\bullet(V_1 \langle 1 \rangle)$ and $S_q^\bullet(V_1)$, where $\langle 1 \rangle$ denotes a shift by one in the $\mathbb{Z}/2\mathbb{Z}$ -parity, and we define in this way a $U_q(\mathfrak{gl}_{m|n})$ -module algebra structure on

$$A_{k|l} = (S_q^\bullet(V_1 \langle 1 \rangle))^{\otimes k} \otimes (S_q^\bullet(V_1))^{\otimes l}, \quad (4.4)$$

where $\langle 1 \rangle$ denotes a shift by one in the $\mathbb{Z}/2\mathbb{Z}$ -degree. As in (Lehrer et al., 2011, Theorem 2.8), it follows that $A_{k|l}$ is a flat deformation of the symmetric algebra $S((V_1 \langle 1 \rangle)^{\oplus k} \oplus V_1^{\oplus l})$, i.e. of the exterior algebra $\bigwedge(V_1^{\oplus k} \oplus (V_1 \langle 1 \rangle)^{\oplus l})$. This means that the graded dimensions of $A_{k|l}$ and of $\bigwedge(V_1^{\oplus k} \oplus (V_1 \langle 1 \rangle)^{\oplus l})$ coincide.

For all $a = 1, \dots, m+n$ and $i = 1, \dots, k+l$ let $X_{ai} = 1 \otimes \dots \otimes 1 \otimes v_a \otimes 1 \otimes \dots \otimes 1 \in A_{k|l}$, the entry v_a being at position i . It is easy to see that the X_{ai} 's generate $A_{k|l}$ as an algebra (this follows for example from (Lehrer et al., 2011, Lemma 2.9)), and by construction they are subject to the relations

$$X_{aj}X_{bi} = -(-1)^{|i||j|} \sum_{a',b'=1}^{m+n} \check{R}_{a'a,b'b}^{-1} X_{a'i}X_{b'j} \quad \text{for all } i < j. \quad (4.5)$$

Here $\check{R}_{a'a,b'b}^{-1}$ denotes the matrix coefficient of the inverse of the braiding \check{R} , and the sign $-(-1)^{|i||j|}$ appears since we shifted the first k copies of $S_q^\bullet V_1$ by one in the $\mathbb{Z}/2\mathbb{Z}$ -degree.

Lemma 4.1. *The map $z_{ai} \mapsto X_{ai}$ defines an algebra isomorphism $\bigwedge_q^\bullet(V_1 \otimes V_2) \rightarrow A_{k|l}$. For all $N \geq 0$, a basis of $\bigwedge_q^N(V_1 \otimes V_2)$ is given by the elements*

$$\left\{ z_{a_1 b_1} \wedge_q \cdots \wedge_q z_{a_N b_N} \left| \begin{array}{l} (a_1, b_1) \leq (a_2, b_2) \leq \cdots \leq (a_N, b_N) \\ \text{in the anti-lexicographic order, and} \\ \text{if } (a_i, b_i) = (a_{i+1}, b_{i+1}) \text{ then } |z_{a_i b_i}| = 1 \end{array} \right. \right\}. \quad (4.6)$$

We point out that this isomorphism does not preserve the $\mathbb{Z}/2\mathbb{Z}$ -parity. However, one gets a parity-preserving isomorphism $\bigwedge_q^\bullet(V_1 \otimes V_2) \rightarrow \mathbb{P}A_{k|l}$, see remark 3.2.

Proof. First, let us notice that the relations (4.2) hold for the X_{ai} 's, and hence the map $z_{ai} \mapsto X_{ai}$ is well-defined. Indeed, the first two relations hold simply because they hold in $\bigwedge_q^\bullet(V_1)$, while the other relations are directly obtained from (4.5) (being very careful with signs).

It is a straightforward computation using the relations (4.2) to show that (4.6) span $\bigwedge_q^N(V_1 \otimes V_2)$. Since the X_{ai} 's generate $A_{k|l}$ as an algebra, the map $z_{ai} \mapsto X_{ai}$ is surjective and the images of the elements (4.6) span $A_{k|l}$. On the other side, one checks directly that the cardinality of the set (4.6) coincides with the dimension of $\bigwedge^N(V_1^{\oplus k} \oplus (V_1 \langle 1 \rangle)^{\oplus l})$, which by construction is the same as the dimension of the graded part of $A_{k|l}$ of degree N . It follows that the images of the elements (4.6) must be linearly independent and the map $z_{ai} \mapsto X_{ai}$ is an isomorphism of algebras. \square

The following result is the quantum version of super skew Howe duality. The corresponding non-quantized result can be found in (Cheng and Wang, 2012, Chapter 5). In the non-super case, the result has already been proven in (Cautis et al., 2014, Theorem 4.2.2). A generalization to the super case, but only from one side, already appeared in (Wu and Zhang, 2009, Theorem 2.2).

Theorem 4.2 (Quantized super skew Howe duality). *For all $N > 0$ the two actions*

$$U_q(\mathfrak{gl}_{m|n}) \curvearrowright \bigwedge_q^N(\mathbb{C}_q^{m|n} \otimes \mathbb{C}_q^{k|l}) \curvearrowright U_q(\mathfrak{gl}_{k|l}) \quad (4.7)$$

commute and generate each other's centralizer. Moreover, we have

$$\bigwedge_q^N(\mathbb{C}_q^{m|n} \otimes \mathbb{C}_q^{k|l}) \cong \bigoplus_{\lambda \in H(m|n) \cap H(l|k)} L_{q,m|n}(\lambda) \otimes L_{q,k|l}(\lambda^T) \quad (4.8)$$

as $U_q(\mathfrak{gl}_{m|n}) \otimes U_q(\mathfrak{gl}_{k|l})$ -modules.

Proof. It is clear that the two actions in (4.7) commute with each other. It is then enough to prove the decomposition (4.8), which implies the first assertion. We remark that in the non-quantized case, the analogue of (4.8) holds (see (Cheng and Wang, 2012, Theorem 5.18 and Remark 5.20)).

The space $W = \bigwedge_q^N (\mathbb{C}_q^{m|n} \otimes \mathbb{C}_q^{k|l})$, being a quotient of $\bigotimes^N (\mathbb{C}_q^{m|n} \otimes \mathbb{C}_q^{k|l})$ which is a semisimple $(U_q(\mathfrak{gl}_{m|n}) \otimes U_q(\mathfrak{gl}_{k|l}))$ -module by proposition 3.1, is semisimple. Moreover it must decompose as $\bigoplus_{\lambda \in H_{m|n}, \mu \in H_{k|l}} (L_{q,m|n}(\lambda) \otimes L_{q,k|l}(\mu))^{\oplus \kappa_{\lambda,\mu}}$. As in finite-dimensional representation theory of semisimple Lie algebras, it is very easy to argue that the multiplicities $\kappa_{\lambda,\mu}$ are uniquely determined by the dimensions of the weight spaces of W , that is by the character of W . The character of W coincides with the character of $\bigwedge^N (\mathbb{C}^{m|n} \otimes \mathbb{C}^{k|l})$, hence the multiplicities $\kappa_{\lambda,\mu}$ are the same as in the non-quantized case, and (4.8) follows. \square

Proposition 4.3. *As a $U_q(\mathfrak{gl}_{m|n})$ -representation, the module (4.7) decomposes as*

$$\bigwedge_q^N (\mathbb{C}_q^{m|n} \otimes \mathbb{C}_q^{k|l}) \cong \bigoplus_{\substack{\mathbf{a} \in \text{Comp}(N_1, k), \\ \mathbf{b} \in \text{Comp}(N_2, l), \\ N_1 + N_2 = N}} \bigwedge_q^{\mathbf{a}} \mathbb{C}_q^{m|n} \otimes \bigwedge_q^{\mathbf{b}} (\mathbb{C}_q^{m|n} \langle 1 \rangle), \quad (4.9)$$

where $\text{Comp}(M, h) = \{\mathbf{a} = (a_1, \dots, a_h) \mid a_i \in \mathbb{Z}_{\geq 0}, a_1 + \dots + a_h = M\}$ denotes the set of all compositions of M with h parts, and for $\mathbf{a} = (a_1, \dots, a_k)$ set

$$\bigwedge_q^{\mathbf{a}} \mathbb{C}_q^{m|n} = \bigwedge_q^{a_1} \mathbb{C}_q^{m|n} \otimes \dots \otimes \bigwedge_q^{a_k} \mathbb{C}_q^{m|n}. \quad (4.10)$$

When considered as a $U_q(\mathfrak{gl}_{k|l})$ -module, (4.9) is the weight space decomposition, and $\bigwedge_q^{\mathbf{a}} \mathbb{C}_q^{m|n} \otimes \bigwedge_q^{\mathbf{b}} (\mathbb{C}_q^{m|n} \langle 1 \rangle)$ is the weight space of weight $(\mathbf{a}, \mathbf{b}) = a_1 \varepsilon_1 + \dots + a_k \varepsilon_k + b_1 \varepsilon_{k+1} + \dots + b_l \varepsilon_{k+l}$.

We recall that $\bigwedge_q^{\mathbf{b}} (\mathbb{C}_q^{m|n} \langle 1 \rangle)$ is just the symmetric braided algebra, but with shifted $\mathbb{Z}/2\mathbb{Z}$ -parity, see Remark 3.2.

Proof. The isomorphism (4.9) is just the isomorphism of $\bigwedge_q^N (\mathbb{C}_q^{m|n} \otimes \mathbb{C}_q^{k|l})$ with the graded part of $A_{k|l}$ of degree N . It is straightforward to check that

this isomorphism is $U_q(\mathfrak{gl}_{m|n})$ -equivariant, and that the last assertion on the weight spaces holds. \square

From now on we set $l = 0$ and consider the case of $V_2 = \mathbb{C}_q^k$. We will mainly be interested in the action of $U_q(\mathfrak{gl}_k)$ on the $U_q(\mathfrak{gl}_{m|n})$ -representation $\bigoplus_{\mathbf{a}} \bigwedge_q^{\mathbf{a}} \mathbb{C}_q^{m|n}$. Hence from now on the symbols E_i, F_i will denote the generators of $U_q(\mathfrak{gl}_k)$, and will act on $\bigoplus_{\mathbf{a}} \bigwedge_q^{\mathbf{a}} \mathbb{C}_q^{m|n}$.

We identify the weight lattice \mathbf{P} of \mathfrak{gl}_k with \mathbb{Z}^k . We can then reinterpret skew Howe duality by saying that for all $\mathbf{a} \in \mathbb{N}^k$ we have morphisms

$$E_i: \bigwedge_q^{\mathbf{a}} \mathbb{C}_q^{m|n} \longrightarrow \bigwedge_q^{\mathbf{a}+\alpha_i} \mathbb{C}_q^{m|n}, \quad (4.11)$$

$$F_i: \bigwedge_q^{\mathbf{a}} \mathbb{C}_q^{m|n} \longrightarrow \bigwedge_q^{\mathbf{a}-\alpha_i} \mathbb{C}_q^{m|n}, \quad (4.12)$$

$$\mathbf{q}^h: \bigwedge_q^{\mathbf{a}} \mathbb{C}_q^{m|n} \longrightarrow \bigwedge_q^{\mathbf{a}} \mathbb{C}_q^{m|n}, \quad (4.13)$$

satisfying the defining relations of $U_q(\mathfrak{gl}_k)$, where we set $\bigwedge_q^{\mathbf{a}} \mathbb{C}_q^{m|n} = 0$ in case $a_i < 0$ for some $i = 1, \dots, k$. By the last claim of proposition 4.3, the element \mathbf{q}^h acts on $\bigwedge_q^{\mathbf{a}} \mathbb{C}_q^{m|n}$ by $q^{\langle h, \mathbf{a} \rangle}$.

We notice that the action given by skew Howe duality is local, in the following sense:

Proposition 4.4. *Fix $k \geq 2$, let $\mathbf{a} = (a_1, \dots, a_k)$ and consider $\bigwedge_q^{\mathbf{a}} \mathbb{C}_q^{m|n}$. Then we have*

$$E_i = \text{id}^{\otimes(i-1)} \otimes E \otimes \text{id}^{\otimes(k-i-1)} \quad (4.14)$$

as morphisms $\bigwedge_q^{\mathbf{a}} \mathbb{C}_q^{m|n} \mapsto \bigwedge_q^{\mathbf{a}+\alpha_i} \mathbb{C}_q^{m|n}$, where

$$E: \bigwedge_q^{a_i} \mathbb{C}_q^{m|n} \otimes \bigwedge_q^{a_{i+1}} \mathbb{C}_q^{m|n} \rightarrow \bigwedge_q^{a_i+1} \mathbb{C}_q^{m|n} \otimes \bigwedge_q^{a_{i+1}-1} \mathbb{C}_q^{m|n} \quad (4.15)$$

is the generator E_1 of $U_q(\mathfrak{gl}_2)$. The analogous statement holds for the F_i 's.

Proof. This is straightforward. \square

Moreover, we state the following easy result:

Lemma 4.5. *Let $k = 2$, and consider skew Howe duality applied to $\Lambda_q^N(\mathbb{C}_q^{m|n} \otimes \mathbb{C}_q^2)$. Then*

$$E_1^{(N)}: \Lambda_q^0(\mathbb{C}_q^{m|n}) \otimes \Lambda_q^N(\mathbb{C}_q^{m|n}) \rightarrow \Lambda_q^N(\mathbb{C}_q^{m|n}) \otimes \Lambda_q^0(\mathbb{C}_q^{m|n}) \quad \text{and} \quad (4.16)$$

$$F_1^{(N)}: \Lambda_q^N(\mathbb{C}_q^{m|n}) \otimes \Lambda_q^0(\mathbb{C}_q^{m|n}) \rightarrow \Lambda_q^0(\mathbb{C}_q^{m|n}) \otimes \Lambda_q^N(\mathbb{C}_q^{m|n}) \quad (4.17)$$

are the identity map after the natural identification of both spaces with $\Lambda_q^N(\mathbb{C}_q^{m|n})$.

5. The category \mathbf{Sp}^+

Following [Cautis et al. \(2014\)](#), we introduce the category \mathbf{Sp}^+ which gives a graphical interpretation of skew Howe duality.

The category $\widetilde{\mathbf{Sp}}^+$

First, we are going to define a monoidal category $\widetilde{\mathbf{Sp}}^+$. Objects of $\widetilde{\mathbf{Sp}}^+$ are sequences of non-negative integers. The tensor product is given by concatenation of sequences. Morphisms are monoidally generated by identities and by the two elements

$$\mathbf{E}^{(r)} = \mathbf{E}^{(r)} \mathbf{1}_{a \otimes b}: a \otimes b \longrightarrow a + r \otimes b - r \quad (5.1)$$

$$\mathbf{F}^{(r)} = \mathbf{F}^{(r)} \mathbf{1}_{a \otimes b}: a \otimes b \longrightarrow a - r \otimes b + r \quad (5.2)$$

for all $a, b \in \mathbb{Z}_{\geq 0}$ and $1 \leq r \leq b$ such that $a + r, b - r \in \mathbb{Z}_{\geq 0}$ or $a - r, b + r \in \mathbb{Z}_{\geq 0}$, respectively. By abuse of notation, we omit sometimes to write $\mathbf{1}_{a \otimes b}$ on the right of $\mathbf{E}^{(r)}$ and $\mathbf{F}^{(r)}$.

Let us denote by \emptyset the empty sequence, which is the unit of the monoidal structure. We also want to have an isomorphism $0 \cong \emptyset$ which allows us to identify 0 with the monoidal unit. We will implicitly use this isomorphism to let 0's appear and disappear. We impose moreover that $\mathbf{E}^{(r)} \mathbf{1}_{0 \otimes r}: r \cong 0 \otimes r \rightarrow r \otimes 0 \cong r$ and $\mathbf{F}^{(r)} \mathbf{1}_{r \otimes 0}: r \cong r \otimes 0 \rightarrow 0 \otimes r \cong r$ are the identity morphisms.

Graphically, we represent the identity morphism $a \rightarrow a$ and the morphisms (5.1) and (5.2) by

$$\text{id}_a = \begin{array}{c} a \\ | \\ a \end{array}, \quad E^{(r)} \mathbf{1}_{a \otimes b} = \begin{array}{c} a+r \quad b-r \\ | \quad | \\ \swarrow \quad \searrow \\ | \quad | \\ a \quad b \end{array} \quad \text{and} \quad F^{(r)} \mathbf{1}_{a \otimes b} = \begin{array}{c} a-r \quad b+r \\ | \quad | \\ \swarrow \quad \searrow \\ | \quad | \\ a \quad b \end{array} \quad (5.3)$$

respectively. Notice that we read the pictures always from the bottom to the top. The tensor product of two morphisms is given by horizontal juxtaposition, while composition of morphisms is given by stacking them one on top of the other.

We adopt the following graphical convention: we use thick strands (as in (5.3)) when they are labeled by an integer $a > 0$, we use thin strands when they are labeled by 1, and we use dashed strands when they are labeled by 0. In particular, $E \mathbf{1}_{a \otimes b} = E^{(1)} \mathbf{1}_{a \otimes b}$ and $F \mathbf{1}_{a \otimes b} = F^{(1)} \mathbf{1}_{a \otimes b}$ are represented by the diagrams

$$\begin{array}{c} a+1 \quad b-1 \\ | \quad | \\ \swarrow \quad \searrow \\ | \quad | \\ a \quad b \end{array} \quad \text{and} \quad \begin{array}{c} a-1 \quad b+1 \\ | \quad | \\ \swarrow \quad \searrow \\ | \quad | \\ a \quad b \end{array}. \quad (5.4)$$

We enlarge the category $\widetilde{\mathbf{Sp}}^+$ by taking as new homomorphism spaces the \mathbb{k} -vector spaces spanned by the old ones, and by taking then the additive envelope of the resulting category. In this way $\widetilde{\mathbf{Sp}}^+$ becomes an additive \mathbb{k} -linear category. For brevity in the following, when we will write $\mathbf{a} \in \widetilde{\mathbf{Sp}}^+$, we will mean a sequence of non-negative integers (although a generic object of $\widetilde{\mathbf{Sp}}^+$ is a formal direct sum of such sequences). We will use a similar convention for the categories \mathbf{Sp}^+ , \mathbf{Sp} and $\mathbf{Sp}(\beta)$ which will be introduced later on.

The category \mathbf{Sp}^+

We define a category \mathbf{Sp}^+ by imposing the graphical versions of the relations of the generators of $U_q(\mathfrak{gl}_k)$:

Definition 5.1 (Cautis et al. (2014)). We define the category \mathbf{Sp}^+ to be the quotient of the category $\widetilde{\mathbf{Sp}}^+$ modulo the following relations:

- Far away elements commute:

$$\begin{array}{c}
 a-r \quad b+r \quad c-s \quad d+s \quad a-r \quad b+r \quad c-s \quad d+s \\
 \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} \cdots \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} = \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} \cdots \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} \\
 \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} \quad \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} \\
 a \quad b \quad c \quad d \quad a \quad b \quad c \quad d
 \end{array} \quad (5.5a)$$

$$\begin{array}{c}
 a+r \quad b-r \quad c+s \quad d-s \quad a+r \quad b-r \quad c+s \quad d-s \\
 \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} \cdots \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} = \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} \cdots \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} \\
 \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} \quad \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} \\
 a \quad b \quad c \quad d \quad a \quad b \quad c \quad d
 \end{array} \quad (5.5b)$$

- The morphisms $E^{(r)}$ and $F^{(r)}$ commute also when they are near to each other:

$$\begin{array}{c}
 a-r \quad b+r+s \quad c-s \quad a-r \quad b+r+s \quad c-s \\
 \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} = \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} \\
 \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} \quad \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} \\
 a \quad b \quad c \quad a \quad b \quad c
 \end{array} \quad \text{and} \quad \begin{array}{c}
 a+r \quad b-r-s \quad c+s \quad a+r \quad b-r-s \quad c+s \\
 \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} = \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} \\
 \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} \quad \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} \\
 a \quad b \quad c \quad a \quad b \quad c
 \end{array} \quad (5.5c)$$

- Divided powers multiply as usual:

$$\begin{array}{c}
 a-r-s \quad b+r+s \quad a-r-s \quad b+r+s \\
 \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} = \begin{bmatrix} r+s \\ r \end{bmatrix} \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} \\
 \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} \quad \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} \\
 a \quad b \quad a \quad b
 \end{array} \quad \text{and} \quad \begin{array}{c}
 a+r+s \quad b-r-s \quad a+r+s \quad b-r-s \\
 \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} = \begin{bmatrix} r+s \\ r \end{bmatrix} \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} \\
 \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} \quad \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} \begin{array}{|c|} \hline \uparrow \\ \hline \end{array} \\
 a \quad b \quad a \quad b
 \end{array} \quad (5.5d)$$

- The commuting relation between E and F is

$$\begin{array}{c} a & b \\ | & | \\ \swarrow 1 & \searrow \\ | & | \\ \swarrow 1 & \searrow \\ | & | \\ a & b \end{array} - \begin{array}{c} a & b \\ | & | \\ \swarrow 1 & \searrow \\ | & | \\ \swarrow 1 & \searrow \\ | & | \\ a & b \end{array} = [a - b] \begin{array}{c} a & b \\ | & | \\ | & | \\ | & | \\ | & | \\ a & b \end{array}. \quad (5.5e)$$

- And finally, we have the Serre relations:

$$\begin{array}{c} a+2 & b-1 & c-1 \\ | & | & | \\ \swarrow 2 & \searrow & \\ | & | & | \\ \swarrow 1 & \searrow & \\ | & | & | \\ a & b & c \end{array} - \begin{array}{c} a+2 & b-1 & c-1 \\ | & | & | \\ \swarrow 1 & \searrow & 1 \\ | & | & | \\ \swarrow 1 & \searrow & \\ | & | & | \\ a & b & c \end{array} + \begin{array}{c} a+2 & b-1 & c-1 \\ | & | & | \\ | & | & | \\ \swarrow 2 & \searrow & \\ | & | & | \\ | & | & | \\ a & b & c \end{array} = 0, \quad (5.5f)$$

$$\begin{array}{c} a+1 & b+1 & c-2 \\ | & | & | \\ | & | & | \\ \swarrow 1 & \searrow & \\ | & | & | \\ | & | & | \\ a & b & c \end{array} - \begin{array}{c} a+1 & b+1 & c-2 \\ | & | & | \\ \swarrow 1 & \searrow & 1 \\ | & | & | \\ \swarrow 1 & \searrow & \\ | & | & | \\ a & b & c \end{array} + \begin{array}{c} a+1 & b+1 & c-2 \\ | & | & | \\ | & | & | \\ \swarrow 1 & \searrow & \\ | & | & | \\ | & | & | \\ a & b & c \end{array} = 0, \quad (5.5g)$$

$$\begin{array}{c} a-2 & b+1 & c+1 \\ | & | & | \\ \swarrow 2 & \searrow & \\ | & | & | \\ \swarrow 1 & \searrow & \\ | & | & | \\ a & b & c \end{array} - \begin{array}{c} a-2 & b+1 & c+1 \\ | & | & | \\ \swarrow 1 & \searrow & 1 \\ | & | & | \\ \swarrow 1 & \searrow & \\ | & | & | \\ a & b & c \end{array} + \begin{array}{c} a-2 & b+1 & c+1 \\ | & | & | \\ | & | & | \\ \swarrow 2 & \searrow & \\ | & | & | \\ | & | & | \\ a & b & c \end{array} = 0, \quad (5.5h)$$

$$\begin{array}{c} a-1 & b-1 & c+2 \\ | & | & | \\ | & | & | \\ \swarrow 1 & \searrow & \\ | & | & | \\ | & | & | \\ a & b & c \end{array} - \begin{array}{c} a-1 & b-1 & c+2 \\ | & | & | \\ \swarrow 1 & \searrow & 1 \\ | & | & | \\ \swarrow 1 & \searrow & \\ | & | & | \\ a & b & c \end{array} + \begin{array}{c} a-1 & b-1 & c+2 \\ | & | & | \\ | & | & | \\ \swarrow 1 & \searrow & \\ | & | & | \\ | & | & | \\ a & b & c \end{array} = 0. \quad (5.5i)$$

We define now a monoidal functor $\mathcal{G}_{m|n}^+ : \widetilde{\mathbf{Sp}}^+ \rightarrow \mathbf{Rep}_{m|n}^+$. On objects, $\mathcal{G}_{m|n}^+$ is given by $\mathcal{G}_{m|n}^+(a) = \bigwedge_q^a \mathbb{C}_q^{m|n}$, while on the morphisms (5.1) and (5.2) we set

$$\mathcal{G}_{m|n}^+(E^{(r)}) = E^{(r)} : \bigwedge_q^a \mathbb{C}_q^{m|n} \otimes \bigwedge_q^b \mathbb{C}_q^{m|n} \longrightarrow \bigwedge_q^{a+r} \mathbb{C}_q^{m|n} \otimes \bigwedge_q^{b-r} \mathbb{C}_q^{m|n}, \quad (5.6)$$

$$\mathcal{G}_{m|n}^+(F^{(r)}) = F^{(r)} : \bigwedge_q^a \mathbb{C}_q^{m|n} \otimes \bigwedge_q^b \mathbb{C}_q^{m|n} \longrightarrow \bigwedge_q^{a-r} \mathbb{C}_q^{m|n} \otimes \bigwedge_q^{b+r} \mathbb{C}_q^{m|n}, \quad (5.7)$$

where E and F , the generators of $U_q(\mathfrak{gl}_2)$, act on $\bigoplus_{a+b=N} \bigwedge^a \mathbb{C}_q^{m|n} \otimes \bigwedge^b \mathbb{C}_q^{m|n}$ by skew Howe duality. It follows from lemma 4.5 (and the fact that the relations from Definition 5.1 mimic the quantum group relations) that the functor is well-defined.

We then get automatically as a consequence of theorem 4.2:

Theorem 5.2 (See also Cautis et al. (2014)). *The functor $\mathcal{G}_{m|n}^+$ descends to a full functor $\mathcal{G}_{m|n}^+ : \mathbf{Sp}^+ \rightarrow \mathbf{Rep}_{m|n}^+$.*

In the non-super case there is actually an easy way to turn $\mathcal{G}_{m|n}^+$ into an equivalence of categories by passing to a quotient of \mathbf{Sp}^+ , as proven in Cautis et al. (2014) (and as also follows by the presentation of the Schur algebra given in Doty and Giaquinto (2002)):

Definition 5.3. *We define $\mathbf{Sp}_{\leq m}^+$ to be the quotient of the category \mathbf{Sp}^+ modulo all objects $a > m$ (formally, we quotient out all morphisms that factor through a tensor product of objects with at least one tensor factor greater than m).*

Then:

Theorem 5.4 (See (Cautis et al., 2014, Theorem 4.4.1)). *In case $n = 0$, the functor $\mathcal{G}_{m|0}^+$ descends to an equivalence of categories $\mathcal{G}_{m|0}^+ : \mathbf{Sp}_{\leq m}^+ \rightarrow \mathbf{Rep}_m^+$.*

For a sequence $\mathbf{a} = (a_1, \dots, a_r) \in \mathbf{Sp}^+$ we let $|\mathbf{a}| = a_1 + \dots + a_r$. Notice that, by construction, $\mathrm{Hom}_{\mathbf{Sp}^+}(\mathbf{a}, \mathbf{b}) \neq 0$ if and only if $|\mathbf{a}| = |\mathbf{b}|$. Hence we have:

Lemma 5.5. *Let $\mathbf{a}, \mathbf{b} \in \mathbf{Sp}^+$, and let $m \geq |\mathbf{a}|$. Then the quotient functor $\mathbf{Sp}^+ \rightarrow \mathbf{Sp}_{\leq m}^+$ induces an isomorphism $\mathrm{Hom}_{\mathbf{Sp}^+}(\mathbf{a}, \mathbf{b}) \cong \mathrm{Hom}_{\mathbf{Sp}_{\leq m}^+}(\mathbf{a}, \mathbf{b})$.*

We recall also the following lemma:

Lemma 5.6. *Let $\mathbf{a} \in \mathbf{Sp}^+$ and let $N = |\mathbf{a}|$. There are a monomorphism $\iota_{\mathbf{a}} : \mathbf{a} \hookrightarrow 1^{\otimes N}$ and an epimorphism $\pi_{\mathbf{a}} : 1^{\otimes N} \twoheadrightarrow \mathbf{a}$ with $\pi_{\mathbf{a}} \circ \iota_{\mathbf{a}} = \mathrm{id}_{\mathbf{a}}$.*

Proof. It is sufficient to prove the claim for an object $a \in \mathbb{Z}_{\geq 0}$. Let

$$\iota_a = F_{a-1}^{(1)} \cdots F_2^{(a-2)} F_1^{(a-1)} : a \cong a \otimes 0^{\otimes a-1} \rightarrow 1^{\otimes a} \quad (5.8)$$

and

$$\pi_a = E_1^{(a-1)} \cdots E_{a-2}^{(r)} E_{a-1}^{(1)} : 1^{\otimes a} \rightarrow a \otimes 0^{\otimes a-1} \cong a. \quad (5.9)$$

It follows from the relation (5.5e) that $\pi_a \circ \iota_a$ is a multiple of the identity, whence the claim. \square

As a consequence, for all objects $\mathbf{a}, \mathbf{b} \in \mathbf{Sp}^+$ with $|\mathbf{a}| = |\mathbf{b}| = N$ we can factor the identity of $\text{Hom}_{\mathbf{Sp}^+}(\mathbf{a}, \mathbf{b})$ as

$$\text{Hom}_{\mathbf{Sp}^+}(\mathbf{a}, \mathbf{b}) \hookrightarrow \text{Hom}_{\mathbf{Sp}^+}(1^{\otimes N}, 1^{\otimes N}) \twoheadrightarrow \text{Hom}_{\mathbf{Sp}^+}(\mathbf{a}, \mathbf{b}). \quad (5.10)$$

The functor \mathcal{J}^+

Set in \mathbf{Sp}^+

$$c_{1,1} = (q^{-1} - \text{FE}) \mathbf{1}_{1 \otimes 1} = q^{-1} \begin{array}{c} \begin{array}{cc} 1 & 1 \\ \uparrow & \uparrow \\ 1 & 1 \end{array} - \begin{array}{c} \begin{array}{cc} 1 & 1 \\ \uparrow & \uparrow \\ \downarrow & \downarrow \end{array} \end{array}, \quad (5.11)$$

$$c_{1,1}^{-1} = (q - \text{FE}) \mathbf{1}_{1 \otimes 1} = q \begin{array}{c} \begin{array}{cc} 1 & 1 \\ \uparrow & \uparrow \\ 1 & 1 \end{array} - \begin{array}{c} \begin{array}{cc} 1 & 1 \\ \uparrow & \uparrow \\ \downarrow & \downarrow \end{array} \end{array}. \quad (5.12)$$

The notation is motivated by the fact (easy to check) that $c_{1,1}^{-1}$ is the inverse of $c_{1,1}$.

We define a monoidal functor $\mathcal{J}^+ : \mathbf{H} \rightarrow \mathbf{Sp}^+$. On objects, we set $\mathcal{J}^+(1) = 1$. On morphisms, we set

$$\mathcal{J}^+ \left(\begin{array}{c} \begin{array}{cc} 1 & 1 \\ \uparrow & \uparrow \\ \downarrow & \downarrow \\ 1 & 1 \end{array} \end{array} \right) = c_{1,1} \quad \text{and} \quad \mathcal{J}^+ \left(\begin{array}{c} \begin{array}{cc} 1 & 1 \\ \uparrow & \uparrow \\ \downarrow & \downarrow \\ 1 & 1 \end{array} \end{array} \right) = c_{1,1}^{-1}. \quad (5.13)$$

It is easy to check that \mathcal{J}^+ is well-defined, i.e. that $c_{1,1}$ satisfies the defining relations (2.11) of the Hecke algebra. We will use the pictures in (5.13) to denote the morphisms $c_{1,1}$ and $c_{1,1}^{-1}$ of \mathbf{Sp}^+ .

REMARK 5.7. Alternatively, one could use lemma 5.8 below to check that \mathcal{J}^+ is well-defined by comparing, for each fixed $m \geq 0$, the composition of the functor \mathcal{J}^+ restricted to the full subcategory \mathcal{H}_m and composed with the quotient functor $\mathbf{Sp}^+ \rightarrow \mathbf{Sp}_{\leq m}^+$, to the composition $(\mathcal{G}_{m|0}^+)^{-1} \circ \mathcal{RT}_{m|0}$, where $(\mathcal{G}_{m|0}^+)^{-1}$ is an essential inverse to the equivalence of categories $\mathcal{G}_{m|0}^+ : \mathbf{Sp}_{\leq m}^+ \rightarrow \mathbf{Rep}_m^+$.

Lemma 5.8. *The composition $\mathcal{G}_{m|n}^+ \circ \mathcal{J}^+$ gives the Reshetikhin-Turaev invariant $\mathcal{RT}_{m|n}$.*

Proof. It suffices to check that the image of the generators of the Hecke algebra under the Reshetikhin-Turaev functor (that descends to \mathcal{H}) agree with the image under $\mathcal{G}_{m|n}^+ \circ \mathcal{J}^+$. This amounts to checking the following equality in $\mathbf{Rep}_{m|n}^+$:

$$\mathcal{G}_{m|n}^+ \left(\begin{array}{c} \begin{array}{cc} 1 & 1 \\ \downarrow & \downarrow \\ \text{---} & \text{---} \\ \uparrow & \uparrow \\ 1 & 1 \end{array} \end{array} \right) = - \mathcal{RT}_{m|n} \left(\begin{array}{c} \begin{array}{cc} 1 & 1 \\ \uparrow & \uparrow \\ \text{---} & \text{---} \\ \downarrow & \downarrow \\ 1 & 1 \end{array} \end{array} \right) + q^{-1} \mathcal{RT}_{m|n} \left(\begin{array}{c} \begin{array}{cc} 1 & 1 \\ \uparrow & \uparrow \\ \uparrow & \uparrow \\ 1 & 1 \end{array} \end{array} \right) \quad (5.14)$$

Let $x_a \otimes x_b \in \mathbb{C}_q^{m|n} \otimes \mathbb{C}_q^{m|n}$ be a standard basis vector. Under the isomorphism (4.9) it corresponds to $z_{a1} \wedge_q z_{b2} \in \bigwedge_q^2(\mathbb{C}_q^{m|n} \otimes \mathbb{C}_q^2)$. We can now compute

$$EF(z_{a1} \wedge_q z_{b2}) = E(z_{a2} \wedge_q z_{b2}) = z_{a2} \wedge_q z_{b1} + q z_{a1} \wedge_q z_{b2}. \quad (5.15)$$

We have now four cases:

- if $a = b$ and $|a| = 0$, then $z_{a2} \wedge_q z_{a1} = -q z_{a1} \wedge_q z_{a2}$ and hence (5.15) gives 0;
- if $a = b$ and $|a| = 1$, then $z_{a2} \wedge_q z_{a1} = q^{-1} z_{a1} \wedge_q z_{a2}$ and hence (5.15) gives $[2] z_{a1} \wedge_q z_{b2}$;
- if $a < b$, then $z_{a2} \wedge_q z_{b1} = -(-1)^{|a||b|} z_{b1} \wedge_q z_{a2} + (q^{-1} - q) z_{a1} \wedge_q z_{b2}$ and hence (5.15) is equal to $-(-1)^{|a||b|} z_{b1} \wedge_q z_{a2} + q^{-1} z_{a1} \wedge_q z_{b2}$;
- if $a > b$, then $z_{a2} \wedge_q z_{b1} = -(-1)^{|a||b|} z_{b1} \wedge_q z_{a2}$ and hence (5.15) is equal to $-(-1)^{|a||b|} z_{b1} \wedge_q z_{a2} + q z_{a1} \wedge_q z_{b2}$.

An immediate comparison with the formulas (3.12) for the R -matrix proves the claim. \square

Proposition 5.9. *The functor \mathcal{J}^+ is fully faithful.*

Proof. We must show that for all $r \geq 0$ the functor \mathcal{J}^+ induces an isomorphism $\mathcal{H}_r \cong \text{End}_{\text{Sp}^+}(1^{\otimes r})$. Choose $m \geq r$ and $n = 0$. Then, by Schur-Weyl duality, the composition $\mathcal{G}_{m|0}^+ \circ \mathcal{J}^+$ induces an isomorphism $\mathcal{H}_r \cong \text{End}_{\text{Rep}^+}((\mathbb{C}_q^m)^{\otimes r})$. By theorem 5.4 together with lemma 5.5, on the other side, $\mathcal{G}_{m|0}^+$ induces an isomorphism $\text{End}_{\text{Sp}^+}(1^{\otimes r}) \cong \text{End}_{\text{Rep}^+}((\mathbb{C}_q^m)^{\otimes r})$, and the claim follows. \square

We recall the following immediate consequence:

Lemma 5.10. *The homomorphism space $\text{End}_{\text{Sp}^+}(1^{\otimes N})$ is monoidally generated by $c_{1,1}$.*

Braided structure

As a reference for the definition of a braided category, we refer to (Turaev, 1994, §1.2), from where we also copy the notation.

We recall the following observation, which we will use in the proof of the next proposition.

Lemma 5.11 (See also (Cautis et al., 2014, §6.2)). *A braided structure on Sp^+ is uniquely determined by the braiding of $1 \otimes 1$. In other words, suppose we have two braided structures on Sp^+ , with braidings $c_{\mathbf{a},\mathbf{b}}$ and $c'_{\mathbf{a},\mathbf{b}}$ respectively. If $c_{1,1} = c'_{1,1}$ then the two braided structures coincide.*

Proof. We need to show that $c_{\mathbf{a},\mathbf{b}} = c'_{\mathbf{a},\mathbf{b}}$ for all objects $\mathbf{a}, \mathbf{b} \in \text{Sp}^+$. Since the braiding is compatible with the monoidal structure (see (Turaev, 1994, (1.2.b) and (1.2.c))), it is sufficient to prove that $c_{a,b} = c'_{a,b}$ for all $a, b \in \mathbb{Z}_{\geq 0}$. Using naturality (Turaev, 1994, (1.2.d)) we get

$$\begin{aligned} c_{a,b} &= c_{a,b}(\pi_a \otimes \pi_b)(\iota_a \otimes \iota_b) = (\pi_a \otimes \pi_b)c_{1 \otimes N, 1 \otimes N'}(\iota_a \otimes \iota_b) \\ &= (\pi_a \otimes \pi_b)c'_{1 \otimes N, 1 \otimes N'}(\iota_a \otimes \iota_b) = c'_{a,b}(\pi_a \otimes \pi_b)(\iota_a \otimes \iota_b) = c'_{a,b} \end{aligned} \quad (5.16)$$

and we are done. \square

The category Sp^+ is a sort of inverse limit of the categories $\text{Sp}_{\leq m}^+$. Each one of the categories $\text{Sp}_{\leq m}^+$, being equivalent to Rep_m^+ , is braided. This allows to induce a braided structure on Sp^+ :

Proposition 5.12. *The category \mathbf{Sp}^+ is a braided category, with braiding on $1 \otimes 1$ given by $c_{1,1}$.*

Proof. Given objects $\mathbf{a}, \mathbf{b} \in \mathbf{Sp}^+$, choose $m \geq |\mathbf{a}| + |\mathbf{b}|$. In the category $\mathbf{Sp}_{\leq m}^+$, which is braided since it is equivalent to \mathbf{Rep}_m^+ , there is a braiding morphism, which via the isomorphism given by lemma 5.5 gives an element $c_{\mathbf{a}, \mathbf{b}} \in \text{Hom}_{\mathbf{Sp}^+}(\mathbf{a}, \mathbf{b})$. This element does not depend on the choice of m (this is a consequence of the naturality of the construction, and can be proved using (5.11) and lemma 5.11). It follows that the elements $c_{\mathbf{a}, \mathbf{b}}$ define a braided structure (the axioms hold since they hold in $\mathbf{Sp}_{\leq m}^+$ for all big enough m). \square

Actually, we can also give a direct proof, which does not use the braided structure of \mathbf{Rep}_m^+ (and we could even recover the braided structure of \mathbf{Rep}_m^+ from it).

Alternative proof. We define the braiding on $1 \otimes 1$ using (5.11). By lemma 5.11, there is a unique possible way to extend it to a candidate braiding on the whole category, using (5.16). We then need to check that such extension is natural, i.e. that for $\mathbf{a}, \mathbf{a}', \mathbf{b}, \mathbf{b}' \in \mathbf{Sp}^+$ and $f: \mathbf{a} \rightarrow \mathbf{a}'$, $g: \mathbf{b} \rightarrow \mathbf{b}'$ we have

$$(f \otimes g)c_{\mathbf{a}, \mathbf{b}} = c_{\mathbf{a}', \mathbf{b}'}(f \otimes g). \quad (5.17)$$

We can suppose $|\mathbf{a}| = |\mathbf{a}'| = M$ and $|\mathbf{b}| = |\mathbf{b}'| = N$ (otherwise the claim is trivial). We then have

$$c_{\mathbf{a}', \mathbf{b}'}(f \otimes g) = (\pi_{\mathbf{a}'} \otimes \pi_{\mathbf{b}'})c_{1^{\otimes M}, 1^{\otimes N}}(\iota_{\mathbf{a}'} \otimes \iota_{\mathbf{b}'})(f \otimes g)(\pi_{\mathbf{a}} \otimes \pi_{\mathbf{b}})(\iota_{\mathbf{a}} \otimes \iota_{\mathbf{b}}). \quad (5.18)$$

If we set $\tilde{f} = \iota_{\mathbf{a}'} f \pi_{\mathbf{a}}$ and $\tilde{g} = \iota_{\mathbf{b}'} g \pi_{\mathbf{b}}$ and we suppose that we can prove naturality for $c_{1^{\otimes M}, 1^{\otimes N}}$, then we get $c_{1^{\otimes M}, 1^{\otimes N}}(\tilde{f} \otimes \tilde{g}) = (\tilde{f} \otimes \tilde{g})c_{1^{\otimes M}, 1^{\otimes N}}$ and (5.17) follows.

So we reduced the problem to the case $\mathbf{a} = \mathbf{a}' = 1^{\otimes M}$ and $\mathbf{b} = \mathbf{b}' = 1^{\otimes N}$. By lemma 5.10, we can then assume that f and g are just $c_{1,1}$ tensored with identities, and the claim then follows immediately from the braid relation for $c_{1,1}$. \square

Up to rescaling, our formula for $c_{1,1}$ coincides with Lusztig's symmetry (Lusztig, 1993, 5.2.1) (cf. also Queffelec (2015)). In (Cautis et al., 2014, §6.1) a braided structure on \mathbf{Sp}^+ is constructed using these Lusztig's symmetries. Then by lemma 5.11 we can deduce that the braiding defined in

proposition 5.12 is closely related to the one in (Cautis et al., 2014, §6.1), and we have the following explicit formula:

$$c_{a,b} = (-1)^{a+ab} q^{-a} \sum_{-k+l-m=a-b} q^{-km+l} E^{(k)} F^{(l)} E^{(m)} \mathbf{1}_{a \otimes b}. \quad (5.19)$$

It can be shown (Cautis et al., 2014, Lemma 6.1.1) that this formula is equivalent to the following ones:

$$c_{a,b} = \begin{cases} (-1)^{b(a+1)} q^{-b} \sum_{s \geq 0} (-q)^s F^{(s+a-b)} E^{(s)} \mathbf{1}_{a \otimes b} & \text{if } a - b \geq 0, \\ (-1)^{a(b+1)} q^{-a} \sum_{s \geq 0} (-q)^s E^{(s-a+b)} F^{(s)} \mathbf{1}_{a \otimes b} & \text{if } a - b \leq 0. \end{cases} \quad (5.20)$$

Lusztig's symmetries have been widely studied, in particular in Lusztig (1993). We recall some formulas from (Lusztig, 1993, Section 37.1.3) in our conventions:

$$\begin{array}{c} \text{Diagram 1} \\ a \quad b \end{array} = (-q)^{a-b} \begin{array}{c} \text{Diagram 2} \\ a \quad b \end{array}, \quad \begin{array}{c} \text{Diagram 3} \\ a \quad b \end{array} = (-q)^{b-a-2} \begin{array}{c} \text{Diagram 4} \\ a \quad b \end{array}, \quad (5.21)$$

$$\begin{array}{c} \text{Diagram 5} \\ a \quad b \end{array} = (-q)^{b-a+2} \begin{array}{c} \text{Diagram 6} \\ a \quad b \end{array}, \quad \begin{array}{c} \text{Diagram 7} \\ a \quad b \end{array} = (-q)^{a-b} \begin{array}{c} \text{Diagram 8} \\ a \quad b \end{array}. \quad (5.22)$$

6. Graphical calculus for duals

The category \mathbf{Sp}

We define now a new monoidal category \mathbf{Sp} by adding to \mathbf{Sp}^+ left and right duals. In detail, objects of \mathbf{Sp} are (formal direct sums of) sequences of integers and the tensor product of objects is given by concatenation of sequences. We set $a^* = -a$ for all $a \in \mathbb{Z}$. Morphisms are monoidally generated by the identity morphisms id_a for $a \in \mathbb{Z}$, by the morphisms of \mathbf{Sp}^+ (subject to the relations of \mathbf{Sp}^+) and by adjunction morphisms turning a^* into a left and right dual of a for all $a \in \mathbb{Z}_{>0}$.

We use orientations to distinguish in our pictures between objects $a > 0$ and their duals. Graphically, we represent the identity morphism $a^* \rightarrow a^*$ for $a > 0$ by

$$\begin{array}{c} a^* \\ \downarrow \\ a^* \end{array} \quad (6.1)$$

As usual, we represent the adjunction morphisms for $a > 0$ by the pictures

$$\begin{array}{c} \text{---} \curvearrowright \text{---} \\ a \quad a^* \end{array} \quad \begin{array}{c} \text{---} \curvearrowleft \text{---} \\ a^* \quad a \end{array} \quad \begin{array}{c} a^* \quad a \\ \text{---} \cup \text{---} \end{array} \quad \begin{array}{c} a \quad a^* \\ \text{---} \cup \text{---} \end{array} \quad (6.2)$$

Saying that they turn a^* into a left and right dual of a amounts to imposing the duality relations

$$\begin{array}{c} a \\ \downarrow \\ \text{---} \cup \text{---} \\ \uparrow \\ a \end{array} = \begin{array}{c} a \\ \downarrow \\ \text{---} \cup \text{---} \\ \uparrow \\ a \end{array} = \begin{array}{c} a \\ \downarrow \\ \text{---} \\ \uparrow \\ a \end{array} \quad \text{and} \quad \begin{array}{c} a^* \\ \downarrow \\ \text{---} \cup \text{---} \\ \uparrow \\ a^* \end{array} = \begin{array}{c} a^* \\ \downarrow \\ \text{---} \cup \text{---} \\ \uparrow \\ a^* \end{array} = \begin{array}{c} a^* \\ \downarrow \\ \text{---} \\ \uparrow \\ a^* \end{array} \quad (6.3)$$

. Notice that every object $\mathbf{a} = (a_1, \dots, a_\ell) \in \mathbf{Sp}$ has a left and right dual $\mathbf{a}^* = (a_\ell^*, \dots, a_1^*)$, with adjunction morphisms obtained by nesting (graphically) the generating adjunction morphisms (6.2). Hence, by adjunction, we have right and left duals of the generating morphisms $\mathbf{E}^{(r)}$ and $\mathbf{F}^{(r)}$. We impose that these left and right duals coincide. In pictures:

$$\begin{array}{c} (a+r)^* \quad (b-r)^* \\ \downarrow \quad \downarrow \\ \text{---} \cup \text{---} \\ \downarrow \quad \downarrow \\ a^* \quad b^* \end{array} = \begin{array}{c} (a+r)^* \quad (b-r)^* \\ \downarrow \quad \downarrow \\ \text{---} \cup \text{---} \\ \downarrow \quad \downarrow \\ a^* \quad b^* \end{array} \quad (6.4)$$

for $E^{(r)}\mathbf{1}_{(b-r, a+r)}$ and

$$(6.5)$$

for $F^{(r)}\mathbf{1}_{(b+r, a-r)}$. Since $E^{(r)}$ and $F^{(r)}$, together with identities and the adjunction morphisms, generate the morphisms of \mathbf{Sp} , it follows that the left and right duals of *every* morphism coincide.

We denote

$$E^{(r)}\mathbf{1}_{a^* \otimes b^*} = (F^{(r)}\mathbf{1}_{a+r \otimes b-r})^* \quad \text{and} \quad F^{(r)}\mathbf{1}_{a^* \otimes b^*} = (E^{(r)}\mathbf{1}_{a-r \otimes b+r})^* \quad (6.6)$$

and we use the pictures

$$(6.7)$$

for (6.4) and (6.5), respectively.

Since the category $\mathbf{Rep}_{m|n}$ is ribbon, we have

Proposition 6.1. *The functor $\mathcal{G}: \mathbf{Sp}^+ \rightarrow \mathbf{Rep}_{m|n}^+$ extends to a functor $\mathcal{G}: \mathbf{Sp} \rightarrow \mathbf{Rep}_{m|n}$.*

Proof. This is almost straightforward. It is sufficient to map the adjunction morphisms (6.2) to the duality morphisms of the ribbon structure of $\mathbf{Rep}_{m|n}$ under the functor \mathcal{G} . It is then a general consequence of the fact that the category $\mathbf{Rep}_{m|n}$ is ribbon that left and right duals of morphisms agree, which is expressed diagrammatically by relations (6.4) and (6.5). \square

Proposition 6.2. *The functor $\mathcal{G}: \mathbf{Sp} \rightarrow \mathbf{Rep}_{m|n}$ is full.*

Proof. Let $\mathbf{a} = (a_1, \dots, a_\ell)$, $\mathbf{b} = (b_1, \dots, b_\kappa)$ be two objects of \mathbf{Sp} , and suppose without loss of generality that $a_i, b_j \neq 0$ for all i, j . Suppose first that the following condition is satisfied:

- (*) there are indexes $1 \leq \ell' \leq \ell$ and $1 \leq \kappa' \leq \kappa$ such that $a_i > 0$ if and only if $i \leq \ell'$ while $b_j > 0$ if and only if $j > \kappa'$.

Let $\mathbf{a}' = (a_1, \dots, a_{\ell'})$ and $\mathbf{a}'' = (a_{\ell'+1}, \dots, a_\ell)$, and $\mathbf{b}' = (b_1, \dots, b_{\kappa'})$ and $\mathbf{b}'' = (b_{\kappa'+1}, \dots, b_\kappa)$. Let us define sequences \mathbf{c} and \mathbf{d} of objects of \mathbf{Sp}^+ by $\mathbf{c} = (\mathbf{b}')^* \otimes \mathbf{a}'$ and $\mathbf{d} = \mathbf{b}'' \otimes (\mathbf{a}'')^*$. Then we have an isomorphism

$$\begin{aligned} \mathrm{Hom}_{\mathbf{Rep}_{m|n}}(\bigwedge_q^{\mathbf{a}}, \bigwedge_q^{\mathbf{b}}) &\longrightarrow \mathrm{Hom}_{\mathbf{Rep}_{m|n}^+}(\bigwedge_q^{\mathbf{c}}, \bigwedge_q^{\mathbf{d}}) \\ \gamma &\longmapsto \hat{\gamma} = (\mathrm{ev}_{\mathbf{b}'^*} \otimes \mathrm{id}_{\mathbf{d}})(\mathrm{id}_{\mathbf{b}'^*} \otimes \gamma \otimes \mathrm{id}_{\mathbf{a}''^*})(\mathrm{id}_{\mathbf{c}} \otimes \mathrm{coev}_{\mathbf{a}''}) \end{aligned} \quad (6.8)$$

whose inverse is given by

$$\begin{aligned} \mathrm{Hom}_{\mathbf{Rep}_{m|n}^+}(\bigwedge_q^{\mathbf{c}}, \bigwedge_q^{\mathbf{d}}) &\longrightarrow \mathrm{Hom}_{\mathbf{Rep}}(\bigwedge_q^{\mathbf{a}}, \bigwedge_q^{\mathbf{b}}) \\ \gamma &\longmapsto \tilde{\gamma} = (\mathrm{id}_{\mathbf{b}} \otimes \mathrm{ev}_{\mathbf{a}''^*})(\mathrm{id}_{\mathbf{b}'} \otimes \gamma \otimes \mathrm{id}_{\mathbf{a}''})(\mathrm{coev}_{\mathbf{b}'} \otimes \mathrm{id}_{\mathbf{a}}), \end{aligned} \quad (6.9)$$

where for simplicity we omitted the symbol \bigwedge_q in the subscripts.

Let now $\gamma: \bigwedge_q^{\mathbf{a}} \rightarrow \bigwedge_q^{\mathbf{b}}$ be a morphism in $\mathbf{Rep}_{m|n}$. Since $\mathcal{G}_{m|n}^+$ is full, there is an element $\varphi \in \mathrm{Hom}_{\mathbf{Sp}^+}(\mathbf{c}, \mathbf{d})$ such that $\mathcal{G}_{m|n}^+(\varphi) = \hat{\gamma}$. Let

$$\tilde{\varphi} = (\mathrm{id}_{\mathbf{b}} \otimes \mathrm{ev}_{\mathbf{a}''^*})(\mathrm{id}_{\mathbf{b}'} \otimes \varphi \otimes \mathrm{id}_{\mathbf{a}''})(\mathrm{coev}_{\mathbf{b}'} \otimes \mathrm{id}_{\mathbf{a}}) \in \mathbf{Sp}, \quad (6.10)$$

where now $\mathrm{ev}_{\mathbf{a}''^*}$ and $\mathrm{coev}_{\mathbf{b}'}$ denote the adjunction morphisms in \mathbf{Sp} . Then $\mathcal{G}_{m|n}(\tilde{\varphi}) = \tilde{\hat{\gamma}} = \gamma$, hence γ is in the image of $\mathcal{G}_{m|n}$.

Let now \mathbf{a} and \mathbf{b} be general, and let $\gamma \in \mathrm{Hom}_{\mathbf{Rep}}(\bigwedge_q^{\mathbf{a}}, \bigwedge_q^{\mathbf{b}})$. Choose permutations w and z such that $w\mathbf{a}$ and $z\mathbf{b}$ satisfy (*). Corresponding to some reduced expressions of w and z there are isomorphisms $\Psi_1: \bigwedge_q^{\mathbf{a}} \rightarrow \bigwedge_q^{w\mathbf{a}}$ and $\Psi_2: \bigwedge_q^{\mathbf{b}} \rightarrow \bigwedge_q^{z\mathbf{b}}$ which are given by the braiding of \mathbf{Rep} . Although the category \mathbf{Sp} is not braided, it is straightforward to see that these isomorphisms are in the image of $\mathcal{G}_{m|n}$, since the braiding morphisms of $\mathbf{Rep}_{m|n}$ are obtained by taking the duals (graphically “rotating”) of the braiding morphisms of $\mathbf{Rep}_{m|n}^+$ (cf. also the pictures (6.21) and (6.22) below). So there exist morphisms $\widetilde{\Psi}_1$

and $\widetilde{\Psi_2^{-1}}$ in \mathbf{Sp} such that $\mathcal{G}(\widetilde{\Psi_1}) = \Psi_1$ and $\mathcal{G}(\widetilde{\Psi_2^{-1}}) = \Psi_2^{-1}$. By the first part of the proof, there is a morphism φ in \mathbf{Sp} such that $\mathcal{G}(\varphi) = \Psi_2 \circ \gamma \circ \Psi_1^{-1}$. Hence $\mathcal{G}(\widetilde{\Psi_2^{-1}} \circ \varphi \circ \widetilde{\Psi_1}) = \gamma$, and we are done. \square

REMARK 6.3. Notice that proposition 6.2 is a special case of the following more general statement. Consider a full strong monoidal functor $\mathcal{G}: \mathcal{M}^+ \rightarrow \mathcal{B}$ from a monoidal category \mathcal{M}^+ to a ribbon category \mathcal{B} . Suppose that the images of objects of \mathcal{M}^+ together with their duals in \mathcal{B} generate the objects of \mathcal{B} . Suppose also that the category \mathcal{M}^+ is a subcategory of a rigid category \mathcal{M} and the functor \mathcal{G} extends to a functor $\tilde{\mathcal{G}}: \mathcal{M} \rightarrow \mathcal{B}$. Then $\tilde{\mathcal{G}}$ is also full.

We state also the following result, which seems actually to be known, but we have not been able to find a reference in full generality. The analogous statement in the non-quantized setting can be found in (Brundan and Stroppel, 2012, Theorem 7.8).

Proposition 6.4 (Super mixed quantum Schur-Weyl duality). *The Reshetikhin-Turaev functor $\mathcal{RT}_{m|n}$ induces a full functor $\mathbf{Br}(d) \rightarrow \mathbf{Rep}_{m|n}$. In particular, for any $r, s \geq 0$ we have a surjective map*

$$\mathbf{Br}_{\eta_{r,s}}(d) \longrightarrow \mathbf{End}_{\mathbf{Rep}_{m|n}} \left((\mathbb{C}_q^{m|n})^{\otimes r} \otimes (\mathbb{C}_q^{m|n})^{*\otimes s} \right). \quad (6.11)$$

Moreover, if $(m+1)(n+1) > r+s$ then this map is also injective.

Proof. The fullness of the functor $\mathbf{Br}(d) \rightarrow \mathbf{Rep}_{m|n}$ follows from the fullness of the functor $\mathbf{H} \rightarrow \mathbf{Rep}_{m|n}^+$ exactly as in the proof of proposition 6.2 (cf. also remark 6.3). If $(m+1)(n+1) > r+s$ then it follows that (6.11) is faithful by comparing dimensions, since $\dim \mathbf{Br}_{\eta_{r,s}}(d) = (r+s)! = \dim \mathcal{H}_{r+s}$ and

$$\dim \mathbf{End}_{\mathbf{Rep}_{m|n}} \left((\mathbb{C}_q^{m|n})^{\otimes r} \otimes (\mathbb{C}_q^{m|n})^{*\otimes s} \right) = \dim \mathbf{End}_{\mathbf{Rep}_{m|n}} \left((\mathbb{C}_q^{m|n})^{\otimes r+s} \right) = \dim \mathcal{H}_{r+s}. \quad (6.12)$$

The last equation follows by super Schur-Weyl duality (proposition 3.1), since all partitions of $r+s$ are in $H_{m|n}$ if $(m+1)(n+1) > r+s$. \square

We define also the following morphisms:

$$\begin{aligned}
 E^{(r)} \mathbf{1}_{a \otimes b^*} &= \begin{array}{c} \begin{array}{cc} a+r & (b+r)^* \\ \downarrow & \downarrow \\ \text{---} r \text{---} \\ \downarrow & \downarrow \\ a & b^* \end{array} \\ = \begin{array}{cc} a+r & (b+r)^* \\ \downarrow & \downarrow \\ \text{---} r \text{---} \\ \downarrow & \downarrow \\ a & b^* \end{array} \end{array}, & F^{(r)} \mathbf{1}_{a \otimes b^*} = \begin{array}{c} \begin{array}{cc} a-r & (b-r)^* \\ \downarrow & \downarrow \\ \text{---} r \text{---} \\ \downarrow & \downarrow \\ a & b^* \end{array} \\ = \begin{array}{cc} a-r & (b-r)^* \\ \downarrow & \downarrow \\ \text{---} r \text{---} \\ \downarrow & \downarrow \\ a & b^* \end{array} \end{array}, \quad (6.13)
 \end{aligned}$$

$$\begin{aligned}
 E^{(r)} \mathbf{1}_{a^* \otimes b} &= \begin{array}{c} \begin{array}{cc} (a-r)^* & b-r \\ \downarrow & \downarrow \\ \text{---} r \text{---} \\ \downarrow & \downarrow \\ a^* & b \end{array} \\ = \begin{array}{cc} (a-r)^* & b-r \\ \downarrow & \downarrow \\ \text{---} r \text{---} \\ \downarrow & \downarrow \\ a^* & b \end{array} \end{array}, & F^{(r)} \mathbf{1}_{a^* \otimes b} = \begin{array}{c} \begin{array}{cc} (a+r)^* & b+r \\ \downarrow & \downarrow \\ \text{---} r \text{---} \\ \downarrow & \downarrow \\ a^* & b \end{array} \\ = \begin{array}{cc} (a+r)^* & b+r \\ \downarrow & \downarrow \\ \text{---} r \text{---} \\ \downarrow & \downarrow \\ a^* & b \end{array} \end{array}. \quad (6.14)
 \end{aligned}$$

The category $\mathbf{Sp}(\beta)$

As we have seen, via the functor $\mathcal{G}_{m|n}$ the category \mathbf{Sp} gives all morphisms of $\mathbf{Rep}_{m|n}$. However, \mathbf{Sp} is somehow too big (any non-trivial homomorphism space is infinite dimensional, since it contains the diagram consisting of a disjoint union of an arbitrary number of bubbles). We hence define a quotient $\mathbf{Sp}(\beta)$, depending on the choice of our element q^β .

Definition 6.5. *We define the category $\mathbf{Sp}(\beta)$ to be the quotient of the category \mathbf{Sp} modulo the relations*

$$\begin{array}{c} 1 \\ \downarrow \\ \text{---} 1 \text{---} \\ \downarrow \\ 1 \end{array} = [\beta - 1] \begin{array}{c} 1 \\ \downarrow \\ 1 \end{array}, \quad \begin{array}{c} 1 \\ \downarrow \\ \text{---} 1 \text{---} \\ \downarrow \\ 1 \end{array} = [\beta - 1] \begin{array}{c} 1 \\ \downarrow \\ 1 \end{array}, \quad (6.15a)$$

and

$$1 \bigcirc = 1 \bigcirc = [\beta], \quad (6.15b)$$

$$\begin{array}{c} 1 \quad 1^* \\ \downarrow \quad \downarrow \\ \text{---} \quad \text{---} \\ \downarrow \quad \downarrow \\ 1 \quad 1^* \end{array} - \begin{array}{c} 1 \quad 1^* \\ \downarrow \quad \downarrow \\ \text{---} 1 \text{---} \\ \downarrow \quad \downarrow \\ 1 \quad 1^* \end{array} = [2 - \beta] \begin{array}{c} 1 \quad 1^* \\ \downarrow \quad \downarrow \\ 1 \quad 1 \end{array}, \quad (6.15c)$$

$$\begin{array}{c} 1^* \quad 1 \\ \downarrow \quad \uparrow \\ \text{---} \end{array} - \begin{array}{c} 1^* \quad 1 \\ \downarrow \quad \uparrow \\ \text{---} \\ \downarrow \quad \uparrow \\ 1^* \quad 1 \end{array} = [2 - \beta] \begin{array}{c} 1^* \quad 1 \\ \downarrow \quad \uparrow \\ \text{---} \\ \downarrow \quad \uparrow \\ 1^* \quad 1 \end{array}. \quad (6.15d)$$

Recall that we set $d = m - n$. We then have:

Theorem 6.6. *The functor $\mathcal{G}_{m|n}: \mathbf{Sp} \rightarrow \mathbf{Rep}_{m|n}$ descends to a full functor*

$$\mathcal{G}_{m|n}: \mathbf{Sp}(d) \rightarrow \mathbf{Rep}_{m|n}. \quad (6.16)$$

Proof. We need to check that relations (6.15) are satisfied by morphisms in $\mathbf{Rep}_{m|n}$. Relation (6.15b) is (3.21). Let us check the first one of (6.15a) (the second one is analogous). Using (5.14) we have in $\mathbf{Rep}_{m|n}$:

$$\begin{array}{c} 1 \\ \uparrow \\ \text{---} \\ \downarrow \\ 1 \end{array} \begin{array}{c} 1 \\ \uparrow \\ \text{---} \\ \downarrow \\ 1 \end{array} = - \begin{array}{c} 1 \\ \uparrow \\ \text{---} \\ \downarrow \\ 1 \end{array} \begin{array}{c} 1 \\ \uparrow \\ \text{---} \\ \downarrow \\ 1 \end{array} + q^{-1} \begin{array}{c} 1 \\ \uparrow \\ \text{---} \\ \downarrow \\ 1 \end{array} \begin{array}{c} 1 \\ \uparrow \\ \text{---} \\ \downarrow \\ 1 \end{array} = -q^{-d} + q^{-1}[d] \begin{array}{c} 1 \\ \uparrow \\ \text{---} \\ \downarrow \\ 1 \end{array} = [d-1] \begin{array}{c} 1 \\ \uparrow \\ \text{---} \\ \downarrow \\ 1 \end{array}. \quad (6.17)$$

We now check (6.15c). Again using (5.14) we have in $\mathbf{Rep}_{m|n}$:

$$\begin{array}{c} 1 \quad 1^* \\ \uparrow \quad \downarrow \\ \text{---} \\ \downarrow \quad \uparrow \\ 1 \quad 1^* \end{array} = \begin{array}{c} 1 \quad 1^* \\ \uparrow \quad \downarrow \\ \text{---} \\ \downarrow \quad \uparrow \\ 1 \quad 1^* \end{array} - q^{-1} \begin{array}{c} 1 \quad 1^* \\ \uparrow \quad \downarrow \\ \text{---} \\ \downarrow \quad \uparrow \\ 1 \quad 1^* \end{array} - q \begin{array}{c} 1 \quad 1^* \\ \uparrow \quad \downarrow \\ \text{---} \\ \downarrow \quad \uparrow \\ 1 \quad 1^* \end{array} + \begin{array}{c} 1 \quad 1^* \\ \uparrow \quad \downarrow \\ \text{---} \\ \downarrow \quad \uparrow \\ 1 \quad 1^* \end{array} \\
= \begin{array}{c} 1 \quad 1^* \\ \uparrow \quad \downarrow \\ \text{---} \\ \downarrow \quad \uparrow \\ 1 \quad 1^* \end{array} - q^{-1+d} \begin{array}{c} 1 \quad 1^* \\ \uparrow \quad \downarrow \\ \text{---} \\ \downarrow \quad \uparrow \\ 1 \quad 1^* \end{array} - q^{1-d} \begin{array}{c} 1 \quad 1^* \\ \uparrow \quad \downarrow \\ \text{---} \\ \downarrow \quad \uparrow \\ 1 \quad 1^* \end{array} + [d] \begin{array}{c} 1 \quad 1^* \\ \uparrow \quad \downarrow \\ \text{---} \\ \downarrow \quad \uparrow \\ 1 \quad 1^* \end{array} \quad (6.18)$$

and the claim follows since $-q^{-1+d} - q^{1-d} + [d] = [d-2]$. Analogously one can check (6.15d). \square

Proposition 6.7. *The functor $\mathcal{I}^+: \mathbf{H} \rightarrow \mathbf{Sp}^+$ extends to a fully faithful functor $\mathcal{I}_\beta: \mathbf{Br}(\beta) \rightarrow \mathbf{Sp}(\beta)$.*

Proof. Since the oriented Brauer category $\mathbf{Br}(\beta)$ is generated by the braiding from \mathbf{H} together with the adjunction maps (cups and caps), there is a unique natural way to extend the functor \mathcal{J}^+ to $\mathbf{Br}(\beta)$. We should check that the defining relations of $\mathbf{Br}(\beta)$ hold in the image. Relation (2.9a) is a direct consequence of (6.15b), while relations (2.9b) can be showed by the same computation as in (6.17).

Now, let us first prove that \mathcal{J}_d is full. The main ingredient will be the fact that $\mathcal{J}: \mathbf{H} \rightarrow \mathbf{Sp}^+$ is fully faithful (proposition 5.9). Consider a diagram element $\varphi \in \text{Hom}_{\mathbf{Sp}(\beta)}(\mathbf{a}, \mathbf{b})$ with \mathbf{a} and \mathbf{b} containing only copies of 1 and 1^* . We slice φ into elementary pieces, as (5.3), (6.2) and (6.7), tensored with identities. Of course the labels that appear in the slices can be greater than 1. We thus replace, at the bottom and the top of each slice, each higher label a (respectively, a^*) by $1^{\otimes a}$ (respectively, $(1^*)^{\otimes a}$) by inserting $\pi_a \iota_a$ (respectively, its dual), where ι_a and π_a are the injection and the projection from lemma 5.6. Moreover, we can suppose that all caps and cups are labeled by 1 using the following trick:

The diagram shows an equality between two expressions. On the left, a curved line (cap) connects two boxes. The left box is labeled π_a and has multiple upward arrows from a label '1'. The right box is labeled ι_a^* and has multiple downward arrows to a label '1*'. On the right, the same cap is shown, but the left box is now labeled ι_a (with a downward arrow to the π_a box) and the right box is labeled π_a (with an upward arrow from the ι_a^* box). The right box also has multiple downward arrows to a label '1*'. The entire equation is labeled (6.19) on the far right.

(and similarly in the case of cups).

Now we have sliced our diagram so that if one slice contains higher labels then it does not contain cups or caps (so if it contains higher labels then there is no interaction between upwards pointing and downwards pointing strands). The slices containing cups or caps lie in the image of \mathcal{J}_β , obviously. The other slices lie also in the image of \mathcal{J}_β , since \mathcal{J}^+ is full. Hence \mathcal{J}_β is full.

In order to prove that \mathcal{J}_β is faithful, consider first the case $\beta = d$. For any objects $\mathbf{a}, \mathbf{b} \in \mathbf{Br}(d)$ we can choose m, n big enough so that the Reshetikhin-Turaev functor $\mathcal{RT}_{m|n}$ induces an isomorphism $\text{Hom}_{\mathbf{Br}(d)}(\mathbf{a}, \mathbf{b}) \rightarrow \text{Hom}_{\text{Rep}_{m|n}}(\bigwedge_q^{\mathbf{a}}, \bigwedge_q^{\mathbf{b}})$ (this is an immediate generalization of proposition 6.4). Since this map factors through $\text{Hom}_{\mathbf{Br}(d)}(\mathbf{a}, \mathbf{b}) \rightarrow \text{Hom}_{\mathbf{Sp}(d)}(\mathbf{a}, \mathbf{b})$, and we just proved that this is surjective, it follows that this is injective. The case β generic follows by specialization (formally, one needs to define the category $\mathbf{Sp}(\beta)$ over $\mathbb{C}(q)[q^\beta, q^{-\beta}]$,

and then the argument before shows that $\text{Hom}_{\mathbf{Sp}(d)}(\mathbf{a}, \mathbf{b})$ is free of the same rank as for $\beta = d$. \square

By construction, there is an obvious functor $i: \mathbf{Sp}^+ \rightarrow \mathbf{Sp}$. We will denote also by $i: \mathbf{Sp}^+ \rightarrow \mathbf{Sp}(\beta)$ its composition with the quotient functor $\mathbf{Sp} \rightarrow \mathbf{Sp}(\beta)$.

Proposition 6.8. *The functor $i: \mathbf{Sp}^+ \rightarrow \mathbf{Sp}(\beta)$ is fully faithful.*

Proof. We shall prove that i induces an isomorphism $\text{Hom}_{\mathbf{Sp}^+}(\mathbf{a}, \mathbf{b}) \cong \text{Hom}_{\mathbf{Sp}(\beta)}(\mathbf{a}, \mathbf{b})$ for all objects $\mathbf{a}, \mathbf{b} \in \mathbf{Sp}^+$. First, we claim that it is enough to consider the case $\mathbf{a} = \mathbf{b} = 1^{\otimes N}$. Indeed, using (5.10) we have a commutative diagram

$$\begin{array}{ccccc}
 \text{Hom}_{\mathbf{Sp}^+}(\mathbf{a}, \mathbf{b}) & \hookrightarrow & \text{Hom}_{\mathbf{Sp}^+}(1^{\otimes N}, 1^{\otimes N}) & \twoheadrightarrow & \text{Hom}_{\mathbf{Sp}^+}(\mathbf{a}, \mathbf{b}) \\
 \downarrow & & \downarrow & & \downarrow \\
 \text{Hom}_{\mathbf{Sp}(\beta)}(\mathbf{a}, \mathbf{b}) & \hookrightarrow & \text{Hom}_{\mathbf{Sp}(\beta)}(1^{\otimes N}, 1^{\otimes N}) & \twoheadrightarrow & \text{Hom}_{\mathbf{Sp}(\beta)}(\mathbf{a}, \mathbf{b})
 \end{array} \tag{6.20}$$

Then supposing that the middle vertical map is an isomorphism, the left square gives the injectivity and the right square gives the surjectivity of the map $\text{Hom}_{\mathbf{Sp}^+}(\mathbf{a}, \mathbf{b}) \rightarrow \text{Hom}_{\mathbf{Sp}(\beta)}(\mathbf{a}, \mathbf{b})$, which hence is an isomorphism.

Se we can restrict to the case of $1^{\otimes N}$. Since the functors \mathcal{I}^+ and \mathcal{I}_β are fully faithful, we have isomorphisms $\text{End}_{\mathbf{Sp}^+}(1^{\otimes N}) \cong \mathcal{H}_N \cong \text{End}_{\mathbf{Br}(\beta)}(1^{\otimes N}) \cong \text{End}_{\mathbf{Sp}(\beta)}(1^{\otimes N})$, and we are done. \square

As a corollary of proposition 6.7 or alternatively of proposition 6.8 we have the following important fact:

Corollary 6.9. *In $\mathbf{Sp}(\beta)$ we have $\text{End}_{\mathbf{Sp}(\beta)}(a) = \mathbb{C}_q$ for all $a \in \mathbb{Z}_{\geq 0}$.*

Ribbon structure

In $\mathbf{Sp}(\beta)$ let us define

$$c_{1^*,1} = \begin{array}{c} 1 \quad 1^* \\ \nearrow \quad \searrow \\ \searrow \quad \nearrow \\ 1^* \quad 1 \end{array} = \begin{array}{c} 1 \quad 1^* \\ \nearrow \quad \searrow \\ \nearrow \quad \searrow \\ 1^* \quad 1 \end{array}, \quad c_{1^*,1}^{-1} = \begin{array}{c} 1 \quad 1^* \\ \nearrow \quad \searrow \\ \searrow \quad \nearrow \\ 1^* \quad 1 \end{array} = \begin{array}{c} 1 \quad 1^* \\ \nearrow \quad \searrow \\ \searrow \quad \nearrow \\ 1^* \quad 1 \end{array}, \quad (6.21)$$

$$c_{1,1^*}^{-1} = \begin{array}{c} 1^* \quad 1 \\ \nearrow \quad \searrow \\ \searrow \quad \nearrow \\ 1 \quad 1^* \end{array} = \begin{array}{c} 1^* \quad 1 \\ \nearrow \quad \searrow \\ \nearrow \quad \searrow \\ 1 \quad 1^* \end{array}, \quad c_{1,1^*} = \begin{array}{c} 1^* \quad 1 \\ \nearrow \quad \searrow \\ \searrow \quad \nearrow \\ 1 \quad 1^* \end{array} = \begin{array}{c} 1^* \quad 1 \\ \nearrow \quad \searrow \\ \searrow \quad \nearrow \\ 1 \quad 1^* \end{array}. \quad (6.22)$$

The notation we use is motivated by the following lemma:

Lemma 6.10. *We have*

$$\begin{array}{c} 1 \quad 1^* \\ \nearrow \quad \searrow \\ \searrow \quad \nearrow \\ 1 \quad 1^* \end{array} = \begin{array}{c} 1 \quad 1^* \\ \nearrow \quad \searrow \\ \nearrow \quad \searrow \\ 1 \quad 1^* \end{array} = \text{id}_{1 \otimes 1^*} \quad \text{and} \quad \begin{array}{c} 1^* \quad 1 \\ \nearrow \quad \searrow \\ \searrow \quad \nearrow \\ 1^* \quad 1 \end{array} = \begin{array}{c} 1^* \quad 1 \\ \nearrow \quad \searrow \\ \searrow \quad \nearrow \\ 1^* \quad 1 \end{array} = \text{id}_{1^* \otimes 1}. \quad (6.23)$$

Proof. Let us just check one of the four equalities, the other ones being similar (actually, one of the four even implies the other three using the properties of the braiding $c_{1,1}$ and of the duality, see [Ohtsuki \(2002\)](#) and [Turaev \(1994\)](#)).

We compute:

$$\begin{aligned}
& \text{Diagram 1} = \text{Diagram 2} - q \text{Diagram 3} - q^{-1} \text{Diagram 4} + \text{Diagram 5} \\
& = ([\beta] - [2][\beta - 1]) \text{Diagram 6} + [2 - \beta] \text{Diagram 7} + \text{Diagram 8} .
\end{aligned}
\tag{6.24}$$

Let us expand the second summand:

$$\text{Diagram 7} = [2] \text{Diagram 9} = [2][\beta - 1] \text{Diagram 10} .
\tag{6.25}$$

Let us now expand the last summand:

$$\text{Diagram 8} = \text{Diagram 11} = 3 \text{Diagram 12} + \text{Diagram 13}
\tag{6.26}$$

$$\begin{aligned}
&= \text{Diagram 1} + [\beta - 1][\beta - 1] \text{Diagram 2} \\
&= [\beta - 2] \text{Diagram 3} + [\beta - 1][\beta - 1] \text{Diagram 4} \\
&= [\beta - 2] \text{Diagram 5} + ([\beta - 2][\beta - 2] + [\beta - 1][\beta - 1]) \text{Diagram 6}.
\end{aligned}$$

The diagrams are as follows:

- Diagram 1: A vertical line with a loop on the left. The loop has three crossings and is labeled with '3' inside. The top and bottom of the loop are labeled '1' and '1*' respectively. The main vertical line is also labeled '1' and '1*' at the top and bottom.
- Diagram 2: Two parallel vertical lines, both labeled '1' and '1*' at the top and bottom.
- Diagram 3: A vertical line with a loop on the left. The loop has two crossings and is labeled with '1' and '1*' at the top and bottom. The main vertical line is also labeled '1' and '1*' at the top and bottom.
- Diagram 4: Two parallel vertical lines, both labeled '1' and '1*' at the top and bottom.
- Diagram 5: A vertical line with a loop on the left. The loop has one crossing and is labeled with '1' and '1*' at the top and bottom. The main vertical line is also labeled '1' and '1*' at the top and bottom.
- Diagram 6: Two parallel vertical lines, both labeled '1' and '1*' at the top and bottom.

Above we passed from the second line to the third one by applying an analogue of (6.15a) with color 2 on the boundary points. Such an equation can be obtained from (6.15a) by first applying (5.5e) to reduce the colors.

Putting all together, the claim follows since

$$[\beta] - [2][\beta - 1] + [\beta - 2] = -[\beta - 2] + [\beta - 2] = 0 \quad (6.27)$$

and

$$[2 - \beta][2][\beta - 1] + [\beta - 2][\beta - 2] + [\beta - 1][\beta - 1] = [2 - \beta][\beta] + [\beta - 1][\beta - 1] = 1 \quad (6.28)$$

(using the formulas from section 2). \square

We define more generally candidate braiding elements c_{a,b^*} , c_{a,b^*}^{-1} , $c_{a^*,b}$ and $c_{a^*,b}^{-1}$ exactly as we did for $a = b = 1$. It follows from lemma 6.10 and from the proof of lemma 5.11, and in particular from (5.16), that they are inverse to each other. In addition, we define elements c_{a^*,b^*} and c_{a^*,b^*}^{-1} by taking the duals of the elements $c_{a,b}$ and $c_{a,b}^{-1}$, respectively.

We then define recursively

$$c_{a,b \otimes b'} = (\text{id}_a \otimes c_{a,b'})(c_{a,b} \otimes \text{id}_b), \quad (6.29)$$

$$c_{a \otimes a',b} = (c_{a,b} \otimes \text{id}_{a'})(\text{id}_a \otimes c_{a',b}). \quad (6.30)$$

Notice that this is well-defined, i.e. when defining $c_{\mathbf{a} \otimes \mathbf{a}', \mathbf{b} \otimes \mathbf{b}'}$ it does not matter in which order we use (6.29) and (6.30) above.

Lemma 6.11. *The family of maps $\{c_{\mathbf{a}, \mathbf{b}} \mid \mathbf{a}, \mathbf{b} \in \mathbf{Sp}(\beta)\}$ is natural.*

Proof. Let $f: \mathbf{a} \rightarrow \mathbf{a}'$, $g: \mathbf{b} \rightarrow \mathbf{b}'$. We need to show that $(g \otimes f)c_{\mathbf{a}, \mathbf{b}} = c_{\mathbf{a}', \mathbf{b}'}(f \otimes g)$. Since morphisms in $\mathbf{Sp}(\beta)$ are generated by the morphisms $\mathbf{E}^{(r)}$ and $\mathbf{F}^{(r)}$ of \mathbf{Sp}^+ and the adjunction morphisms (6.2), it suffices to consider such morphisms. For $\mathbf{E}^{(r)}$ and $\mathbf{F}^{(r)}$ the statement follows by the naturality of the braiding of \mathbf{Sp}^+ (see proposition 5.12). Hence we only need to consider the adjunction morphisms. For them, the claim follows immediately from the fact that $c_{\mathbf{a}, \mathbf{b}}^{-1}$ is the inverse of $c_{\mathbf{a}, \mathbf{b}}$, as illustrated below:

□

We will now define a family of twists $\{\theta_{\mathbf{a}}\}_{\mathbf{a} \in \mathbf{Sp}(\beta)}$ compatible with the braiding $\{c_{\mathbf{a}, \mathbf{b}}\}_{\mathbf{a}, \mathbf{b} \in \mathbf{Sp}(\beta)}$. For $a \in \mathbb{Z}_{\geq 0}$ set $\theta_a \in \text{End}_{\mathbf{Sp}(\beta)}(a)$ to be multiplication by $q^{-a\beta + a(a-1)}$. In pictures

$$\begin{array}{c} a \\ \uparrow \\ \boxed{\theta_a} \\ \downarrow \\ a \end{array} = q^{-a\beta + a(a-1)} \begin{array}{c} a \\ \downarrow \\ a \end{array}. \quad (6.31)$$

The choice of the scalar is made in such a way that the functor $\mathcal{G}_{m|n}$ sends θ_a to the twist of $\mathbf{Rep}_{m|n}$, cf. (3.20). Moreover we let θ_{a^*} be the dual of θ_a , that is

$$\begin{array}{c} a^* \\ \downarrow \\ \boxed{\theta_{a^*}} \\ \downarrow \\ a^* \end{array} = \begin{array}{c} a^* \\ \downarrow \\ \boxed{\theta_a} \\ \uparrow \\ a^* \end{array} = \begin{array}{c} a^* \\ \uparrow \\ \boxed{\theta_a} \\ \downarrow \\ a^* \end{array} = q^{-a\beta + a(a-1)} \begin{array}{c} a^* \\ \downarrow \\ a^* \end{array}. \quad (6.32)$$

More generally for $\mathbf{a} \in \mathbf{Sp}(\beta)$ define $\theta_{\mathbf{a}}$ recursively:

$$\theta_{\mathbf{a} \otimes \mathbf{a}'} = c_{\mathbf{a}', \mathbf{a}} c_{\mathbf{a}, \mathbf{a}'} (\theta_{\mathbf{a}} \otimes \theta_{\mathbf{a}'}) = (\theta_{\mathbf{a}'} \otimes \theta_{\mathbf{a}}) c_{\mathbf{a}', \mathbf{a}} c_{\mathbf{a}, \mathbf{a}'}. \quad (6.33)$$

The equality between the two expressions follows by the naturality of the braiding. Again using the naturality of the braiding it is easy to show that this gives a definition of $\theta_{\mathbf{a}}$ independent of the order in which we perform the recursion.

Lemma 6.12. *The family of maps $\{\theta_{\mathbf{a}} \mid \mathbf{a} \in \mathbf{Sp}(\beta)\}$ is natural.*

Proof. Let $\varphi: \mathbf{a} \rightarrow \mathbf{a}'$. We need to show that $\varphi\theta_{\mathbf{a}} = \theta_{\mathbf{a}'}\varphi$. Again, it is sufficient to consider generating morphisms of $\mathbf{Sp}(\beta)$. First, consider the case $\varphi = \mathbf{E}^{(r)}: a \otimes b \mapsto a + r \otimes b - r$ for $a, b \in \mathbb{Z}_{\geq 0}$. By (5.5d) it is enough to consider the case $r = 1$. We have:

$$\begin{aligned}
& \begin{array}{c} a+1 \quad b-1 \\ \uparrow \quad \uparrow \\ \diagdown \quad \diagup \\ \boxed{\theta_{a \otimes b}} \\ \downarrow \quad \downarrow \\ a \quad b \end{array} = \begin{array}{c} a+1 \quad b-1 \\ \uparrow \quad \uparrow \\ \diagdown \quad \diagup \\ \boxed{\theta_a} \quad \boxed{\theta_b} \\ \downarrow \quad \downarrow \\ a \quad b \end{array} = q^{-(a+b)\beta + a(a-1) + b(b-1)} \begin{array}{c} a+1 \quad b-1 \\ \uparrow \quad \uparrow \\ \diagdown \quad \diagup \\ \text{twist} \\ \downarrow \quad \downarrow \\ a \quad b \end{array} \\
& = q^{-(a+b)\beta + a(a-1) + b(b-1) + 2(a-b) + 2} \begin{array}{c} a+1 \quad b-1 \\ \uparrow \quad \uparrow \\ \text{twist} \\ \diagdown \quad \diagup \\ \downarrow \quad \downarrow \\ a \quad b \end{array} = \begin{array}{c} a+1 \quad b-1 \\ \uparrow \quad \uparrow \\ \boxed{\theta_{a+1}} \quad \boxed{\theta_{b-1}} \\ \downarrow \quad \downarrow \\ a \quad b \end{array} = \begin{array}{c} a+1 \quad b-1 \\ \uparrow \quad \uparrow \\ \boxed{\theta_{a+1 \otimes b-1}} \\ \downarrow \quad \downarrow \\ a \quad b \end{array}. \quad (6.34)
\end{aligned}$$

Here, the first equality is a consequence of the definition (6.33) of the twist on a tensor product, the second one is the definition (6.32) of the twist on a single strand, the third one is a consequence of (5.21) and (5.22), and the last two equalities are symmetric to the two first ones. A similar computation yields the same result for $\mathbf{F}^{(r)}: a \otimes b \rightarrow a - r \otimes b - r$ for $a, b \in \mathbb{Z}_{\geq 0}$.

Second, we need to consider the case in which $\varphi: \emptyset \rightarrow a \otimes a^*$ is the duality map. By writing $\text{id}_a = \pi_a \otimes \iota_a$ (see lemma 5.6), since we already showed that the twist commutes with π_a and ι_a , it is enough to consider the case $a = 1$.

We have

$$\begin{array}{c} 1 \quad 1^* \\ \uparrow \quad \downarrow \\ \boxed{\theta_{1 \otimes 1^*}} \\ \downarrow \end{array} = \begin{array}{c} 1 \quad 1^* \\ \downarrow \quad \uparrow \\ \boxed{\theta_1} \quad \boxed{\theta_{1^*}} \\ \downarrow \quad \uparrow \\ \downarrow \end{array} \quad (6.35)$$

and the claim follows since

$$\begin{array}{c} 1 \\ \downarrow \\ \downarrow \quad \uparrow \\ \uparrow \\ 1 \end{array} = q^\beta \begin{array}{c} 1 \\ \downarrow \\ \downarrow \\ \uparrow \\ 1 \end{array} = \begin{array}{c} 1 \\ \downarrow \\ \boxed{\theta_1^{-1}} \\ \uparrow \\ 1 \end{array}. \quad (6.36) \quad \square$$

Altogether we get:

Theorem 6.13. *The category $\mathbf{Sp}(\beta)$ is ribbon.*

The technical computation below will be useful in the proof of Proposition 6.15.

Lemma 6.14. *We have:*

$$\begin{array}{c} a \\ \downarrow \\ \downarrow \quad \uparrow \\ \uparrow \\ a \end{array} = q^{-a\beta + a(a-1)} \begin{array}{c} a \\ \downarrow \\ \downarrow \\ \uparrow \\ a \end{array} = \begin{array}{c} a \\ \downarrow \\ \boxed{\theta_a} \\ \uparrow \\ a \end{array}, \quad (6.37)$$

$$\begin{array}{c} a \\ \downarrow \\ \downarrow \quad \uparrow \\ \uparrow \\ a \end{array} = q^{a\beta - a(a-1)} \begin{array}{c} a \\ \downarrow \\ \downarrow \\ \uparrow \\ a \end{array} = \begin{array}{c} a \\ \downarrow \\ \boxed{\theta_a^{-1}} \\ \uparrow \\ a \end{array}. \quad (6.38)$$

Proof. We check only the first equation, the second being similar. Using the definition of the braiding, it is easy to deduce:

$$\begin{array}{c} a \\ \downarrow \\ \downarrow \quad \uparrow \\ \uparrow \\ a \end{array} = q^{-a} \sum_{s=0}^a (-q)^s \begin{bmatrix} \beta - s \\ a \end{bmatrix} \begin{bmatrix} a \\ s \end{bmatrix} \begin{array}{c} a \\ \downarrow \\ \downarrow \\ \uparrow \\ a \end{array}. \quad (6.39)$$

First, let us notice that for any $k \in \mathbb{N}$ we have

$$\begin{aligned}
\sum_{s=0}^a (-q)^s \begin{bmatrix} a \\ s \end{bmatrix} \begin{bmatrix} \beta - s - k \\ a \end{bmatrix} &= \sum_{s=0}^a (-q)^s \begin{bmatrix} \beta - s - k \\ a \end{bmatrix} \left(q^{a-s} \begin{bmatrix} a-1 \\ s-1 \end{bmatrix} + q^{-s} \begin{bmatrix} a-1 \\ s \end{bmatrix} \right) \\
&= \sum_{s=0}^{a-1} (-1)^s \begin{bmatrix} a-1 \\ s \end{bmatrix} \left(\begin{bmatrix} \beta - s - k \\ a \end{bmatrix} - q^a \begin{bmatrix} \beta - s - k - 1 \\ a \end{bmatrix} \right) \\
&= q^{-\beta+k+a} \sum_{s=0}^{a-1} (-q)^s \begin{bmatrix} a-1 \\ s \end{bmatrix} \begin{bmatrix} \beta - s - k - 1 \\ a-1 \end{bmatrix}. \quad (6.40)
\end{aligned}$$

Applying this formula recursively we can compute the r.h.s. of (6.39)

$$\begin{aligned}
q^{-a} \sum_{s=0}^a (-q)^s \begin{bmatrix} \beta - s \\ a \end{bmatrix} \begin{bmatrix} a \\ s \end{bmatrix} &= q^{-a} q^{-\beta+a} \sum_{s=0}^{a-1} (-q)^s \begin{bmatrix} a-1 \\ s \end{bmatrix} \begin{bmatrix} \beta - s - 1 \\ a-1 \end{bmatrix} \quad (6.41) \\
&= q^{-a} q^{-\beta+a} q^{-\beta+(a-1)+1} \dots q^{-\beta+1+(a-1)} \sum_{s=0}^0 (-q)^s \begin{bmatrix} 0 \\ s \end{bmatrix} \begin{bmatrix} \beta - s - a \\ 0 \end{bmatrix} \\
&= q^{-a\beta+a(a-1)},
\end{aligned}$$

whence the claim. \square

The category $\mathbf{Sp}(d)$ allows us to recover the Reshetikhin-Turaev invariant:

Proposition 6.15. *There is an obvious braided functor $\mathcal{Q}_\beta: \mathbf{Tangles}_q^{\text{lab}} \rightarrow \mathbf{Sp}(\beta)$. For $\beta = d$, the composition $\mathcal{G}_{m|n} \circ \mathcal{Q}_d$ gives the Reshetikhin-Turaev functor $\mathcal{RT}_{m|n}$.*

Proof. One has to check that this functor is well-defined, i.e. that the relations of $\mathbf{Tangles}$ are satisfied in the image of \mathcal{Q}_β . We refer to (Ohtsuki, 2002, Figure 3.6) for the explicit relations of $\mathbf{Tangles}^{\text{lab}}$, which we now check. Relations (Ohtsuki, 2002, (3.2) and (3.3)) about height changes and far away commutation are clear, the duality relation (Ohtsuki, 2002, (3.10)) is (6.3) and (Ohtsuki, 2002, (3.11)) is implied by (6.4) and (6.5). The framed Reidemeister I move described by Relation (Ohtsuki, 2002, (3.16)) is implied by lemma 6.14. Reidemeister moves II and III described in relations (Ohtsuki, 2002, (3.13) and (3.15)) follow since \mathbf{Sp}^+ is braided, and the version of the Reidemeister II move from relation (Ohtsuki, 2002, (3.14)) is (6.23). The last claim is clear. \square

Equivalence of categories

We constructed a full functor $\mathcal{G}: \mathbf{Sp}(d) \rightarrow \mathbf{Rep}_{m|n}$. A natural problem is to ask for additional relations on $\mathbf{Sp}(d)$ such that this functor becomes an equivalence of categories. The following result tells us that it is enough to look for relations on \mathbf{Sp}^+ :

Proposition 6.16. *Suppose that S is a set of relations on \mathbf{Sp}^+ such that the functor $\mathcal{G}_{m|n}^+$ descends to an equivalence of categories $\mathbf{Sp}^+/S \rightarrow \mathbf{Rep}_{m|n}^+$. Then the functor $\mathcal{G}_{m|n}$ restricts to an equivalence of categories $\mathbf{Sp}(d)/S \rightarrow \mathbf{Rep}_{m|n}$.*

Proof. We already know that the functor $\mathcal{G}_{m|n}$ is essentially surjective and full. Hence it is enough to show that it is faithful. This follows by comparing dimensions of homomorphism spaces. Let $\mathbf{a}, \mathbf{b} \in \mathbf{Sp}(d)/S$ be two objects: we want to show that $\dim \mathrm{Hom}_{\mathbf{Sp}(d)/S}(\mathbf{a}, \mathbf{b}) = \dim \mathrm{Hom}_{\mathbf{Rep}_{m|n}}(\bigwedge^{\mathbf{a}}, \bigwedge^{\mathbf{b}})$. We proceed by induction on the total number D of duals in the tensor product expansion of \mathbf{a} and \mathbf{b} . If $D = 0$ then the claim holds by hypothesis, since \mathbf{a}, \mathbf{b} are objects of \mathbf{Sp}^+ . Hence suppose $D > 0$. Without loss of generality, suppose that \mathbf{a} contains a dual object. Since both the categories $\mathbf{Sp}^+(d)/S$ and $\mathbf{Rep}_{m|n}$ are braided, we can suppose without loss of generality that the dual object appears at the last place in the tensor product decomposition of \mathbf{a} , that is $\mathbf{a} = \mathbf{a}' \otimes a_r^*$. Then we have

$$\begin{aligned} \dim \mathrm{Hom}_{\mathbf{Sp}(d)/S}(\mathbf{a}' \otimes a_r^*, \mathbf{b}) &= \dim \mathrm{Hom}_{\mathbf{Sp}(d)/S}(\mathbf{a}', \mathbf{b} \otimes a_r) \\ &= \dim \mathrm{Hom}_{\mathbf{Rep}_{m|n}}(\bigwedge_q^{\mathbf{a}'}, \bigwedge_q^{\mathbf{b}} \otimes \bigwedge_q^{a_r}) = \dim \mathrm{Hom}_{\mathbf{Rep}_{m|n}}(\bigwedge_q^{\mathbf{a}'} \otimes \bigwedge_q^{a_r^*}, \bigwedge_q^{\mathbf{b}}), \end{aligned} \tag{6.42}$$

where the middle equality follows by induction, and we are done. \square

In particular, let us define the category $\mathbf{Sp}(d)_{\leq m}$ to be the quotient of the category $\mathbf{Sp}(d)$ modulo objects $a > m$ (compare with definition 5.3). Then we have the following corollary:

Corollary 6.17. *The functor $\mathcal{G}_{m|0}$ descends to an equivalence of categories*

$$\mathcal{G}_{m|0}: \mathbf{Sp}(d)_{\leq m} \xrightarrow{\sim} \mathbf{Rep}_{m|0}. \tag{6.43}$$

Similarly, combining proposition 6.16 with the main result from Grant (2016), we obtain:

Corollary 6.18. *The category $\text{Rep}_{1|1}$ is equivalent to the quotient of $\text{Sp}(0)$ modulo the relation*

$$[a+1][a][b][b-1] \begin{array}{c} \uparrow \quad \uparrow \\ a \quad b \end{array} - [2][a+1][b-1] \begin{array}{c} \uparrow \quad \uparrow \\ \swarrow \quad \searrow \\ a \quad b \end{array} + [2]^2 \begin{array}{c} \uparrow \quad \uparrow \\ \swarrow \quad \searrow \\ a \quad b \end{array} = 0 \quad (6.44)$$

for all $a, b \in \mathbb{Z}_{\geq 0}$.

REMARK 6.19. In principle, it is possible to describe a full set of relations for any m, n . Namely, it follows by Schur-Weyl duality (proposition 3.1) that the kernel of the action of the Hecke algebra is generated by the Young symmetrizer Y_b corresponding to the partition b given by one box of size $(n+1) \times (m+1)$. A standard argument (see the proof of (Sartori, 2014, Theorem 3.3.12)) shows that Y_b , considered as a morphism in Sp^+ , generates the kernel of the functor $\mathcal{G}_{m|n}^+$. By proposition 6.16 it follows that Y_b generates the kernel of $\mathcal{G}_{m|n}: \text{Sp}(d) \rightarrow \text{Rep}_{m|n}$. Anyway, although explicit, the relation Y_b is not handy (already in the case of $\mathfrak{gl}_{1|1}$ it involves four strands, while relation (6.44) above only involves two), and one would like to find nicer relations, involving a minimal number of strands.

More relations in $\text{Sp}(\beta)$

We compute in this subsection some further relations which are implied from the defining relations of the categories Sp and $\text{Sp}(\beta)$.

Lemma 6.20. *In Sp we have*

$$\begin{array}{c} a+r+s \quad (b+r+s)^* \\ \begin{array}{c} \uparrow \quad \uparrow \\ \swarrow \quad \searrow \\ a \quad b^* \end{array} \end{array} = \begin{bmatrix} r+s \\ r \end{bmatrix} \begin{array}{c} \uparrow \quad \uparrow \\ \swarrow \quad \searrow \\ a \quad b^* \end{array} \quad \text{and} \quad \begin{array}{c} a+r+s \quad (b+r+s)^* \quad (a-r-s)^* \quad b-r-s \\ \begin{array}{c} \uparrow \quad \uparrow \\ \swarrow \quad \searrow \\ a^* \quad b \end{array} \end{array} = \begin{bmatrix} r+s \\ r \end{bmatrix} \begin{array}{c} \uparrow \quad \uparrow \\ \swarrow \quad \searrow \\ a^* \quad b \end{array} . \quad (6.45)$$

Proof. The proof is similar in both cases, so we only detail the first one. We have:

$$\begin{array}{c} \begin{array}{|c|} \hline s \\ \hline \end{array} \\ \begin{array}{|c|} \hline r \\ \hline \end{array} \\ a \quad b^* \end{array} = \begin{array}{c} \begin{array}{|c|} \hline s \\ \hline \end{array} \\ \begin{array}{|c|} \hline r \\ \hline \end{array} \\ a \quad b^* \end{array} = \begin{array}{c} \begin{array}{|c|} \hline s \\ \hline \end{array} \\ \begin{array}{|c|} \hline r+s \\ \hline \end{array} \\ a \quad b^* \end{array} = \begin{bmatrix} r+s \\ r \end{bmatrix} \begin{array}{c} \begin{array}{|c|} \hline s \\ \hline \end{array} \\ \begin{array}{|c|} \hline r+s \\ \hline \end{array} \\ a \quad b^* \end{array}. \quad (6.46)$$

The first equality follows from the relation

$$\begin{array}{c} a+r+s \\ \begin{array}{|c|} \hline s \\ \hline \end{array} \\ \begin{array}{|c|} \hline r \\ \hline \end{array} \\ a \quad r \quad s \end{array} = \begin{array}{c} a+r+s \\ \begin{array}{|c|} \hline r+s \\ \hline \end{array} \\ \begin{array}{|c|} \hline s \\ \hline \end{array} \\ a \quad r \quad s \end{array}, \quad (6.47)$$

which is the graphical version of a higher order Serre relation, and follows from the relations defining Sp^+ . \square

Lemma 6.21. *In Sp we have*

$$\begin{array}{c} a+2 \quad b+1 \quad (c+1)^* \\ \begin{array}{|c|} \hline 2 \\ \hline \end{array} \\ a \quad b \quad c^* \end{array} - \begin{array}{c} a+2 \quad b+1 \quad (c+1)^* \\ \begin{array}{|c|} \hline 1 \\ \hline \end{array} \\ a \quad b \quad c^* \end{array} + \begin{array}{c} a+2 \quad b+1 \quad (c+1)^* \\ \begin{array}{|c|} \hline 2 \\ \hline \end{array} \\ a \quad b \quad c^* \end{array} = 0 \quad (6.48)$$

as well as

$$\begin{array}{c} (a-1)^* \quad b+1 \quad c-2 \\ \begin{array}{|c|} \hline 1 \\ \hline \end{array} \\ a^* \quad b \quad c \end{array} - \begin{array}{c} (a-1)^* \quad b+1 \quad c-2 \\ \begin{array}{|c|} \hline 1 \\ \hline \end{array} \\ a^* \quad b \quad c \end{array} + \begin{array}{c} (a-1)^* \quad b+1 \quad c-2 \\ \begin{array}{|c|} \hline 1 \\ \hline \end{array} \\ a^* \quad b \quad c \end{array} = 0. \quad (6.49)$$

Proof. As the two formulas are similar, we only detail the first one. We can rewrite the first summand of (6.48) as

$$\begin{array}{c} \text{Diagram 1} \end{array} = \begin{array}{c} \text{Diagram 2} \end{array} = \begin{array}{c} \text{Diagram 3} \end{array}, \quad (6.50)$$

the second summand as

$$\begin{array}{c}
\begin{array}{ccc}
\begin{array}{c} \text{Diagram 1: Three vertical lines } a, b, c. \text{ Line } a \text{ has two upward arrows. Line } b \text{ has two upward arrows. Line } c \text{ has two upward arrows. A horizontal line connects line } a \text{ to line } b \text{ at the top, labeled } 1. \text{ Another horizontal line connects line } a \text{ to line } b \text{ at the bottom, labeled } 1. \end{array}
& = &
\begin{array}{c} \text{Diagram 2: Similar to Diagram 1, but the horizontal lines are curved. The top curve is labeled } 1. \text{ The bottom curve is labeled } 1. \end{array}
\\
a & b & c
\end{array}
\end{array}
\quad (6.51)$$

and the third summand as

$$\begin{array}{c}
\text{Diagram 1: Three vertical lines labeled } a, b, c \text{ at the bottom. Line } a \text{ has an upward arrow. Line } b \text{ has a downward arrow. Line } c \text{ has a downward arrow. A line segment labeled '2' connects the top of line } a \text{ to the top of line } b. A curved line labeled '1' connects the top of line } b \text{ to the top of line } c.} \\
= \\
\begin{array}{c}
\text{Diagram 2: Three vertical lines labeled } a, b, c \text{ at the bottom. Line } a \text{ has an upward arrow. Line } b \text{ has a downward arrow. Line } c \text{ has a downward arrow. A line segment labeled '2' connects the top of line } a \text{ to the top of line } b. A dashed vertical line is between } b \text{ and } c. A curved line labeled '1' connects the top of line } b \text{ to the top of line } c, passing through the dashed line.}
\end{array}
\end{array} \quad (6.52)$$

Now the claim follows by the Serre relation

$$\begin{array}{c} \text{Diagram 1} \end{array} - \begin{array}{c} \text{Diagram 2} \end{array} + \begin{array}{c} \text{Diagram 3} \end{array} = 0. \quad (6.53)$$

Lemma 6.22. *In \mathbf{Sp} we have*

$$\begin{array}{c} a+1 \quad b+1 \quad (c+2)^* \\ \diagdown \quad \diagup \\ \text{---} \quad \text{---} \quad \text{---} \\ \diagup \quad \diagdown \\ a \quad b \quad c^* \end{array} - \begin{array}{c} a+1 \quad b+1 \quad (c+2)^* \\ \diagdown \quad \diagup \\ \text{---} \quad \text{---} \quad \text{---} \\ \diagup \quad \diagdown \\ a \quad b \quad c^* \end{array} + \begin{array}{c} a+1 \quad b+1 \quad (c+2)^* \\ \diagdown \quad \diagup \\ \text{---} \quad \text{---} \quad \text{---} \\ \diagup \quad \diagdown \\ a \quad b \quad c^* \end{array} = 0 \quad (6.54)$$

as well as

$$\begin{array}{c} (a-2)^* \quad b+1 \quad c-1 \\ \begin{array}{|c|} \hline \text{Diagram 1} \\ \hline \end{array} \end{array} - \begin{array}{c} (a-2)^* \quad b+1 \quad c-1 \\ \begin{array}{|c|} \hline \text{Diagram 2} \\ \hline \end{array} \end{array} + \begin{array}{c} (a-2)^* \quad b+1 \quad c-1 \\ \begin{array}{|c|} \hline \text{Diagram 3} \\ \hline \end{array} \end{array} = 0. \quad (6.55)$$

Proof. The proof is similar to the proof of Lemma 6.21, and we omit it. \square

Lemma 6.23. *In Sp we have*

$$\begin{array}{c} (a-1)^* \quad b+1 \quad (c+2)^* \\ \begin{array}{|c|} \hline \text{Diagram 1} \\ \hline \end{array} \end{array} - \begin{array}{c} (a-1)^* \quad b+1 \quad (c+2)^* \\ \begin{array}{|c|} \hline \text{Diagram 2} \\ \hline \end{array} \end{array} + \begin{array}{c} (a-1)^* \quad b+1 \quad (c+2)^* \\ \begin{array}{|c|} \hline \text{Diagram 3} \\ \hline \end{array} \end{array} = 0 \quad (6.56)$$

as well as

$$\begin{array}{c} (a-2)^* \quad b-1 \quad (c+1)^* \\ \begin{array}{|c|} \hline \text{Diagram 1} \\ \hline \end{array} \end{array} - \begin{array}{c} (a-2)^* \quad b-1 \quad (c+1)^* \\ \begin{array}{|c|} \hline \text{Diagram 2} \\ \hline \end{array} \end{array} + \begin{array}{c} (a-2)^* \quad b-1 \quad (c+1)^* \\ \begin{array}{|c|} \hline \text{Diagram 3} \\ \hline \end{array} \end{array} = 0. \quad (6.57)$$

Proof. We again omit the proof. \square

REMARK 6.24. By swapping all orientations of the uprights, we obtain (with analogous proofs) six more relations from lemmas 6.21, 6.22, and 6.23. Moreover, by replacing all morphisms E_i with F_i we obtain, again with analogous proofs, other twelve relations. For space reasons, we do not write all of them.

Lemma 6.25. *In $\text{Sp}(\beta)$ we have*

$$\begin{array}{c} a \quad b^* \\ \begin{array}{|c|} \hline \text{Diagram 1} \\ \hline \end{array} \end{array} - \begin{array}{c} a \quad b^* \\ \begin{array}{|c|} \hline \text{Diagram 2} \\ \hline \end{array} \end{array} = [a + b - \beta] \begin{array}{c} a \quad b^* \\ \begin{array}{|c|} \hline \text{Diagram 3} \\ \hline \end{array} \end{array} \quad (6.58)$$

and

$$\begin{array}{c} a^* \quad b \\ \downarrow \quad \downarrow \\ \text{1} \quad \leftarrow \\ \downarrow \quad \downarrow \\ \text{1} \quad \rightarrow \\ \downarrow \quad \downarrow \\ a^* \quad b \end{array} - \begin{array}{c} a^* \quad b \\ \downarrow \quad \downarrow \\ \text{1} \quad \leftarrow \\ \downarrow \quad \downarrow \\ \text{1} \quad \rightarrow \\ \downarrow \quad \downarrow \\ a^* \quad b \end{array} = [\beta - a - b] \begin{array}{c} a^* \quad b \\ \downarrow \quad \downarrow \\ \downarrow \quad \downarrow \\ \downarrow \quad \downarrow \\ \downarrow \quad \downarrow \\ a^* \quad b \end{array}. \quad (6.59)$$

Proof. We check only the first relation, the second one being similar. We have

$$\begin{aligned}
& \text{Diagram 1} = \text{Diagram 2} \\
& = \text{Diagram 3} - [a-1] \text{Diagram 4} - [b-1] \text{Diagram 5} + [a-1][b-1] \text{Diagram 6} \\
& = \text{Diagram 7} - [a-1] \text{Diagram 8} - [b-1] \text{Diagram 9} + [a-1][b-1][\beta] \text{Diagram 10} \\
& = \text{Diagram 11} - [2-\beta] \text{Diagram 12} + [a-1][1-\beta] \text{Diagram 13} \\
& \quad + [b-1][1-\beta] \text{Diagram 14} + [a-1][b-1][\beta] \text{Diagram 15},
\end{aligned}
\tag{6.60}$$

where we used relations (6.15a), (6.15c) and (6.15d). The claim now follows

since

$$\begin{aligned}
& [2 - \beta][a][b] - [a - 1][1 - \beta][b] - [b - 1][1 - \beta][a] - [a - 1][b - 1][\beta] \quad (6.61) \\
&= [b]([2 - \beta][a] - [1 - \beta][a - 1]) + [b - 1]([\beta - 1][a] - [\beta][a - 1]) \\
&= [b][1 - \beta + a] + [b - 1][\beta - a] = [b][-\beta + a + 1] - [b - 1][-\beta + a] \\
&= [a + b - \beta].
\end{aligned}$$

□

7. Mixed skew Howe duality and the idempotent quantum group

We now introduce quotients of the idempotent version of $U_q(\mathfrak{gl}_k)$, depending on the choice of a sequence of signs, which are the natural targets for $\mathfrak{gl}_{m|n}$ link invariants. We constructed a first example of this in [Queffelec and Sartori \(2018\)](#), here we generalize it further.

Doubled Schur algebras

Let us fix a sequence $\boldsymbol{\eta} = (\eta_1, \dots, \eta_k) \in \{\pm 1\}^k$ of signs. Let $\mathbf{P}_{\boldsymbol{\eta}}$ be the set of sequences $\lambda = (\lambda_1, \dots, \lambda_{k+l})$ with $\lambda_i \in \mathbb{Z}_{\geq 0}$ if $\eta_i = 1$ and $\lambda_i \in \beta - \mathbb{Z}_{\geq 0}$ if $\eta_i = -1$. We let also $\alpha_i = (0, \dots, 0, 1, -1, 0, \dots, 0)$, the entry 1 being at position i .

Definition 7.1. We define $\dot{\mathbf{U}}_q(\mathfrak{gl}_{\boldsymbol{\eta}})_{\beta}$ to be the additive $\mathbb{C}(q, q^{\beta})$ -linear category with:

- **objects:** formal direct sums of $\mathbf{1}_{\lambda}$ for $\lambda \in \mathbf{P}_{\boldsymbol{\eta}}$;
- **morphisms:** generated by identity endomorphisms $\mathbf{1}_{\lambda}$ in $\text{Hom}(\mathbf{1}_{\lambda}, \mathbf{1}_{\lambda})$, and morphisms $E_i \mathbf{1}_{\lambda} = \mathbf{1}_{\lambda + \alpha_i} E_i \in \text{Hom}(\mathbf{1}_{\lambda}, \mathbf{1}_{\lambda + \alpha_i})$, $F_i \mathbf{1}_{\lambda} = \mathbf{1}_{\lambda - \alpha_i} F_i \in \text{Hom}(\mathbf{1}_{\lambda}, \mathbf{1}_{\lambda - \alpha_i})$. We will often abbreviate by omitting some of the symbols $\mathbf{1}_{\lambda}$. The morphisms are subject to the following relations:

$$(DS1) \quad [E_i, F_j] \mathbf{1}_{\lambda} = \delta_{i,j} [\lambda_i - \lambda_{i+1}] \mathbf{1}_{\lambda},$$

$$\begin{aligned}
(DS2) \quad E_i^2 E_j \mathbf{1}_{\lambda} - (q + q^{-1}) E_i E_j E_i \mathbf{1}_{\lambda} + E_j E_i^2 \mathbf{1}_{\lambda} &= 0 \quad \text{if } j = i \pm 1, \\
F_i^2 F_j \mathbf{1}_{\lambda} - (q + q^{-1}) F_i F_j F_i \mathbf{1}_{\lambda} + F_j F_i^2 \mathbf{1}_{\lambda} &= 0 \quad \text{if } j = i \pm 1,
\end{aligned}$$

$$(DS3) \quad E_i E_j \mathbf{1}_{\lambda} = E_j E_i \mathbf{1}_{\lambda} \text{ and } F_i F_j \mathbf{1}_{\lambda} = F_j F_i \mathbf{1}_{\lambda} \text{ if } |i - j| > 1.$$

We indifferently use: $\mathbf{1}_{\lambda+\alpha_i}E_i\mathbf{1}_\lambda = E_i\mathbf{1}_\lambda = \mathbf{1}_{\lambda+\alpha_i}E_i$, since knowing the source or the target of the 1-morphism is enough to determine the other one. We use the following convention: if the symbol $\mathbf{1}_\lambda$ appears and $\lambda \notin \mathbf{P}_\eta$, then $\mathbf{1}_\lambda = 0$.

For the sequence of signs $\boldsymbol{\eta}_{r,s}$ (see (2.10)), the category $\dot{\mathbf{U}}_q(\mathfrak{gl}_{r+s})_{\boldsymbol{\eta}_{r,s}}$ was already introduced in Queffelec and Sartori (2018). In the special case $s = 0$ the category $\dot{\mathbf{U}}_q(\mathfrak{gl}_r)_{\boldsymbol{\eta}_{r,0}}$ was already defined in Cautis et al. (2014), and denoted by $\dot{\mathbf{U}}_q^{\geq 0}(\mathfrak{gl}_r)$.

Let us define a functor $\Phi_\eta: \dot{\mathbf{U}}_q(\mathfrak{gl}_\eta)_\beta \rightarrow \mathbf{Sp}(\beta)$ by setting on the generators

$$\mathbf{1}_\lambda \mapsto \mathbf{a}, \quad (7.1)$$

$$E_i\mathbf{1}_\lambda \mapsto (\mathrm{id}^{\otimes(i-1)} \otimes E \otimes \mathrm{id}^{\otimes(k-i-1)})\mathbf{1}_\mathbf{a}, \quad (7.2)$$

$$F_i\mathbf{1}_\lambda \mapsto (\mathrm{id}^{\otimes(i-1)} \otimes F \otimes \mathrm{id}^{\otimes(k-i-1)})\mathbf{1}_\mathbf{a}, \quad (7.3)$$

where \mathbf{a} is determined by

$$a_i = \begin{cases} \lambda_i & \text{if } \eta_i = +1, \\ (-\lambda_i + \beta)^* & \text{if } \eta_i = -1, \end{cases} \quad (7.4)$$

Notice that depending on the index i , the morphisms E and F in the equations above can be either (5.4), (6.7), (6.13) or (6.14).

Proposition 7.2. *The functor Φ_η is well-defined.*

Proof. We shall check that the relations defining $\dot{\mathbf{U}}_q(\mathfrak{gl}_\eta)_\beta$ are satisfied by the images of the E_i 's and F_i 's. The commuting relation (DS3) is straightforward. Also (DS1) is straightforward in case $|i - j| > 1$. Since the other relations are local, we only need to consider the case in which two neighbor indices $i, i + 1$ are involved such that $\eta_i \neq \eta_{i+1}$ (otherwise the relations hold because they hold in \mathbf{Sp}^+ by definition).

Relation (DS1) for $i = j \pm 1$ is very easy to check, and follows from relation (5.5c). In the case $i = j$, the relation (DS1) is proved in lemma 6.25. The Serre relations (DS2) are proven in lemmas 6.21, 6.22 and 6.23 (cf. also remark 6.24). \square

We denote by $\mathbf{Sp}(\beta)_\eta$ the additive full subcategory of $\mathbf{Sp}(\beta)$ with objects (a_1, \dots, a_k) such that $a_i \in \mathbb{Z}_{\geq 0}$ if $\eta_i = 1$ and $a_i = b_i^*$ for some $b_i \in \mathbb{Z}_{\geq 0}$ if

$\eta_i = -1$. Then Φ_η has values in $\mathbf{Sp}(\beta)_\eta$ and, indeed, we will prove that this is an equivalence of categories. But first, we want to investigate what happens when we swap two signs of the sequence η .

Let also $1 \leq i \leq k$ be an index with $\eta_i = 1$, $\eta_{i+1} = -1$ and let s be the simple transposition $(i, i+1)$. Denote by $\eta' = s\eta$ the swapped sequence. Using the braiding in $\mathbf{Sp}(\beta)$ it is immediate to show that $\mathbf{Sp}(\beta)_\eta$ and $\mathbf{Sp}(\beta)_{\eta'}$ are equivalent. Pulling back this braiding² we can construct an equivalence $\dot{\mathbf{U}}_q(\mathfrak{gl}_\eta)_\beta \rightarrow \dot{\mathbf{U}}_q(\mathfrak{gl}_{\eta'})_\beta$, which is explicitly given by some analogue of Lusztig's symmetries:

Lemma 7.3. *We have an isomorphism of categories $\mathsf{T}_i: \dot{\mathbf{U}}_q(\mathfrak{gl}_\eta)_\beta \rightarrow \dot{\mathbf{U}}_q(\mathfrak{gl}_{\eta'})_\beta$ given by the map*

$$\begin{aligned}
\mathbf{1}_\lambda &\mapsto \mathbf{1}_{s\lambda}, \\
E_{i-1}\mathbf{1}_\lambda &\mapsto (-1)^{\lambda_{i+1}+1}(qE_{i-1}E_i - E_iE_{i-1})\mathbf{1}_{s\lambda}, \\
F_{i-1}\mathbf{1}_\lambda &\mapsto (-1)^{\lambda_{i+1}+1}(q^{-1}F_iF_{i-1} - F_{i-1}F_i)\mathbf{1}_{s\lambda}, \\
E_{i+1}\mathbf{1}_\lambda &\mapsto (-1)^{\lambda_i+1}(qE_{i+1}E_i - E_iE_{i+1})\mathbf{1}_{s\lambda}, \\
F_{i+1}\mathbf{1}_\lambda &\mapsto (-1)^{\lambda_{i+1}+1}(q^{-1}F_iF_{i+1} - F_{i+1}F_i)\mathbf{1}_{s\lambda}, \\
E_i\mathbf{1}_\lambda &\mapsto (-q)^{\lambda_i - \lambda_{i+1}}F_i\mathbf{1}_{s\lambda}, \\
F_i\mathbf{1}_\lambda &\mapsto (-q)^{-\lambda_i + \lambda_{i+1} + 2}E_i\mathbf{1}_{s\lambda}.
\end{aligned} \tag{7.5}$$

Moreover, we have a commutative diagram

$$\begin{array}{ccc}
\dot{\mathbf{U}}_q(\mathfrak{gl}_\eta)_\beta & \xrightarrow{\Phi_\eta} & \mathbf{Sp}(\beta)_\eta \\
\mathsf{T}_i \downarrow & & \downarrow \mathsf{T}_i \\
\dot{\mathbf{U}}_q(\mathfrak{gl}_{\eta'})_\beta & \xrightarrow{\Phi_{\eta'}} & \mathbf{Sp}(\beta)_{\eta'}
\end{array} \tag{7.6}$$

where the right vertical arrow is defined on objects by applying the transposition s and on morphisms by $\mathsf{T}_i(x) = T_i x T_i^{-1}$, where

$$T_i = \mathrm{id}_{(a_1, \dots, a_{i-1})} \otimes c_{a_i, a_{i+1}} \otimes \mathrm{id}_{(a_{i+2}, \dots, a_k)}. \tag{7.7}$$

²More explicitly, the formulas in Lemma 7.3 are obtained by conjugating the images of the generators of $\dot{\mathbf{U}}_q(\mathfrak{gl}_\eta)_\beta$ in $\mathbf{Sp}(\beta)$ by the up/down braidings, and finding equivalent diagrams that describe these elements without using up/down crossings and can be pulled back to $\dot{\mathbf{U}}_q(\mathfrak{gl}_{\eta'})_\beta$.

Proof. The proof is a lengthy but straightforward calculation. One shows explicitly that (7.5) give a well-defined map and that the diagram (7.6) commutes. Formulas analogous to (7.5) define explicitly the inverse of T_i . \square

As a consequence, we get:

Corollary 7.4. *If $\boldsymbol{\eta}, \boldsymbol{\eta}'$ are two sequences of signs with the same number of +1 and -1 then $\dot{\mathsf{U}}_q(\mathfrak{gl}_{\boldsymbol{\eta}})_{\beta}$ and $\dot{\mathsf{U}}_q(\mathfrak{gl}_{\boldsymbol{\eta}'})_{\beta}$ are isomorphic.*

We can now prove:

Proposition 7.5. *The functor $\Phi_{\boldsymbol{\eta}}: \dot{\mathsf{U}}_q(\mathfrak{gl}_{\boldsymbol{\eta}})_{\beta} \rightarrow \mathsf{Sp}(\beta)$ is full.*

Proof. By lemma 7.3, and in particular by (7.6), we can restrict to the case $k = r + s$ and $\boldsymbol{\eta} = \boldsymbol{\eta}_{r,s}$.

Let $\lambda = (1, \dots, 1, \beta - 1, \dots, \beta - 1)$ and $\mathbf{a} = \Phi_{\boldsymbol{\eta}}(\mathbf{1}_{\lambda}) = 1^{\otimes r} \otimes (1^*)^{\otimes s}$, and let us prove that the image of $\Phi_{\boldsymbol{\eta}}$ contains $\mathsf{End}_{\mathsf{Sp}(\beta)}(\mathbf{a})$. Indeed, $\mathsf{End}_{\mathsf{Sp}(\beta)}(\mathbf{a})$ is just the walled Brauer algebra $\mathsf{Br}_{\boldsymbol{\eta}}(\beta)$, which is generated by upward or downward pointing crossings and cup-caps (cf. for example Brundan and Stroppel (2012)). Hence $\mathsf{End}_{\mathsf{Sp}(\beta)}(\mathbf{a})$ is generated by

$$\mathrm{id}^{\otimes i} \otimes c_{1,1} \otimes \mathrm{id}^{\otimes (r+s-i-2)} \quad \text{for } i = 0, \dots, r-2, \quad (7.8)$$

$$\mathrm{id}^{\otimes i} \otimes c_{1^*,1^*} \otimes \mathrm{id}^{\otimes (r+s-i-2)} \quad \text{for } i = r+1, \dots, r+s-2, \quad (7.9)$$

$$\mathrm{id}^{\otimes (r-1)} \otimes (\mathsf{EF} \mathbf{1}_{1 \otimes 1^*}) \otimes \mathrm{id}^{\otimes (s-1)}. \quad (7.10)$$

These elements are clearly in the image of $\dot{\mathsf{U}}_q(\mathfrak{gl}_{\boldsymbol{\eta}_{r,s}})_{\beta}$. Similarly, we can show that $\Phi_{\boldsymbol{\eta}_{r,s}}: \mathsf{Hom}_{\dot{\mathsf{U}}_q(\mathfrak{gl}_{\boldsymbol{\eta}_{r,s}})_{\beta}}(\mathbf{1}_{\lambda}, \mathbf{1}_{\mu}) \rightarrow \mathsf{Hom}_{\mathsf{Sp}(\beta)}(\Phi_{\boldsymbol{\eta}_{r,s}}(\mathbf{1}_{\lambda}), \Phi_{\boldsymbol{\eta}_{r,s}}(\mathbf{1}_{\mu}))$ is surjective in the case $\mathbf{1}_{\lambda} = \mathbf{a}' \otimes \mathbf{a}''$ and $\mathbf{1}_{\mu} = \mathbf{b}' \otimes \mathbf{b}''$, where \mathbf{a}', \mathbf{b}' are tensor products of 1's and 0's and $\mathbf{a}'', \mathbf{b}''$ are tensor products of 1^* 's and 0^* 's.

Let us now consider the case of two general elements $\mathbf{1}_{\lambda}, \mathbf{1}_{\mu} \in \mathsf{P}_{\boldsymbol{\eta}_{r,s}}$, so that $\Phi_{\boldsymbol{\eta}_{r,s}}(\mathbf{1}_{\lambda}) = \mathbf{a} = \mathbf{a}' \otimes \mathbf{a}''$ and $\Phi_{\boldsymbol{\eta}_{r,s}}(\mathbf{1}_{\mu}) = \mathbf{b} = \mathbf{b}' \otimes \mathbf{b}''$ with $\mathbf{a}', \mathbf{b}' \in \mathsf{Sp}^+$ and $\mathbf{a}'', \mathbf{b}''$ duals of objects from Sp^+ , and let us show that $\mathsf{Hom}_{\mathsf{Sp}(\beta)}(\mathbf{a}, \mathbf{b})$ is in the image of $\Phi_{\boldsymbol{\eta}}$. The strategy of the proof is the following: first, we prove that this is true up to enlarging r and s (pictorially, we allow more horizontal space for more strands) and then we show that this enlarging was not necessary.

Let also $r' \geq r, s' \geq s$ and let $j: \dot{\mathbf{U}}_q(\mathfrak{gl}_{\eta_{r,s}})_\beta \hookrightarrow \dot{\mathbf{U}}_q(\mathfrak{gl}_{\eta_{r',s'}})_\beta$ be the inclusion which maps $X_i \mapsto X_{i+r'-r}$ for $X_i = E_i, F_i$. Notice that the following diagram commutes by construction

$$\begin{array}{ccc} \dot{\mathbf{U}}_q(\mathfrak{gl}_{\eta_{r,s}})_\beta & \xrightarrow{\Phi_{\eta_{r,s}}} & \mathrm{Sp}(\beta) \\ j \downarrow & & \downarrow \\ \dot{\mathbf{U}}_q(\mathfrak{gl}_{\eta_{r',s'}})_\beta & \xrightarrow{\Phi_{\eta_{r',s'}}} & \mathrm{Sp}(\beta) \end{array} \quad (7.11)$$

where the vertical map on the right is given by tensoring with $0^{\otimes(r'-r)}$ and $0^{\otimes(s'-s)}$ on the left and on the right, respectively. We claim that there are some r' and s' such that $\mathrm{Hom}_{\mathrm{Sp}(d)}(\mathbf{a}, \mathbf{b})$ is in the image of $\Phi_{\eta_{r',s'}}$. Indeed, we can write any morphism $\varphi: \mathbf{a} \rightarrow \mathbf{b}$ as $(\pi_{\mathbf{b}'} \otimes \pi_{\mathbf{b}''}) \circ \varphi' \circ (\iota_{\mathbf{a}'} \otimes \iota_{\mathbf{a}''})$ for $\varphi' \in \mathrm{Hom}(1^{\otimes|\mathbf{a}'|} \otimes 1^{\otimes|\mathbf{a}''|}, 1^{\otimes|\mathbf{b}'|} \otimes 1^{\otimes|\mathbf{b}''|})$. Choose $r' = \max\{|\mathbf{a}'|, |\mathbf{b}'|\}$ and $s' = \max\{|\mathbf{a}''|, |\mathbf{b}''|\}$. Then, by the first paragraph of the proof, φ' is in the image of $\Phi_{\eta_{r',s'}}$. Since $\pi_{\mathbf{b}'} \otimes \pi_{\mathbf{b}''}$ and $\iota_{\mathbf{a}'} \otimes \iota_{\mathbf{a}''}$ are also in the image of $\Phi_{\eta_{r',s'}}$, the morphism φ is, too.

Choose now $r' \geq r$ and $s' \geq s$ minimal such that $\mathrm{Hom}_{\mathrm{Sp}(\beta)}(\mathbf{a}, \mathbf{b})$ is in the image of $\Phi_{\eta_{r',s'}}$. We want to show by contradiction that $r' = r$ and $s' = s$. Suppose, on the contrary, that $r' > r$ (the case $s' > s$ being analogous). Choose $\varphi \in \mathrm{Hom}_{\mathrm{Sp}(\beta)}(\mathbf{a}, \mathbf{b})$ and pick $x \in \dot{\mathbf{U}}_q(\mathfrak{gl}_{\eta_{r',s'}})_\beta$ with $\varphi = \Phi_{\eta_{r',s'}}(x)$. It follows from the PBW Theorem applied to the quantum group $\mathbf{U}_q(\mathfrak{gl}_{r'+s'})$ that we can write x as linear combination $x = \sum_h \gamma_h x_h$ of monomials x_h in the generators E_i, F_i such that in each x_h the generators F_i appear to the right of the E_i 's. For each x_h we have now three possibilities:

- (i) The generator F_1 appears at least once in x_h , and we can assume that there are no E_1 's on the right of this F_1 . Then $\Phi_{\eta_{r',s'}}(x_h) = 0$, since this F_1 is sent to some $(\mathbf{F} \otimes \mathrm{id}^{r'+s'-2})\mathbf{1}_c$, where $c_1 = 0$. Hence we can remove the term $\gamma_h x_h$ from x without changing the value of $\Phi_{\eta_{r',s'}}(x)$.
- (ii) The generator E_1 appears at least once in x_h , and there are no F_1 's. Similarly as before, by looking at the first entry of the sequences (which are zero both at the bottom and at the top, since $r' > r$) we conclude that this monomial x_h acts by zero. Hence we can remove $\gamma_h x_h$ from x .
- (iii) The generators E_1 and F_1 do not appear in x_h .

So we proved that we do not need E_1 and F_1 , so r' was not minimal, and this is a contradiction. \square

Mixed skew Howe duality

Let us now set $\beta = d = m - n$ and fix an integer N . Let

$$\Xi_{\eta,N} = \{a_1^{\eta_1} \otimes \cdots \otimes a_k^{\eta_k} \mid a_i \in \mathbb{Z}_{\geq 0}, \eta_1 a_1 + \cdots + \eta_k a_k + dp_{\eta} = N\} \quad (7.12)$$

where $p_{\eta} = \#\{i \mid \eta_i = -1\}$, and we use the convention $a^{-1} = a^*$ for $a \in \mathbb{Z}_{\geq 0}$.

Consider the space

$$W = \bigoplus_{\alpha \in \Xi_{\eta,N}} \bigwedge_{\mathbb{C}_q}^{\alpha} \mathbb{C}_q^{m|n}. \quad (7.13)$$

Composing the map Φ_{η} and the functor $\mathcal{G}_{m|n}$ we then have:

Proposition 7.6. *The assignment $x \mapsto \mathcal{G}_{m|n} \circ \Phi_{\eta}(x)$ defines a map*

$$\Psi_{\eta,N}: \dot{\mathbf{U}}_q(\mathfrak{gl}_{\eta})_{\beta} \rightarrow \text{End}_{U_q(\mathfrak{gl}_{m|n})}(W). \quad (7.14)$$

As a direct consequence of proposition 7.5 we obtain:

Theorem 7.7 (Mixed skew Howe duality). *For all $m, n \geq 0$, $N \in \mathbb{Z}$ and for all sequences of signs η the map $\Psi_{\eta,N}$ is surjective. In particular, $\dot{\mathbf{U}}_q(\mathfrak{gl}_{\eta})_d$ acts $U_q(\mathfrak{gl}_{m|n})$ -equivariantly on the $U_q(\mathfrak{gl}_{m|n})$ -module W and generates the full centralizer.*

We expect a double centralizing property to hold, i.e. we expect the action of $U_q(\mathfrak{gl}_{m|n})$ to generate the full centralizer of the $\dot{\mathbf{U}}_q(\mathfrak{gl}_{\eta_{r,s}})_d$ -action.

REMARK 7.8. By the theorem, the space W inherits an action of $\dot{\mathbf{U}}_q(\mathfrak{gl}_{\eta})_{\beta}$. Since the latter is an idempotent version of $U_q(\mathfrak{gl}_k)$, this induces an action of $U_q(\mathfrak{gl}_k)$ on W . In particular, we have commuting actions

$$U_q(\mathfrak{gl}_k) \curvearrowright \bigoplus_{\alpha \in \Xi_{\eta,N}} \bigwedge_{\mathbb{C}_q}^{\alpha} \mathbb{C}_q^{m|n} \curvearrowright U_q(\mathfrak{gl}_{m|n}). \quad (7.15)$$

However, since there are infinitely many \mathfrak{gl}_k weights involved, the projections onto the \mathfrak{gl}_k -weight spaces are not in the image of the map $U_q(\mathfrak{gl}_k) \rightarrow \text{End}_{U_q(\mathfrak{gl}_{m|n})}(W)$, which is therefore not surjective. But the image of $U_q(\mathfrak{gl}_k)$ together with the projections onto the weight spaces generate all intertwiners.

REMARK 7.9. Consider the special case $\boldsymbol{\eta} = \boldsymbol{\eta}_{r,s}$. Then we have commuting actions of $U_q(\mathfrak{gl}_{r+s})$ and $U_q(\mathfrak{gl}_{m|n})$ on

$$\bigoplus_{\substack{a_1 + \dots + a_r - a_{r+1} \\ \dots - a_{r+s} + (m-n)s = N}} \bigwedge_q^{a_1} \mathbb{C}_q^{m|n} \otimes \dots \otimes \bigwedge_q^{a_r} \mathbb{C}_q^{m|n} \otimes \bigwedge_q^{a_{r+1}} (\mathbb{C}_q^{m|n})^* \otimes \dots \otimes \bigwedge_q^{a_{r+s}} (\mathbb{C}_q^{m|n})^*. \quad (7.16)$$

Notice that for $s = 0$ this gives back the usual skew Howe duality. As a $U_q(\mathfrak{gl}_{r+s})$ -module, (7.16) is a weight module and is $U_q(\mathfrak{gl}_r) \otimes U_q(\mathfrak{gl}_s)$ -finite. Moreover, the action of the lower triangular part $U_q(\mathfrak{n}^-)$ is locally finite. In general, however, it is not finitely generated. This is true in the case $n = 1$ (since the dimension of the exterior powers of $\mathbb{C}^{m|1}$ is bounded) and then we can conclude that (7.16) is in the parabolic category $\mathcal{O}^p(U_q(\mathfrak{gl}_{r+s}))$ (however with respect to the negative Borel) corresponding to the parabolic subalgebra $\mathfrak{p} \subseteq \mathfrak{gl}_{r+s}$ with Levi subalgebra $\mathfrak{gl}_r \oplus \mathfrak{gl}_s$.

EXAMPLE 7.10. Let us illustrate the example $m = n = 1$, $r = s = 1$ and $N = 0$. Then we have two commuting actions

$$U_q(\mathfrak{gl}_{1|1}) \curvearrowright \bigoplus_{a=0}^{\infty} \bigwedge_q^a \mathbb{C}_q^{1|1} \otimes \bigwedge_q^a (\mathbb{C}_q^{1|1})^* \curvearrowright U_q(\mathfrak{gl}_2). \quad (7.17)$$

Notice that $\bigwedge_q^0 \mathbb{C}_q^{1|1} \otimes \bigwedge_q^0 (\mathbb{C}_q^{1|1})^* \cong \mathbb{C}_q$ (the trivial representation), while all other $\bigwedge_q^a \mathbb{C}_q^{1|1} \otimes \bigwedge_q^a (\mathbb{C}_q^{1|1})^*$ for $a \geq 1$ are four-dimensional, indecomposable and isomorphic to each other (see Sartori (2015)). The two commuting actions (7.17) can be pictured as in figure 2. In the picture, each dot corresponds to a basis vector. The vertical arrows denote the action of the generators E and F of $U_q(\mathfrak{gl}_{1|1})$, while the horizontal double arrows denote the action of the generators E (solid) and F (dashed, although not the whole action is shown) of $U_q(\mathfrak{gl}_2)$. Of course, the actions have coefficients which are not indicated. From the picture, one sees that (7.17), as an $U_q(\mathfrak{gl}_2)$ -module, decomposes as $M^-(1, -1) \oplus M^-(1, -1) \oplus P^-(1, -1)$, where $M^-(1, -1)$ denotes the (opposite) Verma module with lowest weight $(1, -1)$ and $P^-(1, -1)$ denotes the (opposite) indecomposable projective module with head isomorphic to $M^-(1, -1)$. In particular, one can see explicitly that the action of $\dot{U}_q(\mathfrak{gl}_2)$, together with the projections onto the weight spaces, gives the full centralizer of the $U_q(\mathfrak{gl}_{1|1})$ -action.




Figure 2: The two commuting actions of (7.17).

Equivalence of categories

We conclude this section proving that the functors $\Phi_{\eta_{r,s}}$ glue together giving an equivalence of categories between the direct limit of $\dot{\mathbf{U}}_q(\mathfrak{gl}_{\eta_{r,s}})_\beta$ for $r, s \rightarrow \infty$ and $\mathbf{Sp}(\beta)$.

Proposition 7.11. *The functor $\Phi_\eta: \dot{\mathbf{U}}_q(\mathfrak{gl}_\eta)_\beta \rightarrow \mathbf{Sp}(\beta)$ is faithful.*

Proof. We will only sketch the idea of the proof, leaving the details, which are anyway straightforward, to the interested reader. As in the previous proof, by lemma 7.3 we can restrict to the case $\eta = \eta_{r,s}$. In the case $s = 0$ faithfulness can be proven as usual, by reducing to the Hecke algebra case, or see (Cautis et al., 2014, Theorem 4.4.1). For $s > 0$, if the target categories $U_q(\mathfrak{gl}_{r+s})_{\eta_{r,s}}$ were ribbon, a standard argument would allow to reduce to the upwards case. Although this is not exactly the case, one can adapt the argument.

Let $\varphi \in \text{Hom}_{U_q(\mathfrak{gl}_{\eta_{r,s}})_\beta}(\mathbf{1}_\lambda, \mathbf{1}_\mu)$ and suppose $\Phi_{\eta_{r,s}}(\varphi) = 0$. One can define a map $\dot{\mathbf{U}}_q(\mathfrak{gl}_{r+s})_{\eta_{r,s}} \rightarrow \dot{\mathbf{U}}_q(\mathfrak{gl}_{r'+s})_{\eta_{r',s}}$ for r' big enough, given graphically by

$$\begin{array}{c} \uparrow \dots \uparrow \downarrow \dots \downarrow \\ \boxed{\varphi} \\ \uparrow \dots \uparrow \downarrow \dots \downarrow \end{array} \mapsto \varphi' = \begin{array}{c} \uparrow \dots \uparrow \downarrow \dots \downarrow \\ \begin{array}{c} \text{curved lines} \\ \downarrow \\ \boxed{\varphi} \end{array} \\ \uparrow \dots \uparrow \downarrow \dots \downarrow \end{array} . \quad (7.18)$$

Using an argument similar to the proof of proposition 7.5 (cf. also (Queffelec and Sartori, 2018, Lemma 4.1 and Remark 4.2)), it follows that φ' is in the

image of the obvious map $\dot{\mathbf{U}}_q(\mathfrak{gl}_{\eta_{r',0}})_\beta \rightarrow \dot{\mathbf{U}}_q(\mathfrak{gl}_{\eta_{r',s}})_\beta$, hence it has a preimage $\tilde{\varphi}' \in \dot{\mathbf{U}}_q(\mathfrak{gl}_{\eta_{r',0}})_\beta = \dot{\mathbf{U}}_q^{\geq 0}(\mathfrak{gl}_{r'})$. Since its image in $\mathbf{Sp}(\beta)$ is zero, we have $\tilde{\varphi}' = 0$, hence $\varphi' = 0$. Since the construction (7.18) can be inverted (by a similar picture) it follows that also $\varphi = 0$. \square

In particular, we can define an infinite version $\dot{\mathbf{U}}_q(\mathfrak{gl}_{\eta_{\infty,\infty}})_\beta$ as the direct limit of $\dot{\mathbf{U}}_q(\mathfrak{gl}_{\eta_{r,s}})_\beta$ for $r, s \rightarrow \infty$. Propositions 7.5 and 7.11 then give:

Corollary 7.12. *The functors $\Phi_{\eta_{r,s}}$ induce an equivalence of categories between $\dot{\mathbf{U}}_q(\mathfrak{gl}_{\eta_{\infty,\infty}})_\beta$ and $\mathbf{Sp}(\beta)$.*

8. Link invariants

In this final section, we explain how colored link invariants of type A can be interpreted inside the categories $\mathbf{Sp}(\beta)$ or $\dot{\mathbf{U}}_q(\mathfrak{gl}_{\eta_{\infty,\infty}})_\beta$.

The $\mathfrak{gl}_{m|n}$ link invariant and the HOMFLY-PT polynomial

By proposition 6.15 we have an invariant of tangles $\mathcal{Q}_\beta: \mathbf{Tangles}_q^{\text{lab}} \rightarrow \mathbf{Sp}(\beta)$ with values in $\mathbf{Sp}(\beta)$. Moreover, for $\beta = d = m - n$ the diagram

$$\begin{array}{ccc} & & \mathbf{Sp}(d) \\ & \nearrow \mathcal{Q}_d & \downarrow \mathfrak{G}_{m|n} \\ \mathbf{Tangles}_q^{\text{lab}} & \xrightarrow{\mathcal{RT}_{m|n}} & \mathbf{Rep}_{m|n} \end{array} \quad (8.1)$$

commutes. Hence the invariant \mathcal{Q}_d gives rise to all such \mathfrak{gl}_{m-n} Reshetikhin-Turaev invariants, for which $m - n = d$.

Let $T \in \text{Hom}_{\mathbf{Tangles}_q^{\text{lab}}}(\mathbf{a}, \mathbf{b})$ be a labeled tangle. Then the lifted invariant \mathcal{Q}_d assigns to it an element of $\text{Hom}_{\mathbf{Sp}(d)}(\mathbf{a}, \mathbf{b})$. Since this space is in general bigger than $\text{Hom}_{\mathbf{Rep}_{m|n}}(\bigwedge_q^{\mathbf{a}} \mathbb{C}_q^{m|n}, \bigwedge_q^{\mathbf{b}} \mathbb{C}_q^{m|n})$, the lifted invariant \mathcal{Q}_d is finer than the Reshetikhin-Turaev invariant $\mathcal{RT}_{m|n}$.

We are mostly interested in the special case of a link L (hence $L \in \text{End}_{\mathbf{Tangles}_q^{\text{lab}}}(\emptyset)$, where \emptyset is the empty sequence). We will denote by $P_\beta(L)$ the Reshetikhin-Turaev invariant of links $\mathcal{Q}_\beta(L)$. By corollary 6.9, the endomorphism space

$\text{End}_{\text{Sp}(\beta)}(\emptyset)$ is one-dimensional, and if $\beta = d = m - n$ then this endomorphism space is canonically isomorphic to $\text{End}_{\text{Rep}_{m|n}}(\mathbb{C}_q)$. Hence in the case of links the lifted invariant \mathcal{Q}_d gives the same link polynomial as the classical Reshetikhin-Turaev invariant. Since \mathcal{Q}_d does not depend on m and n , but only on their difference $d = m - n$, we deduce the following well-known observation:

Proposition 8.1. *The Reshetikhin-Turaev invariant of links labeled by exterior powers of the vector representation of $\mathfrak{gl}_{m|n}$ only depends on the difference $m - n$.*

REMARK 8.2. This is more generally true for links labeled by partitions, see for example [Queffelec and Sartori \(2015\)](#).

Notice that also in the case β generic $P_\beta(L)$ defines an invariant of oriented framed links, which is a polynomial in two variables q and q^β . Then $P_\beta(L)$ is the colored HOMFLY-PT polynomial of the link L (see [Queffelec and Sartori \(2018, 2015\)](#)).

Cutting strands and the Alexander polynomial

In the case where $\beta = 0$, it is a well-known phenomenon that $P_0(L)$ is trivial, since (6.15b) sends trivial circles to zero. In order to obtain an interesting invariant in the $\beta = 0$ case, one has to cut open one strand (see for example [Sartori \(2015\)](#)). We hence define for $a \in \mathbb{Z}_{\geq 0}$ the link invariant

$$\tilde{P}_{\beta,a} : \text{Links}^{\text{lab}} \longrightarrow \mathbb{C}_q \quad (8.2)$$

given by assigning to $L \in \text{End}_{\text{Tangles}^{\text{lab}}}(\emptyset)$ the element $\mathcal{Q}_\beta(\tilde{L}) \in \text{End}_{\text{Sp}(\beta)}(a)$, where \tilde{L} is obtained by L by cutting one strand labeled by a . By the following proposition 8.3, this does not depend on the strand cut and is therefore a well-defined link invariant.

In the case β generic, one obtains the reduced HOMFLY-PT polynomial. The two specializations $\beta = 0$ and $\beta = m$ yield respectively the Alexander and the reduced \mathfrak{sl}_m Reshetikhin-Turaev polynomial (see [Queffelec and Sartori \(2018, 2015\)](#) for our normalizations of these invariants and for details).

Proposition 8.3. *Let $a \in \mathbb{Z}_{\geq 0}$ and let $\tau \in \text{End}_{\text{Sp}(\beta)}(a \otimes a)$. Then*

$$\begin{array}{c} a \\ \downarrow \\ \boxed{\tau} \\ \uparrow \\ a \end{array} \text{ (with a loop) } = \begin{array}{c} \uparrow \\ a \\ \boxed{\tau} \\ a \\ \downarrow \end{array} \text{ (with a loop) } . \quad (8.3)$$

Proof. It is enough to check the statement for τ a basis element. Since $\text{End}_{\text{Sp}(\beta)}(a \otimes a) \cong \text{End}_{\text{Sp}^+}(a \otimes a)$, it is easy to check that the elements $\{E^{(k)}F^{(k)} \mid 0 \leq k \leq a\}$ give a basis. We have

$$\begin{array}{c} \uparrow \\ a \\ \text{ (diagonal lines) } \\ a \\ \downarrow \end{array} = \begin{array}{c} \uparrow k \\ \text{ (crossing) } \\ \downarrow k \end{array} = \begin{array}{c} \uparrow k \\ \text{ (parallel lines) } \\ \downarrow k \end{array} = \begin{array}{c} \uparrow k \\ \text{ (diagonal lines) } \\ \downarrow k \end{array} , \quad (8.4)$$

where the first equality is a consequence of relations (5.21) and (5.22) and the third one follows by an iterated use of (5.5e). \square

In light of the inclusion of categories $\Phi_{\boldsymbol{\eta}}$, it is possible to regard all these invariants of tangles also as having values in $\dot{\mathbf{U}}_q(\mathfrak{gl}_{\boldsymbol{\eta}})_{\beta}$ for some sequence of signs $\boldsymbol{\eta}$ long enough. For a more detailed description of link invariants with values in $\dot{\mathbf{U}}_q(\mathfrak{gl}_{\boldsymbol{\eta}})_{\beta}$ we refer to [Queffelec and Sartori \(2018\)](#). We point out that working in $\dot{\mathbf{U}}_q(\mathfrak{gl}_{\boldsymbol{\eta}})_{\beta}$ is particularly handy for computation purposes, since $\dot{\mathbf{U}}_q(\mathfrak{gl}_{\boldsymbol{\eta}})_{\beta}$ has a more rigid structure than $\text{Sp}(\beta)$. Moreover, we believe this can give more insight for developing a categorification.

REMARK 8.4. ³ In this paper, we studied *skew* Howe duality, generalizing [Cautis et al. \(2014\)](#), hence we obtain the category $\text{Sp}(d)$ which describes intertwining operators between exterior powers of the natural representation of $U_q(\mathfrak{gl}_{m|n})$. In this setting, the easiest case where we can explicitly describe all relations is when $n = 0$ and we consider the category $\text{Rep}_{m|0}$ (see corollary 6.17).

³Since this paper was written, the program outlined in this remark has been partly carried out by Tubbenhauer, Vaz and Wedrich in [Tubbenhauer et al. \(2017\)](#). Interestingly, their spider categories also allow to mix symmetric and exterior powers.

However, one could also be interested in replacing *skew* Howe duality by *symmetric* Howe duality, and define analogous (and similar) categories $\mathbf{Sp}_{\text{sym}}^+(\beta)$ of “symmetric” spiders, which would describe intertwining operators between symmetric powers of $\mathbb{C}_q^{m|n}$ (see also Rose and Tubbenhauer (2015)).

It is interesting to notice that symmetric powers actually appear already in our picture: indeed, $\bigwedge_q^a(\mathbb{C}_q^{m|n})$ is isomorphic to $(S_q^a \mathbb{C}_q^{n|m})\langle a \rangle$, where $\langle a \rangle$ denotes a shift by a in the $\mathbb{Z}/2\mathbb{Z}$ -degree. This suggests that there is a strong similarity between the “skew categories” \mathbf{Sp}^+ , $\mathbf{Sp}(\beta)$ and the “symmetric categories” $\mathbf{Sp}_{\text{sym}}^+$, $\mathbf{Sp}_{\text{sym}}(\beta)$. Indeed, one can fix the conventions so that \mathbf{Sp}^+ and $\mathbf{Sp}_{\text{sym}}^+$ are the same. However, the parity shift above introduces some sign differences between $\mathbf{Sp}(\beta)$ and $\mathbf{Sp}_{\text{sym}}(\beta)$ which prevent an isomorphism on the nose.

Alexander, J. W., 1928. Topological invariants of knots and links. Trans. Amer. Math. Soc. 30 (2), 275–306.

URL <http://dx.doi.org/10.2307/1989123>

Berenstein, A., Zwicknagl, S., 2008. Braided symmetric and exterior algebras. Trans. Amer. Math. Soc. 360 (7), 3429–3472.

URL <http://dx.doi.org/10.1090/S0002-9947-08-04373-0>

Bernstein, J., Frenkel, I., Khovanov, M., 1999. A categorification of the Temperley-Lieb algebra and Schur quotients of $U(\mathfrak{sl}_2)$ via projective and Zuckerman functors. Selecta Math. (N.S.) 5 (2), 199–241.

URL <http://dx.doi.org/10.1007/s000290050047>

Blanchet, C., 2000. Hecke algebras, modular categories and 3-manifolds quantum invariants. Topology 39 (1), 193–223.

URL [http://dx.doi.org/10.1016/S0040-9383\(98\)00066-4](http://dx.doi.org/10.1016/S0040-9383(98)00066-4)

Brundan, J., Stroppel, C., 2012. Gradings on walled Brauer algebras and Khovanov’s arc algebra. Adv. Math. 231 (2), 709–773.

URL <http://dx.doi.org/10.1016/j.aim.2012.05.016>

Cautis, S., Kamnitzer, J., Licata, A., 2010. Categorical geometric skew Howe duality. Invent. Math. 180 (1), 111–159.

URL <http://dx.doi.org/10.1007/s00222-009-0227-1>

Cautis, S., Kamnitzer, J., Morrison, S., 2014. Webs and quantum skew Howe duality. Math. Ann. 360 (1-2), 351–390.

URL <http://dx.doi.org/10.1007/s00208-013-0984-4>

- Chari, V., Pressley, A., 1994. A guide to quantum groups. Cambridge University Press, Cambridge.
- Cheng, S.-J., Lam, N., Zhang, R. B., 2004. Character formula for infinite-dimensional unitarizable modules of the general linear superalgebra. *J. Algebra* 273 (2), 780–805.
URL [http://dx.doi.org/10.1016/S0021-8693\(03\)00538-6](http://dx.doi.org/10.1016/S0021-8693(03)00538-6)
- Cheng, S.-J., Wang, W., 2012. Dualities and representations of Lie superalgebras. Vol. 144 of Graduate Studies in Mathematics. American Mathematical Society, Providence, RI.
- Dipper, R., Doty, S., Stoll, F., 2014. The quantized walled Brauer algebra and mixed tensor space. *Algebr. Represent. Theory* 17 (2), 675–701.
URL <http://dx.doi.org/10.1007/s10468-013-9414-2>
- Doty, S., Giaquinto, A., 2002. Presenting Schur algebras. *Int. Math. Res. Not.* (36), 1907–1944.
URL <http://dx.doi.org/10.1155/S1073792802201026>
- Frenkel, I., Khovanov, M., Stroppel, C., 2006. A categorification of finite-dimensional irreducible representations of quantum \mathfrak{sl}_2 and their tensor products. *Selecta Math. (N.S.)* 12 (3-4), 379–431.
URL <http://dx.doi.org/10.1007/s00029-007-0031-y>
- Freyd, P., Yetter, D., Hoste, J., Lickorish, W. B. R., Millett, K., Ocneanu, A., 1985. A new polynomial invariant of knots and links. *Bull. Amer. Math. Soc. (N.S.)* 12 (2), 239–246.
URL <http://dx.doi.org/10.1090/S0273-0979-1985-15361-3>
- Geer, N., 2005. The Kontsevich integral and quantized Lie superalgebras. *Algebr. Geom. Topol.* 5, 1111–1139.
URL <http://dx.doi.org/10.2140/agt.2005.5.1111>
- Geer, N., Patureau-Mirand, B., 2008. On the colored HOMFLY-PT, multi-variable and Kashaev link invariants. *Commun. Contemp. Math.* 10 (suppl. 1), 993–1011.
URL <http://dx.doi.org/10.1142/S0219199708003125>
- Geer, N., Patureau-Mirand, B., 2010. Multivariable link invariants arising from Lie superalgebras of type I. *J. Knot Theory Ramifications* 19 (1),

- 93–115.
 URL <http://dx.doi.org/10.1142/S0218216510007784>
- Gorsky, E., Gukov, S., Stosic, M., Apr. 2013. Quadruply-graded colored homology of knots. ArXiv e-prints.
- Grant, J., 2016. A generators and relations description of a representation category of $U_q(\mathfrak{gl}(1|1))$. *Algebraic & Geometric Topology* 16 (1), 509–539.
 URL <http://dx.doi.org/10.2140/agt.2016.16.509>
- Howe, R., 1989. Remarks on classical invariant theory. *Trans. Amer. Math. Soc.* 313 (2), 539–570.
 URL <http://dx.doi.org/10.2307/2001418>
- Howe, R., 1995. Perspectives on invariant theory: Schur duality, multiplicity-free actions and beyond. In: *The Schur lectures (1992) (Tel Aviv)*. Vol. 8 of *Israel Math. Conf. Proc.* Bar-Ilan Univ., Ramat Gan, pp. 1–182.
 URL <http://dx.doi.org/10.1007/BF02771542>
- Jones, V. F. R., 1985. A polynomial invariant for knots via von Neumann algebras. *Bull. Amer. Math. Soc. (N.S.)* 12 (1), 103–111.
 URL <http://dx.doi.org/10.1090/S0273-0979-1985-15304-2>
- Kamnitzer, J., Tingley, P., 2009. The crystal commutor and Drinfeld’s unitarized R -matrix. *J. Algebraic Combin.* 29 (3), 315–335.
 URL <http://dx.doi.org/10.1007/s10801-008-0137-0>
- Kassel, C., 1995. *Quantum groups*. Vol. 155 of *Graduate Texts in Mathematics*. Springer-Verlag, New York.
- Kauffman, L. H., Saleur, H., 1991. Free fermions and the alexander-conway polynomial. *Communications in mathematical physics* 141 (2), 293–327.
 URL <https://doi.org/10.1007/BF02101508>
- Kazhdan, D., Lusztig, G., 1979. Representations of Coxeter groups and Hecke algebras. *Invent. Math.* 53 (2), 165–184.
 URL <http://dx.doi.org/10.1007/BF01390031>
- Khovanov, M., 2000. A categorification of the Jones polynomial. *Duke Math. J.* 101 (3), 359–426.
 URL <http://dx.doi.org/10.1215/S0012-7094-00-10131-7>

- Khovanov, M., 2002. A functor-valued invariant of tangles. *Algebr. Geom. Topol.* 2, 665–741.
URL <http://dx.doi.org/10.2140/agt.2002.2.665>
- Khovanov, M., 2004. $\mathfrak{sl}(3)$ link homology. *Algebr. Geom. Topol.* 4, 1045–1081.
URL <http://dx.doi.org/10.2140/agt.2004.4.1045>
- Khovanov, M., 2006. An invariant of tangle cobordisms. *Trans. Amer. Math. Soc.* 358 (1), 315–327.
URL <http://dx.doi.org/10.1090/S0002-9947-05-03665-2>
- Khovanov, M., 2007. Triply-graded link homology and Hochschild homology of Soergel bimodules. *Internat. J. Math.* 18 (8), 869–885.
URL <http://dx.doi.org/10.1142/S0129167X07004400>
- Khovanov, M., Lauda, A. D., 2009. A diagrammatic approach to categorification of quantum groups. I. *Represent. Theory* 13, 309–347.
URL <http://dx.doi.org/10.1090/S1088-4165-09-00346-X>
- Khovanov, M., Lauda, A. D., 2010. A categorification of quantum $\mathfrak{sl}(n)$. *Quantum Topol.* 1 (1), 1–92.
URL <http://dx.doi.org/10.4171/QT/1>
- Khovanov, M., Lauda, A. D., 2011. A diagrammatic approach to categorification of quantum groups II. *Trans. Amer. Math. Soc.* 363 (5), 2685–2700.
URL <http://dx.doi.org/10.1090/S0002-9947-2010-05210-9>
- Khovanov, M., Rozansky, L., 2008. Matrix factorizations and link homology. *Fund. Math.* 199 (1), 1–91.
URL <http://dx.doi.org/10.4064/fm199-1-1>
- Kuperberg, G., 1996. Spiders for rank 2 Lie algebras. *Comm. Math. Phys.* 180 (1), 109–151.
URL <http://projecteuclid.org/getRecord?id=euclid.cmp/1104287237>
- Kwon, J.-H., 2014. Crystal bases of q -deformed Kac modules over the quantum superalgebras $U_q(\mathfrak{gl}(m|n))$. *Int. Math. Res. Not. IMRN* (2), 512–550.
- Lauda, A. D., Queffelec, H., Rose, D. E. V., 2015. Khovanov homology is a skew Howe 2-representation of categorified quantum $\mathfrak{sl}(m)$. *Algebraic &*

- Geometric Topology 15 (5), 2517–2608.
 URL <http://dx.doi.org/10.2140/agt.2015.15.2517>
- Lehrer, G. I., Zhang, H., Zhang, R. B., 2011. A quantum analogue of the first fundamental theorem of classical invariant theory. *Comm. Math. Phys.* 301 (1), 131–174.
 URL <http://dx.doi.org/10.1007/s00220-010-1143-3>
- Lusztig, G., 1993. Introduction to quantum groups. Vol. 110 of *Progress in Mathematics*. Birkhäuser Boston Inc., Boston, MA.
- Mackaay, M., Webster, B., Feb. 2015. Categorified skew Howe duality and comparison of knot homologies. *ArXiv e-prints*.
- Mackaay, M., Yonezawa, Y., 2013. $\mathfrak{sl}(n)$ -Web categories. *ArXiv e-prints*.
- Manolescu, C., Ozsváth, P., Szabó, Z., Thurston, D., 2007. On combinatorial link Floer homology. *Geom. Topol.* 11, 2339–2412.
 URL <http://dx.doi.org/10.2140/gt.2007.11.2339>
- Mitsuhashi, H., 2006. Schur-Weyl reciprocity between the quantum superalgebra and the Iwahori-Hecke algebra. *Algebr. Represent. Theory* 9 (3), 309–322.
 URL <http://dx.doi.org/10.1007/s10468-006-9014-5>
- Morton, H. R., 2002. Power sums and homfly skein theory. *Geometry & Topology Monographs* 4, 235–244.
- Murakami, H., Ohtsuki, T., Yamada, S., 1998. Homfly polynomial via an invariant of colored plane graphs. *Enseign. Math.* (2) 44 (3-4), 325–360.
- Murakami, J., 1992. The multi-variable Alexander polynomial and a one-parameter family of representations of $\mathcal{U}_q(\mathfrak{sl}(2, \mathbb{C}))$ at $q^2 = -1$. In: *Quantum groups*. Springer Berlin Heidelberg, pp. 350–353.
- Ohtsuki, T., 2002. Quantum invariants. Vol. 29 of *Series on Knots and Everything*. World Scientific Publishing Co. Inc., River Edge, NJ, a study of knots, 3-manifolds, and their sets.
- Przytycki, J. H., 1999. Fundamentals of Kauffman bracket skein modules. *Kobe J. Math.* 16 (1), 45–66, [arXiv:9809113](https://arxiv.org/abs/9809113).

- Przytycki, J. H., Traczyk, P., 1987. Conway algebras and skein equivalence of links. *Proc. Amer. Math. Soc.* 100 (4), 744–748.
URL <http://dx.doi.org/10.2307/2046716>
- Queffelec, H., 2015. Skein modules from skew Howe duality and affine extensions. *SIGMA* 11 (030).
URL <https://doi.org/10.3842/SIGMA.2015.030>
- Queffelec, H., Rose, D. E. V., 2016. The \mathfrak{sl}_n foam 2-category: a combinatorial formulation of Khovanov-Rozansky homology via categorical skew Howe duality. *Advances in Mathematics* 302, 1251–1339.
URL <https://doi.org/10.1016/j.aim.2016.07.027>
- Queffelec, H., Sartori, A., Jun. 2015. A note on the $\mathfrak{gl}_{m|n}$ link invariants and the HOMFLY-PT polynomial. ArXiv e-prints.
- Queffelec, H., Sartori, A., 2018. HOMFLY-PT and Alexander polynomials from a doubled Schur algebra. *Quantum Topol* 9, 323–347.
URL <https://doi.org/10.4171/QT/109>
- Reshetikhin, N. Y., Turaev, V. G., 1990. Ribbon graphs and their invariants derived from quantum groups. *Comm. Math. Phys.* 127 (1), 1–26.
URL <http://projecteuclid.org/euclid.cmp/1104180037>
- Rose, D. E. V., Tubbenhauer, D., 2015. Symmetric webs, Jones-Wenzl recursions and q -Howe duality. *International Mathematics Research Notices* 2016 (17), 5249–5290.
URL <https://doi.org/10.1093/imrn/rnv302>
- Rouquier, R., Dec. 2008. 2-Kac-Moody algebras. ArXiv e-prints.
- Sartori, A., 2014. Categorification of tensor powers of the vector representation of $U_q(\mathfrak{gl}(1|1))$. Ph.D. thesis, Universität Bonn, uRN: [urn:nbn:de:hbz:5n-36359](https://nbn-resolving.org/urn:nbn:de:hbz:5n-36359).
- Sartori, A., 2015. The Alexander polynomial as quantum invariant of links. *Ark. Mat.* 53 (1), 177–202.
URL <http://dx.doi.org/10.1007/s11512-014-0196-5>
- Sartori, A., 2016. Categorification of tensor powers of the vector representation of $U_q(\mathfrak{gl}(1|1))$. *Selecta Mathematica* 22 (2), 669–734.
URL <https://doi.org/10.1007/s00029-015-0202-1>

- Scheunert, M., 1992. Serre-type relations for special linear lie superalgebras. *letters in mathematical physics* 24 (3), 173–181.
URL <http://dx.doi.org/10.1007/BF00402892>
- Soergel, W., 1997. Kazhdan-Lusztig-Polynome und eine Kombinatorik für Kipp-Moduln. *Represent. Theory* 1, 37–68.
URL <http://dx.doi.org/10.1090/S1088-4165-97-00006-X>
- Tubbenhauer, D., 2014. \mathfrak{sl}_n -webs, categorification, and Khovanov-Rozansky homologies. *ArXiv e-prints*.
- Tubbenhauer, D., Vaz, P., Wedrich, P., 2017. Super q-howe duality and web categories. *Algebraic & geometric topology* 17 (6), 3703–3749.
URL <https://doi.org/10.2140/agt.2017.17.3703>
- Turaev, V., 1991. Skein quantization of Poisson algebras of loops on surfaces. *Ann. Sci. École Norm. Sup. (4)* 24 (6), 635–704.
URL http://www.numdam.org/item?id=ASENS_1991_4_24_6_635_0
- Turaev, V. G., 1994. Quantum invariants of knots and 3-manifolds. Vol. 18 of *de Gruyter Studies in Mathematics*. Walter de Gruyter & Co., Berlin.
- Viro, O. Y., 2006. Quantum relatives of the Alexander polynomial. *Algebra i Analiz* 18 (3), 63–157.
URL <http://dx.doi.org/10.1090/S1061-0022-07-00956-9>
- Weimer, F., 2005. A tangle analogon for Leduc’s deformation of the walled Brauer algebra. *Ph.D. thesis*, University of Stuttgart.
- Wu, Y., Zhang, R. B., 2009. Unitary highest weight representations of quantum general linear superalgebra. *J. Algebra* 321 (11), 3568–3593.
URL <http://dx.doi.org/10.1016/j.jalgebra.2008.02.040>
- Zhang, H., 2009. The quantum general linear supergroup, canonical bases and Kazhdan-Lusztig polynomials. *Sci. China Ser. A* 52 (3), 401–416.
URL <http://dx.doi.org/10.1007/s11425-008-0150-8>
- Zhang, R., 1998. Structure and representations of the quantum general linear supergroup. *Communications in mathematical physics* 195 (3), 525–547.
URL <https://doi.org/10.1007/s002200050401>