



HAL
open science

First Record of *Echinoparyphium recurvatum* (Trematoda, Echinostomatidae) in Pakistan with New Avian Definitive Host, *Vanellus leucurus*

Nadir Ali Birmani, Saima Naz, Gabriel Mouahid

► **To cite this version:**

Nadir Ali Birmani, Saima Naz, Gabriel Mouahid. First Record of *Echinoparyphium recurvatum* (Trematoda, Echinostomatidae) in Pakistan with New Avian Definitive Host, *Vanellus leucurus*. Archives on Veterinary Science and Technology, 2019, 10.29011/2637-9988/100049 . hal-02064162

HAL Id: hal-02064162

<https://hal.science/hal-02064162v1>

Submitted on 11 Mar 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Review Article

First Record of *Echinoparyphium recurvatum* (Trematoda, Echinostomatidae) in Pakistan with New Avian Definitive Host, *Vanellus leucurus*

Nadir Ali Birmani¹, Saima Naz,¹ Gabriel Mouahid^{2*}¹Department of Zoology, University of Sindh, Jamshoro, Sindh, Pakistan²IHPE UMR 5244 CNRS/UPVD/IFREMER/UM/ University of Perpignan, Perpignan, France

*Corresponding author: Gabriel Mouahid, IHPE UMR 5244 CNRS/UPVD/IFREMER/UM/ University of Perpignan, 58 avenue Paul Alduy, 66860 Perpignan, France. Tel: +33-468662184; Fax: +33-468662281; Email: mouahid@univ-perp.fr

Citation: Birmani NA, Naz S, Mouahid G (2019) First Record of *Echinoparyphium recurvatum* (Trematoda, Echinostomatidae) in Pakistan with New Avian Definitive Host, *Vanellus leucurus*: 149. DOI: 10.29011/2637-9988/100049

Received Date: 18January, 2019; Accepted Date: 07February, 2019; Published Date: 15February, 2019

Abstract

Parasitological examinations of White-tailed Lapwing *Vanellus leucurus* (Charadriiformes: Charadriidae) collected from Dadu district (Sindh Province, Southern Pakistan) were carried out at the Department of Zoology, University of Sindh, Jamshoro. A total of 64 trematodes belonging to the genus *Echinoparyphium* [1], were collected from the small intestine of two individual hosts. On the basis of the number and varying size of collar spines, number and size of ventral spines, body shape, arrangement of genital organs, distribution of vitellaria and other diagnostic characters, these trematodes were identified as *Echinoparyphium recurvatum* [2]. Our study provides the first Pakistan record of the trematode *E. recurvatum* with also the first record of *V. leucurus* as avian definitive host.

Keywords: Charadriiformes; Charadriidae; *Echinoparyphium recurvatum*; Sindh Province; Southern Pakistan; *Vanellus leucurus*

Introduction

Genus *Echinoparyphium* is an important taxon in the family Echinostomatidae and has considerable importance in medical and veterinary sciences [3]. It has previously been described as *Distomum* by von Linstow [2] using specimens collected from the small intestine of *Fuligula manila*. Dietz [1] carried out a revision of Echinostomatidae and proposed different genus including the genus *Echinoparyphium*. Kanev [4] published a checklist with 151 species of *Echinoparyphium* using freshwater snails as first and second intermediate hosts, tadpoles and fish as a second intermediate hosts, and birds and mammals as definitive hosts. Since the creation of the genus, most of the published reports are based on the adequate descriptions of adult specimens. The more differentiating character is the morphometry of collar spines in adult stage, specifically the total number of spines (29-45), their sharply pointed shape and their arrangement in double row, with dorsal aboral spines noticeably longer than dorsal oral ones.

In the present paper, we report for the first time the presence of *E. recurvatum* recovered from the small intestine of the White-tailed Lapwing *Vanellus leucurus* (Charadriiformes: Charadriidae) in the Sindh province (Southern Pakistan) and describe the mor-

phology of the adult. The White-tailed Lapwing is inhabitant of freshwater marshy areas and lake shores. They more frequently seek food in shallow water. They are omnivorous and their food includes insects (chiefly beetles, grasshoppers, caterpillars, and fly larvae), mollusks, worms and crustaceans [5] which are potential sources of developmental stages of a variety of helminth parasites.

Materials And Methods

Four live White-tailed Lapwings *Vanellus leucurus* were collected from the Dadu city (GPS coordinates: 26°44'2.76"N; 67°46'46.20"E) located in the North-Western Sindh province, Pakistan. Birds were examined for the presence of trematode parasites in the Parasitology laboratory of the department of Zoology, University of Sindh, Jamshoro, in January 2016. A total of 64 trematodes belonging to the genus *Echinoparyphium*, were collected from the small intestine of two birds and were processed according to the method given by Garcia and Ash [6]. Line drawings were made with the help of a camera Lucida. All measurements were based on 13 mature specimens and are given in micrometer (µm) followed by range in parenthesis. For practical reasons, we have chosen to give the value of the minor axis and the major axis for any spherical or subspherical organ. Length and width are kept for elongated structures such as the body or its various parts, oesophagus, head collar spines, etc. Identification was made with the help of keys given by Jones *et al.* [7] and Yamaguti [8]. Specimens

are available at the Museum of Natural History in Paris under the voucher reference: MNHN HEL682.

Taxonomic Part

Description (based on 13 mature specimens; (Figures 1-3; Table 1)




Figure 1: Morphology of *Echinoparyphium recurvatum*: Lateral view of entire adult worm.




Figure 2: Morphology of *Echinoparyphium recurvatum*: anterior end of the adult with collar spined arrangement (45 spines), Oral Sucker (OS), Prepharynx (Pph), Pharynx (Ph) and Oesophagus (Oe).




Figure 3: Morphology of *Echinoparyphium recurvatum*: genital pore region with Ventral Sucker (VS), Uterus (U) with Eggs (E), end part of the uterus or Metraterm (M), Genital Atrium (GA) and cirrus sac with Seminal Vesicle (SV), Pars Prostatica (PP), Ejaculatory Duct (ED) and Genital Pore (GP).

Body small, spinose, elongate, tapering posteriorly, 3868 (3150-3905) length by 684 (521-690) width. Maximum width at level of ventral sucker. Forebody small, measuring 657 (600-682) length ventrally bent slightly. Hindbody measuring 2736 (2051-2818) length. Integument armed with very small spines which extend from the head crown to the testicular region. Head collar reniform, well developed, measuring 305 (288-310) by 263 (220-

271) and armed with 45 spines arranged in double rows. Corner oral spines 6 on each side, 53.5 (35-70) length by 15.5 (9-22) width; lateral spines 39 (30-46) length by 10 (9-13) width, dorsal spines on oral sucker region 75.5 (40-90) length by 12.5 (9-14) width; dorsal spines away from oral sucker 98.5 (80-120) length by 39 (30-50) width. Oral sucker subterminal, pear shaped in most of the specimens, measuring 118 (110-125) by 110 (100-118). Pre-pharynx very small, 14.5 (14-18) length. Pharynx muscular, elongate-oval, measuring 110 (106-120) by 78 (70-80). Oesophagus small, tubular, measuring 394 (368-422) length. Intestinal bifurcation in front of ventral sucker. Ventral sucker large, muscular, cup-shaped, measuring 473 (425-500) by 421 (392-430). Testes tandem, elongate-oval, smooth, contiguous, in some specimens slightly lobed, located in posterior half of the body. Anterior testicle measuring 394 (385-405) by 315 (302-320) located in the 3rd quarter of the body, whereas, a major part of posterior testicle present in the 3rd quarter and very small portion of it extend to last 4th quarter of body. Posterior testicle measuring 447 (430-455) by 315 (310-320). Post-testicular space measuring 815 (810-820) length. Cirrus sac muscular, elongate, located at antero-lateral side of the ventral sucker, measuring 440 (410-450) by 196 (180-210). Genital pore preacetabular in position, funnel like in appearance. Ovary subspherical, median, measuring 223 (220-235) by 184 (180-206). Distance between ventral sucker and ovary, 789 (700-800) length. Mehlis gland distinct. Uterine seminal receptacle immediately between anterior testicle and the acetabulum, laterally elongate, measuring 210 (200-220) by 105 (100-110). Uterus without loops, filled with eggs, covering an area between ovary and ventral sucker. Eggs operculated, measuring 79 (60-98) length by 51 (35-65) width. Vitellaria composed of large, irregular follicles, distributed in lateral fields of the body, commencing from mid-level of the uterus, extending backward reaching up to posterior extremity, confluent in post-testicular area in few specimens. Excretory vesicle tubular. Excretory pore terminal.

Discussion

The genus *Echinoparyphium* is distinct from other genera of Echinostomatidae in possessing a well-developed head collar with 29 to 45 spines in a double row, with small spines covering the tegument extending to the testicular level ventrally and to the anterior margin of the ventral sucker dorsally. Large testes are tandem,

elongated-oval, smooth or irregular, contiguous, post-equatorial and located in center of the hind body. Ovary small, spherical, median or submedian, pre-equatorial [9]. The genus *Echinoparyphium* counts many parasitic species of the intestine of many groups of birds and some species of mammals and the Type species is *E. elegans* [3,10,11]. Several species of genus *Echinoparyphium* have been found to infect the gastrointestinal tract of aquatic birds and mammals worldwide [12,13]. They are known to infect a large number of different taxa of birds (Anseriformes, Charadriiformes, Ciconiiformes, Columbiformes, Falconiformes, Galliformes, Gruiformes, Passeriformes, Podicipediformes, Strigiformes) and few taxa of mammals (Muridae (*Hydromys chrysogaster*, *Rattus norvegicus*, *Rattus colletti* and *Rattus rattus*), Cricetidae (*Ondatra zibethicus*), Canidae (*Alopex lagopus*) and Hominidae (*Homo sapiens*) [3,14].

According to the morphological description (Figure 1-3) and to the measurements (Table 1), our specimens belong to the genus *Echinoparyphium* and to the species *recurvatum*. It is the first time that this species is described in Pakistan and the first time that the avian host White-tailed Lapwing *Vanellus leucurus* is found to be a definitive host. Khan *et al.* [15] reported two species of *Echinoparyphium* in Pakistan: (i) *Echinoparyphium ellisi* [16] from *Anas crecca* in Nowshera (Khyber Pakhtunkhwa, one of the four provinces of Pakistan located in the northwestern Region) which belongs to the “recurvatum” group according to the phylogenetic study of Kostadinova *et al.* [17] and differs from our specimens in having a very long body (6200 x 1048 µm compared to our specimens 3868 x 684µm); (ii) *Echinoparyphium recurvatum* from *Anas crecca* in Peshawar (Khyber Pakhtunkhwa) which differs from our specimens in having 37 collar spines instead of 45 as it is well known for this species. The species known to have 37 collar spines is *E. serratum* described by Howell [18] and not *E. recurvatum*.

An updated list of natural avian hosts of *E. recurvatum* along with geographical distribution is given (Table 2). It shows that *E. recurvatum* is cosmopolitan species with a wide geographical distribution associated to a wide range of avian definitive host taxa including 7 orders and 10 families. The lack of specificity towards the definitive hosts is assigned to a low specificity towards the first and second intermediate hosts [19].

Citation: Birmani NA, Naz S, Mouahid G (2019) First Record of *Echinoparyphium recurvatum* (Trematoda, Echinostomatidae) in Pakistan with New Avian Definitive Host, *Vanellus leucurus*: 149. DOI: 10.29011/ 2637-9988/100049

Characters	Lee <i>et al.</i> [20]	Sohn [21]	Sereno-Uribe <i>et al.</i> [9]	Present study
Body (L)	3500-4700	2010-3090 (2760)	2750-3220 (2940)	3150-3905 (3868)
Body (W)	500-650	460-610 (550)	400-550 (460)	521-690 (684)
Head collar (Ma)	290-340	260-316 (283)	190-300 (269)	288-310 (305)
Head collar (Mi)	ng	173-214 (188)	170-220 (200)	220-271 (263)
Corner oral spines (L)	ng	53-70	49-66 (57)	39-68 (53)
Corner oral spines (W)	ng	ng	9-12 (10)	11-20 (16)
Dorsal oral spines (L)	ng	ng	45-62 (54)	58-93 (65)
Dorsal oral spines (W)	ng	ng	6-12 (9)	11-14 (12)
Oral sucker (Ma)	120-150	92-122 (109)	78-120 (96)	110-125 (118)
Oral sucker (Mi)	120-150	92-112 (99)	93-110 (100)	100-118 (110)
Pharynx (L)	110-130	61-87 (77)	100-108 (90)	106-120 (110)
Pharynx (W)	90-110	36-71 (46)	50-60 (60)	70-80 (78)
Oesophagus (L)	160-190	204-377 (315)	270-430 (350)	368-422 (394)
Cirrus sac (L)	300-400	209-337 (263)	190-260 (230)	410-450 (440)
Cirrus sac (W)	130-180	82-143 (117)	90-150 (150)	180-210 (196)
Ventral sucker (Ma)	320-400	306-383 (337)	260-380 (320)	425-500 (473)
Ventral sucker (Mi)	320-390	296-367 (333)	240-320 (290)	392-430 (421)
Ovary (Ma)	130-220	92-153 (122)	110-140 (120)	220-235 (223)
Ovary (Mi)	130-200	82-133 (112)	100-130 (120)	180-206 (184)
Anterior testis (L)	280-420	235-377 (329)	200-360 (300)	385-405 (394)
Anterior testis (W)	210-280	153-214 (188)	90-220 (150)	302-320 (315)
Posterior testis (L)	390-480	265-408 (355)	280-450 (370)	430-455 (447)
Posterior testis (W)	180-290	163-204 (187)	110-180 (140)	310-320 (315)
Egg (L)	82-97	96-105 (102)	70-96 (85)	65-93 (79)
Egg (W)	54-59	64-71 (68)	40-56 (47)	37-65 (51)

Table 1: Comparative morphometrics (in μm) of adult worms of *Echinoparyphium recurvatum* obtained in different studies; length (L), width (W), major axis (Ma), minor axis (Mi), ng: not given and mean values given in parentheses after ranges.

Host birds	Order and family	Geographical distribution	Reference
<i>Phasianus colchicus</i>	Galliformes, Phasianidae	Texas, USA	Pence <i>et al.</i> [22]
<i>Meleagris gallopavo</i>	Galliformes, Phasianidae	Eastern Kansas, USA	McJunkin <i>et al.</i> [23]
<i>Anas clypeata</i> , <i>Gallus gallus</i>	Anseriformes, Anatidae, Galliformes, Phasianidae	Mexico	Sereno-Uribe <i>et al.</i> [9]

<i>Gallinula chloropus galeata, Phimosus infuscatus</i>	Gruiformes, Rallidae, Pelecaniformes, Threskiornithidae	Buenos Aires Province Argentina	Lunaschi <i>et al.</i> [24]
<i>Charadrius hiaticula</i>	Charadriiformes, Charadriidae	Slovak Republic	Macko <i>et al.</i> [25]
<i>Buteo buteo</i>	Falconiformes, Accipitridae	Czech Republic	Sitko [26]
<i>Anser anser</i>	Anseriformes, Anatidae	Gilan province, Iran	Hosseini <i>et al.</i> [27]
<i>Gallinago gallinago, Tadorna ferruginea</i>	Charadriiformes, Scolopacidae, Anseriformes, Anatidae	India	Mehra [28]
<i>Vanellus leucurus</i>	Charadriiformes, Charadriidae	Sindh Province, Pakistan	Current paper
<i>Larus dominicanus</i>	Charadriiformes, Laridae	Otago Peninsula New Zealand	Latham and Poulin [19]
<i>Chroicocephalus novaeseelandia</i>	Charadriiformes, Laridae	Otago Peninsula New Zealand	Fredensborg <i>et al.</i> [29]
<i>Aythya novaeseelandia</i>	Anseriformes, Anatida	Lake Wanaka and the Waitaki River watershed, New Zealand	Davis [30]
<i>Anas spp., Anser anser, Aythya novaeseelandiae, Botaurus poy-ciloptilus, Branta canadensis, Carduelis chloris, Cygnus atratus, Larus dominicanus, Tadorna variegata,</i>	Anseriformes, Anatidae, Anseriformes, Anatidae, Anseriformes, Anatidae, Pelecaniformes, Ardeidae, Anseriformes, Anatidae, Passeriformes, Fringillidae, Anseriformes, Anatidae, Charadriiformes, Laridae, Anseriformes, Anatidae	New Zealand	McKenna [31]

Table 2: Natural avian hosts of *Echinoparyphium recurvatum* and their geographical location.

References

- Dietz E (1909) Die Echinostomiden der Vogel. ZoologischerAnzeiger Leipzig 6: 180-192.
- Linstow von O (1873) On the development of *Distomum nodulosum*. The Annals and Magazine of Natural History 12: 345-347.
- Huffman JE, Fried B (2012) The biology of *Echinoparyphium* (Trematoda, Echinostomatidae). Acta Parasitologica 57: 199-210.
- Kanev I (1990) A checklist of the helminth parasites of *Echinis, Echinostoma*, Echinostomatidae (Trematoda) with references for their re-naming, replacement and reclassify. Publishing House of the Bulgarian Academy of Sciences, Sofia 1990.
- Roberts TJ (1992) The Birds of Pakistan: Passeriformes. Volume 2. Oxford University Press. Karachi 598.
- Garcia LA, Ash LR (Ed.). (1979) **Diagnostic Parasitology: Clinical laboratory manual**. 2nd Eds. The CV Mosby Company. 11830 Westline Industrial Drive, St Louis, Missouri 63141, USA 174.
- Jones A, Bray RA, Gibson DI (2005) Keys to the Trematoda. Volume 2. CABI Publishing and The Natural History Museum, London, UK 745.
- Yamaguti S (1971) Synopsis of digenetic trematodes of vertebrates. Volume I and II. Keigaku Publishing Co. Tokyo, Japan 1575.
- Sereno-Urbe AL, Pinacho-Pinacho CD, Sanchéz-Cordero V, Garcia-Varela M (2015) Morphological and molecular analyses of larval and adult stages of *Echinoparyphium recurvatum* von Linstow, 1873 (Digenea: Echinostomatidae) from central Mexico. Journal of Helminthology 89: 458-464.
- Looss A (1899) Weitere Beitreige zur Kenntniss der Trematoden-Fauna Aegyptens, zugleich Versuch einer natiiirlichen Gliederung des Genus *Distomum* Retzius. Zoologische Jahrbücher, Abteilung Systematik, 12: 521-784.
- Mouahid A, Moné H (1988) *Echinoparyphium elegans* (Looss, 1899) (Digenea: Echinostomatidae): the life cycle and redescription of the adult with a revision of the 43-spined members of the genus *Echinoparyphium*. Systematic Parasitology 12: 149-157.
- McCarthy AM (1990) Speciation of Echinostomes; evidence for the existence of two sympatric sibling species in the complex *Echinoparyphium recurvatum* Von Linstow, 1873 (Digenea: Echinostomatidae). Parasitology 101: 35-42.
- Kostadinova A, Gibson DI (2000) The systematics of the Echinostomatidae. In: Fried B, Graczyk T (ed.). Echinostomes as experimental

- models for biological research. Kluwer Academic Publishers, Dordrecht, Netherlands 2000: 31-57.
14. Esteban JG, Toledo R, Sanchez L, Muñoz-Antol C (1997) Life-cycle of *Euparyphium albuferensis* n. sp. (Trematoda: Echinostomatidae) from rats in Spain. *Systematic Parasitology* 38: 211-219.
 15. Khan AJ, Khan SW, Riaz S (1983) Helminth parasites of Wild Duck (*Anas crecca*) from N.W.F.P., Peshawar, Pakistan. *Bulletin of Zoology (Peshawar)* 1: 57-62.
 16. Johnston TH, Simpson ER (1944) Larval trematodes from Australian freshwater mollusks. Part IX. Transactions of the Royal Society of South Australia. lxxviii, 125-132.
 17. Kostadinova A, Herniou EA, Barrett J, Littlewood DT (2003) Phylogenetic relationships of *Echinostoma Rudolphi*, 1809 (Digenea: Echinostomatidae) and related genera re-assessed via DNA and morphological analyses. *Systematic Parasitology* 54: 159-176.
 18. Howell MJ (1968) The life-cycle of *Echinoparyphium serratum* sp. nov. (Digenea: Echinostomatidae). *Parasitology* 58: 573-582.
 19. Latham ADM, Poulin R (2002) New records of gastrointestinal helminths from the southern black-backed gull (*Larus dominicanus*) in New Zealand. *New Zealand Journal of Zoology* 29: 253-257.
 20. Lee SH, Sohn WM, Chai JY (1990) *Echinostoma revolutum* and *Echinoparyphium recurvatum* recovered from house rats in Yangyang-gun, Kangwon-do. *Korean Journal of Parasitology* 28: 235-240.
 21. Sohn WM (1998) Life history of *Echinoparyphium recurvatum* (Trematoda: Echinostomatidae) in Korea. *Korean Journal of Parasitology* 36: 91-98.
 22. Pence DB, Young VE, Guthery FS (1980) Helminths of the ringnecked pheasant, *Phasianus colchicus* (Gmelin) (Phasianidae) from the Texas Panhandle. *Proceedings of the Helminthological Society of Washington* 47: 144-147.
 23. McJunkin JW, Applegate RD, Zelmer DA (2003) Enteric helminths of juvenile and adult wild turkeys (*Meleagris gallopavo*) in eastern Kansas. *Avian Diseases* 47: 1481-1485.
 24. Lunaschi LI, Cremonte F, Drago FB (2007) Checklist of digenean parasites of birds from Argentina. *Zootaxa*, 1403: 1-36.
 25. Macko KJ, Hanzelova V, Mackova A (2004) Platyhelminthes of three species of shorebirds (*Charadriiformes*) in the Slovak Republic. *Helminthologia* 41: 151-159.
 26. Sitko J (1998) Trematodes of birds of prey (Falconiformes) in Czech-Republic. *Helminthologia* 35: 131-146.
 27. Hosseini SH, Saifuri P, Eslami A, Nabieian S (2001) Parasitic infections of graylag goose (*Anser anser*) in Gilan province, Iran. *Iranian Journal of Veterinary Medicine* 56: 56-60.
 28. Mehra HR (1982) The fauna of India and the adjacent countries. Platyhelminths. Vol. I. Trematoda. Zoological Survey of India. Eka Press. Calcutta 1982: 65-72.
 29. Fredensborg BL, Latham AD, Poulin R (2004) New records of gastrointestinal helminths from the red-billed gull (*Larus novaehollandiae scopulinus*). *New Zealand Journal of Zoology* 31: 75-80.
 30. Davis NE (2006) A survey of waterfowl for echinostomes and schistosomes from Lake Wanaka and the Waitaki River watershed, New Zealand. *Journal of Helminthology* 80: 33-40.
 31. McKenna PB (2010) An updated checklist of helminth and protozoan parasites of birds in New Zealand. *WebmedCentral Parasitology* 1.