



HAL
open science

Role of an opportunistic pathogen in the decline of stressed oak trees

Benoit Marçais, N. Breda

► **To cite this version:**

Benoit Marçais, N. Breda. Role of an opportunistic pathogen in the decline of stressed oak trees. Journal of Ecology, 2006, 94 (6), pp.1214-1223. 10.1111/j.1365-2745.2006.01173.x . hal-02064818

HAL Id: hal-02064818

<https://hal.science/hal-02064818>

Submitted on 12 Mar 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Role of an opportunistic pathogen in the decline of stressed oak trees

Marçais B.¹ and Bréda N.²

¹ UMR 1136 Interactions arbres/microorganismes, équipe de Pathologie Forestière, INRA-Nancy, 54280 Champenoux, France, marçais@nancy.inra.fr

² UMR 1137 Ecologie et Ecophysiologie Forestières, INRA-Nancy, 54280 Champenoux, France, breda@nancy.inra.fr

SUMMARY

1 The importance of opportunistic pathogens, in particular of *Armillaria* species, in forest decline has often been open to debate.

2 In order to assess the role of *Armillaria gallica* in the decline of oak trees, 60 *Quercus robur* trees with high (HIP trees) or low (LIP trees) level of *A. gallica* inoculum, as measured by the density of epiphytic rhizomorphs on the root collar, were artificially defoliated for two years. Half of the HIP trees were treated when first defoliated with boric acid to reduce the *A. gallica* inoculum potential (BHIP trees). The ability of *in situ* rhizomorphs to colonize plant material was similar for LIP and BHIP, but was lower than in HIP trees, indicating that the boric acid treatment reduced the level of *A. gallica* inoculum.

3 Tree growth was similar between treatments as determined by dendrochronological comparisons. Although defoliation greatly reduced both tree growth and sapwood starch reserves at the beginning of autumn, growth response to defoliation and sapwood starch concentration at the beginning of autumn were similar for LIP, BHIP and HIP trees.

4 HIP trees suffered considerably greater crown deterioration and mortality following defoliation than either BHIP or LIP trees (62 %, 32 % and 5 % mortality rates, respectively). The trees that died had very low sapwood starch concentrations. In addition, at similar levels of sapwood starch, HIP trees were much more likely to die than LIP or BHIP trees.

5 Two other factors influenced tree mortality: past stress that reduced the tree growth a few years prior to the start of the experiment was shown to alter the tree's ability to cope with defoliation. Oak mildew selectively infected the defoliated trees and increased the severity of the defoliation stress.

6 Thus, trees subjected to high level of *A. gallica* inoculum had a lower ability to overcome the defoliation stress. These findings supports the forest decline models developed by Manion (1991) and shows that it is important to take in to account the role of opportunistic pathogens in tree mortality processes.

Key word: forest decline, *Armillaria gallica*, opportunistic pathogens, *Quercus robur*, starch, carbohydrate reserves, defoliation, time-lag effect, community interactions

INTRODUCTION

Tree mortality remains a poorly understood process that is often difficult to predict (Franklin et al, 1987, Pedersen, 1998). Pathogens are important agents of tree mortality that may regulate host demography and strongly alter the structure of plant communities (Hansen, 1999; Gilbert, 2002; Møller, 2005). Several examples of forest declines linked to severe epidemics have been described (Anagnostakis, 1987; Weste and Marks, 1987; Gibbs *et al*, 1999). However, in many cases of forest decline, the role of pathogens is less significant as the process appears to be multifactorial with the involvement of just opportunistic parasites, i.e. organisms unable to colonize a host unless it has been first weakened as a result of another stress. Manion (1991) developed a conceptual model of forest decline that postulates a conjunction of 3 different types of factors that must occur for the onset of a decline: predisposing factors act over the long term to weaken the trees, while inciting factors are short term stresses that trigger the decline, and contributing factors, mostly opportunistic organisms, act on the weakened trees to increase or to speed up the level of decline and mortality. However, the importance of opportunistic organisms in forest decline has been controversial, in part because they tend to be late invaders of trees that are about to die. While they have some times been considered to play an important part in the decline, greatly aggravating the problem (Guillaumin *et al*, 1985; Houston, 1992; Wargo and Harrington, 1991), they have also been considered as minor components of declines which were mainly related to tree aging, location on inadequate sites, or pollution (Becker and Lévy, 1982, Mueller-Dombois, 1992; Landmann *et al*, 1993; Thomas *et al*, 2002, Frey et al, 2004). Nevertheless, forest dieback results from ecosystem processes that both results from and induces community imbalances.

We investigated the decline of oak trees to assess the importance of opportunistic pathogens in forest decline. Oak decline has been an episodic problem in Europe during the 20th century and typically involves the action of several biotic and abiotic factors operating in sequence (Thomas *et al*, 2002). Such declines have been a growing concern, in particular because of the possible relation with climatic change. In oak decline, fungal pathogens colonizing either the root system (*Armillaria* sp.) or the bole bark (*Biscogniauxia mediterranea*) and bark insects such as *Agilus* species have been reported as important opportunistic parasites, contributing to tree mortality. As in most cases of forest decline, the importance of those opportunistic pathogens is controversial (Thomas *et al*, 2002).

A. gallica, one of the most frequently found *Armillaria* species on declining oaks, is a wood decay root rotting fungus and an aggressive colonizer of oak stumps. From colonized wood, it forms in forest soil a network of rhizomorphs, perennial cord-like organs, that are capable of colonizing the tree bases of most oak trees in the forest. Rhizomorphs develop epiphytically on the root collar (Redfern and Filip, 1991, Marçais and Caël, 2006). *A. gallica* is an opportunistic pathogen with a low level of aggressiveness and is unable to colonize vigorously growing hosts (Wargo and Harrington, 1991). However, the fungus often invades trees weakened by insect defoliation or drought. Inoculum potential, defined by Garrett (1956) as a combination of rhizomorph abundance at the host surface and the vitality of those propagules has been considered an important feature determining the ability of *Armillaria* to invade an hosts. The importance of inoculum potential has mainly been documented for aggressive *Armillaria* species. In forests of NW America, *A. ostoyae* was shown to be able to rapidly invade tree stumps created by selective logging from quiescent lesions and subsequently develop an increased inoculum potential in the vicinity, resulting in increased infection in surrounding trees (Cruickshank et al, 1997; Morrison et al 2001). Comparatively, little information exists on the importance of inoculum potential for opportunistic *Armillaria* species such as *A. gallica*.

In healthy mature oaks, the highest concentration of total non-structural carbohydrates are found in the stem, in October, just prior to leaf fall (Barbaroux and Bréda, 2002). Artificial defoliation has been found to induce a reduction in sugars and in amino acids in

the roots of oak seedlings (Parker and Patton, 1975), while the level of starch is affected only when trees are defoliated sufficiently severely to cause refoliation in the same season (Wargo *et al.*, 1972). Several studies have shown that physiological imbalances in starch compared to sugars in declining trees may decrease their resistance to insect (Dunn *et al.*, 1990) and fungal organisms (Wargo, 1981). Renaud and Mauffette (1991) reported that crown dieback in Sugar Maple was associated with reduced concentrations of carbohydrates, and suggested that such an imbalance in sugar/starch compounds may decrease the resistance of the trees to biotic and abiotic stresses. That is, tree carbohydrate reserves may be used as a physiological marker for the trees ability to overcome stress. Nevertheless, no clear threshold of total non-structural carbohydrates leading to a increased tree mortality risk has been established.

Our aim is to determine the effect of *A. gallica* and leaf loss on the survival of young, pedunculate oak trees. This soil pathogen is very common and it is difficult to identify comparable trees colonized or not by epiphytic rhizomorphs. However, *A. gallica* inoculum potential can show very high within-stand heterogeneity levels (Marçais and Caël, 2006) and comparing the response to defoliation of trees subjected to high or low inoculum potential is possible. The hypothesis was that trees with a dense root collar colonization by epiphytic *A. gallica* rhizomorphs would exhibit a greater decline or mortality following stress (i.e. leaf loss), than trees with less root collar rhizomorph colonization. Stress was induced by defoliation and physiological consequences were quantified by determining the end of the season, carbohydrate reserve levels.

MATERIAL AND METHODS

Study plot and experimental design

The study plot was set up in a 20-year-old stand of 8-12 cm diameter at 1.3 m from soil level *Quercus robur*, naturally regenerated in the Champenoux communal forest in NE France. Oaks are the dominant tree species with an understory of *Carpinus betulus*. The soil was homogeneous across the stand consisting of a hydromorphic clay loam with a calcareous clay layer, 30-45 cm below the soil surface. The humus layer was a eutrophic mull (pH 4.7). A previous study (Marçais and Caël, 2006) showed that the *Armillaria gallica* inoculum potential was highly heterogeneous within this stand, presenting an aggregated pattern with a range of 10 m that could be related to the colonization pattern of tree stumps of the previous stand by *Armillaria*.

Fifteen blocks of 5 trees, with approximately 8-12 m between the furthest trees within a block, were selected in the Spring of 2000 (Fig 1). Five blocks consisted of trees selected to have less than 1.2 mg.cm⁻² epiphytic *Armillaria* rhizomorphs on the bark of the collar area (referred to as "low inoculum potential trees - LIP") while 10 blocks consisted of trees selected with a collar density of rhizomorphs between 5.5-9 mg.cm⁻² (referred to as "high inoculum potential trees - HIP"). Rhizomorphs on tree collars were measured using the method of Marçais and Caël, 2006. Briefly, a small section of the collar was exposed and the density of epiphytic rhizomorph on the tree collar estimated by rhizomorph counts on a grid. Five randomly selected blocks containing HIP trees were treated with boric acid at the beginning of July 2000, in order to reduce the *Armillaria* inoculum potential (referred to as the BHIP treatment). The humus layer was brushed away up to 1m from the tree base and a litre of 3 % boric acid sprayed on to the soil (Bauce and Allen, 1992).

Four trees per block were artificially defoliated at the end of June 2000 and again at the beginning of July 2001 to mimic the action of late insect defoliators such as *Lymantria dispar* or *Thaumetopoea procesionea*. The tree crowns were bent to the soil by pulling the upper trunk with a rope and all the leaves were cut at the petiole with scissors. Trees were then lifted to recover their former position. A control tree in each block was bent but not defoliated.




Fig. 1. Experimental design: 5 trees were selected within each of the 15 blocks. The blocks with shading were treated with boric acid to reduce *A. gallica* inoculum potential. Epiphytic *Armillaria* rhizomorph density on tree collar: +, 1.1 mg.cm⁻²; ★, 1.1-2.3 mg.cm⁻²; ▲, 2.3-3.4 mg.cm⁻²; ▲, 3.4-6.0 mg.cm⁻²; ▼, > 6.0 mg.cm⁻².

Measuring tree growth

To determine radial growth, each tree was cored to the pith at 1.3m from soil level (one core per tree) during the winter 2001-2002. Tree radial growth was not observed after 2001 because severe mortality as a consequence of the treatment occurred in the autumn of 2001 and would have led to a strong bias in the growth results. Each early-wood, late-wood and total ring width was measured microscopically to the nearest 0.01 mm. Following the measurements, individual ring-width series were cross-dated to ensure dating accuracy (Becker, 1989). The effect of time, defoliation and *Armillaria* inoculum potential treatment, on past tree growth was analyzed with a mixed model using SAS (SAS/STAT 8.1, SAS Institute Inc., Cary, NC). Blocks were treated as random variables and a first order autoregressive covariance structure was assumed for the ring widths of successive years.

Determination of *Armillaria* inoculum potential

In order to monitor the impact of the boron treatment, colonization rate of wood segments by soil rhizomorphs was determined. Fifteen days following the boron treatment, soil at the collar of each of the 75 trees was removed to expose a *Armillaria* rhizomorph, taking care not to disturb the rhizomorph network. A freshly cut pedunculate oak branch, 3-4 cm in diameter and 15 cm long, was attached to the rhizomorph within 10-20 cm of the collar with a rubber band, and the soil replaced. The colonization of the branch was checked after one year. The wood segments were retrieved and the presence of *Armillaria* mycelial fans, beneath the bark was checked. Whenever *Armillaria* fans were present, a sample was taken to determine the species, using Alu I digested rDNA intergenic spacer profiles (Harrington and Wingfield, 1995). The difference in frequency of branch colonization by *A. gallica* between the 3 treatments was analyzed by logistic regression analysis, including a block effect. The density of epiphytic rhizomorphs on the tree collars was also checked at the end of the experimental period, in the summer of 2003 (Marçais and Caël, 2006).

Measuring crown status and carbohydrate reserve

Twelve defoliated trees, (8 LIP + 4 HIP) and 6 control trees (4 LIP + 2 HIP) were monitored in greater detail. The leaves of 12 defoliated trees were collected during the period of artificial defoliation in June 2000 and July 2001, in order to determine each tree's total leaf area. Individual leaf area was measured from sub-samples from both upper- and lower crown positions (20 sun exposed and 20 shadow leaves) for each defoliated tree using a portable area meter coupled to a transparent belt conveyer (LI-3000A and LI-3050A, Lincoln, Nebraska, USA). Samples were oven-dried for 24 hours at 60°C and the specific leaf area was determined as the ratio of leaf area to dry weight ($\text{cm}^2 \cdot \text{g}^{-1}$). This ratio was used to compute individual tree leaf areas from the leaves dry weight.

The eighteen trees (defoliated + controls) were sampled to determine the concentration of total non-structural carbohydrates (TNC, including starch, glucose, fructose and sucrose) in the sapwood of tree trunks in June 2000 and 2001 prior to defoliation and in October 2000 and 2001 after leaf abscission, but prior to the first frost. These June and October dates correspond to the times of minimum and maximum concentrations in TNC respectively, as determined by seasonal analysis of TNC in oak stems (Barbaroux and Bréda, 2002). Two short cores including the whole sapwood were extracted, one from the base of the stem (from a height of 0-1.3 m) and the other from a major root, frozen and stored at -20°C until freeze-dried. Heartwood was removed from the cores and the sapwood and bark were analyzed together. TNC concentration, i.e. starch and soluble sugars, was enzymatically determined according to Barbaroux and Bréda (2002) and Barbaroux et al. (2003). The effect of time and defoliation TNC concentration was analyzed with a mixed model using the "mixed" procedure of SAS. The significance of TNC evolution between June and October of 2000 and 2001 for both defoliated and control trees was tested with contrasts.

In October 2001, the defoliated trees ($n = 57$) were sampled in the bole and analyzed for TNC concentration as previously described. However, no control trees were sampled at this time. The relationship between the starch concentration in the sapwood in autumn 2001 and the *Armillaria* inoculum potential was analyzed by variance analysis using the SAS "mixed" procedure, introducing the block as a random variable. To control for variation caused by oak mildew attack and past stress events, we introduced mildew severity in 2000 and relative growth reduction in 1995 (RGR95) as covariate. Indeed, a severe oak mildew infection developed on the studied trees in the summer of 2000: following defoliation, newly emerged leaves were infected by oak mildew, *Erisiphe alphitoides*, while spring formed leaves of control trees were not. In addition, a past stress event that had an impact on tree growth in 1995 was detected.

The crown status was monitored several times per year over a period of 4 years. The level of infection of oak mildew in the summer of 2000 was rated as: 1 (no visible leaf necrosis, but presence of infection); 2 (5-33% of leaves with necrosis); 3 (more than 33% leaves with necrosis of leaf margin, but a normal leaf size); and 4 (very severe necrosis on the majority of leaves, with greatly reduced leaf size). Ten leaves from the new foliation were collected in September 2000 to quantify total chlorophyll content using a SPAD-502 chlorophyll meter (Minolta, Japan) and to measure leaf area. The calibration used between transmittance T and chlorophyll content was:

$$[\text{chlorophyll}] (\mu\text{mol} \cdot \text{m}^{-2}) = 0.08 * T^2 + 11.597 - T - 98.548$$

In the spring of each year, the crown was rated as either: 0 (healthy); 1 (moderately declining, with sparse foliage over the entire crown but with no major dead limbs); 2 (severely declining, with sparse foliage and also the death of major limbs); 3 (dead trunk and collar). The difference in crown status between trees with the 3 *Armillaria* inoculum potential treatments (LIP, HIP, BHIP) was analyzed by procedure "genmod" of SAS using a multinomial distribution with the cumulative logit link. Two trees, (one HIP and a BHIP), were damaged as a result of the trunk bending during artificial defoliation in 2001 and were

discarded from further analysis. The trunk sapwood starch concentration, as well as the interaction between trunk sapwood starch concentration and treatment, was introduced as a covariate. Trunk sapwood starch concentration was used as a surrogate for the level of stress the trees sustained. The block factor, nested within treatment was introduced as a fixed effect. Differences between the 3 inoculum potential treatments were tested using contrasts. To determine the possible colonization by *Armillaria*, the collar area and major roots of trees that died was checked by looking for mycelial fans beneath the bark in the cambium area. Whenever suspected *Armillaria* mycelial fans were detected, a sample was taken to determine the *Armillaria* species.

Table 1. Effect of defoliation and *Armillaria* treatments [§] on trees growth (radial increment, mm.year⁻¹).

Effect	Num df ‡	Den df ‡	F value	Pr > F
Year	16	171	50.7	<0.001
Inoculum Potential	2	14.2	0.1	0.908
Year * Inoculum Potential	32	172	0.8	0.827
Defoliation	1	45.4	6.3	0.016
Year*Defoliation	16	171	3.8	<0.001
Inoculum Potential * Defoliation	2	45.3	0.2	0.798
Year * Inoculum Potential * Defoliation	32	172	0.9	0.663

[§] The 3 *Armillaria* inoculum potential treatment are low *A. gallica* IP; high *A. gallica* IP treated with boron; and high *A. gallica* IP not treated with boron.

[‡] Degree freedom of numerator and denominator respectively.

The block effect is specified as a random effect and is thus not included in this table of fixed effect. The between block variance is of 0.023 and is not significant ($z = 1.15$, $p = 0.125$).

RESULTS

Past tree radial growth

There was no difference in past growth or in growth reduction following artificial defoliation between the trees exposed to low/high *A. gallica* inoculum potential or between trees treated/un-treated with boron (Fig. 2, table 1). The growth of un-defoliated or defoliated trees did not significantly differ prior to their defoliation in 2000. The artificial defoliation induced a growth reduction of 38 % in 2000 and 80 % in 2001. In 2001, only early wood, (i.e. only one large vessel layer) was produced by defoliated trees.

In the recent past, the trees experienced a severe reduction in radial growth in 1995-96 which could be correlated to an especially rainy spring; we calculated using a soil water balance model (Granier *et al.*, 1999) and climatic data from a nearby weather station an excess of water in 1995 of 492 mm, as compared to an average value of 340 mm/year for that stand; that putatively induced severe waterlogging above the clay layer. The severity of this growth reduction on individual trees was measured by the relative growth reduction in 1995 ($RGR_{95} = [\text{mean radial growth } 1990-94 - \text{radial growth } 1995] / \text{mean radial growth } 1990-94$). The RGR_{95} was not significantly correlated with epiphytic rhizomorph density at the tree collar ($r = 0.052$, $p = 0.69$). It was slightly higher for BHIP trees (0.64 ± 0.07) than for LIP and HIP trees (0.56 ± 0.04 and 0.58 ± 0.07 , respectively).




Fig. 2. Past radial growth of the studied trees. —○—, HIP trees; —△—, LIP trees; —●—, BHIP trees; —◆—, non defoliated control (HIP, LIP and BHIP). The arrows indicate the years of artificial defoliation. The standard error represented by the bar was estimated from the residual variance of the mixed model.

Impact of boric acid treatment

The boric acid treatment had a significant impact on the epiphytic rhizomorph density. While at the start of the experiment, the initial epiphytic rhizomorph dry weight on tree collars was 1.0 ± 0.2 , 5.4 ± 0.2 , 5.6 ± 0.3 mg.cm⁻² for trees of low inoculum potential (LIP), boron treated high inoculum potential (BHIP) and non treated high inoculum potential (HIP) groups respectively, it was 1.5 ± 0.3 , 1.0 ± 0.2 , 4.3 ± 0.4 mg.cm⁻² respectively, in 2003. In addition, the frequency of branch segment colonization by *Armillaria* was significantly reduced by the boron treatment, with colonization rates respectively of 62.5 %, 52.2 %, and 87% for trees of the LIP, BHIP and HIP groups ($\chi^2 = 6.66$, $p = 0.036$). The *Armillaria* species that colonized the segment was determined in 40 cases and only *A. gallica* was detected.

Refoliation of defoliated trees

The trees' leaf area in June 2000 ranged from 9 to 20 m². After the first defoliation, new leaves developed in approximately 2 weeks. As a consequence to the severe oak mildew infection in 2000, the newly formed leaves had a reduced leaf area (18.6 ± 1.5 cm² for defoliated trees versus 48.9 ± 8.6 cm² for controls, $t = 5.9$, $p < 0.001$) and chlorophyll content (234 ± 12 μmol.m⁻² for defoliated trees versus 448 ± 19 for controls, $t = 19.9$, $p = 0.003$) relative to leaves from trees that were not defoliated. The level of infection was extremely severe on 42% of the trees with almost a second defoliation induced by the oak mildew. There was no significant difference in mildew infection between LIP, BHIP and HIP trees ($\chi^2 = 0.68$, $p = 0.713$). In July 2001, just prior to the second artificial defoliation, the average tree' total leaf area was 5.8 m², compared to 13.5 m² in June 2000, prior to the first defoliation (a reduction of 56 %, paired t-test = 9.5, $p < 0.001$). The impact of this was mainly on the leaf area (paired t-test = 2.4, $p = 0.035$), while the number of leaves per tree decreased slightly, but not significantly. No *E. alphitoïdes* infection occurred on the newly formed leaves following the 2001 defoliation.

Carbohydrate reserve and mortality

Very similar results were obtained for the analysis of trunk and root sapwood starch concentrations; only the results for trunk starch sapwood concentrations are presented. The variance analysis showed that defoliation, time and the defoliation x time interaction all significantly influence the trunk starch sapwood concentrations (result not shown, Fig. 3). While the starch concentration increased from late June 2000 to October 2000 in the control trees ($F = 19.36$, $p < 0.001$), it decreased for the defoliated trees ($F = 178.30$, $p < 0.001$). Similarly, in 2001, the sapwood starch concentration increased between June to October in the control trees ($F = 27.58$, $p < 0.001$) while it remained constant for the defoliated trees ($F = 0.70$, $p = 0.408$). In October 2001, although the defoliated trees had significantly lower starch concentrations in the lower bole sapwood than the non-defoliated controls (Fig. 3), they had similar concentrations of higher soluble sugars.

The starch concentration measured in trunk sapwood in June 2000 were 4.1 ± 1.1 and 3.5 ± 1.6 for LIP and HIP, t -test = 0.71, $p = 0.489$, respectively. In addition, in October 2001, no significant difference in the trunk sapwood starch concentration existed between the 3 treatments, LIP, HIP and BHIP (table 2), although there was a tendency for LIP trees to have higher starch reserves than HIP trees (1.6 ± 0.4 , 1.0 ± 0.4 , 0.8 ± 0.4 respectively for LIP, BHIP and HIP trees). Trees that experienced the highest relative growth reduction in 1995 (RGR95) or that were more severely infected by oak mildew following the 2000 defoliation, had a significantly lower level of starch reserve in the trunk sapwood in October 2001 (table 2). In June 2000, RGR95 and starch concentration of the lower bole sapwood were not significantly correlated ($r = -0.18$, $p = 0.464$).




Fig. 3. Carbohydrate reserves concentration (g/100 g DW) in the lower bole of trees not defoliated (open bars) or defoliated (closed bars). Soluble sugars include glucose, fructose and sucrose.

Table 2. Effect of tree status and history on starch reserves in trunk sapwood (October 2001) of artificially defoliated trees.

Sources	Numerator df	Denominator df	F value	P value
RGR95 [§]	1	39	6.28	0.014
Mildew severity in 2000	1	39	5.05	0.039
Inoculum Potential [†]	2	39	2.36	0.119

[§] Relative growth reduction in 1995 (see section “past tree growth” of the results).

[†] The three inoculum potential treatments are low *A. gallica* IP, high *A. gallica* IP treated with boron and high *A. gallica* IP not treated with boron.

The block effect is specified as a random effect and is thus not included in this table of fixed effect. The between block variance is of 0.03 and is not significant ($z = 0.24$, $p = 0.406$).

Only one tree died in the winter 2000-01, in the HIP treatment. The mortality was greater in 2001. Most of the mortality occurred between October and December (15 trees out of 19), while some trees died in the spring of 2002 with some additional mortality in 2004. The dieback always started in the upper crown and progressed downwards. No *Agrilus bilineatus* colonization was detected in the upper crown; this insect is an opportunistic invader of the bark of stressed oak trees. The last step of the process was the invasion of the collar and lower bole by *Armillaria*. Seventeen of the 19 dead trees were invaded by *Armillaria* and all tested isolates were *A. gallica*. Three trees were uprooted for a more detailed investigation of symptoms. The root system was completely invaded by *Armillaria*. An isolate from all the lesions that could have resulted from independent sources were checked for species identification and only *A. gallica* was detected ($n = 8$).




Fig. 4. Decline status of the trees crown in summer 2004. healthy; moderately declining; severely declining; dead. There was 20 defoliated trees per treatment and all together 15 controls trees (not defoliated). LIP, HIP, low and high *Armillaria* inoculum potential; BHIP, high *Armillaria* inoculum potential treated with boric acid.

The mortality was very different in the 3 treatments, with 62 % mortality for HIP trees, 32 % for BHIP trees and 5 % for LIP trees. The decline status of the crown was also very different between the treatments (Table 3, Fig. 4), the crown of the HIP trees being in a significantly worse status compared to both the LIP and BHIP trees as revealed by contrast analysis (χ^2 of 6.14, $p = 0.013$ and 5.26, $p = 0.022$ respectively), while the LIP and the BHIP trees were not different from each other ($\chi^2 = 0.04$, $p = 0.850$). Trees showing a low level of starch reserves in the trunk sapwood in the autumn of 2001 experienced greater crown deterioration and mortality than trees with a high sapwood starch concentrations (Table 3, Fig. 5). Most of the trees that died had a very low sapwood starch concentration. In addition, trees that experienced severe growth reduction in 1995, i.e. a high RGR95 ($\chi^2 = 4.33$, $p = 0.037$) and trees that were more severely infected by oak mildew following the 2000 defoliation ($\chi^2 = 4.89$, $p = 0.027$), showed a greater deterioration of crown status in 2004 and a higher mortality level. However, when these 2 effects were introduced in a model already containing the trees' trunk sapwood starch level, they did not add significant information (results not shown).

Table 3. Effect of tree history (defoliation and oak mildew infection) on the crown status of the trees in the summer of 2004: result of the multinomial analysis.

Sources			df	Likelihood χ^2	P value
Starch	Sapwood	Concentration (SSC)	1	7.44	0.006
Inoculum Potential §			2	7.83	0.020
SSC * Inoculum Potential			2	0.57	0.751
Block(Inoculum Potential)			12	15.50	0.215

§ The three inoculum potential treatments are low *A. gallica* IP, high *A. gallica* IP treated with boron and high *A. gallica* IP not treated with boron.




Fig. 5. Mortality probability in 2001-04 of artificially defoliated trees and starch concentrations of trunk sapwood in October 2001. —, Low *A. gallica* inoculum potential trees (LIP); , High inoculum potential trees treated with boric acid (BHIP); - - -, High inoculum potential trees not treated with boric acid (HIP). The vertical lines on the x-axis, top and bottom, indicate the starch concentration for trees that died and trees that survived, . The mortality probability is derived from the multinomial analysis ($1-p(\text{CDS} \leq 3)$, with CDS, the crown decline status).

However, one major difference to Pedersen's (1997) model is that the predisposing and inciting stress were not sufficient to induce significant mortality under our conditions. The results indicate that *Armillaria gallica*, although unable to attack vigorous trees and despite a very late intervention in the decline process, was actively involved in the trees decline following defoliation. Trees subjected to a high *A. gallica* inoculum potential, with both a high rhizomorph density on the root collar and the presence of rhizomorphs with a high colonizing capacity, experienced 10 fold increase in post-defoliation mortality compared with trees with a low *A. gallica* inoculum potential. However, as the experiment was conducted in an uncontrolled environment, we cannot rule out interference from undetermined factors that might influence these results. Nevertheless, treating trees exposed to high *A. gallica* inoculum potential with boric acid (BHIP trees) both reduced the inoculum potential and the mortality level, suggesting that the observed lower mortality rate might indeed be caused by the decreased presence of *A. gallica*. The status of the crown in BHIP trees was found to be intermediate between HIP and LIP trees and was linked to an increased level of the 1995 stress with a higher value of RGR95 for these trees, and lower levels of starch reserves in the autumn of 2001. When this was accounted for, the difference in crown status between LIP and BHIP trees was not significant. Possibly, the trees with low starch reserves that were exposed to high *A. gallica* inoculum potential were unable to both defend themselves against the pathogen and mobilize sufficient soluble carbohydrates to adequately harden themselves against the action of frost. In agreement with this hypothesis, it has been demonstrated that trees with very low starch reserves (under 5 mg/g dry weight) are prone to attack by opportunistic organisms (Wargo & Harrington, 1991, Dunn *et al*, 1987). According to the opportunistic pathogen concept, trees subjected to the same level of stress should have a higher likelihood of decline/mortality if they are faced with a high level of *A. gallica* inoculum (Gregory *et al*, 1991). Although we applied a uniform defoliation stress level, the trees exhibited differences as a consequence of past events (RGR95) and an unexpected pathogen, oak mildew, infected the trees with different levels of severity during the course of the experiment. An integrated method to determine the level of stress experienced by the trees is to compare trees that had similar levels of carbohydrate reserves at the end of the 2001 growing season. These results show that for similar levels of carbohydrate reserve, trees supporting high levels of *A. gallica* inoculum indeed experienced a greater decline/mortality.

Infection of new leaves by oak mildew following defoliation appears to have amplified the negative effects of defoliation stress and to have been a significant factor leading to the decline of these trees. Usually, oak mildew attacks are not severe because the pathogen is only able to infect expanding leaves and is not present at the beginning of the season when the foliage of mature oak trees develops. The impact of mildew on trees whose phenology is disturbed by defoliation has previously been documented (Thomas *et al*, 2004); however, its importance may have been under-appreciated. In 2000, many of the trees in this study were nearly defoliated for a second time as a result of the oak mildew infection. The importance of this infection on the new leaves of previously defoliated trees has been previously demonstrated by protecting the refoliation of oak trees (using a chemical treatment) following an infection by *Thaumetopoea processionea* against oak mildew (Marçais, unpublished results). These findings stress the importance of taking into account the interaction between parasites/pathogens. Pathogens that on their own may not have a strong impact can by interacting with other parasites, have a significantly greater impact on their host and on the surrounding plant community. Such interaction between pathogen / parasites were documented for mortality processes leading to either succession or to an altered dominance between two competing plants (De Rooij-Van der Goes, 1995; Holah and Alexander, 1999).

Tree mortality during this study appeared as a complex process fitting Manion's (1991) model of forest decline, i.e. involving the action of several factors acting on different time scales and the interaction of different factors such as defoliation/oak mildew/*Armillaria*.

Attempts to assign the mortality of forest trees to one or more causes might thus be incorrect particularly in the case of decline or background mortality and might explain why this process has often been difficult to predict (Bigler *et al*, 2004, Suarez *et al*, 2004). More generally, it has been recently stressed that more attention should be paid to indirect mechanisms, such as increased predation likelihood, by which parasites can regulate host populations (Møller, 2005). Such an indirect mechanism is reported in this study: *A. gallica* did not have any direct impact on host morbidity, but the pathogen affected the ability to cope with acute stress, such as defoliation. If *A. gallica* has such an impact on a tree's ability to withstand chronic stress as a result of, for example competition, this pathogen could have an impact on tree density, an important forest attribute.

Acknowledgements

The authors are especially grateful to C. Barbaroux, J. Biedermann and L. Lhoste for their contribution to the carbohydrate measurements, to François Géréma who contributed the dendrochronological work; to Olivier Caël who helped in the field work and to Everett Hansen for reviewing the manuscript. Finally, to the members of the forest pathology and phytoecology laboratories who helped with the manual defoliation.

LITERATURE CITED

- Ameglio, T., Ewers, F.W., Cochard, H., Martignac, M., Vandame, M., Bodet, C. & Cruiziat P. (2001) Winter stem xylem pressure in walnut trees: effects of carbohydrates, cooling and freezing. *Tree Physiology*, **21**, 387-394
- Anagnostakis, S. L. (1987). Chestnut blight: the classical problem of an introduced pathogen. *Mycologia* **79**, 23–37
- Barbaroux, C., & Bréda, N. (2002) Contrasting distribution and seasonal dynamics of carbohydrate reserves in stem wood of adult-ring porous sessile oak and diffuse-porous beech trees. *Tree Physiology*, **22**, 1201-1210.
- Barbaroux, C., Bréda, N. & Dufrêne, E. (2003) Distribution of above-ground and below-ground carbohydrate reserves in adult trees of two contrasting broad-leaved species (*Quercus petraea* and *Fagus sylvatica*). *New Phytologist*, **157**, 605-615
- Bauce, E., & Allen, D.C. (1992) Role of *Armillaria calvescens* and *Glycobius speciosus* in a sugar maple decline. *Canadian Journal of Forest Research*, **22**, 549-552.
- Becker, M. (1989) The role of climate on present and past vitality of silver fir forests in the Vosges mountains of northeastern France. *Canadian Journal of Forest Research*, **19**, 1110-1117.
- Becker, M., & Lévy, G. (1982) Le dépérissement du chêne en forêt de Tronçais : les causes écologiques. *Annales des Sciences Forestières*, **39**, 439-444
- Bigler, C., Gričar, J., Bugmann, H. & Čufar K. (2004) Growth patterns as indicators of impending tree death in silver fir. *Forest Ecology and Management*, **199**, 183-190
- Cherubini, P., Fontana, G. Rigling, D., Bobbertin, M. Brang, P., & Innes, J.L. (2002) Tree-life history prior to death: two fungal pathogens affect tree-ring growth differently. *Journal of Ecology*, **90**, 839-850
- Cruickshank M.G., Morrison, D.J. & Punja, Z.K. (1997) Incidence of *Armillaria* species in precommercial thinning stumps and spread of *Armillaria ostoyae* to adjacent Douglas-fir trees. *Canadian Journal of Forest Research*, **27**, 481-490
- De Rooij-Van der Goes PCEM (1995) Then role of plant-parasitic nematodes and soil-borne fungi in the decline of *Ammophila arenaria* (L.) Link. *New Phytologist* **129**, 661-669
- Dunn, J.P., Kimmerer, T.W. & Potter, D.A. (1987) Winter starch reserves of white oaks as a predictor of attack by the two lined chestnut borer, *Agilus bilineatus* (Weber) (*Coleoptera: Buprestidae*). *Oecologia*, **74**, 352-355
- Dunn, J.P., Potter, D.A. & Kimmerer, T.W. (1990) Carbohydrate reserves, radial growth, and mechanisms of resistance of oak trees to phloem-boring insects. *Oecologia*, **83**, 458-468
- Franklin, J.F., Shugart, H.H., & Harmon, M.E. (1987) Tree death as an ecological process. *BioScience*, **27**, 259-288

- Frey, B.R., Lieffers, V.L., Hogg, E.H. & Landhäusser, S.M. (2004) Predicting landscape patterns of aspen dieback: mechanisms and knowledge gaps. *Canadian Journal of Forest Research*, **34**, 1379-1390
- Garrett, S. D. (1956) Rhizomorph behaviour in *Armillaria mellea*, (Vahl) Quél. II. Logistics of infection. *Ann. Bot.* **78**, 193-209.
- Gibbs, J.N., Lipscombe, M.A. & Peace, A.J. (1999) The impact of *Phytophthora* disease on riparian populations of common alder (*Alnus glutinosa*) in southern Britain. *European Journal of Forest Pathology*, **29**, 39-50.
- Gilbert, G. S. (2002). Evolutionary ecology of plant diseases in natural ecosystems. *Annual Review of Phytopathology*, **40**, 13-44
- Granier A., Bréda N., Biron P., Viville S. (1999) A lumped water balance model to evaluate duration and intensity of drought constraints in forest stands. *Ecological Modelling*, **116**, 269-283
- Gregory, R.A., Williams, M.W., Wong, B.L. & Hawley, G.J. (1986) Proposed scenario for dieback and decline of *Acer saccharum* in northeastern USA and southeastern Canada. *IAWA Bulletin*, **7**, 357-369.
- Gregory, S.C., Rishbeth, J. & Shaw, C.G. III (1991) Pathogenicity and virulence. *Armillaria root disease* (eds C.G Shaw III, & G.A. Kile), pp 76-87. Agriculture Handbook No. 691, Forest Service USDA, Washington.
- Guillaumin, J.J., Bernard, C., Delatour, C. & Belgrand, M. (1985) Contribution à l'étude du dépérissement du chêne : pathologie racinaire en forêt de Tronçais. *Annales des Sciences Forestières*, **42**, 1-22.
- Hansen, E. M. (1999) Disease and diversity in forest ecosystems. *Australasian Plant Pathology*, **28**, 313-319
- Harrington, T.C. & Wingfield, B.D. (1995) A PCR-based identification method for species of *Armillaria*. *Mycologia*, **87**, 280-288.
- Holah, J .C. & Alexander, H. M. (1999). Soil pathogenic fungi have the potential to affect the co-existence of two tallgrass prairie species. *Journal of Ecology*, **87**, 598-608
- Houston, D.R. (1992) A host-saprogen model for forest dieback-decline diseases. *Forest decline concepts*, eds P.D. Manion & D. Lachance), pp 3-25. APS Press, St Paul.
- Landmann, G., Becker, M., Delatour, C., Dreyer, E. & Dupouey, J.L. (1993) Oak dieback in France: historical and recent records, possible causes, current investigations. (Ed *Rundgespräche der Kommission für Ökologie*). pp 97-114. Bd 5 'Zustand und Gefährdung der Laubwälder'.
- Manion, P.D. (1991) *Tree disease concepts*. Prentice Hall, Englewood Cliffs, NJ.
- Marçais, B. & Caël, O. (2006) Spatial pattern of *Armillaria* epiphytic rhizomorphs density on the collar of oak trees at the stand level. *Forest Pathology*, **36**, 32-40
- Møller, A.P. (2005) Parasitism and the regulation of host populations. *Parasitism & Ecosystems* (eds F. Thomas, F. Renaud & J.F. Guégan) pp 43-53. Oxford University press, Oxford
- Morrison, D.J., Pellow, K.W., Nemec, A.F.L. & Norris, D.J. (2001) Effects of selective cutting on the epidemiology of *Armillaria* root disease in the southern interior of British Columbia. *Canadian Journal of Forest Research*, **31**, 59-70.
- Mueller-Dombois, D. (1992) A natural dieback theory, cohort senescence as an alternative to the decline disease theory. *Forest decline concepts* (eds P.D. Manion & D. Lachance), pp 26-37. APS Press, St Paul.
- Parker, J. & Patton, R.L. (1975) Effects of repeated defoliation on some metabolites in roots of black oak seedlings. *Canadian Journal of Forest Research*, **5**, 457-463
- Pedersen, B.S. (1997) The role of stress in the mortality of midwestern oaks as indicated by growth prior to death. *Ecology*, **79**, 79-93.
- Pedersen, B.S. (1998) Modeling tree mortality in response to short- and long-term environmental stresses. *Ecological modelling*, **105**, 347-351.
- Redfern, D.B., & Filip, G.M. (1991) Inoculum and Infection. *Armillaria root disease* (eds C.G Shaw III, & G.A. Kile), pp 48-60. Agriculture Handbook No. 691, Forest Service USDA.
- Renaud, J.P. & Mauffette, Y. (1991) The relationships of crown dieback with carbohydrate content & growth of sugar maple (*Acer saccharum*). *Canadian Journal of Forest Research*, **21**, 1111-1118.

- Suarez, M.L., Ghermandi, L. & Kitzberger, T. (2004) Factor predisposing episodic drought-induced tree mortality in *Nothofagus* – site, climate and growth trends. *Journal of Ecology*, **92**, 954-966
- Thomas, F.M., Blank, R. & Hartmann, G. (2002) Abiotic and biotic factors and their interactions as causes of oak decline in Central Europe. *Forest Pathology*, **32**, 277-307.
- Thomas, F.M., Meyer, G. & Popp, M. (2004) Effects of defoliation on the frost hardiness and the concentrations of soluble sugars and cyclitols in the bark tissue of pedunculate oak (*Quercus robur* L.). *Annals of Forest Science*, **61**, 455-463.
- Wargo, P.M. (1981) Defoliation, Dieback and Mortality. *The Gypsy moth: research toward integrated pest management* (eds C.C. Doane & M.M. Mc Manus), pp 240-248. USDA For Ser Tech Bull 1584, Washington.
- Wargo, P.M., Parker, J. & Houston, D.R. (1972) Starch content in roots of defoliated sugar maple. *Forest Science*, **18**, 203-204
- Wargo, P.M., & Harrington, T. C. (1991) Host stress and Susceptibility. *Armillaria root disease* (eds C.G Shaw III, & G.A. Kile), pp 88-101. Agriculture Handbook No. 691, Forest Service USDA, Washington.
- Weste, G. & Marks G. C. (1987) The Biology of *Phytophthora Cinnamomi* in Australasian Forests *Annu. Rev. Phytopathol.* **25**, 207-229