



Multifractal analysis of different hydrological products of X-band radar

I Skouri-Plakali, Igor Paz, A. Ichiba, Auguste Gires, I Tchiguirinskaia, D Schertzer

► To cite this version:

I Skouri-Plakali, Igor Paz, A. Ichiba, Auguste Gires, I Tchiguirinskaia, et al.. Multifractal analysis of different hydrological products of X-band radar. European Geophysics Union (EGU), Apr 2017, Vienne, Austria. 2017. hal-02067081

HAL Id: hal-02067081

<https://hal.science/hal-02067081>

Submitted on 14 Mar 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Introduction.

The project aimed to compare the data between X-band and C-band radar to study the impact of small-scale rainfall variability over the Bièvre catchment (in SW of Paris) which is 40km away from the radar (see I. Paz et al. X4.145 on Fri 28 Apr)^[5]. The purpose of this work is to analyse and inter-compare 5 (where was available) different products of X-band radar with pulsed emissions for simple and dual polarization.

The signal of KDP is noisy so the use of a smoothed signal is used to compute KDP. Two different filters are used: the Finite Impulse Response (FIR) and the median one. The data were processed with standard Rainbow software.

Product	Dual Polarization	Data used for products	Required parameters	Equation used	Filter
DPSRI	✓	dBZ, ZDR, KDP	a=150 b=1.3	$Z=aR^b$	FIR
DPSRI	✓	dBZ, ZDR, KDP	a=200 b=1.6	$Z=aR^b$	FIR
DPSRI-SP	✓	dBZ, ZDR, KDP	a=150 b=1.3	$Z=aR^b$	Median
DPSRI-PS	✓	dBZ, ZDR, KDP	a=150 b=1.3	$Z=aR^b$	FIR
SRI	x	dBZ	a=200 b=1.6	$Z=aR^b$	FIR

Methodology

a. ENPC polarimetric X-band radar

- Oblate drops -> differential phase shift (ΦDP) between the horizontal and vertical axis. The specific differential phase shift (=grad ΦDP) is used for rainfall rate computation.
 - Rainfall estimations
 - Strong KDP ($> 1^\circ/\text{km}$) -> directly $R = c \left(\frac{KDP}{f} \right)^d$
 - Low KDP -> Marshall-Palmer relation $Z = aR^b$

b. Spectral analysis: the first indication of the scaling behaviour of the field

Spectral slope β calculated $E(k) \approx k^{-\beta}$, where
k:wavelength

c. Universal Multifractals parameters fully define statistics across scales^[2,3,4]

- $\alpha \in [0,2]$ (multifractality index): the variability of intermittency with respect to intensity level
 - C_1 (mean intermittency): mean inhomogeneity of the field at large and small scales ($C_1=0$ for homogeneous fields)
 - H : the degree of non-conservation, the scale dependency of the mean field ($H=0$ considered in our case) $\langle \left(R_{\lambda}^{(\eta)} \right)^q \rangle \approx \lambda^{K(q,\eta)}$ $K(q,\eta) = \eta^{\alpha} K(q)$

C_1 and α estimated with the **Double Trace Moment** (DTM) technique [1]

Values in italics correspond to a scale break at 4km, while beige background correspond to a scale break at 2km. The events E7 and E6 are long so in order to calculate the parameters we had to divide them into shorter equal events. The E7 is divided in three and the E6 in two. The first line of every product shows the values for small scales and the second for large scales.


Data

The signal of KDP is noisy so the use of a smoothed signal is used to compute KDP. Two different filters are used: the Finite Impulse Response (FIR) and the median one. The data were processed with standard Rainbow software.

Table 2: Rainfall events and their characteristics

Event	Date	Start time (UTC)	End time (UTC)	Duration (in h)
E1	12-13/09/2015	06:05:00	18:55:00	36.83
E2	12/09/2015	06:05:00	01:15:00	20.25
E3	13/09/2015	05:45:00	18:55:00	13.17
E4	16/09/2015	00:05:00	16:50:00	16.75
E5	05-06/10/2015	09:10:00	16:05:00	30.92
E6	21-23/05/2016	12:25:00	23:50:00	59.42
E7	28-31/05/2016	08:25:00	00:00:00	87.58
E8	17/06/2016	00:00:00	21:15:00	21.25

Table 3: Figures of cumulative rainfall of the selected events as it is presented in the report for Veolia^[6]



Conclusions _

- With the X-band radar, a scale break can be seen in 2km. It means that small scale behaviour is not accessible with C-band radars
 - The scale break is less pronounced for some events (i.e. 28-31/05/2016)
 - SRI seems to produce data with higher homogeneity
 - α stronger for SRI and DPSRI $a=200$ $b=1.6$

Future work

- Analysing more events of lower and higher intensity
 - Comparing with other types of radar data and with rain gauge measurements
 - Use them in urban hydrological models (Multi Hydro) to find out which filter is better to use

Acknowledgements

The authors greatly acknowledge financial support from the Chair “Hydrology for Resilient Cities” (endowed by Veolia) of Ecole des Ponts ParisTech.

References

- [1]Levallée et al. 1993, Nonlinear variability and landscape topography: analysis and simulation. In L. De Cola & N. Lam, eds. *Fractals in geography*. Prentice-Hall, pp. 171–205.
 - [2]Schertzer and Lovejoy 1987, Physical modeling and Analysis of Rain and Clouds by Anisotropic Scaling Multiplicative Processes. *Journal of Geophysical Research, D* 8(8), pp. 9693–9714.
 - [3]Schertzer and Lovejoy 1991, Nonlinear geodynamical variability: multiple singularities, universality and observables. In *Nonlinear variability in Geophysics Scaling and Fratals*, pp. 41-82, Kluwer, Dordrecht
 - [4]Schertzer and Lovejoy 1997, Universal Multifractals do Exist! *Journal of Applied Meteorology*, 36, pp.1296–1303.
 - [5] Paz, I. S. R. et al. (2017). Comparison between rain gauges, C-band and X-band radar data, and their hydrological impacts on a Paris region catchment. *Water Resources Research* (submitted).
 - [6]Paz, I. S.R., Skouri-Plakali I. et al (2016). Comparison of the rainfall data resulting from C-band and X-band measurements to study the impact of small-scale rainfall cariability over the SIAVB catchment

