



Growth potential can affect timing of maturity in a long-lived semelparous fish

K. Yokouchi, Françoise Daverat, N. Fukuda, R. Sudo, M. J. Miller, K. Tsukamoto, Pierre Elie, R. Poole

► To cite this version:

K. Yokouchi, Françoise Daverat, N. Fukuda, R. Sudo, M. J. Miller, et al.. Growth potential can affect timing of maturity in a long-lived semelparous fish. *Biology Letters*, 2018, 14 (7), pp.4. 10.1098/rsbl.2018.0269 . hal-02074327

HAL Id: hal-02074327

<https://hal.science/hal-02074327>

Submitted on 20 Mar 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

BIOLOGY LETTERS

Future growth potential can affect female size at maturation of a long-lived semelparous fish

Journal:	<i>Biology Letters</i>
Manuscript ID	Draft
Article Type:	Research
Date Submitted by the Author:	n/a
Complete List of Authors:	Yokouchi, Kazuki; Irstea Bordeaux; FRA, National Research Institute of Fisheries Science Daverat, Francoise; Irstea Bordeaux Fukuda, Nobuto; FRA, National Research Institute of Fisheries Science Sudo, Ryusuke; FRA, National Research Institute of Aquaculture Miller, Michael; Nihon University Tsukamoto, Katsumi; Nihon University, Elie, Pierre; Irstea Bourdeaux Poole , Russell; Marine Institute
Subject:	Ecology < BIOLOGY
Categories:	Population Ecology
Keywords:	<i>Anguilla anguilla</i>, growth trajectory, otolith, probability of maturation, European eel

SCHOLARONE™
Manuscripts

Future growth potential can affect female size at maturation of a long-lived semelparous fish

Kazuki Yokouchi^{1,2*}, Françoise Daverat¹, Nobuto Fukuda², Ryusuke Sudo³, Michael J. Miller⁴, Katsumi Tsukamoto⁴, Pierre Elie¹, Russell Poole⁵

¹Unit EPBX, IRSTEA, 50 Avenue de Verdun, F-33612 Cestas Cedex, France

²National Research Institute of Fisheries Science, FRA, Yokosuka, 238-0316, Japan

³National Research Institute of Aquaculture, FRA, Minami-ise, 516-0193, Japan

⁴College of Bioresource Sciences, Nihon University, Fujisawa, 252-0880, Japan

⁵Marine Institute, Furnace, Newport, Co. Mayo., Ireland

*kazuki.yokouchi@affrc.gp.jp

Summary

The relationship between female size at maturity and individual growth trajectories of a long-lived semelparous species of fish was investigated using the European eel. A Bayesian model was applied to 338 individual growth trajectories of maturing migration-stage females from France, Ireland, Netherlands and Hungary. This clearly showed that when prospects for further growth were low, the onset of silvering process would be triggered for the eels to leave growth habitats and migrate to the spawning area. Therefore, female eels tended to attain larger body size when growth prospect was high enough to risk spending extra time in their growth habitats.

Keywords: *Anguilla anguilla*; European eel; otolith; growth; probability of maturation

29 1. Introduction

30 For most animals, the set of factors determining the timing of reproduction are
31 important for their fitness. Long-lived species are thought to respond to environmental
32 constraints by adjusting their effort for reproduction [1]. Once an individual is able to
33 mature, thereafter the processes can be controlled by long-term predictive cues such as
34 photoperiod, food supply and temperature. Factors such the frequency of reproduction
35 referred to as iteroparity and semelparity may influence the body size at maturation [2].
36 Age and size at maturation are key life-history traits that affect growth rate, survival and
37 fecundity [3-4].

38 The relation between size and the probability of maturing has been intensively
39 studied using the probabilistic maturation reaction norm (PMRN) [5] with respect to
40 phenotypic changes related to evolutionary responses. The PMRN could be affected
41 by the individual growth histories, but how growth affects size and age at maturation is
42 not fully understood yet [6].

43 The European eel *Anguilla anguilla* is a long-lived semelparous fish species that
44 undertakes exceptionally long catadromous transoceanic migrations both at the larval
45 stage while dispersing to their juvenile growth habitats in rivers, lakes, and lagoons in
46 Europe and again as “silver eels” that mature as they migrate to the spawning area in the
47 Sargasso Sea. Silver eels migrate downstream predominantly during high river-flow
48 periods and new-moon phases [7], and androgen hormones appear to be important to
49 facilitate the initiation of the migrations as the silvering process begins [8]. Large
50 variations in the size and age at silvering of European eels [9] indicate though, that a
51 range of factors affect the timing of the start of the silvering process that differ among
52 individuals or geographic locations. Although it’s possible to evaluate the proximate
53 environmental or endocrine conditions when maturation and the silver eel spawning
54 migrations begin, what triggers the activation of this process itself is not yet known.

55 The 6,000 km or longer migration of European eels requires that enough energy is
56 stored in their bodies to be able to complete the long journey, so there must be a
57 threshold size before eels would start to mature and become silver eels. Maturation of
58 females is thought to be responsible for a size-maximizing strategy [10], because
59 fecundity is entirely determined by body size. Enlargement of the body is
60 advantageous for females, whereas the growth phase over a decade or more can
61 increases the risk of mortality. This tradeoff between size at maturity and growth rate

suggests that females should start the process of silvering when growth is slowing down and no better growth prospects are expected.

We investigated the link between maturation probabilities and the body size of European eels. We tested the hypothesis that the probability to become a maturing silver eel was linked to growth patterns of eels, with poor growth prospects leading to a higher probability of silvering being triggered at smaller sizes.

2. Materials and methods

Data about body size and age of silver-phase female European eels ($n = 338$) were used from Europe (Ireland, France, Netherlands and Hungary) for the years 2000–2007 to overview these traits widely from a population. Data were extracted from EU-FP5th EELREP database (Loire, Ste. Eulalie, Grevelingen, Balaton, Certes, Nive and Rhine river systems) and additional data from the Burrishoole and Corrib rivers were obtained (R. Poole unpublished).

Individual growth trajectories were acquired using the data of otoliths (calcium carbonate structures of the inner ear of fishes). Back-calculation analysis was undertaken using a relationship between body size and otolith radius. Measurements were made of the radius (mm) of the i th annuli (R_i), which is the distance from the otolith mark at recruitment to the continental habitats to the i th annuli, and of the radius (mm) of the otolith at capture (R), which is the distance from the mark at recruitment to the otolith edge. The L_T of the fish at age i yrs (L_i , mm) was estimated using the following formula: $L_i = L_r + (L - L_r) R_i R^{-1}$, where L_r is the mean L_T of glass eels when they recruit to coasts [11], and L is the L_T at capture (mm). Annual body increment (G_i : mm/year) was calculated as $G_i = L_i - L_{i-1}$.

The average growth rate of a particular period (differential of individual size-age relationship) and growth acceleration/deceleration (second-order differential of individual size-age relationship) were selected as explanatory variables. Logistic regression models describing the probability of silvering were set up where we tested if body size and growth histories over the year preceding silvering were a significant proximate cue(s) for silvering. The basic form for these logistic models was $\text{logit}(p) = \log_e[p / (1 - p)] = c_0 + c_1 L_{t,i} + c_2 L'_{t,i} + c_3 L''_{t,i} + \text{Inds}$, where p is the probability of silvering (early maturation), c_0 is a constant, c_1 is the coefficient for the size effect ($L_{t,i}$) at various age t of individual i , c_2 is the coefficient for the individual growth ($L'_{t,i}$) from

age $t-4$ to age t of individual i , c_3 is the coefficient for the acceleration/deceleration of growth ($L''_{t,i}$) between age $t-4 \sim t$ and age $t-5 \sim t-1$ of individual i , and $Inds$ is the random-effect of individuals.

Silvering probability was fitted to a sigmoid curve using the Bernoulli distribution. We deployed constant prior as the priors and hyperpriors and used Markov chain Monte Carlo to draw samples from the distributions of interest. Three chains were initiated at the maximum likelihood estimates and were run for 5000 iterations as a burn in, after which every fifth iteration was recorded to remove autocorrelation, until 1000 samples had been obtained. To assess the significance of the parameters, the estimated probability distributions of parameters were confirmed to not include zero within the range of the distribution. Convergence of the parameters estimated was confirmed by the iteration figures and that R-hat was close to 1. The random effect was assessed by the deviance information criterion (DIC) for the significance in the model. The programs R (R2Winbugs package) and WinBugs were used for the Bayesian analysis.

3. Results

Silver eel females from Ireland, France, Netherlands and Hungary had a wide range of total lengths at silvering that was from 436 to 982 mm, except for one smaller eel that was 377 mm ($n = 338$, figure 1b). The size distribution showed that for the onset of silvering maturation for eels to occur, there must first be an enlargement of the body beyond a certain minimum size, which appears to be about 430 mm, with only one outlier. The ages and growth rates of silver eel females ranged widely from 4 to 44 yrs-old and from 12.1 to 148.2 mm yr⁻¹, respectively. Annual body increment (G_i) of females was relatively stable but with a large variance until age 10, and then it decreased until 20 yrs-old when it stabilized again (figure 1a).

Size ($L_{t,i}$), average growth ($L'_{t,i}$) and the acceleration/deceleration of growth ($L''_{t,i}$) had a significant effect on the probability of silvering for females in the model. The significant improvement of the random-effect model fit was confirmed by the lower DIC (1554) of the model than that of a model having only fixed effects (1619). Convergence of parameters was confirmed visually and the R-hats ranged from 1.02 to 1.15 (table 1). Estimated distributions of parameters $L_{t,i}$, positively, and $L'_{t,i}$ and $L''_{t,i}$ negatively affected the silvering probability of eels (figure 2) and the distributions of parameters did not include zero within the ranges (table 1), indicating that lower growth

128 in recent years and/or a large growth deceleration leads to a higher probability of
129 silvering.

130

131 **4. Discussion**

132 This study demonstrated that when the prospect for further growth in their present
133 habitats was low, eels tended to start the silvering maturation process and leave those
134 growth habitats to start their spawning migrations. This indicates that constant high
135 growth leads to a larger size at silvering, whereas poor or decreasing growth results in
136 eels silvering earlier at smaller sizes. Such a strategy would be adaptive if mortality
137 during the growth phase is low enough to take the risk to spend more time in the growth
138 stage as a long-lived semelparous fish.

139 Our model is in agreement with the dynamics of endocrine factors during silvering.
140 In silver eels, growth-related physiological traits are reduced whereas
141 maturation-related physiological traits are enhanced [12-13]. From an
142 endocrinological point of view, the silvering corresponds to a conversion of the somatic
143 growth mode into a maturation mode. A deceleration of growth may be a cause or a
144 consequence of this conversion.

145 Our results suggested that the silvering size was linked with deceleration of growth
146 in addition to growth rates. In fact, our model is likely to apply well to the case of the
147 north and south distribution areas of eels, such as old larger eels in Ireland and small
148 younger eels in Italian lagoons [14]. Steady growth of larger and older female silver
149 eels in Ireland [15] suggested that the prospect of further stable moderate growth within
150 present habitats could result in the postponement of the onset of silvering. In addition,
151 larger eels might have relatively small risks for staying in northern-latitude
152 low-productivity habitats because they are one of the top predators in their aquatic
153 ecosystems. In the high-productivity ecosystems adjacent to the Mediterranean Sea,
154 eels would face higher mortality compared to that in the North. If the mortality or
155 variability of mortality were relatively high in a region, a trade-off between somatic
156 growth and the risk of dying before reproduction would lead to a female tactic shifting
157 to maturing at the earliest possible opportunity [9]. Traits related to the reproductive
158 migration could be affected by large-scale environmental changes among long-lived
159 migratory species like sea birds and salmon [e.g. 16-17]. For a large panmictic
160 population of eels, silvering being triggered by a reduction of growth is likely an

161 evolutionary adaptation that allows eels to maximize the benefits of inhabiting a wide
162 range of habitats with various levels of growth potential.

163

164 **Acknowledgements**

165 We thank Patrick Lambert and Eric Rochard for their valued suggestions.

166

167 **References**

- 168 1. Erikstad KE, Fauchald P, Tveraa T, Steen H. 1998 On the cost of reproduction in
169 long-lived birds: the influence of environmental variability. *Ecology* **79**, 1781-1788.
170 (doi:10.1890/0012-9658(1998)079[1781:OTCORI]2.0.CO;2)
- 171 2. Kozłowski J, Wieget RG. 1986 Optimal allocation of energy to growth and
172 reproduction. *Theor. Popul. Biol.* **29**, 16-37. (doi:10.1016/0040-5809(86)90003-1)
- 173 3. Roff DA. 1992 *The evolution of life histories: theory and analysis*. Berlin,
174 Germany: Springer.
- 175 4. Stearns SC. 1992 *The evolution of life histories*. New York: Oxford University
176 Press.
- 177 5. Heino M, Dieckmann U, Godø, OR. 2002 Measuring probabilistic reaction norms
178 for age and size at maturation. *Evolution* **56**, 669–678.
179 (doi.org/10.1111/j.0014-3820.2002.tb01378.x)
- 180 6. Kuparinen A, O'Hara RB, Merilä J. 2008 The role of growth history in determining
181 age and size at maturation in exploited fish populations. *Fish Fish.* **9**, 201–207.
182 (doi.org/10.1111/j.1467-2979.2008.00284.x)
- 183 7. Haro A. 2003 Downstream migration of silver-phase anguillid eels. In *Eel Biology*,
184 pp. 215-222. Tokyo, Japan: Springer.
- 185 8. Sudo R, Tsukamoto K. 2015 Migratory restlessness and the role of androgen for
186 increasing behavioral drive in the spawning migration of the Japanese eel. *Sci. Rep.*
187 **5**, 17430. (doi:10.1038/srep17430)
- 188 9. van den Thillart G, Rankin JC, Dufour S. (eds) 2009 *Spawning migration of the*
189 *European eel*. New York, NY: Springer.
- 190 10. Helfman GS, Facey DE, Hales Jr. LS, Bozeman EL. 1987 Reproductive ecology of
191 the American eel. In *Common strategies of anadromous and catadromous fishes*.
192 pp. 42–56. Xx: American Fisheries Society Publication.

- 193 11. Svedäng H, Neuman E, Wickström H. 1996 Maturation patterns in female
194 European eel: age and size at the silver eel stage. *J. Fish Biol.* **48**, 342–351.
195 (doi:10.1111/j.1095-8649.1996.tb01432.x)
- 196 12. Aroua S, Schmitz M, Baloché S, Vidal B, Rousseau K, Dufour, S. 2005 Endocrine
197 evidence that silvering, a secondary metamorphosis in the eel, is a pubertal rather
198 than a metamorphic event. *Neuroendocrinol.* **82**, 221–232.
199 (doi:10.1159/000092642)
- 200 13. van Ginneken V, Durif C, Balm SP, Boot R, Verstegen MWA, Antonissen E, van
201 den Thillart G. 2007 Silvering of European eel (*Anguilla anguilla* L.): seasonal
202 changes of morphological and metabolic parameters. *Anim. Biol.* **57**, 63–77. (doi:
203 10.1163/157075607780002014)
- 204 14. Daverat F et al. 2012 One century of eel growth: changes and implications. *Ecol.*
205 *Freshwat. Fish* **21**, 325–336. (doi:10.1111/j.1600-0633.2011.00541.x)
- 206 15. Poole RW, Reynolds JD. 1996 Growth rate and age at migration of *Anguilla*
207 *anguilla*. *J. Fish Biol.* **48**, 633–642. (doi:10.1111/j.1095-8649.1996.tb01458.x)
- 208 16. Cubaynes S, Doherty PF, Schreiber EA, Gimenez O. 2011 To breed or not to breed:
209 a seabird's response to extreme climatic events. *Biol. Lett.* **7**, 303–306.
210 (doi:10.1098/rsbl.2010.0778)
- 211 17. Kinnison MT, Quinn TP, Unwin MJ. 2011 Correlated contemporary evolution of
212 life history traits in New Zealand Chinook salmon, *Oncorhynchus tshawytscha*.
213 *Heredity* **106**, 448–459. (doi:10.1038/hdy.2010.162)
- 214

215 **Figure captions**

216 **Figure 1.** (a) The estimated mean body increment (growth in 1 year) and standard
217 deviation (bars) and (b) size and age at silvering maturation of individual female
218 European eels.

219
220 **Figure 2.** Probability curves of silvering maturation as a function of body size (total
221 length) in the Bayesian model for (a) recent 5-yrs growth ($L'_{t,i}$) and (b) the deceleration
222 of growth ($L''_{t,i}$) with several values of growth or trends.

223
224 **Table 1.** Parameter estimates with 95% confidence intervals from the best probability
225 of silvering maturation model for female European eels with random effect of
226 individuals.

parameter	estimate \pm SD	95% CI	R hat
body size (L)	0.016 ± 0.003	$0.011 \sim 0.021$	1.15
recent 5-yrs growth (L')	-0.015 ± 0.003	$-0.021 \sim -0.009$	1.13
inclination of growth (L'')	-0.032 ± 0.005	$-0.041 \sim -0.023$	1.02

227
228 **Short title:** Growth prospect affect size to mature

229
230 **Competing interests**

231 We have no competing interests.

232
233 **Authors' contributions**

234 K.Y., F.D., K.T., P.E. and R.P. conceived the idea. K.Y. and F.D. designed the study and
235 conducted data analyses. K.Y., F.D., N.F., R.S., M.J.M. drafted the manuscript. All
236 authors contributed to revise it and gave final approval for publication.

237
238 **Funding**

239 K.Y. was supported by a Research Fellowship for Young Scientists (No. 245842) from
240 the Japan Society for Promotion of Science. K.Y., F.D. and P.E. were supported by the
241 CPER programs of Region Aquitaine, France.




Figure 1. (a) The estimated mean body increment (growth in 1 year) and standard deviation (bars) and (b) size and age at silvering maturation of individual female European eels.

figure 1
156x219mm (300 x 300 DPI)




Figure 2. Probability curves of silvering maturation as a function of body size (total length) in the Bayesian model for (a) recent 5-yrs growth (L' , i) and (b) the deceleration of growth (L'' , i) with several values of growth or trends.

figure 2
 100x83mm (300 x 300 DPI)