



HAL
open science

q-differential equations for q-classical polynomials and q-Jacobi-Stirling numbers

Ana Loureiro, Jiang Zeng

► **To cite this version:**

Ana Loureiro, Jiang Zeng. q-differential equations for q-classical polynomials and q-Jacobi-Stirling numbers. *Mathematical News / Mathematische Nachrichten*, 2016, 289 (5-6), pp.693-717. 10.1002/mana.201400381 . hal-02075487

HAL Id: hal-02075487

<https://hal.science/hal-02075487v1>

Submitted on 6 Feb 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

q -DIFFERENTIAL EQUATIONS FOR q -CLASSICAL POLYNOMIALS AND q -JACOBI-STIRLING NUMBERS

ANA F. LOUREIRO AND JIANG ZENG

ABSTRACT. We introduce, characterise and provide a combinatorial interpretation for the so-called q -Jacobi-Stirling numbers. This study is motivated by their key role in the (reciprocal) expansion of any power of a second order q -differential operator having the q -classical polynomials as eigenfunctions in terms of other even order operators, which we explicitly construct in this work. The results here obtained can be viewed as the q -version of those given by Everitt *et al.* and by the first author, whilst the combinatorics of this new set of numbers is a q -version of the Jacobi-Stirling numbers given by Gelineau and the second author.

1. INTRODUCTION AND MAIN RESULTS

The present paper concerns with the explicit construction of even order q -differential operators having the so-called q -classical polynomials as eigenfunctions. Motivated by this goal, we introduce, develop and give a combinatorial interpretation to the so-called q -Jacobi-Stirling numbers. Among other things, this work consists of a q -version of the studies in [2, 6, 7, 8, 11, 12, 26], while assembling the information in a coherent framework. To our best knowledge the results here presented are new in the theory.

At the centre is the q -derivative operator or *Jackson-derivative*, here denoted as D_q and defined for any polynomial f as follows

$$(D_q f)(x) := \frac{f(qx) - f(x)}{(q-1)x} \quad \text{if } x \neq 0, \quad (1.1)$$

and $(D_q f)(x) = f'(0)$ if $x = 0$, where q is a complex number such that $q \neq 0$ and $|q| \neq 1$. Throughout the text, the symbols j, k, m and n will be allocated for integer values and we will write $n \geq k$ to mean all the integers n greater than or equal to k . As $q \rightarrow 1$, the q -derivative operator D_q reduces to the derivative operator, $D_q \rightarrow D$, and its action on the sequence of monomials $\{x^n\}_{n \geq 0}$ is given by

$$(D_q \zeta^n)(x) = \begin{cases} [n]_q x^{n-1} & \text{if } n \geq 1, \\ 0 & \text{if } n = 0, \end{cases}$$

where $[a]_q := \frac{q^a - 1}{q - 1}$ for any complex number a .

Recalling the classical q -notations found on [10, 20], the q -shifted factorial (also known as the q -Pochhammer symbol) is defined by $(x; q)_k := \prod_{i=0}^{k-1} (1 - xq^i)$, for $k \geq 1$, and $(x; q)_0 := 1$, whilst the q -binomial coefficients are defined by

$$\begin{bmatrix} n \\ k \end{bmatrix}_q = \frac{(q; q)_n}{(q; q)_k (q; q)_{n-k}} \quad \text{if } 0 \leq k \leq n,$$

and $\begin{bmatrix} n \\ k \end{bmatrix}_q = 0$ otherwise. In addition, for practical reasons, we introduce the supplementary notation

$$[a; q]_n := \prod_{k=0}^{n-1} [a+k]_q \quad \text{and} \quad [n]_q! := [1; q]_n.$$

Hence, $[a; q]_n := (1 - q)^{-n} (q^a; q)_n$, for any $n \geq 0$.

Throughout the text, we consider sequences of polynomials of successive degrees starting at zero and leading coefficient equals to one, which we refer to as *monic polynomial sequences* (MPS in short). An MPS forms a basis of the vector space of polynomials with complex coefficients, denoted by \mathcal{P} , and we will mainly deal with the so-called q -classical polynomial sequences. These are monic orthogonal polynomial sequences (soon abbreviated to MOPS) possessing Hahn's property [15], that goes as follows:

Definition 1. An MOPS $\{P_n\}_{n \geq 0}$ is called q -classical whenever the sequence of its q -derivatives, $\{P_n^{[1]}\}_{n \geq 0}$, where $P_n^{[1]}(x) := \frac{1}{[n+1]_q} (D_q P_{n+1})(x)$, is also an MOPS.

Relevant properties about orthogonality and the q -classical character of a sequence are summarised in §3.1, but we highlight the fact that an MOPS $\{P_n\}_{n \geq 0}$ is q -classical if and only if there exist two polynomials Φ (of degree at most 2) and Ψ (of degree 1) such that

$$\mathcal{L}_q[P_n](x) := \left(\Phi(x) D_q \circ D_{q^{-1}} - \Psi(x) D_{q^{-1}} \right) P_n(x) = \chi_n P_n(x), \quad n \geq 0, \quad (1.2)$$

where

$$\chi_n = \begin{cases} -[n]_{q^{-1}} \Psi'(0) & \text{if } \deg \Phi = 0, 1, \\ [n]_{q^{-1}} ([n-1]_q - \Psi'(0)) & \text{if } \deg \Phi = 2, \quad n \geq 0. \end{cases} \quad (1.3)$$

In other words, the elements of a q -classical MOPS are eigenfunctions of the second-order q -differential operator \mathcal{L}_q of q -Sturm-Liouville type (see Proposition 2 for more details).

Regarding their importance in various fields of mathematics, there is a considerable bibliography on the subject – without any attempt for completion, we refer to [5, 10, 16, 17, 19, 20, 22, 23, 33]. In particular, they appear in the literature in the framework of discretisations of hypergeometric second order differential operators in q -lattices (see [20, 34]). Structural properties of such polynomials have been intensively studied in [1, 19, 31] following the program developed for P. Maroni for the classical ones by using an algebraic approach. From an historical point of view, in [15], q -classical orthogonal polynomial sequences appear for the first time as those sequences of orthogonal polynomials $\{p_n\}_{n \geq 0}$ such that the sequence of polynomials $\{q_n(x) := D_q p_{n+1}(x)\}_{n \geq 0}$ is also orthogonal. The q -classical polynomials include the q -polynomials of Al-Salam and Carlitz, the Stieltjes-Wigert, the q -Laguerre, the Little q -Jacobi, and an exhaustive list can be found in Table 1 – see [19] for a detailed survey.

When q tends to 1, we recover the classical polynomials of Hermite, Laguerre, Bessel or Jacobi. These sequences are the only orthogonal polynomial sequences whose elements are eigenfunctions of the second-order linear differential operator $\mathcal{L} := \mathcal{L}_1$, known as the Bochner differential operator, named after [3]. Krall [21] showed that if the elements of a classical polynomial sequence are eigenfunctions of a linear differential operator, then it must be of even order. In fact, as observed in [30, Theorem 1], there exists a polynomial P so that any even order differential operator having classical polynomials as eigenfunctions can be written as $P(\mathcal{L})$. Thus, if $\mathcal{O} = \sum_{k=0}^{2n} a_k(x) D^k$, where $a_k(x)$ are polynomials of degree at most k , then a natural question is to state the connection between the polynomials $a_k(x)$ and the polynomial coefficients Φ and Ψ in the differential operator \mathcal{L} . This question has been addressed in [26] and partly in [7, 8]. More precisely, explicit expressions for any integral power of \mathcal{L} were obtained for each of the classical polynomial families, orthogonal with respect to a positive-definite measure in a series of papers by Everitt *et al.* (see [7, 8] and the references therein). Meanwhile, in [26] a similar study was carried out in a more coherent framework gathering the properties of all the classical families (not restricting to the positive-definite case, and therefore including, for example, the Bessel case). The key to attain such expressions were the Stirling numbers in the study of the cases of Hermite and Laguerre polynomials, whereas the cases of Bessel and Jacobi polynomials required the introduction of a new pair of numbers: the so-called α -modified Stirling numbers $\widehat{s}_\alpha(n, k)$ and $\widehat{S}_\alpha(n, k)$. These are best known as Jacobi-Stirling numbers (as called in [7]), which can be defined as follows

$$\prod_{i=0}^{n-1} (x - i(\alpha + i)) = \sum_{k=0}^n \widehat{s}_\alpha(n, k) x^k \quad \text{and} \quad x^n = \sum_{k=0}^n \widehat{S}_\alpha(n, k) \prod_{i=0}^{k-1} (x - i(\alpha + i)), \quad n \geq 0,$$

where α represents a complex number. Since their introduction, several properties of the Jacobi-Stirling numbers and its companions including combinatorial interpretations have been established, see [2, 6, 11, 12, 24, 32].

In this paper, we obtain explicit expressions for any positive integer power of the q -differential operator \mathcal{L}_q given in (1.2) and the key to achieve this is via the introduction of a new set of numbers, that we call as q -Jacobi-Stirling numbers, to which we obtain several properties as well as a combinatorial interpretation. The results here obtained are the q -version to those in [26] (and also in [7, 8]), as well as to those in [2, 6, 11, 12, 30, 32], since we provide here combinatorial interpretations to the arisen coefficients and eigenvalues. This study has the merit of addressing all the q -classical polynomial sequences as whole in a coherent framework that brings together generalisations of the q -differential equation (1.2) and associated combinatorial interpretations.

Our starting point is Theorem 1, which gives an explicit expression of q -differential operators of arbitrary even order, $\mathcal{L}_{k;q}$ say, having the q -classical polynomials as eigenfunctions. Obviously, this sequence of operators have \mathcal{L}_q as a particular (and simplest) case.

Theorem 1. *Let $\{P_n\}_{n \geq 0}$ be a q -classical MOPS fulfilling (1.2). For any integer $k \geq 0$, the polynomial $y(x) = P_n(x)$ also fulfils*

$$\mathcal{L}_{k;q}[y](x) := \sum_{\nu=0}^k \Lambda_{k,\nu}(x; q) \left(D_{q^{-1}}^{k-\nu} \circ D_q^k y \right) (q^{-\nu} x) = \Xi_n(k; q) y(x), \quad (1.4)$$

where, for each $\nu = 0, 1, \dots, k$,

$$\Lambda_{k,\nu}(x; q) = \frac{q^{-(k-\nu)(\nu+1)}}{[k-\nu]_q!} \left(\prod_{\sigma=1}^{\nu} \chi_{\sigma}^{[k-\sigma]} \right) \left(\prod_{\sigma=0}^{k-\nu-1} q^{-\sigma \deg \Phi} \Phi(q^{\sigma} x) \right) D_q^{k-\nu} P_k(x),$$

with $\chi_n^{[k]} = q^{-k \deg \Phi - 1} [n]_{q^{-1}} \left([n + 2k]_q \frac{\Phi''(0)}{2} + z \right)$, $z = -\left(\frac{\Phi''(0)}{2} + q\Psi'(0) \right)$ and

$$\Xi_n(k; q) = \prod_{\sigma=0}^{k-1} \chi_{n-\sigma}^{[\sigma]} = \begin{cases} \left(-q^{\frac{(k-1)(1-\deg \Phi)}{2}} \Psi'(0) \right)^k \prod_{\sigma=0}^{k-1} ([n]_{q^{-1}} - [\sigma]_{q^{-1}}) & \text{if } \deg \Phi = 0, 1, \\ q^{-\frac{k(k+1)}{2}} \prod_{\sigma=0}^{k-1} ([n]_{q^{-1}} (z + [n]_q) - [\sigma]_{q^{-1}} (z + [\sigma]_q)) & \text{if } \deg \Phi = 2. \end{cases} \quad (1.5)$$

Observe that by taking $q \rightarrow 1$ in Theorem 1, we recover the results obtained in [26, Theorems 2.1 and 2.2].

In some cases, the orthogonality measures associated to certain q -classical sequences $\{P_n\}_{n \geq 0}$ can be expressed via a weight function, so that there exists a function $W_q(x)$ possessing all the necessary properties with support in (a, b) , with $a < b$, possibly infinite, so that

$$\int_a^b P_n(x) P_m(x) W_q(x) dx = N_n \delta_{n,m}, \quad \text{with } N_n \neq 0, n, m \geq 0,$$

where the symbol $\delta_{n,m}$ stands for the *Kronecker delta symbol*. If a q -classical sequence $\{P_n\}_{n \geq 0}$, orthogonal with respect to the weight function W_q , fulfils the q -differential equation (1.2), then $W_q(x)$ satisfies

$$q^{-1} D_{q^{-1}} (\Phi(x) W_q(x)) + \Psi(x) W_q(x) = \lambda g(x), \quad (1.6)$$

where λ is an arbitrary complex number and $g \neq 0$ is a locally integrable function with rapid decay representing the null form, as it is, for instance, the case of the Stieltjes function $g(x) = \exp(-x^{1/4}) \sin(x^{1/4}) \mathbb{1}_{[0, +\infty)}(x)$ where $\mathbb{1}_A(x)$ stands for the characteristic function of the set A : it equals 1 if $x \in A$ and 0 otherwise. (For more details we refer to [19, p.81-82].) If this is the case, a q -self-adjoint version of Theorem 1 is obtained:

Corollary 1. *With the same assumptions as those in Theorem 1, if the q -classical sequence $\{P_n\}_{n \geq 0}$ is orthogonal with respect to the weight function $W_q(x)$, then the operator in (1.4) can be written as*

$$\mathcal{L}_{k;q}[y](x) := q^{-k} (W_q(x))^{-1} D_{q^{-1}}^k \left(\left(\prod_{\sigma=0}^{k-1} q^{-\sigma \deg \Phi} \Phi(q^\sigma x) \right) W_q(x) (D_q^k y(x)) \right), \quad (1.7)$$

for any polynomial y , and has $P_n(x)$ as eigenfunctions.

Further to this, we link integral powers of the operator \mathcal{L}_q in (1.2) with $\mathcal{L}_{k;q}$, via a set of numbers that we introduce and characterise, the so-called q -Jacobi-Stirling numbers (see Section 2), together with the q -Stirling numbers. The characterisation of the q -Jacobi-Stirling numbers in Section 2 includes, among other things, a combinatorial interpretation. The reason for this lies on the fact that the bridge between powers of the operator \mathcal{L}_q with $\mathcal{L}_{k;q}$ is tied to the relation between their corresponding eigenvalues $(\chi_n)^k$ and $\Xi_n(k; q)$, respectively. Observe that χ_n are linear in n and $\Xi_n(k; q)$ are essentially q -shifted factorials when $\deg \Phi$ equals 0 or 1. Hence, the pair of q -Stirling numbers of first and second kind $(c_q(k, j), S_q(k, j))_{j, k \geq 0}$ allows us to bridge them. However, the eigenvalues χ_n become quadratic in n and $\Xi_n(k; q)$ a product of quadratic polynomials in n when $\deg \Phi = 2$. In order to link powers of χ_n to $\Xi_n(k; q)$ we introduce a new pair of numbers – the q -Jacobi-Stirling numbers of first and second kind, denoted as $(Jc_k^j(z; q^{-1}), JS_k^j(z; q^{-1}))_{j, k \geq 0}$, both depending on a complex valued parameter z . With this we are able to show that:

Theorem 2. *Let k be a positive integer and \mathcal{L}_q the linear differential operator defined in (1.2). The identities*

$$\mathcal{L}_q^k[f](x) = \begin{cases} \sum_{j=0}^k S_{q^{-1}}(k, j) q^{(\deg \Phi - 1) \frac{j(j-1)}{2}} (-\Psi'(0))^{k-j} \mathcal{L}_{j;q}[f](x) & \text{if } \deg \Phi = 0, 1, \\ \sum_{j=0}^k JS_k^j(z; q^{-1}) q^{\frac{j(j+1)}{2} - k} \mathcal{L}_{j;q}[f](x) & \text{if } \deg \Phi = 2, \end{cases} \quad (1.8)$$

are valid for any polynomial f , where $\mathcal{L}_{k;q}$ is the operator given in (1.4) and $z = -(1 + q\Psi'(0))$.

Here $\mathcal{L}_q^k[f](x) := \mathcal{L}_q[\mathcal{L}_q^{k-1}[f]](x)$ for $k = 1, 2, \dots$, with the convention $\mathcal{L}_q^0[f](x) := f(x)$. The identities (1.8) can be reversed to obtain the following pair of reciprocal relations.

Corollary 2. *For any integer k and any polynomial f we have*

$$\mathcal{L}_{k;q}[f](x) = \begin{cases} q^{(1 - \deg \Phi) \frac{k(k-1)}{2}} \sum_{j=0}^k c_{q^{-1}}(k, j) (-\Psi'(0))^{k-j} \mathcal{L}_q^j[f](x) & \text{if } \deg \Phi = 0, 1, \\ \sum_{j=0}^k Jc_k^j(z; q^{-1}) q^{j - \frac{k(k+1)}{2}} \mathcal{L}_q^j[f](x) & \text{if } \deg \Phi = 2. \end{cases} \quad (1.9)$$

The outline of the manuscript reads as follows. We begin with the characterisation of the q -Jacobi-Stirling numbers in Section 2, providing their properties in §2.1, which are also set in comparison to those of the q -Stirling numbers and of the Jacobi-Stirling. For this reason, the proofs are either brief or omitted. A combinatorial interpretation is given in §2.2 and, at last, §2.3 is devoted to symmetric generalisations on this new pair of numbers. In Section 3 we revise the core properties of the q -classical polynomials and we prove Theorem 1, Corollary 1, Theorem 2 and Corollary 2 given above. Finally in Section 4, we illustrate these results with a detailed analysis of the cases of the Al-Salam-Carlitz, the Stieltjes-Wigert and the Little q -Jacobi polynomials.

2. THE q -JACOBI-STIRLING NUMBERS

We address the problem of relating powers of a (complex) number x and a corresponding q -factorial, and therefore the introduction of q -analogs of the Stirling numbers along with their modifications (also known as Jacobi-Stirling numbers).

The sequence of factorials $\left\{ \prod_{i=0}^{n-1} (x - [i]_q) \right\}_{n \geq 0}$, alike the sequence of monomials $\{x^n\}_{n \geq 0}$, forms a basis of the vector space of polynomials, here denoted as \mathcal{P} . Thus, there is a pair of sequences of numbers bridging each of these q -factorials of a complex number x and its corresponding powers – these are the so-called q -Stirling numbers, here denoted as $(c_q(n, k), S_q(n, k))_{0 \leq k \leq n}$. Precisely, for each non-negative integer n , the pair $(c_q(n, k), S_q(n, k))_{0 \leq k \leq n}$ (see [13]) is such that

$$\prod_{i=0}^{n-1} (x - [i]_q) = \sum_{k=0}^n (-1)^{n-k} c_q(n, k) x^k, \quad (2.1)$$

$$x^n = \sum_{k=0}^n S_q(n, k) \prod_{i=0}^{k-1} (x - [i]_q). \quad (2.2)$$

A recurrence relation fulfilled by this pair of q -Stirling numbers can be readily deduced from the identity $x \prod_{i=0}^{n-1} (x - [i]_q) = \prod_{i=0}^n (x - [i]_q) + [n]_q \prod_{i=0}^{n-1} (x - [i]_q)$, which goes as follows:

$$c_q(n+1, k+1) = c_q(n, k) + [n]_q c_q(n, k+1), \quad (2.3)$$

$$S_q(n+1, k+1) = S_q(n, k) + [k+1]_q S_q(n, k+1), \quad (2.4)$$

for $0 \leq k \leq n$ with $c_q(n, k) = S_q(n, k) = 0$, if $k \notin \{1, \dots, n\}$, and $c_q(0, 0) = S_q(0, 0) = 1$. Naturally, the standard pair of Stirling numbers is recovered from $(c_q(n, k), S_q(n, k))_{0 \leq k \leq n}$ when $q \rightarrow 1$.

As it will soon become clear the connection between the operators $\mathcal{L}_{k;q}$ and \mathcal{L}_q^k will be transferred to the problem of establishing inverse relations between the following sequence of factorials, depending on the prescribed parameter z (possibly complex),

$$\left\{ \prod_{i=0}^{n-1} (x - [i]_q (z + [i]_{q^{-1}})) \right\}_{n \geq 0} \quad (2.5)$$

and the sequence of monomials. Straightforwardly, the set (2.5) is a basis of \mathcal{P} . For this reason, there exists a pair of coefficients, $(\mathbf{Jc}_n^k(z; q), \mathbf{JS}_n^k(z; q))_{0 \leq k \leq n}$ say, performing the change of basis to $\{z^n\}_{n \geq 0}$. We will refer to this pair of coefficients as the q -Jacobi-Stirling numbers defined through

$$\prod_{i=0}^{n-1} (x - [i]_q (z + [i]_{q^{-1}})) = \sum_{k=0}^n (-1)^{n-k} \mathbf{Jc}_n^k(z; q) x^k, \quad n \geq 0, \quad (2.6)$$

$$x^n = \sum_{k=0}^n \mathbf{JS}_n^k(z; q) \prod_{i=0}^{k-1} (x - [i]_q (z + [i]_{q^{-1}})), \quad n \geq 0. \quad (2.7)$$

Remark 1. A y -version of the q -Stirling numbers was introduced in [18, Definition 3] in order to compute the moments of some rescaled Al-Salam-Chihara polynomials. It is easy to see that the latter generalized Stirling numbers are a rescaled version of the Jacobi-Stirling numbers. For example, the y -version of the q -Stirling numbers of the second kind $S_q(n, k, y)$ are defined by

$$x^n = \sum_{k=0}^n S_q(n, k, y) \prod_{j=0}^{k-1} (x - [j]_q (1 - yq^{-j})). \quad (2.8)$$

It follows from (2.7) and (2.8) that $\mathbf{JS}_n^k(z; q) = S_q(n, k, y^{-1}) y^{n-k} (1 - q^{-1})^k$ with $z = \frac{y-1}{1-q^{-1}}$.

Bearing in mind the identity fulfilled by the factorials in (2.5)

$$\begin{aligned} x \prod_{i=0}^{k-1} (x - [i]_q(z + [i]_{q^{-1}})) &= \prod_{i=0}^k (x - [i]_q(z + [i]_{q^{-1}})) \\ &\quad + [k]_q(z + [k]_{q^{-1}}) \prod_{i=0}^{k-1} (x - [i]_q(z + [i]_{q^{-1}})), \end{aligned}$$

we readily deduce the following triangular recurrence relations fulfilled by these two sets of numbers

$$\mathbf{Jc}_{n+1}^{k+1}(z; q) = \mathbf{Jc}_n^k(z; q) + [n]_q(z + [n]_{q^{-1}}) \mathbf{Jc}_n^{k+1}(z; q), \quad (2.9)$$

$$\mathbf{JS}_{n+1}^{k+1}(z; q) = \mathbf{JS}_n^k(z; q) + [k+1]_q(z + [k+1]_{q^{-1}}) \mathbf{JS}_n^{k+1}(z; q), \quad (2.10)$$

for $0 \leq k \leq n$, with $\mathbf{Jc}_n^k(z; q) = \mathbf{JS}_n^k(z; q) = 0$, if $k \notin \{1, \dots, n\}$, and $\mathbf{Jc}_0^0(z; q) = \mathbf{JS}_0^0(z; q) = 1$.

The pair (2.6)-(2.7) represents a pair of inverse relations and we have

$$\sum_{j \geq 0} \mathbf{Jc}_n^j(z; q) \mathbf{JS}_j^k(z; q) = \sum_{j \geq 0} \mathbf{JS}_n^j(z; q) \mathbf{Jc}_j^k(z; q) = \delta_{n,k}, \quad n, k \geq 0,$$

as it can be derived directly from their definition.

2.1. Properties of the q -Jacobi-Stirling numbers. An explicit expression for the q -Jacobi-Stirling numbers of second kind $\mathbf{Jc}_n^k(z; q)$ can be obtained directly from the Newton interpolation formula for a polynomial f that we next recall.

Lemma 1 (Newton's interpolation formula). *Let b_0, b_1, \dots, b_{n-1} be distinct numbers. Then, for any polynomial f of degree less than or equal to n we have*

$$f(x) = \sum_{j=0}^n \left(\sum_{r=0}^j \frac{f(b_r)}{\prod_{0 \leq k \leq j, k \neq r} (b_r - b_k)} \right) \prod_{i=0}^{j-1} (x - b_i). \quad (2.11)$$

From (2.11), with $f(x) = x^n$ and $b_k = [k]_q([k]_{q^{-1}} + z)$, and (2.7) we derive immediately the following result:

Proposition 1. *For $0 \leq j \leq n$ we have*

$$\mathbf{JS}_n^j(z; q) = \sum_{r=0}^j (-1)^{j-r} \frac{q^{-\binom{r}{2} - r(j-r)} ([r]_q([r]_{q^{-1}} + z))^n}{[r]_q! [j-r]_q! \prod_{0 \leq k \leq j, k \neq r} (z + [k+r]_{q^{-1}})}. \quad (2.12)$$

Remark 2. *By letting $q \rightarrow 1$, the pair $(\mathbf{Jc}_n^k(z; q), \mathbf{JS}_n^k(z; q))_{0 \leq k \leq n}$ reduces to the Jacobi-Stirling numbers in [7, 11, 32] and the relation (2.12) reduces to [7, (4.4)].*

For each integer $k \geq 1$, it follows from the recurrence relation (2.10) that

$$\sum_{n \geq k} \mathbf{JS}_n^k(z; q) x^n = \frac{x}{1 - [k]_q([k]_{q^{-1}} + z)} \sum_{n \geq k-1} \mathbf{JS}_n^{k-1}(z; q) x^n$$

and therefore

$$\sum_{n \geq k} \mathbf{JS}_n^k(z; q) x^n = \prod_{i=1}^k \frac{x}{1 - [i]_q([i]_{q^{-1}} + z)}.$$

Theorem 3. *Let n, k be positive integers with $n \geq k$. The Jacobi-Stirling numbers $JS_n^k(z, q)$ and $Jc_n^k(z, q)$ are polynomials in z of degree $n - k$ with coefficients in $\mathbb{N}[q, q^{-1}]$. Moreover, if*

$$JS_n^k(z; q) = a_{n,k}^{(0)}(q) + a_{n,k}^{(1)}(q)z + \cdots + a_{n,k}^{(n-k)}(q)z^{n-k}, \quad (2.13)$$

$$Jc_n^k(z; q) = b_{n,k}^{(0)}(q) + b_{n,k}^{(1)}(q)z + \cdots + b_{n,k}^{(n-k)}(q)z^{n-k}, \quad (2.14)$$

then

$$a_{n,k}^{(n-k)} = S_q(n, k), \quad b_{n,k}^{(n-k)} = c_q(n, k).$$

Proof. This follows from (2.9) and (2.10) by induction on n . \square

Remarks 1. (1) *The leading coefficient of $JS_n^k(z; q)$ and $Jc_n^k(z; q)$ in (2.13) are the q -Stirling numbers of second kind $S_q(n, k)$ and the q -Stirling numbers of first kind $c_q(n, k)$, respectively. Hence, we have*

$$\lim_{z \rightarrow \infty} \frac{JS_n^k(z; q)}{z^{n-k}} = S_q(n, k) \text{ whilst } \lim_{z \rightarrow \infty} \frac{Jc_n^k(z; q)}{z^{n-k}} = c_q(n, k).$$

(2) *Observe that $x - [i]_{q^{-1}} \left(\frac{1}{q-1} + [i]_q \right) = \frac{q}{q-1} \left(\frac{q-1}{q} x - [i]_q \right)$, for $i = 0, 1, 2, \dots$. Therefore, from the pair of identities (2.1)–(2.2) and (2.6)–(2.7), it follows that*

$$JS_n^k\left(\frac{1}{q-1}; q^{-1}\right) = \left(\frac{q}{q-1}\right)^{n-k} S_q(n, k) \quad \text{and} \quad Jc_n^k\left(\frac{1}{q-1}; q^{-1}\right) = \left(\frac{q}{q-1}\right)^{n-k} c_q(n, k), \quad (2.15)$$

for $k = 0, 1, \dots, n$.

(3) *When $z = 0$, by analogy with the ordinary central factorial numbers (cf. [11]), we can define the corresponding q -Jacobi-Stirling numbers as q -central factorial numbers of even indices :*

$$U_q(n, k) = JS_n^k(0; q), \quad V_q(n, k) = Jc_n^k(0; q). \quad (2.16)$$

Therefore, the following recurrences hold true :

$$U_q(n+1, k+1) = U_q(n, k) + [k+1]_q [k+1]_{q^{-1}} U_q(n, k+1), \quad 0 \leq k \leq n, \quad (2.17)$$

$$V_q(n+1, k+1) = V_q(n, k) + [n]_q [n]_{q^{-1}} V_q(n, k+1), \quad 0 \leq k \leq n, \quad (2.18)$$

with $U_q(n, k) = V_q(n, k) = 0$, if $k \notin \{1, \dots, n\}$, and $U_q(0, 0) = V_q(0, 0) = 1$, $n \geq 0$.

Some value of the q -Jacobi-Stirling numbers of first kind are as follows:

$$Jc_n^1(z; q) = \prod_{k=1}^{n-1} [k]_q (z + [k]_{q^{-1}}), \quad Jc_n^{(n)}(z, q) = 1,$$

$$Jc_3^2(z; q) = (3 + q + q^{-1}) + (2 + q)z,$$

$$Jc_4^2(z; q) = (q^{-3} + 5q^{-2} + 11q^{-1} + q^3 + 11q + 5q^2 + 15)$$

$$+ (2q^3 + 14q + 8q^2 + 2q^{-2} + 7q^{-1} + 15)z + (4q + q^3 + 3 + 3q^2)z^2,$$

$$Jc_4^3(z; q) = (3q^{-1} + 6 + q^2 + 3q + q^{-2}) + (3 + 2q + q^2)z.$$

Some values of the q -Jacobi-Stirling numbers of second kind are as follows:

$$JS_n^1(z; q) = (1 + z)^{n-1}, \quad JS_n^{(n)}(z; q) = 1,$$

$$JS_3^2(z; q) = (3 + q + q^{-1}) + (2 + q)z,$$

$$JS_4^2(z; q) = (9 + q^{-2} + q^2 + 5q + 5q^{-1}) + (11 + 3q^{-1} + 2q^2 + 8q)z + (3q + 3 + q^2)z^2.$$

2.2. Combinatorial interpretation of the q -Jacobi-Stirling numbers. For any positive integer n , we consider the set of two copies of the integers:

$$[n]_2 = \{1_1, 1_2, \dots, n_1, n_2\}.$$

Definition 2. A Jacobi-Stirling k -partition of $[n]_2$ is a partition of $[n]_2$ into $k + 1$ subsets B_0, B_1, \dots, B_k of $[n]_2$ satisfying the following conditions:

- (1) there is a zero block B_0 , which may be empty and cannot contain both copies of any $i \in [n]$,
- (2) $\forall j \in [k]$, each nonzero block B_j is not empty and contains the two copies of its smallest element and does not contain both copies of any other number.

We shall denote the zero block by $\{\dots\}_0$. For example, the partition

$$\pi = \{\{2_2, 5_1\}_0, \{1_1, 1_2, 2_1\}, \{3_1, 3_2, 4_2\}, \{4_1, 5_2\}\}$$

is not a Jacobi-Stirling 3-partition of $[5]_2$, while

$$\pi' = \{\{2_2, 5_1\}_0, \{1_1, 1_2, 2_1\}, \{3_1, 3_2\}, \{4_1, 4_2, 5_2\}\}$$

is a Jacobi-Stirling 3-partition of $[5]_2$. We order the blocks of a partition in increasing order of their minimal elements. By convention, the zero block is at the first position.

Definition 3. An inversion of type 1 of π is a pair (b_1, B_j) , where $b_1 \in B_i$ for some i ($1 \leq i < j$) and $b_1 > c_1$ for some $c_1 \in B_j$. An inversion of type 2 of π is a pair (b_2, B_j) , where $b_2 \in B_i$ for some i ($0 \leq i < j$) and $b_2 > c_2$ for some $c_2 \in B_j$ and $b_1 \notin B_j$, where a_i means integer a with subscript $i = 1, 2$. Let $\text{inv}_i(\pi)$ be the number of inversions of π of type $i = 1, 2$ and set $\text{inv}(\pi) = \text{inv}_2(\pi) - \text{inv}_1(\pi)$. Let $\Pi(n, k, i)$ denote the set of Jacobi-Stirling k -partitions of $[n]_2$ such that the zero-block contains i numbers with subscript 1.

For example, for the Jacobi-Stirling 2-partitions of $[3]_2$ we have the following tableau

JS 2-partitions of $[3]_2$	inv_1	inv_2	inv
$\{\}_0, \{1_1, 1_2, 3_2\}, \{2_1, 2_2, 3_1\}$	0	0	0
$\{\}_0, \{1_1, 1_2, 3_1\}, \{2_1, 2_2, 3_2\}$	1	0	-1
$\{3_2\}_0, \{1_1, 1_2, 3_1\}, \{2_1, 2_2\}$	1	1	0
$\{3_2\}_0, \{1_1, 1_2\}, \{2_1, 2_2, 3_1\}$	0	1	1
$\{2_2\}_0, \{1_1, 1_2, 2_1\}, \{3_1, 3_2\}$	0	0	0
$\{2_1\}_0, \{1_1, 1_2, 2_2\}, \{3_1, 3_2\}$	0	0	0
$\{3_1\}_0, \{1_1, 1_2, 3_2\}, \{2_1, 2_2\}$	0	1	1
$\{3_1\}_0, \{1_1, 1_2\}, \{2_1, 2_2, 3_2\}$	0	0	0

Thus,

$$\sum_{\pi \in \Pi(3,2,0)} q^{\text{inv}(\pi)} = 3 + q + q^{-1} \quad \text{and} \quad \sum_{\pi \in \Pi(3,2,1)} q^{\text{inv}(\pi)} = 2 + q.$$

Theorem 4. For any positive integers n and k and $0 \leq i \leq n - k$ we have

$$a_{n,k}^{(i)}(q) = \sum_{\pi \in \Pi(n,k,i)} q^{\text{inv}(\pi)}.$$

Proof. We prove by induction on $n \geq 1$. The identity is clearly true for $n = 1$. Assume $n > 1$. We divide $\Pi(n, k, i)$ into three subsets as follows.

- $\{n_1, n_2\}$ forms a single block, the enumerative polynomial is $a_{n-1, k-1}^{(i)}(q)$.
- n_1 is in the zero-block and n_2 is in a non zero-block, the enumerative polynomial is $(1 + q + \dots + q^{k-1})a_{n-1, k}^{(i-1)}(q)$.

- n_1 is in a non zero-block B_j , so (n_1, B_i) is an inversion of type 1 for any $i = j + 1, \dots, k$, and n_2 is in any other block B_l ($l \neq j$), so (n_2, B_i) is an inversion for any $i = l + 1, \dots, k$ and $l \neq j$, so the enumerative polynomial is

$$\left(\sum_{j=1}^k q^{-(k-j)} \left(\sum_{l=0}^{j-1} q^{k-l-1} + \sum_{l=j+1}^k q^{k-l} \right) \right) a_{n-1,k}^{(i)}(q) = [k]_q [k]_{q^{-1}} a_{n-1,k}^{(i)}(q).$$

Summing up, we have

$$a_{n,k}^{(i)}(q) = a_{n-1,k-1}^{(i)}(q) + [k]_q a_{n-1,k}^{(i-1)}(q) + [k]_q [k]_{q^{-1}} a_{n-1,k}^{(i)}(q).$$

This is equivalent to (2.10). □

For a permutation σ of $[n]_0 := [n] \cup \{0\}$ (resp. $[n]$) and for $j \in [n]_0$ (resp. $[n]$), denote by $\text{Orb}_\sigma(j) = \{\sigma^\ell(j) : \ell \geq 1\}$ the orbit of j and $\min(\sigma)$ the set of its positive cyclic minima, i.e.,

$$\min(\sigma) = \{j \in [n] : j = \min(\text{Orb}_\sigma(j) \cap [n])\}.$$

Definition 4. Given a word $w = w(1) \dots w(\ell)$ on the finite alphabet $[n]$, a letter $w(j)$ is a record of w if $w(k) > w(j)$ for every $k \in \{1, \dots, j-1\}$. We define $\text{rec}(w)$ to be the number of records of w and $\text{rec}_0(w) = \text{rec}(w) - 1$.

For example, if $w = 574862319$, then the records are 5, 4, 2, 1. Hence $\text{rec}(w) = 4$.

Definition 5. Let $\mathcal{P}(n, k, i)$ be the set of all pairs (σ, τ) such that σ is a permutation of $[n]_0$, τ is a permutation of $[n]$, and both have k cycles.

- i) 1 and 0 are in the same cycle in σ ;
- ii) among their nonzero entries, σ and τ have the same cycle minima;
- iii) $\text{rec}_0(w) = i$, where $w = \sigma(0)\sigma^2(0) \dots \sigma^l(0)$ with $\sigma^{l+1}(0) = 0$.

As Foata and Han [9], we define the B -code of a permutation σ of $[n]$ based on the decomposition of each permutation as a product of disjoint cycles. For a permutation $\sigma = \sigma(1)\sigma(2) \dots \sigma(n)$ and each $i = 1, 2, \dots, n$ let $k := k(i)$ be the smallest integer $k \geq 1$ such that $\sigma^{-k}(i) \leq i$. Then, define

$$\text{B-code}(\sigma) = (b_1, b_2, \dots, b_n) \quad \text{with} \quad b_i := \sigma^{-k(i)}(i) \quad (1 \leq i \leq n).$$

We define the sorting index for permutation σ of $[n]$ by $\text{Sor}(\sigma) = \sum_{i=1}^n (i - b_i)$, while for a permutation σ of $[n]_0$ we define the modified sorting index by $\text{Sor}_0(\sigma) = \sum_{i=1}^n (i - b'_i)$, where $b'_i = b_i$ if $\sigma^{-1}(i) \neq 0$ and $b'_i = i$ if $\sigma^{-1}(i) = 0$. Finally, for any pair (σ, τ) in $\mathcal{P}(n, k, i)$ we define the statistic

$$\text{Sor}(\sigma, \tau) = \text{Sor}(\tau) - \text{Sor}_0(\sigma).$$

Theorem 5. We have $b_{n,k}^{(i)}(q) = \sum_{(\sigma, \tau) \in \mathcal{P}(n, k, i)} q^{\text{Sor}(\sigma, \tau)}$.

Proof. We proceed by induction on $n \geq 1$. The case $n = 1$ is clear. Assume that $n > 1$. We divide $\mathcal{P}(n, k, i)$ into three parts:

- (i) the pairs (σ, τ) such that $\sigma^{-1}(n) = n$. Then n forms a cycle in both σ and τ and the enumerative polynomial is clearly $b_{n-1,k-1}^{(i)}(q)$.
- (ii) the pairs (σ, τ) such that $\sigma^{-1}(n) = 0$. We can construct such pairs starting from a pair (σ', τ') in $\mathcal{P}(n-1, k, i-1)$ as follows: we insert n in σ' as image of 0, i.e., $\sigma(0) = n$, $\sigma(n) = \sigma'(0)$ and $\sigma(i) = \sigma'(i)$ for $i \neq 0, n$, and then we insert n in τ' by choosing an $j \in [n]$ and define $\tau(j) = n$, $\tau(n) = \tau(j)$ and $\tau(l) = \tau'(l)$ for $l \neq j, n$. Clearly, the enumerative polynomial is $[n-1]_q b_{n-1,k}^{(i-1)}(q)$.

- (iii) the pairs (σ, τ) such that $\sigma^{-1}(n) \notin \{0, n\}$. We can construct such pairs by first choosing an ordered pair (σ', τ') in $\mathcal{P}(n-1, k, i)$ and then inserting n in σ' and τ' , respectively. Clearly, the corresponding enumerative polynomial is given by $[n-1]_q [n-1]_{q^{-1}} b_{n-1, k}^{(i)}(q)$.

Summing up, we get the following equation:

$$b_{n, k}^{(i)}(q) = b_{n-1, k-1}^{(i)}(q) + [n-1]_q b_{n-1, k}^{(i-1)}(q) + [n-1]_q [n-1]_{q^{-1}} b_{n-1, k}^{(i)}(q). \quad (2.19)$$

By (2.6) and (2.14), it is easy to see that the coefficients $b_{n, k}^{(i)}(q)$ satisfy the same recurrence. \square

For example, for $n = 3$ and $k = 2$ the pairs (σ, τ) and the associated statistics are as follows:

(σ, τ)	$\text{rec}_0(\sigma)$	B_0 -code σ	B -code τ	$\text{Sor}(\tau)$	$\text{Sor}_0(\sigma)$	$\text{Sor}(\sigma, \tau)$
$(01)(23), (1)(23)$	0	$(1, 2, 2)$	$(1, 2, 2)$	1	1	0
$(01)(23), (13)(2)$	0	$(1, 2, 2)$	$(1, 2, 1)$	2	1	1
$(012)(3), (12)(3)$	0	$(1, 1, 3)$	$(1, 1, 3)$	1	1	0
$(013)(2), (13)(2)$	0	$(1, 2, 1)$	$(1, 2, 1)$	2	2	0
$(013)(2), (1)(23)$	0	$(1, 2, 1)$	$(1, 2, 2)$	1	2	-1
$(031)(2), (1)(23)$	1	$(0, 2, 3)$	$(1, 2, 2)$	1	1	0
$(031)(2), (1)(23)$	1	$(0, 2, 3)$	$(1, 1, 3)$	1	1	0
$(021)(3), (1)(23)$	1	$(0, 2, 3)$	$(1, 2, 1)$	2	1	1

Thus,

$$\sum_{(\sigma, \tau) \in \mathcal{P}(3, 2, 0)} q^{\text{Sor}(\sigma, \tau)} = 3 + q + q^{-1}, \quad \sum_{(\sigma, \tau) \in \mathcal{P}(3, 2, 1)} q^{\text{Sor}(\sigma, \tau)} = 2 + q.$$

2.3. A symmetric generalisation of Jacobi-Stirling numbers. As suggested by Richard Askey (private communication in 2013), it is natural to consider the pair of connection coefficients $\{(S_{z, w}(n, k), s_{z, w}(n, k))\}_{n \geq k \geq 0}$ satisfying

$$x^n = \sum_{k=0}^n S_{z, w}(n, k) \prod_{i=0}^{k-1} (x - (i+z)(i+w)),$$

$$\prod_{i=0}^{n-1} (x - (i+z)(i+w)) = \sum_{k=0}^n s_{z, w}(n, k) x^k.$$

It is readily seen that we have the following recurrence relation for $0 \leq k \leq n$

$$S_{z, w}(n+1, k+1) = S_{z, w}(n, k) + (z+k+1)(w+k+1)S_{z, w}(n, k+1),$$

$$s_{z, w}(n+1, k+1) = s_{z, w}(n, k) - (z+n)(w+n)s_{z, w}(n, k+1),$$

with $S_{z, w}(n, k) = s_{z, w}(n, k) = 0$, if $k \notin \{1, \dots, n\}$, and $S_{z, w}(0, 0) = s_{z, w}(0, 0) = 1$.

It is clear that $S_{z, w}(n, k)$ are symmetric polynomials in z and w with non negative integer coefficients and

$$\sum_{n \geq k} S_{z, w}(n, k) x^n = \prod_{i=1}^k \frac{x}{1 - (i+z)(i+w)x}.$$

When $w = 0$ these numbers reduce to Jacobi-Stirling numbers. Mimicking the arguments in [11], we can prove similar results for the symmetric variants. Hence, we will just state the combinatorial interpretations of these numbers and omit the proofs.

Definition 6. A double signed k -partition of $[n]_2 = \{1_1, 1_2, \dots, n_1, n_2\}$ is a partition of $[n]_2$ into $k+2$ subsets $(B_0, B'_0, B_1, \dots, B_k)$ such that

- (1) there are two distinguishable zero blocks B_0 and B'_0 , any of which may be empty;

- (2) there are k indistinguishable nonzero blocks, all nonempty, each of which contains both copies of its smallest element and does not contain both copies of any other number;
- (3) each zero block does not contain both copies of any number and B'_0 may contain only numbers with subscript 2.

Let $\Pi(n, k)$ be the set of double signed k -partitions of $[n]_2$. For $\pi \in \Pi(n, k)$ denote by $s(\pi)$ (resp. $t(\pi)$) the number of integers with subscript 1 (reps. 2) in B_0 (reps. B'_0) of π .

Theorem 6. *The polynomial $S_{z,w}(n, k)$ is the enumerative polynomial of $\Pi(n, k)$ with z enumerating the numbers with subscript 1 in B_0 and w enumerating the numbers with subscript 2 in B'_0 , i.e.,*

$$S_{z,w}(n, k) = \sum_{\pi \in \Pi(n, k)} z^{s(\pi)} w^{t(\pi)}.$$

For example, all the double signed k -partitions of $[2]_2$ ($1 \leq k \leq 2$) with the corresponding weight are

- $k = 0$: $\pi = \{\{1_1, 2_1\}_0, \{1_2, 2_2\}'_0\}$; with weight $z^2 w^2$;
- $k = 1$: $\pi_1 = \{\{2_2\}_0, \{1\}'_0, \{1_1, 1_2, 2_1\}\}$ with weight 1;
- $\pi_2 = \{\{2_1\}_0, \{1\}'_0, \{1_1, 1_2, 2_2\}\}$ with weight z ;
- $\pi_3 = \{\{1\}_0, \{2_2\}'_0, \{1_1, 1_2, 2_1\}\}$ with weight w ;
- $\pi_4 = \{\{1_1\}_0, \{1_2\}'_0, \{2_1, 2_2\}\}$ with weight zw ;
- $\pi_5 = \{\{2_1\}_0, \{2_2\}'_0, \{1_1, 1_2\}\}$ with weight zw .
- $k = 2$: $\pi = \{\{\}_0, \{1\}'_0, \{1_1, 1_2\}, \{2_1, 2_2\}\}$ with weight 1.

Thus, by the Theorem 6, we have

$$x^2 = S_{z,w}(2, 0) + S_{z,w}(2, 1)(x - zw) + S_{z,w}(2, 2)(x - zw)(x - (z + 1)(w + 1)),$$

where $S_{z,w}(2, 0) = z^2 w^2$, $S_{z,w}(2, 1) = 1 + z + w + 2zw$, $S_{z,w}(2, 2) = 1$.

Definition 7. *Let $\mathcal{P}_0(n, k)$ be the set of all pairs (σ, τ) of permutations of $[n]_0$ such that σ and τ both have k cycles and*

- i) *1 and 0 are in the same cycle in σ and τ .*
- ii) *Among their nonzero entries, σ and τ have the same set of cycle minima.*

Let $rec_0(w_\sigma)$ be the number of left-to right minima of the word $w_\sigma = \sigma(0)\sigma^2(0) \dots \sigma^l(0)$ with $\sigma^{l+1}(0) = 0$.

Theorem 7. *The polynomial $(-1)^{n-k} s_{z,w}(n, k)$ is the enumerative polynomial of pairs (σ, τ) in $\mathcal{P}_0(n, k)$ with z enumerating the $rec_0(w_\sigma)$ and w enumerating $rec_0(w_\tau)$, i.e.,*

$$(-1)^{n-k} s_{z,w}(n, k) = \sum_{(\sigma, \tau) \in \mathcal{P}_0(n, k)} z^{rec_0(w_\sigma)} w^{rec_0(w_\tau)}.$$

Let $(a)_n = a(a+1) \dots (a+n-1)$ for $n \geq 1$ and $(a)_0 = 1$. The Wilson polynomials (see [20]) are more natural and simply expanded in the basis of polynomials in x^2 given by $\{(a + ix)_n (a - ix)_n\}_{n \geq 0}$, where $i^2 = -1$. So, it is legitimate to refer to the pair of set of numbers $\{(w(n, k), W(n, k))\}_{0 \leq k \leq n}$ performing the following change of basis

$$(a + ix)_n (a - ix)_n = \sum_{k=0}^n w(n, k) x^{2k}, \tag{2.20}$$

$$x^{2n} = \sum_{k=0}^n W(n, k) (a + ix)_k (a - ix)_k, \tag{2.21}$$

as the *Wilson numbers of first and second kind*. Then, by Newton's interpolation formula we have

$$w(n, k) = e_{n-k}(a^2, (a+1)^2, \dots, (a+n-1)^2), \quad (2.22)$$

$$W(n, k) = \sum_{r=0}^k \frac{(-1)^{n-r}(a+r)^{2n}}{r!(k-r)!(2a+r)_r(2a+2r+1)_{k-r}}, \quad (2.23)$$

where $e_k(x_1, \dots, x_n)$ denotes the k -th elementary symmetric polynomial of x_1, \dots, x_n . It is readily seen that

$$W(n+1, k+1) = W(n, k) - (a+k+1)^2 W(n, k+1), \quad (2.24)$$

$$w(n+1, k+1) = w(n, k) + (a+n)^2 w(n, k+1), \quad (2.25)$$

with $W(n, k) = w(n, k) = 0$, if $k \notin \{1, \dots, n\}$, and $W(0, 0) = w(0, 0) = 1$, $n \geq 0$.

It is easy to derive from Theorems 6 and 7 that

$$(-1)^{n-k} W(n, k) = \sum_{\pi \in \Pi(n, k)} a^{s(\pi)+t(\pi)}, \quad (2.26)$$

$$s_{z,w}(n, k) = \sum_{(\sigma, \tau) \in \mathcal{P}_0(n, k)} a^{\text{reco}(w_\sigma) + \text{reco}(w_\tau)}. \quad (2.27)$$

Remark 3. *It is worth to notice that the change of basis $\{(a+ix)_n(a-ix)_n\}_{n \geq 0}$ and $\{(x^2 + (a-1)^2)^n\}_{n \geq 0}$ is performed by the aforementioned pair of Jacobi-Stirling numbers $\{((-1)^{n+k} j s_{n+1}^{k+1}(2\alpha), JS_{n+1}^{k+1}(2\alpha))\}_{0 \leq k \leq n}$, see [25, Remark 3.9]. Indeed, the latter pair of numbers is associated to the linear change of basis in the vector space of polynomials in the variable x^2 performed by the Kontorovich-Lebedev integral transform after a slight modification of the kernel, as described in [25].*

Of course, we can formally work out a q -version of the numbers $S_{z,w}(n, k)$ and $s_{z,w}(n, k)$ by mimicking our approach to the q -Jacobi-Stirling numbers. However, from analytical point of view, an appropriate q -version of Wilson numbers should be the connection coefficients between the basis $\{x^n\}_{n \geq 0}$ and the Askey-Wilson basis $\{(az, a/z; q)_n\}_{n \geq 0}$ with $x = (z + 1/z)/2$ [17, Chap. 15 and 16] and $(a, b; q)_n = (a; q)_n (b; q)_n$. Thus, the Askey-Wilson numbers of the second and first kind may be defined by

$$x^n = \sum_{k=0}^n W_q(n, k) (az, a/z; q)_k, \quad (2.28)$$

$$(az, a/z; q)_n = \sum_{k=0}^n w_q(n, k) x^k, \quad (2.29)$$

where $x = (z + 1/z)/2$. By Newton's formula we have

$$W_q(n, k) = \frac{1}{2^n} \sum_{j=0}^k q^{k-j^2} a^{-2j} \frac{(q^j a + q^{-j}/a)^n}{(q, q^{-2j+1}/a^2; q)_j (q, q^{2j+1} a^2; q)_{k-j}}. \quad (2.30)$$

Since $(az, a/z; q)_n = \prod_{k=0}^{n-1} (1 - 2axq^k + a^2q^{2k})$, we derive from (2.29) that

$$w_q(n, k) = (-1)^k (2a)^n q^{\binom{n}{2}} e_{n-k} \left(\frac{a^{-1} + a}{2}, \frac{(aq)^{-1} + aq}{2}, \dots, \frac{(aq^{n-1})^{-1} + aq^{n-1}}{2} \right). \quad (2.31)$$

The following recurrence relations follow easily from (2.28) and (2.29):

$$W_q(n+1, k+1) = -\frac{1}{2aq^k} W_q(n, k) + \frac{1+a^2q^{2k}}{2aq^k} W_q(n, k+1), \quad (2.32)$$

$$w_q(n+1, k+1) = -2aq^n w_q(n, k) + (1+a^2q^{2n}) w_q(n, k+1). \quad (2.33)$$

Let $T_q(n, k) = (-2a)^k q^{\binom{k}{2}} W_q(n, k)$. Then $T_q(n, k)$ are polynomials in a and q with nonnegative coefficients satisfying the recurrence

$$T_q(n+1, k+1) = T_q(n, k) + (1 + a^2 q^{2k}) T_q(n, k+1). \quad (2.34)$$

Though these numbers should be interesting because they are related to the *Askey-Wilson operator*, their combinatorics seem to be quite different from that of Wilson numbers or q -Jacobi-Stirling numbers. We plan to investigate this topic in a future work.

3. EVEN ORDER q -DIFFERENTIAL EQUATIONS FULFILLED BY THE q -CLASSICAL POLYNOMIALS

To come up with the explicit expressions of $\mathcal{L}_{k;q}$ given in Theorem 1 we do not make use of the analytical properties of the measures of each of the q -classical sequences, nor of the properties of the basic hypergeometric corresponding series. (These two approaches would require to split the analysis for each of the q -classical sequences). Rather, we base our study on the algebraic approach, within the principles developed in [28] and in the survey [19]. Here, the information on the orthogonality measures is hidden in the corresponding linear functionals. This methodology has the merit of gathering all the properties in coherent and consistent framework amongst all the possible q -classical cases. The q -classical sequences and the corresponding q -classical operators will be treated as whole. Based on the properties of the corresponding eigenvalues, there will be essentially two cases to single out: depending on whether the degree of Φ is less than or equal to 2. Whenever the q -classical linear functional admits an integral representation via a weight function, we are able to write $\mathcal{L}_{k;q}$ in a q -self adjoint manner, as in Corollary 1. Therefore, we start with preliminary results on the q -classical sequences to the sequel.

3.1. Background. The orthogonality measures will be represented by a moment linear functional that satisfies certain properties. The moment linear functionals (also called *forms*, see for instance [28]) are elements of \mathcal{P}' , the dual space of the vector space of polynomials \mathcal{P} . We adopt the standard notation of $\langle u, f \rangle$ to consider the action of $u \in \mathcal{P}'$ over $f \in \mathcal{P}$. The sequence of moments of $u \in \mathcal{P}'$ results from its action on the canonical sequence $\{x^n\}_{n \geq 0}$ and will be denoted as $(u)_n := \langle u, x^n \rangle, n \geq 0$. Any linear operator $T : \mathcal{P} \rightarrow \mathcal{P}$ has a transpose ${}^t T : \mathcal{P}' \rightarrow \mathcal{P}'$ defined by $\langle {}^t T(u), f \rangle = \langle u, T(f) \rangle, u \in \mathcal{P}', f \in \mathcal{P}$. For example, for any nonzero complex number a , any linear functional u and any polynomial g , let $D_q u$ and $g u$ be the forms defined as usual by $\langle g u, f \rangle := \langle u, g f \rangle, \langle h_a u, f \rangle := \langle u, h_a f \rangle = \langle u, f(ax) \rangle$ and

$$\langle D_q u, f \rangle := -\langle u, D_q f \rangle \text{ for any } f \in \mathcal{P}. \quad (3.1)$$

Thus, with some abuse of notation, D_q acting on linear functionals is minus the transpose of D_q on polynomials: ${}^t D_q := -D_q$ (see [19, 28] for more details). For any $f, g \in \mathcal{P}$ and $u \in \mathcal{P}'$, the following properties hold:

$$D_q(fg)(x) = f(qx)(D_q g)(x) + (D_q f)(x)g(x), \quad (3.2)$$

$$D_q(f(x)u) = f(q^{-1}x)(D_q u) + q^{-1}(D_{q^{-1}} f)(x)u. \quad (3.3)$$

Such properties can be generalised, so that from (3.2) we obtain a q -Leibniz formula [15] for the product of two polynomials

$$D_q^n(fg)(x) = \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q (D_q^{n-k} f)(q^k x) (D_q^k g)(x), \quad n \geq 0, \quad (3.4)$$

whilst, from (3.3) we deduce the n th order q -derivative of the product of a polynomial f by a linear functional u :

$$D_q^n(fu) = \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_{q^{-1}} q^{-(n-k)} (D_{q^{-1}}^{n-k} f)(q^{-k} x) (D_q^k u), \quad n \geq 0. \quad (3.5)$$

The dual sequence $\{u_n\}_{n \geq 0}$ of a given MPS $\{P_n\}_{n \geq 0}$ is a subset of \mathcal{P}' whose elements are uniquely defined by $\langle u_n, P_k \rangle := \delta_{n,k}, n, k \geq 0$, where $\delta_{n,k}$ represents the *Kronecker delta* symbol. Any element u of \mathcal{P}' can be

written in a series of any dual sequence $\{u_n\}_{n \geq 0}$ of a MPS $\{P_n\}_{n \geq 0}$ [28]:

$$u = \sum_{n \geq 0} \langle u, P_n \rangle u_n. \quad (3.6)$$

Given a MPS $\{P_n\}_{n \geq 0}$, let us consider the MPS of its k th q -derivatives, $\{P_n^{[k]}\}_{n \geq 0}$, defined by

$$P_n^{[k]}(x) := \frac{1}{[n+1; q]_k} D_q^k P_{n+k}(x), \quad n \geq 0. \quad (3.7)$$

This definition readily implies a relation between the corresponding dual sequences. Precisely, for any integers j and k , with $0 \leq j \leq k$, the dual sequence $\{u_n^{[k]}\}_{n \geq 0}$ of $\{P_n^{[k]}\}_{n \geq 0}$ and the dual sequence $\{u_n^{[k-j]}\}_{n \geq 0}$ of $\{P_n^{[k-j]}\}_{n \geq 0}$ are related by

$$D_q^j(u_n^{[k]}) = (-1)^j [n+1; q]_j u_{n+j}^{[k-j]}, \quad n \geq 0. \quad (3.8)$$

Whenever a form $u \in \mathcal{P}'$ is such that there exists a PS $\{P_n\}_{n \geq 0}$ so that $\langle u, P_n P_m \rangle = k_n \delta_{n,m}$ with $k_n \neq 0$, for $n, m \geq 0$, then u is called a *regular* form [28, 29]. The PS $\{P_n\}_{n \geq 0}$ is then said to be orthogonal with respect to u and we can assume the system (of orthogonal polynomials) to be monic and the original form u is proportional to u_0 . There is a unique MOPS $\{P_n(x)\}_{n \geq 0}$ with respect to the regular form u_0 and it can be characterised by the (popular) second order recurrence relation (see, for instance, [4])

$$\begin{cases} P_0(x) = 1; & P_1(x) = x - \beta_0, \\ P_{n+2}(x) = (x - \beta_{n+1})P_{n+1}(x) - \gamma_{n+1}P_n(x), & n \geq 0, \end{cases} \quad (3.9)$$

where $\beta_n = \frac{\langle u_0, x P_n^2 \rangle}{\langle u_0, P_n^2 \rangle}$ and $\gamma_{n+1} = \frac{\langle u_0, P_{n+1}^2 \rangle}{\langle u_0, P_n^2 \rangle}$ for all $n \geq 0$. In this case, the elements of the corresponding dual sequence $\{u_n\}_{n \geq 0}$ are given by

$$u_n = (\langle u_0, P_n^2 \rangle)^{-1} P_n u_0, \quad n \geq 0. \quad (3.10)$$

When $u \in \mathcal{P}'$ is regular, let Φ be a polynomial such that $\Phi u = 0$, then $\Phi = 0$ [28, 29].

The q -classical polynomials are orthogonal sequences (see Definition 1) that share a number of properties. Among them, we recall:

Proposition 2. [19] *Let $\{P_n\}_{n \geq 0}$ be an MOPS and u_0 be the corresponding regular form. The following statements are equivalent:*

- (a) $\{P_n\}_{n \geq 0}$ is a q -classical sequence.
- (b) There are two polynomials Φ and Ψ with Φ monic, $\deg \Phi \leq 2$ and $\deg \Psi = 1$ such that u_0 fulfills

$$D_q(\Phi u_0) + \Psi u_0 = 0. \quad (3.11)$$

- (c) There exists a sequence of nonzero numbers $\{\chi_n\}_{n \geq 1}$, and two polynomials Φ and Ψ with $\deg \Phi \leq 2$, $\deg \Psi = 1$ and Φ monic such that $\{P_n\}_{n \geq 0}$ fulfills the q -differential equation (1.2).
- (d) There is a monic polynomial Φ with $\deg \Phi \leq 2$ and a sequence of nonzero numbers $\{\vartheta_n\}_{n \geq 0}$ such that

$$P_n u_0 = \vartheta_n D_q^n \left(\left(\prod_{\sigma=0}^{n-1} q^{-\sigma \deg \Phi} \Phi(q^\sigma x) \right) u_0 \right), \quad n \geq 0. \quad (3.12)$$

Whenever $\{P_n\}_{n \geq 0}$ is q -classical, then so is $\{P_n^{[k]}\}_{n \geq 0}$, for any positive integer k and we have:

Corollary 3. [19] *Let k be a positive integer. If $\{P_n\}_{n \geq 0}$ is q -classical, then so is $\{P_n^{[k]}\}_{n \geq 0}$. The corresponding q -classical form $u_0^{[k]}$ fulfills $D_q(\Phi_k u_0^{[k]}) + \Psi_k u_0^{[k]} = 0$ with $\Phi_k(x) = q^{-k \deg \Phi} \Phi(q^k x)$, $\Psi_k(x) =$*

$q^{-k \deg \Phi} \left(\Psi(x) - [k]_q (D_{q^k} \Phi)(x) \right)$ and it is related to u_0 via $u_0^{[k]} = \zeta_k \left(\prod_{\sigma=0}^{k-1} \Phi_\sigma(x) \right) u_0$, where ζ_k is such that $(u_0^{[k]})_0 = 1$, with $\zeta_0 = 1$. Moreover, the sequence $\{P_n^{[k]}\}_{n \geq 0}$ fulfills

$$\Phi_k(x) D_q \circ D_{q^{-1}} \left(P_n^{[k]}(x) \right) - \Psi_k(x) D_{q^{-1}} \left(P_n^{[k]}(x) \right) = \chi_n^{[k]} P_n^{[k]}(x), \quad (3.13)$$

where

$$\chi_n^{[k]} = [n]_{q^{-1}} \left([n-1]_q \frac{\Phi_k'(0)}{2} - \Psi_k'(0) \right) = \begin{cases} -[n]_{q^{-1}} q^{-(\deg \Phi)k} \Psi'(0) & \text{if } \deg \Phi = 0, 1, \\ \frac{[n]_{q^{-1}}}{q^{2k+1}} \left([n+2k]_q - (1+q\Psi'(0)) \right) & \text{if } \deg \Phi = 2, \text{ for } n \geq 0. \end{cases}$$

Observe that the nonzero numbers ϑ_n in the Rodrigues type formula (3.12) are related to the coefficients ζ_k in Corollary 3 via (see [19, p.65] for further details):

$$\vartheta_n = \frac{(-1)^n}{[n]_q!} \langle u_0, P_n^2 \rangle \zeta_n = q^{-\frac{n(n-1)}{2} \deg \Phi} [n]_q! \left(\prod_{\sigma=1}^n \chi_\sigma^{[n-\sigma]} \right)^{-1}, \quad n \geq 0.$$

In addition, we also note that if $\deg \Phi = 2$, then $\chi_{n-k}^{[k]} = q^{k-1} \left([n]_{q^{-1}} (z + [n]_q) - [k]_{q^{-1}} (z + [k]_q) \right)$, for any $0 \leq k \leq n$, where $z = -(1 + q\Psi'(0))$.

3.2. Even order q -differential equations fulfilled by q -classical polynomials. In this section we prove Theorem 1, where we will mainly follow an algebraic approach. Afterwards, we explain the cases where the even order q -differential operator $\mathcal{L}_{k;q}$ admits a representation in a q -self adjoint version, as stated in Corollary 1.

Proof of Theorem 1. In the light of Corollary 3, the q -classical character of the sequence $\{P_n\}_{n \geq 0}$ implies the q -classical character of $\{P_n^{[m]}\}_{n \geq 0}$, no matter the choice of the nonnegative integer m . This encloses the orthogonality of each of the sequences $\{P_n^{[m]}\}_{n \geq 0}$ with respect to $u_0^{[m]}$ and therefore the elements of the corresponding dual sequences, say $\{u_n^{[m]}\}_{n \geq 0}$, can be written as $u_n^{[m]} = \left(\langle u_0^{[m]}, (P_n^{[m]})^2 \rangle \right)^{-1} P_n^{[m]} u_0^{[m]}$ for any integer $n \geq 0$. Consequently, the relation (3.8) becomes

$$D_q^\nu (P_n^{[k]} u_0^{[k]}) = \varpi_n(k, k-\nu) P_{n+\nu}^{[k-\nu]} u_0^{[k-\nu]}, \quad n \geq 0, \quad (3.14)$$

where

$$\varpi_n(k, k-\nu) = (-1)^\nu [n+1; q]_\nu \frac{\langle u_0^{[k]}, (P_n^{[k]})^2 \rangle}{\langle u_0^{[k-\nu]}, (P_{n+\nu}^{[k-\nu]})^2 \rangle}, \quad n \geq 0. \quad (3.15)$$

In this case, the corresponding regular forms $u_0^{[\nu]}$, with $\nu = 0, 1, \dots, k$, satisfy the q -differential equation

$$D_q \left(\Phi_\nu(x) u_0^{[\nu]} \right) + \Psi_\nu(x) u_0^{[\nu]} = 0, \quad (3.16)$$

where Φ_ν and Ψ_ν are given in Corollary 3, respectively. By recalling (3.3), then, based on the fact that $u_0^{[k-1]}$ satisfy (3.16) with $\nu = k-1$, we successively obtain

$$\begin{aligned} \langle \Phi_{k-1}(x) u_0^{[k-1]}, x^m P_n^{[k]} \rangle &= \frac{-1}{[n+1]_q} \langle D_q \left(x^m \Phi_{k-1}(x) u_0^{[k-1]} \right), P_{n+1}^{[k-1]} \rangle \\ &= \frac{-1}{[n+1]_q} \langle u_0^{[k-1]}, \left(-\Psi_{k-1}(x) q^{-m} x^m + q^{-1} \Phi_{k-1}(x) [m]_{q^{-1}} x^{m-1} \right) P_{n+1}^{[k-1]} \rangle, \quad m, n \geq 0. \end{aligned}$$

By virtue of the orthogonality of $\{P_n^{[k-1]}\}_{n \geq 0}$ with respect to $u_0^{[k-1]}$ and because of the degree of the polynomials Φ_{k-1} and Ψ_{k-1} , we deduce

$$\begin{aligned} & \langle \Phi_{k-1}(x) u_0^{[k-1]}, x^m P_n^{[k]} \rangle \\ &= \begin{cases} 0, & m \leq n-1, n \geq 1, \\ -\frac{q^{-n}}{[n+1]_q} \left([n]_q \frac{\Phi_{k-1}''(0)}{2} - \Psi'_{k-1}(0) \right) \langle u_0^{[k-1]}, (P_{n+1}^{[k]})^2 \rangle, & m = n, n \geq 0, \end{cases} \end{aligned}$$

Since $\Phi_{\nu-1}(x) u_0^{[\nu-1]} = \frac{\zeta_{\nu-1}}{\zeta_\nu} u_0^{[\nu]}$, for $\nu \geq 1$, where ζ_ν is such that $(u_0^{[\nu]})_0 = 1$, then we can write

$$\frac{\langle u_0^{[k]}, (P_n^{[k]})^2 \rangle}{\langle u_0^{[k-1]}, (P_{n+1}^{[k-1]})^2 \rangle} = -\frac{\zeta_k}{\zeta_{k-1}} \frac{q^{-n}}{[n+1]_q} \left([n]_q \frac{\Phi_{k-1}''(0)}{2} - \Psi'_{k-1}(0) \right), \quad n \geq 0,$$

which, by finite induction, leads to

$$\frac{\langle u_0^{[k]}, (P_n^{[k]})^2 \rangle}{\langle u_0^{[k-\nu]}, (P_{n+\nu}^{[k-\nu]})^2 \rangle} = (-1)^\nu \frac{\zeta_k}{\zeta_{k-\nu}} \left(\prod_{\sigma=1}^{\nu} \frac{q^{-(n+\sigma-1)}}{[n+\sigma]_q} \left([n+\sigma-1]_q \frac{\Phi_{k-\sigma}''(0)}{2} - \Psi'_{k-\sigma}(0) \right) \right), \quad n \geq 0.$$

Therefore, the coefficients $\varpi_n(k, \nu)$ in (3.15) can also be expressed as follows

$$\varpi_n(k, k-\nu) = \frac{\zeta_k}{\zeta_{k-\nu}} \left(\prod_{\sigma=1}^{\nu} q^{-(n+\sigma-1)} \left([n+\sigma-1]_q \frac{\Phi_{k-\sigma}''(0)}{2} - \Psi'_{k-\sigma}(0) \right) \right), \quad n \geq 0, \quad (3.17)$$

so that from Equation (3.14) we obtain

$$D_q^\nu \left(P_n^{[k]} u_0^{[k]} \right) = \frac{\zeta_k}{\zeta_{k-\nu}} \left(\prod_{\sigma=1}^{\nu} q^{-(n+\sigma-1)} \left([n+\sigma-1]_q \frac{\Phi_{k-\sigma}''(0)}{2} - \Psi'_{k-\sigma}(0) \right) \right) P_{n+\nu}^{[k-\nu]} u_0^{[k-\nu]}, \quad n \geq 0.$$

Substituting $u_0^{[k-\nu]}$ on the right hand side by its expression given in Corollary 3, we obtain

$$D_q^\nu \left(P_n^{[k]} u_0^{[k]} \right) = \zeta_k \left(\prod_{\sigma=1}^{\nu} q^{-(n+\sigma-1)} \left([n+\sigma-1]_q \frac{\Phi_{k-\sigma}''(0)}{2} - \Psi'_{k-\sigma}(0) \right) \right) \left(\prod_{\sigma=0}^{k-\nu-1} \Phi_\sigma \right) P_{n+\nu}^{[k-\nu]}(x) u_0,$$

which can be simplified into

$$D_q^\nu \left((D_q^k P_{n+k})(x) u_0^{[k]} \right) = \zeta_k \left(\prod_{\sigma=1}^{\nu} \chi_{n+\sigma}^{[k-\sigma]} \right) \left(\prod_{\sigma=0}^{k-\nu-1} \Phi_\sigma(x) \right) (D_q^{k-\nu} P_{n+k})(x) u_0, \quad n \geq 0, \quad (3.18)$$

if we take into consideration the definition of $\{P_n^{[\mu]}\}_{n \geq 0}$ given in (3.7) alongside with the expression for $\chi_n^{[k]}$ given in Corollary 3. In particular, we take $n = 0$ in (3.18) to get

$$D_q^\nu \left(u_0^{[k]} \right) = \zeta_k \left(\prod_{\sigma=1}^{\nu} \chi_\sigma^{[k-\sigma]} \right) \left(\prod_{\sigma=0}^{k-\nu-1} \Phi_\sigma \right) P_\nu^{[k-\nu]} u_0, \quad n \geq 0. \quad (3.19)$$

Using the functional version of the q -Leibniz rule (3.5), the left-hand-side of (3.18) can be rewritten as

$$D_q^k \left(D_q^k P_{n+k}(x) u_0^{[k]} \right) = \sum_{\nu=0}^k \begin{bmatrix} k \\ \nu \end{bmatrix}_{q^{-1}} q^{-(k-\nu)} \left(D_{q^{-1}}^{k-\nu} \circ D_q^k P_{n+k} \right) (q^{-\nu} x) (D_q^\nu u_0^{[k]}), \quad n \geq 0,$$

which, after (3.19), can be expressed in terms on u_0 :

$$\begin{aligned} D_q^k \left(D_q^k P_{n+k}(x) u_0^{[k]} \right) &= \sum_{\nu=0}^k \begin{bmatrix} k \\ \nu \end{bmatrix}_{q^{-1}} q^{-(k-\nu)} \zeta_k \left(\prod_{\sigma=1}^{\nu} \chi_\sigma^{[k-\sigma]} \right) \left(\prod_{\sigma=0}^{k-\nu-1} \Phi_\sigma(x) \right) \\ &\quad \times P_\nu^{[k-\nu]}(x) \left(D_{q^{-1}}^{k-\nu} \circ D_q^k P_{n+k} \right) (q^{-\nu} x) u_0, \quad n \geq 0. \end{aligned}$$

Bringing this expression into (3.18), with ν replaced by k , we obtain

$$\begin{aligned} & \sum_{\nu=0}^k \begin{bmatrix} k \\ \nu \end{bmatrix}_{q^{-1}} q^{-(k-\nu)} \left(\prod_{\sigma=1}^{\nu} \chi_{\sigma}^{[k-\sigma]} \right) \left(\prod_{\sigma=0}^{k-\nu-1} \Phi_{\sigma}(x) \right) P_{\nu}^{[k-\nu]} \left(D_{q^{-1}}^{k-\nu} \circ D_q^k P_{n+k} \right) (q^{-\nu} x) u_0 \\ &= \left(\prod_{\sigma=1}^k \chi_{n+\sigma}^{[k-\sigma]} \right) P_{n+k}(x) u_0, \quad n \geq 0. \end{aligned}$$

According to the regularity of u_0 , it follows that $y(x) = P_{n+k}(x)$ satisfies the q -differential equation

$$\sum_{\nu=0}^k \Lambda_{k,\nu}(x; q) \left(D_{q^{-1}}^{k-\nu} \circ D_q^k y \right) (q^{-\nu} x) = \Xi_n(k; q) y(x),$$

where

$$\Lambda_{k,\nu}(x; q) = \begin{bmatrix} k \\ \nu \end{bmatrix}_{q^{-1}} q^{-(k-\nu)} \left(\prod_{\sigma=1}^{\nu} \chi_{\sigma}^{[k-\sigma]} \right) \left(\prod_{\sigma=0}^{k-\nu-1} q^{-\sigma \deg \Phi} \Phi(q^{\sigma} x) \right) P_{\nu}^{[k-\nu]}(x)$$

for $\nu = 0, 1, \dots, k$, and

$$\Xi_n(k; q) = \prod_{\sigma=0}^{k-1} \chi_{n-\sigma}^{[\sigma]}, \quad n \geq 0,$$

with $\chi_n^{[k]}$ given in Corollary 3. Hence, we get the result if we observe that $P_{\nu}^{[k-\nu]}(x) := \frac{1}{[\nu+1; q]_{k-\nu}} D_q^{k-\nu} P_k(x)$, and that $\begin{bmatrix} k \\ \nu \end{bmatrix}_{q^{-1}} \frac{q^{-(k-\nu)}}{[\nu+1; q]_{k-\nu}} = \frac{q^{-(k-\nu)(\nu+1)}}{[k-\nu]_q!}$, for any integers k, ν such that $0 \leq \nu \leq k$. \square

From the proof of Theorem 1, we readily deduce a generalisation of the Rodrigues-type formula (3.12) (which is recovered if we set $n = k$ in the identity (3.20) given below).

Corollary 4. *For any positive integer k , a q -classical polynomial sequence $\{P_n\}_{n \geq 0}$, orthogonal with respect to the q -classical linear functional u_0 , fulfils the relation*

$$D_q^k \left(\left(\prod_{\sigma=0}^{k-1} \Phi_{\sigma}(x) \right) (D_q^k P_n)(x) u_0 \right) = \Xi_n(k; q) P_n(x) u_0, \quad \text{for } n \geq 0, \quad (3.20)$$

where $\Phi_{\sigma}(x)$ are the monic polynomials given in Corollary 3 and $\Xi_n(k; q)$ are the eigenvalues given by (1.5).

Proof. The result is an immediate consequence of the relation (3.18), after setting $\nu = k$ and using the expression for $u_0^{[k]}$ given in Corollary 3. \square

Despite (3.20) is actually a functional relation, in some cases it can be written as a q -differential relation. This occurs whenever u_0 admits an integral representation via a weight function, $W_q(x)$ say, as regular as necessary, so that:

$$\langle u_0, f \rangle = \int_{-\infty}^{+\infty} f(x) W_q(x) dx, \quad \text{for any } f \in \mathcal{P}. \quad (3.21)$$

This weight function W_q is such that the integral on the RHS of the precedent identity must exist for any polynomial f , and must represent the regular form u_0 . For this reason, W_q needs to be continuous at the origin or such that the integral

$$\lim_{\epsilon \rightarrow 0^+} \int_{\epsilon}^1 \frac{W_q(x) - W_q(-x)}{x} dx$$

exists and, in addition, the fact that u_0 fulfils (3.11) implies that W_q must fulfill the q -differential equation (1.6) (a detailed justification can be found in [19, p.81-82]).

Whenever (3.21) is admissible, an MOPS with respect to u_0 may as well be referred to as orthogonal with respect to the weight function W_q . In this case, from (3.20) we deduce a q -self-adjoint version of (1.4):

$$q^{-k} D_{q^{-1}}^k \left(\left(\prod_{\sigma=0}^{k-1} \Phi_\sigma(x) \right) W_q(x) \left(D_q^k P_n \right)(x) \right) = \Xi_n(k; q) W_q(x) P_n(x), \text{ for } n \geq 0. \quad (3.22)$$

Since $\{P_n\}_{n \geq 0}$ forms a basis of \mathcal{P} , we get Corollary 1.

In particular, if we consider $n = k$ case in (3.22), we recover the known Rodrigues' formula (see [20, p.70], [35]):

$$\Xi_n(k; q) P_k(x) = [1; q]_k q^{-k} (W_q(x))^{-1} D_{q^{-1}}^k \left(\left(\prod_{\sigma=0}^{k-1} \Phi_\sigma(x) \right) W_q(x) \right), \quad k \geq 0.$$

3.3. Integral powers of the second-order q -differential operator. The elements of any q -classical sequence $\{P_n\}_{n \geq 0}$ are solutions of the even order q -differential equation in y

$$\mathcal{L}_q^k[y](x) = (\chi_n)^k y(x), \quad n \geq 0. \quad (3.23)$$

This raises the natural question of obtaining an explicit expression for \mathcal{L}_q^k as well as of relating it to the $2k$ th order q -differential operator $\mathcal{L}_{k;q}$ given in (1.4). As the elements of a q -classical polynomial sequence are eigenfunctions of both operators, these problems may be addressed through the comparison between the eigenvalues $(\chi_n)^k$ given by (1.3) and $\Xi_n(k)$ given by (1.5). Observe that $\Psi'(0) \neq [j]_q$ for any integer $j \geq 0$, otherwise the q -classical form u_0 fulfilling (3.11) would have an undetermined sequence of moments, contradicting the assumptions. For this reason, $\Xi_n(k; q) \neq 0$ for $n \geq k$.

Proof of Theorem 2. We begin by relating the k th power of the eigenvalues χ_n given by (1.3) and the eigenvalues $\Xi_n(k; q)$ given by (1.5) for any integer $n \geq 0$. In the light of equalities (2.2) with $x = [n]_{q^{-1}}$, when $\deg \Phi < 2$ we have

$$(\chi_n)^k = \sum_{j=0}^k S_{q^{-1}}(k, j) q^{(\deg \Phi - 1) \frac{j(j-1)}{2}} (-\Psi'(0))^{k-j} \Xi_n(j; q), \quad n \geq 0,$$

whereas, based on the identity (2.7) with $x = [n]_{q^{-1}}(z + [n]_q)$, when $\deg \Phi = 2$ we obtain

$$(\chi_n)^k = \sum_{j=0}^k \text{JS}_k^j(z; q^{-1}) q^{\frac{j(j+1)}{2} - k} \Xi_n(j; q), \quad n \geq 0.$$

According to Proposition 2, the q -classical polynomials are solutions of the $2k$ -order q -differential equation (3.23) where the operator \mathcal{L}_q and the eigenvalues χ_n are those given by (1.2). On the other hand, Theorem 1 ensures the q -classical polynomials to be solutions of the $2k$ -order q -differential equation (1.4), whose corresponding eigenvalues $\Xi_n(k; q)$ are given in Theorem 1. The result now follows from the fact that any q -classical sequence forms a basis of \mathcal{P} . \square

In a similar manner, for a given positive integer k we can express the eigenvalues $\Xi_n(k; q)$ in terms of $(\chi_n)^k$ for any integer $n \geq 0$. Indeed, after $q \rightarrow q^{-1}$, (2.1) and (2.6), with $x = [n]_{q^{-1}}$ and $x = [n]_{q^{-1}}([n]_q - A)$, respectively, imply the following identities

$$\Xi_n(k; q) = \begin{cases} q^{(1-\deg \Phi) \frac{k(k-1)}{2}} \sum_{j=0}^k s_{q^{-1}}(k, j) (-\Psi'(0))^{k-j} (\chi_n)^j & \text{if } \deg \Phi = 0, 1, \\ q^{-\frac{k(k+1)}{2}} \sum_{j=0}^k \text{Jc}_k^j(z; q^{-1})(k, j) q^j (\chi_n)^j & \text{if } \deg \Phi = 2, \end{cases}$$

which hold for any integer $n \geq 0$, where $z = -(1 + q\Psi'(0))$.

By reversing (1.8) using the q -Stirling or q -Jacobi-Stirling numbers of the first kind (or evoking analogous arguments to those in the proof of Theorem 2), we deduce the reciprocal identities of (1.8). More precisely, we obtain Corollary 2.

All the q -classical polynomials were studied in [19]. With the equivalence relation

$$\{P_n(x)\}_{n \geq 0} \sim \{R_n(x)\}_{n \geq 0} \quad \text{if} \quad \exists a \neq 0 \quad \text{such that} \quad R_n(x) = a^{-n} P_n(ax), \quad n \geq 0,$$

several equivalence classes were obtained and these are summarised in Table 1. Since the q -classical polynomials are solutions to the q -differential equation (1.2) and the polynomial coefficient Φ , of degree 2 at most, plays a fundamental role in this work, we have arranged the information according to its properties.

TABLE 1. The q -classical polynomials.

deg Φ	q -classical MOPS
0	Al-Salam Carlitz polynomials · Discrete q -Hermite polynomials
1	Big q -Laguerre · q -Meixner · Wall q -polynomials q -Laguerre polynomials · Little q -Laguerre polynomials q -Charlier I polynomials
2 (with double root)	Alternative q -Charlier polynomials · Stieltjes-Wigert q -polynomials
2 (with 2 single roots)	Little q -Jacobi polynomials · q -Charlier II polynomials Generalized Stieltjes-Wigert q -polynomials · Big q -Jacobi Bi-generalized Stieltjes-Wigert q -polynomials

4. PARTICULAR CASES

Among all the q -classical polynomials, we single out some examples from Table 1 that illustrate the results here obtained. These are mere illustrations, since the results apply to any q -classical family. Among other things, it permits to obtain differently many expressions for even order q -differential equations fulfilled by the q -classical polynomials.

4.1. Al-Salam-Carlitz polynomials. The monic Al-Salam-Carlitz polynomials are

$$P_n(x, a; q) = (-a)^n q^{\binom{n}{2}} \phi_1 \left(\begin{matrix} q^{-n}, x^{-1} \\ 0 \end{matrix}; q, \frac{qx}{a} \right) = \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q (-a)^{n-k} q^{\binom{n-k}{2}} (x^{-1}; q)_k x^k$$

and satisfy a recurrence relation of the type (3.9) with $\beta_n = (1+a)q^n$ and $\gamma_{n+1} = aq^n(q^{n+1} - 1)$, for $n \geq 0$, see [20, p.534], [19]. They are orthogonal with respect to the regular linear functional u_0 fulfilling (3.11) with $\Phi(x) = 1$, $\Psi(x) = (a(q-1))^{-1}(x - (1+a))$, provided that $a \neq 0$. They are positive definite whenever $a < 0$ and $0 < q < 1$ or when $a > 0$ and $q > 1$. The case where $a = -1$ and $0 < q < 1$ corresponds to the so-called Discrete q -Hermite polynomials. We further notice that this MOPS is also Appell with respect to the q -difference operator D_q , so is to say that $P_n^{[k]}(x, a; q) = P_n(x, a; q)$.

In the light of Theorem 1, for any positive integer k , the Al-Salam-Carlitz polynomials are eigenfunctions of the q -differential operator

$$\mathcal{L}_{k;q}[y](x) := \sum_{\nu=0}^k \frac{[1; q^{-1}]_k}{[1; q^{-1}]_{n-k}} \frac{q^{-k}}{(a(q^{-1} - 1))^\nu} P_\nu(x, a; q) \left(D_{q^{-1}}^{k-\nu} \circ D_q^k y \right) (q^{-\nu} x).$$

Following Proposition 2, for each $k \geq 1$, we may write

$$\mathcal{L}_q^k[y](x) = \sum_{j=0}^k S_{q^{-1}}(k, j) q^{-\frac{j(j-1)}{2}} (a(q-1))^{-(k-j)} \mathcal{L}_{j;q}[y](x).$$

When $0 < q < 1$ and $a < 0$, the regular form u_0 admits an integral representation as in (3.21) via the weight function [19, p.87]

$$W_q(x) = K(qx; q)_\infty (a^{-1}qx; q)_\infty \mathbb{1}_{[a^{-1}q^{-1}, q^{-1}]}(x) \quad \text{with} \quad K^{-1} = \int_{a^{-1}q^{-1}}^{q^{-1}} (qt; q)_\infty (a^{-1}qt; q)_\infty dt,$$

where $\mathbb{1}_A(x)$ stands for the characteristic function over the set A (see introduction). Hence, according to Corollary 1, we may alternatively write

$$\mathcal{L}_{k;q}[y](x) := \left(q^k (qx; q)_\infty (a^{-1}qx; q)_\infty \right)^{-1} D_{q^{-1}}^k \left((qx; q)_\infty (a^{-1}qx; q)_\infty (D_q^k y(x)) \right),$$

which is valid for any $y \in \mathcal{P}$. When $k = 1$, the latter reduces to the well known q -Sturm-Liouville equation fulfilled by the Al Salam-Carlitz q -polynomials (see, for instance, [17, p.472]).

4.2. The Stieltjes-Wigert polynomials. The monic Stieltjes-Wigert polynomials are

$$P_n(x; q) = (-1)^n q^{-\frac{n}{2}(2n+1)} {}_1\phi_1 \left(\begin{matrix} q^{-n} \\ 0 \end{matrix}; q, -q^{n+3/2}x \right) = \sum_{k=0}^n \frac{(-1)^{n+k} q^{k(k+\frac{1}{2})-n(n+\frac{1}{2})}}{(q; q)_{n-k}} x^k.$$

They are orthogonal with respect to u_0 fulfilling (3.11) with $\Phi(x) = x^2$ and $\Psi(x) = -(q-1)^{-1}\{x - q^{-3/2}\}$, and they satisfy a recurrence relation of the type (3.9) with $\beta_n = (1+q-q^{n+1})q^{-2n-3/2}$ and $\gamma_{n+1} = (1-q^{n+1})q^{-4(n+1)}$, $n \geq 0$, see [19], [20, p.544]. When $0 < q < 1$, u_0 is positive definite, and admits an integral representation via a weight-function W_q as in (3.21) which is given by

$$W_q(x) = \sqrt{\frac{q}{2\pi \ln q^{-1}}} \exp\left(-\frac{\ln^2 x}{2 \ln q^{-1}}\right) \mathbb{1}_{(0, +\infty)}(x).$$

In the light of Corollary 1, for $0 < q < 1$, the Stieltjes-Wigert q -polynomials are eigenfunctions of

$$\mathcal{L}_{k;q}[y](x) = q^{-k} \exp\left(\frac{\ln^2 x}{2 \ln q^{-1}}\right) D_{q^{-1}}^k \left(x^{2k} \exp\left(-\frac{\ln^2 x}{2 \ln q^{-1}}\right) (D_q^k y(x)) \right).$$

According to Theorem 1, the latter operator can be expanded as follows

$$\mathcal{L}_{k;q}[y](x) = \sum_{\nu=0}^k \begin{bmatrix} k \\ \nu \end{bmatrix}_{q^{-1}} \alpha_{k,\nu} x^{2k-2\nu} P_\nu^{[k-\nu]}(x; q) \left(D_{q^{-1}}^{k-\nu} \circ D_q^k y \right) (q^{-\nu}x),$$

with

$$\alpha_{k,\nu} = q^{-(k-\nu)} ([\nu]_q!) \left(\prod_{\sigma=1}^{\nu} q^{-2k+\sigma} \left([2k-\sigma]_q + \frac{1}{q-1} \right) \right).$$

Finally, by means of the q -Jacobi-Stirling numbers of second kind, for any positive integer k , the k th power of the q -differential operator

$$\mathcal{L}_q := x^2 D_q \circ D_{q^{-1}} + (q-1)^{-1} \{x - q^{-3/2}\} D_{q^{-1}}$$

is given by the formula

$$\mathcal{L}_q^k[y](x) = \sum_{j=0}^k \text{JS}_k^j((q-1)^{-1}; q^{-1}) q^{\frac{j(j+1)}{2}-k} \mathcal{L}_{j;q}[y](x), \quad \forall y \in \mathcal{P},$$

which, after (2.15), can be alternatively written as

$$\mathcal{L}_q^k[y](x) = \sum_{j=0}^k \frac{q^{\frac{j(j-1)}{2}}}{(q-1)^{k-j}} S_q(k, j) \mathcal{L}_{j;q}[y](x), \quad \forall y \in \mathcal{P}.$$

4.3. The Little q -Jacobi polynomials. The monic Little q -Jacobi polynomials are

$$\begin{aligned} P_n(x; a, b|q) &= \frac{(-1)^n q^{\binom{n}{2}} (aq; q)_n}{(abq^{n+1}; q)_n} {}_2\phi_1 \left(\begin{matrix} q^{-n}, abq^{n+1} \\ aq \end{matrix}; q, qx \right) \\ &= \frac{(aq; q)_n}{(abq^{n+1}; q)_n} \sum_{k=0}^n \frac{[n]_q}{[k]_q} \frac{(-1)^{n-k} (abq^{n+1}; q)_k}{(aq; q)_k} q^{\binom{n-k}{2}} x^k \end{aligned}$$

and satisfy the q -differential equation (1.2) with $\Psi(x) = (abq^2(q-1))^{-1} \{(1-abq^2)x + aq - 1\}$ and $\Phi(x) = x(x - b^{-1}q^{-1})$, for $n \geq 0$, see [20, p.482], [19]. Hence, for each positive integer k , we have

$$P_n^{[k]}(x; a, b|q) = P_n(x; aq^k, bq^k|q), \quad n \geq 0.$$

They form an MOPS with respect to the regular linear functional u_0 fulfilling (3.11) provided that the parameters a, b satisfy $a, b, ab \neq q^{-(n+2)}$, for $n \geq 0$. The linear functional u_0 is positive definite when one of the following two conditions is realised [19]

$$\begin{aligned} 0 < q < 1, \quad 0 < a < q^{-1}, \quad b \in]-\infty, q^{-1}[-\{0\}, \\ q > 1, \quad a > q^{-1}, \quad b \in]-\infty, 0[\cup]q^{-1}, +\infty[. \end{aligned}$$

We refer to [19, 20] for further details, but we highlight some results from [19, pp. 100-102].

For certain values of a, b and q , it is possible to get an integral representation for the regular form u_0 via a weight function $W_q(x)$, as described in (3.21). Setting $a := q^{\alpha-1}$ with $\alpha > 0$, then depending on the range of values for b and q , the weight-function $W_q(x)$ can be written as follows:

- For $0 < q < 1$ and $b \in]-\infty, 1[-\{0\}$, we have

$$W_q(x) = K x^{\alpha-1} \frac{(qx; q)_\infty}{(bqx; q)_\infty} \mathbb{1}_{(0, q^{-1})}(x) \quad \text{with } K^{-1} = \int_0^{q^{-1}} t^{\alpha-1} \frac{(qt; q)_\infty}{(bqt; q)_\infty} dt.$$

- For $q > 1$ and $b \geq 1$, we have

$$W_q(x) = K x^{\alpha-1} \frac{(bx; q^{-1})_\infty}{(x; q^{-1})_\infty} \mathbb{1}_{(0, b^{-1})}(x) \quad \text{with } K^{-1} = \int_0^{b^{-1}} t^{\alpha-1} \frac{(bt; q^{-1})_\infty}{(t; q^{-1})_\infty} dt.$$

- For $q > 1$ and $b \leq 0$, we have

$$W_q(x) = K |x|^{\alpha-1} \frac{(bx; q^{-1})_\infty}{(-|x|; q^{-1})_\infty} \mathbb{1}_{(b^{-1}, 0)}(x) \quad \text{with } K^{-1} = \int_0^{|b|^{-1}} t^{\alpha-1} \frac{(|b|t; q^{-1})_\infty}{(-t; q^{-1})_\infty} dt.$$

On the other hand, it is possible to represent u_0 by a discrete measure. Namely, for $0 < q < 1$, we have (see also [20, p.482])

$$u_0 = \frac{(aq; q)_\infty}{(abq^2; q)_\infty} \sum_{k \geq 0} \frac{(bq; q)_k}{(q; q)_k} (aq)^k \delta_{q^k}, \quad \text{with } |aq| < 1, |bq| < 1,$$

whereas, when $q > 1$, we can write

$$u_0 = \frac{(a^{-1}q^{-1}; q^{-1})_\infty}{(a^{-1}b^{-1}q^{-2}; q^{-1})_\infty} \sum_{k \geq 0} \frac{(b^{-1}q^{-1}; q^{-1})_k}{(q^{-1}; q^{-1})_k} (a^{-1}q^{-1})^k \delta_{b^{-1}q^{-(k+1)}}, \quad \text{for } |aq| > 1, |bq| > 1.$$

In the light of Corollary 1, whenever the Little q -Jacobi linear functional u_0 admits an integral representation via a weight function $W_q(x)$, the Little q -Jacobi polynomials $\{P_n(x; a, b|q)\}_{n \geq 0}$ are eigenfunctions of

$$\mathcal{L}_{k;q}[y](x) = q^{-k} (W_q(x))^{-1} D_{q^{-1}}^k \left(\left(\prod_{\sigma=0}^{k-1} x(x - b^{-1}q^{-(\sigma+1)}) \right) W_q(x) (D_q^k y(x)) \right). \quad (4.1)$$

In any of the above mentioned cases, following Theorem 1, the latter operator can be written as follows

$$\mathcal{L}_{k;q}[y](x) = \sum_{\nu=0}^k \begin{bmatrix} k \\ \nu \end{bmatrix}_{q^{-1}} \alpha_{k,\nu}(x) \left(\prod_{\sigma=0}^{k-1} x(x - b^{-1}q^{-(\sigma+1)}) \right) \left(D_{q^{-1}}^{k-\nu} \circ D_q^k y \right) (q^{-\nu} x)$$

with

$$\alpha_{k,\nu}(x) = q^{-(k-\nu)} \left(\prod_{\sigma=1}^{\nu} [\sigma]_{q^{-1}} q^{-2k+\sigma} \left([2k - \sigma]_q + z \right) \right) P_{\nu}(x; aq^{k-\nu}, bq^{k-\nu}|q),$$

and $z = -(1 + q\Psi'(0)) = \frac{1}{q-1} (1 - (abq)^{-1})$.

Finally, by means of the q -Jacobi-Stirling numbers of second kind, for each integer $k \geq 0$, the k th power of the q -differential operator

$$\mathcal{L}_q := x(x - b^{-1}q^{-1})D_q \circ D_{q^{-1}} - ((abq^2(q-1))^{-1} \{ (1 - abq^2)x + aq - 1 \}) D_{q^{-1}} \quad (4.2)$$

is given by the formula

$$\mathcal{L}_q^k[y](x) = \sum_{j=0}^k \text{JS}_k^j(z; q^{-1}) q^{\frac{j(j+1)}{2} - k} \mathcal{L}_{j;q}[y](x),$$

for any $y \in \mathcal{P}$, where $z = -(1 + q\Psi'(0)) = \frac{1}{q-1} (1 - (abq)^{-1})$. According to Corollary 2 the reciprocal relations can also be established:

$$\mathcal{L}_{k;q}[y](x) = \sum_{j=0}^k \text{Jc}_k^j(z; q^{-1}) q^{j - \frac{k(k+1)}{2}} \mathcal{L}_q^j[y](x), \quad \text{for any } y \in \mathcal{P}.$$

Remark 4. Amongst the q -classical sequences it is possible to establish limiting relations between the several q -classical families. (For a catalogue of such relations we refer to [20].) In the same manner, it is also possible to establish limiting relations between two operators, $\mathcal{L}_{k;q}$ and $\widehat{\mathcal{L}}_{k;q}$ (that have P_n and \widehat{P}_n as eigenfunctions, respectively), mirroring the limiting relation that the two corresponding q -classical sequences $\{P_n\}_{n \geq 0}$ and $\{\widehat{P}_n\}_{n \geq 0}$ satisfy.

For example, the monic q -Laguerre polynomial sequence $\{\widehat{P}_n(x; \alpha; q)\}_{n \geq 0}$ is related to the Little q -Jacobi polynomials $\{P_n(x; q^{\alpha}, b|q)\}_{n \geq 0}$, that we have just analysed, via the limiting relation (see [20, p.522]):

$$\begin{aligned} \widehat{P}_n(x; \alpha; q) &= \lim_{b \rightarrow -\infty} ((-bq)^n P_n(-b^{-1}q^{-1}x; q^{\alpha}, b|q)) \\ &= \frac{(-1)^n (q^{\alpha+1}; q)_n}{q^{n(n+\alpha)}} {}_1\phi_1 \left(\begin{matrix} q^{-n} \\ q^{\alpha+1} \end{matrix}; q, -q^{n+\alpha+1}x \right), \quad n \geq 0. \end{aligned}$$

The operator $\widehat{\mathcal{L}}_{q^{-1}} := xD_{q^{-1}} \circ D_q - \frac{q^{\alpha+2}}{q-1} \left(\frac{1}{q^{\alpha+1}} - 1 - x \right) D_q$ has the monic q -Laguerre polynomials $\widehat{P}_n(x; \alpha; q)$ as eigenfunctions, and, more generally, according to Corollary 2, so have the operators

$$\widehat{\mathcal{L}}_{k;q^{-1}}[y](x) = \sum_{j=0}^k c_{q^{-1}}(k, j) (-\Psi'(0))^{k-j} \widehat{\mathcal{L}}_{q^{-1}}^j[y](x),$$

for each $k = 1, 2, \dots$. In fact, the operators $\widehat{\mathcal{L}}_{k;q^{-1}}$ are a limiting case of the operators $\mathcal{L}_{k;q}$ in (4.1) which have the Little q -Jacobi polynomials as eigenfunctions. Indeed, consider the operator $\widetilde{\mathcal{L}}_q = x(x+1)D_q \circ D_{q^{-1}} -$

$\frac{q^{-\alpha-2}(-b(x+1)q^{\alpha+2}+bq+x)}{b(q-1)}D_{q^{-1}}$ obtained from \mathcal{L}_q in (4.2) after the change of variable $x \rightarrow -b^{-1}q^{-1}x$, that is $\tilde{\mathcal{L}}_q[y(t)][-b^{-1}q^{-1}x] = \mathcal{L}_q[y(-b^{-1}q^{-1}t)][x]$, so that

$$\tilde{\mathcal{L}}_q \xrightarrow{b \rightarrow -\infty} x(x+1)D_q \circ D_{q^{-1}} - \frac{-(x+1) + q^{-(\alpha+1)}}{q-1}D_{q^{-1}} = q^{-(\alpha+2)}\widehat{\mathcal{L}}_{q^{-1}}.$$

Besides, recall that $z = \frac{1}{q-1}(1 - (q^\alpha b q)^{-1})$ and therefore $\lim_{b \rightarrow -\infty} z = \frac{1}{q-1}$, so that, by taking into account the identities (2.15), we have for any $y \in \mathcal{P}$

$$\begin{aligned} \lim_{b \rightarrow -\infty} \mathcal{L}_{k;q}[y(-b^{-1}q^{-1}t)](x) &= \lim_{b \rightarrow -\infty} \left(\sum_{j=0}^k Jc_k^j(z; q^{-1}) q^{j - \frac{k(k+1)}{2}} \tilde{\mathcal{L}}_q^j[y](x) \right) \\ &= q^{\binom{k}{2} - k(\alpha+2)} \widehat{\mathcal{L}}_{k;q^{-1}}[y](x). \end{aligned}$$

The detailed analysis done for the q -classical sequences of Al-Salam-Carlitz polynomials, Stieltjes-Wigert polynomials and Little q -Jacobi polynomials, including the limiting relation to the q -Laguerre polynomials, can naturally be extended to all the other q -classical sequences listed in Table 1. The procedure is entirely similar to the ones just described, as the required properties were already studied by several authors and a synthesis of it can be found in [19, 20].

In this paper, we have succeeded in providing the explicit expressions for any integral composite power of a second order q -differential operator having the q -classical polynomials as eigenfunctions. The study required the introduction of the new set of q -Jacobi-Stirling numbers, which we have characterised and provided a combinatorial interpretation. Clearly, by allowing $q \rightarrow 1$ we recover all the classical polynomial related results found in [2, 7, 8, 11, 26, 27, 30] from the viewpoint of both the algebra of these operators and the combinatorial interpretations of the connection coefficients.

Acknowledgements. We thank the referees for careful reading of this manuscript and illuminating comments and remarks.

REFERENCES

- [1] R. Álvarez-Nodarse, J. C. Medem, q -classical polynomials and the q -Askey and Nikiforov-Uvarov tableaux. J. Comput. Appl. Math. 135 (2001), no. 2, 197-223.
- [2] G. E. Andrews, E. S. Egge, W. Gawronski, L. L. Littlejohn, The Jacobi-Stirling numbers. J. Combin. Theory Ser. A 120 (2013), no. 1, 288-303.
- [3] S. Bochner, Über Sturm-Liouvillesche Polynomsysteme, Math. Z. **29** (1929) 730-736.
- [4] T. S. Chihara, An introduction to orthogonal polynomials. Gordon and Breach, NY (1978).
- [5] A. Erdélyi, A. Magnus, F. Oberhettinger, F. Tricomi, Higher Transcendental Functions, McGraw-Hill, New York, vols. 1, 2, 1953, vol. 3, 1955.
- [6] E. S. Egge, Legendre-Stirling permutations, European J. Combin., 31 (2010) 1735-1750.
- [7] W. N. Everitt, L. L. Littlejohn, R. Wellman, G. J. Yoon, Jacobi-Stirling numbers, Jacobi polynomials, and the left-definite analysis of the classical Jacobi differential expression, J. Comput. Appl. Math. 208 (2007) 29-56.
- [8] W. N. Everitt, L. L. Littlejohn, R. Wellman, The left-definite spectral theory for the classical Hermite differential equation, J. Comput. Appl. Math., 121 (2000) 313-330.
- [9] D. Foata, G.-N. Han, New permutation coding and equidistribution of set-valued statistics, Theoret. Comput. Sci., 410 (2009) 3743-3750.
- [10] G. Gasper, M. Rahman, Basic hypergeometric series. With a foreword by Richard Askey. Second edition. Encyclopedia of Mathematics and its Applications, 96. Cambridge University Press, Cambridge, 2004.
- [11] Y. Gelineau, J. Zeng, Combinatorial interpretations of the Jacobi-Stirling numbers. Electron. J. Combin. 17(1) (2010) R70.
- [12] I. M. Gessel, Z. Lin and J. Zeng, Jacobi-Stirling numbers and P -partitions, European J. Combin., 33 (2012) 1987-2000.
- [13] H. W. Gould, The q -Stirling numbers of first and second kinds. Duke Math. J. 28 (1961) 281-289.
- [14] W. Hahn, Über höhere Ableitungen von Orthogonalpolynome, Math. Zeit. 43 (1938)(1), 101.
- [15] W. Hahn, Über Orthogonalpolynome, die q -Differenzgleichungen genügen, Math. Nachr., 2(1-2), (1949), 4-34.
- [16] W. Hahn, Beitrag zur Theorie der Heineschen Reihen, Math. Nachr. 2 (1949), 340-379.

- [17] M. Ismail, Classical and quantum orthogonal polynomials in one variable. With two chapters by Walter Van Assche. With a foreword by Richard A. Askey. Reprint of the 2005 original. Encyclopedia of Mathematics and its Applications, 98. Cambridge University Press, Cambridge, 2009.
- [18] A. Kasraoui, D. Stanton, J. Zeng, The combinatorics of Al-Salam–Chihara q -Laguerre polynomials. *Adv. in Appl. Math.* 47 (2011), no. 2, 216–239.
- [19] L. Khérifi, P. Maroni, The H_q -classical orthogonal polynomials, *Acta Appl. Math.* 71 (2002) 49–115.
- [20] R. Koekoek, P. A. Lesky, R. F. Swarttouw, *Hypergeometric Orthogonal Polynomials and Their q -Analogues*, Springer Monographs in Mathematics (SMM). Springer, 2010.
- [21] H. L. Krall, Certain differential equations for Tchebyshev polynomials, *Duke Math. J.* 4 (1938) 705–718.
- [22] P. A. Lesky, Zweigliedrige Rekursionen für die Koeffizienten von Polynomlösungen Sturm-Liouvillescher q -Differenzgleichungen, *Z. Angew. Math. Mech.* 74 (1994), 497–500.
- [23] P. A. Lesky, Orthogonalpolynome in x und q^{-x} als Lösungen von reellen q -Operatorgleichungen zweiter Ordnung, *Monatsh. Math.* 132 (2001), 123–140.
- [24] Z. Lin, J. Zeng, Positivity properties of Jacobi-Stirling numbers and generalized Ramanujan polynomials. *Adv. in Appl. Math.* 53 (2014), 12–27.
- [25] A. F. Loureiro, S. Yakubovich, The Kontorovich-Lebedev transform as a map between d -orthogonal polynomials, *Stud. Appl. Math.*, 131(3) (2013) 229–265.
- [26] A. F. Loureiro, New results on the Bochner condition about classical orthogonal polynomials, *J. Math. Anal. Appl.*, 364 (2) (2010), 307–323.
- [27] A. F. Loureiro, P. Maroni, P. Z. Rocha, The generalized Bochner condition about classical orthogonal polynomials revisited, *J. Math. Anal. Appl.* 322 (2006) 645–667.
- [28] P. Maroni, Une théorie algébrique des polynômes orthogonaux. Application aux polynômes orthogonaux semi-classiques, C. Brezinski et al. (Eds.), *Orthogonal Polynomials and their Applications*, in: IMACS Ann. Comput. Appl. Math. 9 (1991), 95–130.
- [29] P. Maroni, Variations around classical orthogonal polynomials. Connected problems. *J. Comput. Appl. Math.*, 48 (1993), 133–155.
- [30] L. Miranian, On classical orthogonal polynomials and differential operators, *J. Phys. A: Math Gen.* 38 (2005) 6379–6382.
- [31] J. C. Medem, R. Álvarez-Nodarse, F. Marcellán, On the q -polynomials: a distributional study. *J. Comput. Appl. Math.* 135 (2001), no. 2, 157–196.
- [32] P. Mongelli, Combinatorial interpretations of particular evaluations of complete and elementary symmetric functions, *Electron. J. Combin.*, 19 (2012), R60.
- [33] A. F. Nikiforov, V.B. Uvarov, Polynomials solutions of hypergeometric type difference equations and their classification, *Integral Transforms Spec. Funct.*, 1 (1993), 223–249.
- [34] A. F. Nikiforov, S. K. Suslov, V. B. Uvarov, *Classical orthogonal polynomials of a discrete variable*. Translated from the Russian. Springer Series in Computational Physics. Springer-Verlag, Berlin, 1991.
- [35] M. Weber, A. Erdélyi, On the finite difference analogue of Rodrigues’ formula, *Amer. Math. Monthly* 59 (1952), 163–168.
- [36] J. Zeng, The q -Stirling numbers, continued fractions and the q -Charlier and q -Laguerre polynomials, *J. Comput. Appl. Math.*, t.57 (1995), 413–424.

(Ana F. Loureiro) SCHOOL OF MATHEMATICS, STATISTICS & ACTUARIAL SCIENCE (SMSAS), CORNWALLIS BUILDING, UNIVERSITY OF KENT, CANTERBURY, KENT CT2 7NF (U.K.)

E-mail address, Corresponding author: A.Loureiro@kent.ac.uk

(J. Zeng) UNIVERSITÉ DE LYON, UNIVERSITÉ CLAUDE BERNARD LYON 1, INSTITUT CAMILLE JORDAN, UMR 5208 DU CNRS, 43 BOULEVARD DU 11 NOVEMBRE 1918, 69622 VILLEURBANNE CEDEX (FRANCE)