



HAL
open science

Modelling adhesion-independent cell migration

Gaspard Jankowiak, Diane Peurichard, Anne Reversat, Christian Schmeiser,
Michael Sixt

► **To cite this version:**

Gaspard Jankowiak, Diane Peurichard, Anne Reversat, Christian Schmeiser, Michael Sixt. Modelling adhesion-independent cell migration. 2019. hal-02076901v1

HAL Id: hal-02076901

<https://hal.science/hal-02076901v1>

Preprint submitted on 22 Mar 2019 (v1), last revised 29 Jan 2020 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Modelling adhesion-independent cell migration

Gaspard Jankowiak¹, Diane Peurichard², Anne Reversat³,
Christian Schmeiser⁴, Michael Sixt³

March 22, 2019

1-RICAM-Österreichische Akademie der Wissenschaften, Postgasse 7-9, 1010, Wien, Austria

2-INRIA, 2, rue Simone Iff, 75589 Paris Cedex 12, France

3-Institute of Science and Technology Austria, Am Campus 1, 3400 Klosterneuburg, Austria

4-Faculty of Mathematics, University of Vienna, Oskar-Morgenstern Platz 1, 1090 Vienna, Austria

Abstract

A two-dimensional mathematical model for cells migrating without adhesion capabilities is presented and analyzed. Cells are represented by their cortex, which is modelled as an elastic curve, subject to an internal pressure force. Net polymerization or depolymerization in the cortex is modelled via local addition or removal of material, driving a cortical flow. The model takes the form of a fully nonlinear degenerate parabolic system. An existence analysis is carried out by adapting ideas from the theory of gradient flows. Numerical simulations show that these simple rules can account for the behavior observed in experiments, suggesting a possible mechanical mechanism for adhesion-independent motility.

Keywords: Variational methods; weak solutions; cell motility modelling; cellular cortex; actin polymerization

AMS Subject Classification: 35K40, 35K51, 35Q92, 35A15, 2C17

1 Introduction

One of the most important cellular behaviors is crawling migration. It is observed in many cellular systems both in culture and in vivo [Paluch et al., 2016, Phillipson et al., 2006], and involved in many essential physiological or pathological processes (wound healing, embryonic development, cancer metastasis etc. [Wolf and Friedl, 2006]). Since the works of Abercrombie in 1970 [Abercrombie et al., 1970], which described the multistep model of lamellipodia-based cell migration, numerous authors have studied the mechanisms of actin-based cell migration. As a result, despite some remaining open questions, lamellipodial migration is now well understood and described. This migration mode implies that specific adhesion points transmit intracellular pulling forces from the cytoskeleton to the substrate [Rafelski and Theriot, 2004, Vicente-Manzanares et al., 2009] Actin filaments polymerize below the leading plasma membrane generating pushing forces, and plasma membrane tension resists actin network expansion, pushing back the actin filaments into the cell body. Through adhesion complexes linking the cytoskeleton to the substrate, these retrograde forces are translated into forward locomotion of the cell body [Rafelski and Theriot, 2004].

Yet, recent studies indicate that cell migration can be achieved without specific adhesions in confining three-dimensional environments [Bergert et al., 2015]. Increasing levels of confinement seem to favor adhesion-independent migration in many cell types [Friedl et al., 2001],

and can trigger transitions from integrin-based towards low-adhesive migration modes. However, too strong confinement decreases and even prevents migration [Wolf et al., 2013], due to cell stiffness and nucleus volume. If in the last decade adhesion-independent migration has emerged as a possibly common migration mode, the mechanisms of cell propulsion in this case are still poorly understood. So far in the literature, there is only one known alternative to lamellipodial migration: membrane blebs [Blaser et al., 2006, Charras and Paluch, 2008]. These blebs are cellular extensions free of actin filaments, and generated by intracellular hydrostatic pressure [Tinevez et al., 2009]. Once generated, these blebs grow until a new actin cortex is reassembled inside, which eventually contract and allow the cell to move [Yin Lim et al., 2013]. However, numerous studies display cells migrating in an adhesion-free manner without bleb formation.

Several physical mechanisms have been proposed for force transmission between cell and substrate during migration without focal adhesions: (i) cell migration by swimming (by creation of blebs, [Yin Lim et al., 2013]), (ii) force transmission based on cell-substrate intercalations of lateral protrusions into gaps in the matrix, (iii) chimneying force transmission where cells push against the obstacles [Hawkins et al., 2011, Malawista et al., 2000] or again (iv) flow-friction driven force transmission. In this last case, mechanisms based on non-specific friction between the cell and the substrate have been investigated to account for adhesion-independent migration [Hawkins et al., 2011]. Here, intracellular forces generated by the cytoskeleton are transmitted to the substrate via non-specific friction which has been experimentally measured in [Bergert et al., 2015, Lämmermann et al., 2008]. The molecular origin of nonspecific friction has not been experimentally investigated. Friction could result from interactions between molecules at the cell surface and the substrate, and unveiling the microscopic origin of nonspecific friction will be an important question for future studies.

In this paper, we propose a simplified 2D model for focal adhesion-independent cell migration, based on the mechanisms (iv). We aim to develop a simplified framework to study whether adhesion-free migration could be primarily driven by simple mechanical features.

Our model is focused on the cell's cortex, which is assumed to be an elastic material confined in the horizontal plane. Because of the cytoplasmic pressure, it is subject to outwards pressure forces. Mathematically, this takes the form of a system of parabolic equations, where the polymerization process leads to an advection-type term. This continuous formulation also allows to properly define the reaction forces compensating mass displacement due to membrane renewal. With this very simple model, we are able to trigger cell migration, with speed depending on the geometrical characteristics of the obstacles. Our results seem to be in qualitative agreement with the biological observations. In Section 2, we present the biological observation of leukocyte migrating cells and the experimental setting. Section 3 is concerned with the mathematical model which is analyzed in Section 4. Section 5 presents our numerical results.

2 Leukocyte migration in artificial microchannels

To address the question of adhesion-independent migration of leukocytes, we took advantage of a T cell line, that allows for genetic modifications. In order to remove the link between the adhesion complexes and the substrate, we deleted talin, a cytosolic adaptor protein essential for integrin functionality, using the CRISPR/Cas9 system [Shalem et al., 2014]. In addition, we used well-established microfabricated channels [Vargas et al., 2014] to mimic the confined in vivo environment, coupled to a home-made microfluidic set-up [Reversat et al., 2018]. This set-up allows us to observe that talin KO cells are completely unable to adhere and migrate in

confinement and in smooth-walled channels [Reversat et al., 2018]. Strikingly, their motility is restored in ratchet-baring channels (see Fig. 1).




Figure 1: Leukocyte (in red) migrating from left to right in a ratchet channel. The channel has three sections with wave lengths $6\mu m$, $12\mu m$, and, respectively, $24\mu m$.

3 The mathematical model

Since the mechanisms producing the behavior described in the previous section are not known, we propose a rather simple model for the essential components. The essential idea is that, guided by a chemotactic signal, the cell polarizes with increased actin polymerization near the front end. This is assumed to induce a flow of the cell cortex from front to rear, where depolymerization dominates. The cortex is assumed to be an elastic material with a tendency to equidistribute actin along the cell periphery. This mechanism, together with a constant cytoplasmic excess pressure (actually the pressure difference between cytoplasmic and extracellular pressure), determines the cell shape.

The most critical model ingredient are the forces between the cell and its environment. We assume an unspecific friction with the extracellular liquid, assumed at rest, counteracted by a compensating force, which can be seen as a consequence of the intracellular transport of actin from rear to front. This compensation is chosen such that the cell does not move in an unconstrained environment.




Figure 2: Schematic description of the cell. The cortex is subject to the a number of physical effects: ① pressure forces, ② linear elasticity forces, ③-④ de-/polymerization, ⑤ reaction forces due to transport. Depolymerized actin is transported from the back of the cell (blue region) to the front (green region) where it becomes part of the cortex again via polymerization. The transport inside the cell —in white— is not modeled, but by conservation of the center of mass, it results in a reaction force on the cortex ⑤. The inlet show that at the discrete level (for numerical experiments), on can consider the cortex as a chain of mass points, linked by linear springs.

In accordance with the experimental data we choose a two-dimensional model, which seems reasonable for the experimental situation, where the cell is confined between two flat

surfaces. Fig. 2 illustrates a discrete version of the model, where the cell cortex is described by mass points. We return to this description in Section 5 for simulation purposes, but here we shall formulate a continuous version of the model, where at time $t \geq 0$ the cortex is represented by a Jordan curve

$$\Gamma(t) = \{X(s, t) : s \in \mathbb{T}^1\} \subset \mathbb{R}^2.$$

The one-dimensional torus \mathbb{T}^1 will be represented by intervals of length 1, and the variable s measures the amount of actin material with the total amount normalized to 1. The interior of $\Gamma(t)$ is denoted by $\Omega(t)$, such that $\Gamma(t) = \partial\Omega(t)$. Assuming that $\Gamma(t)$ is smooth enough, we denote by $\tau(t, s)$ and $n(t, s)$ the unit tangent and unit outward normal vectors. Assuming positive orientation of the parametrization, we have

$$\tau = \frac{\partial_s X}{|\partial_s X|}, \quad n = -\tau^\perp,$$

with the convention $(a, b)^\perp = (-b, a)$. The notation is illustrated in Figure 3.




Figure 3: The parameterization and associated vector quantities

The cortex is assumed to be elastic and in equilibrium if $|\partial_s X| = 1$, such that 1 represents the scaled total equilibrium length of the cortex. An elastic resistance against stretching, but not against compression, is described by the potential energy functional

$$E_{\text{el}}(X) = \frac{1}{2} \int_{\mathbb{T}^1} (|\partial_s X| - 1)_+^2 ds. \quad (1)$$

Neglecting resistance against compression can be seen as a convexification of the elastic energy, which facilitates the analysis of Section 4. Actually, we expect the cortex to be always under tension, such that this assumption should not be relevant from a modeling point of view.

This expectation relies on another model ingredient, a cytoplasmic pressure exceeding the extracellular pressure by a constant amount $p > 0$. The associated potential energy contribution is given by

$$E_p(X) = -p |\Omega(t)| = \frac{p}{2} \int_{\mathbb{T}^1} X \cdot \partial_s X^\perp ds. \quad (2)$$

The assumption of a prescribed constant value of p can be seen as a model simplification. Possibly more precise would be the assumption of a fixed prescribed cell volume, measured by the area $|\Omega(t)|$. In this case p would become a time dependent unknown with the mathematical interpretation of a Lagrange multiplier.

The channel walls and other obstacles are modelled by requiring $\Omega(t) \subset \Omega_c$, where the domain $\Omega_c \subset \mathbb{R}^2$ denotes the admissible region. For analysis purposes the obstacles will be softened by introducing the energy contribution

$$E_{\text{obst},\delta}(X) = \int_{\mathbb{T}^1} W_\delta(X) ds, \quad W_\delta = (\delta + \rho_\delta * \mathbb{1}_{\Omega_c})^{-1}, \quad (3)$$

where $\mathbb{1}_{\Omega_c}$ denotes the indicator function of the admissible region, the small parameter $\delta > 0$ measures the softness of the obstacle, and ρ_δ is a positive regularization kernel, approximating the Delta-distribution as $\delta \rightarrow 0$. The formal limit of $E_{\text{obst},\delta}$ as $\delta \rightarrow 0$ takes the value 1, when the constraint $\Omega \subset \Omega_c$ is satisfied, and the value infinity, when it is violated.

The active component of the model comes from actin polymerization and depolymerization in the cortex. We assume that cell polarization manifests itself by local imbalances of this process producing a net increase of actin close to the cell front and a decrease close to the rear of the cell. For simplicity we make the equilibrium assumption that the total amount of actin in the cortex does not change.

We introduce the arclength l , which is given by

$$l(s, t) = \int_0^s |\partial_s X(\sigma, t)| d\sigma. \quad (4)$$

This relation between the arc length l and the Lagrangian variable s can be inverted in terms of the actin density $\rho(l, t)$ per arc length:

$$s(l, t) = \int_0^l \rho(\lambda, t) d\lambda,$$

implying $|\partial_s X| = \rho^{-1}$. We denote by $f(l, t)$ the rate of actin increase ($f > 0$) or decrease ($f < 0$) per time unit and per arc length, so that ρ satisfies

$$\partial_t \rho = f.$$

The above mentioned equilibrium assumption translates to

$$\int_{\Gamma} f dl = \int_{\mathbb{T}^1} f(l(s, t)) |\partial_s X(s, t)| ds = 0, \quad t \geq 0. \quad (5)$$

We then obtain the material derivative for functions of s :

$$\frac{D}{Dt} = \partial_t + \left(\int_0^{l(s,t)} f(\lambda, t) d\lambda \right) \partial_s, \quad (6)$$

which has to be understood relative to the arc length along Γ , measured from the point $X(0, t)$. In particular, the velocity of the cortex relative to the laboratory coordinates is given by DX/Dt .

With the prescription of $f(l, t)$ the total amount of relocated actin is not controlled, since the arclength of the cortex varies with the dynamics. Therefore we shall consider

$$g(s, t) = f(l(s, t), t) |\partial_s X(s, t)|, \quad (7)$$

the rate of change of actin mass per actin mass along the cortex, as given. This roughly assumes growth and decay rates given per actin filament. This leads to

$$\frac{D}{Dt} = \partial_t + \left(\int_0^s g(\sigma, t) d\sigma \right) \partial_s, \quad (8)$$

Friction between the cell surface and the surrounding fluid, which is assumed nonmoving, is modeled by the L^2 -gradient flow for the total energy with the contributions (1), (2), and (3), where the friction force is given in terms of the material derivative:

$$\begin{aligned}\frac{DX}{Dt} &= -E'_{\text{el}}(X) - E'_p(X) - E'_{\text{obst},\delta}(X) \\ &= \partial_s \left((|\partial_s X| - 1)_+ \frac{\partial_s X}{|\partial_s X|} \right) - p \partial_s X^\perp - \nabla W_\delta(X).\end{aligned}$$

Note that the friction coefficient has been eliminated by an appropriate choice of the time scale. This is however not yet the final form of the model. It would predict movement of freely floating cells in the absence of any confinement, as can easily be seen by integrating the equation in the absence of the last term with respect to s :

$$\frac{d}{dt} \int_{\mathbb{T}^1} X ds = - \int_{\mathbb{T}^1} \left(\partial_s X \int_0^s g(\sigma, t) d\sigma \right) ds = \int_{\mathbb{T}^1} X g ds.$$

Since the right hand side will in general be different from zero, the center of mass will move. An explanation is the force, used to move depolymerized G-actin from regions with $g < 0$ to regions with $g > 0$. By the action-reaction principle it seems reasonable to introduce a compensating counterforce with density $\mathcal{F}_{\text{comp}}(s, t)$, acting on the cortex:

$$\partial_t X + \partial_s X \int_0^s g d\sigma = \partial_s \left((|\partial_s X| - 1)_+ \frac{\partial_s X}{|\partial_s X|} \right) - p \partial_s X^\perp - \nabla W_\delta(X) + \mathcal{F}_{\text{comp}}. \quad (9)$$

It can be chosen arbitrarily, except that the total force is fixed:

$$\int_{\mathbb{T}^1} \mathcal{F}_{\text{comp}} ds = - \int_{\mathbb{T}^1} X g ds. \quad (10)$$

We do not make a choice for $\mathcal{F}_{\text{comp}}$ at this point, the precise choice made for our numerical experiments will be discussed in Section 5. One can note that $\mathcal{F}_{\text{comp}}$ is roughly directed from the front towards to back of the cell, since g is negative (respectively positive) where the depolymerization (respectively polymerization).

4 Existence results

4.1 Formulation of the main results

We shall prove a global existence result for the initial value problem for (9) and a convergence result for the limit $\delta \rightarrow 0$ of hard channel walls.

Since the terms describing the cortical flow are chosen to have a vanishing integral with respect to s , we introduce S_{flow} , which is chosen such that

$$\partial_s S_{\text{flow}}[X] = \partial_s X \int_0^s g d\sigma - \mathcal{F}_{\text{comp}}.$$

The initial value problem can then be written in the form

$$\partial_t X + \partial_s(S_{\text{flow}}[X]) = \partial_s \left((|\partial_s X| - 1)_+ \frac{\partial_s X}{|\partial_s X|} \right) - p \partial_s X^\perp - \nabla W_\delta(X), \quad (11)$$

$$X(s, 0) = X^0(s). \quad (12)$$

For function spaces we use the abbreviations $\mathcal{L} := L^2(\mathbb{T}^1)^2$ and $\mathcal{H} := H^1(\mathbb{T}^1)^2$ with the norms $\|\cdot\|_2$ and, respectively, $\|\cdot\|_{1,2}$, where the torus \mathbb{T}^1 is represented by the s -interval $(0, 1)$, and also $\|\cdot\|_p$ for the $L^p(\mathbb{T}^1)$ -norm, $p \geq 1$, and $\langle \cdot, \cdot \rangle_2$ for the \mathcal{L} -scalar product. For the data we shall use the following assumptions:

$$(A1) \quad p < 2\pi,$$

$$(A2) \quad \|S_{\text{flow}}[X]\|_2 \leq c_{\text{flow}}\|X\|_2, \quad \|\partial_s(S_{\text{flow}}[X])\|_2 \leq c_{\text{flow}}\|X\|_{1,2}.$$

$$(A3) \quad W_\delta \text{ is given by (3), where the domain } \Omega_c \subset \mathbb{R}^2 \text{ has a smooth boundary, and } x \cdot \nabla W_\delta(x) \geq 0 \text{ for } x \in \mathbb{R}^2 \text{ and for } \delta \text{ small enough.}$$

$$(A4) \quad X^0 \in \mathcal{H} \text{ and } \{X^0(s) : s \in (0, 1)\} \subset \Omega_c.$$

Assumption (A1) can be motivated by looking at the simplified problem without cortical flow and without obstacles, i.e. $\nabla W_\delta = S_{\text{flow}} = 0$. In this case the expected circular equilibrium state X_{equ} exists only under Assumption (A1) and is given by

$$X_{\text{equ}}(s) = \frac{1}{2\pi - p} (\cos(2\pi s), \sin(2\pi s)).$$

Assumption (A2) is somewhat restrictive compared to the models discussed in Section 3. It can be satisfied for bounded actin growth rate g . The inequality in Assumption (A3) means roughly that the allowed domain Ω_c is star shaped (and that its center has been taken as the origin). Finally, Assumption (A4) on the initial cortex shape implies that the elastic and pressure energies are finite and that the cell lies in the admissible region. It might be noted that we do not assume that X_0 parametrizes a simple curve. We do not refer to this property since our results do not guarantee that it is preserved globally in time.

Theorem 1. *(Global existence for the penalized problem) Let the Assumptions (A1)–(A4) hold and let $\delta > 0$ be small enough. Then there exists a solution X_δ of (11), (12), such that*

$$X_\delta \in H_{\text{loc}}^1(\mathbb{R}_+; \mathcal{L}) \cap L_{\text{loc}}^\infty(\mathbb{R}_+; \mathcal{H}),$$

uniformly with respect to $\delta \rightarrow 0$.

Remark 1. *By the Morrey inequality proved in the appendix, X_δ is Hölder continuous with exponent $\frac{1}{2}$ in terms of s and $\frac{1}{4}$ in terms of t , again uniformly in δ .*

The proof will be carried out in the following three sections. It relies on methods for gradient flows, although they cannot be applied in a straightforward way. The terms on the right hand side of (11) are the L^2 -gradients of the energy functionals E_{el} , E_p , and $E_{\text{obst},\delta}$ (see Section 3), the first of which is convex. The second and third are treated as continuous perturbations. The cortical flow term on the left hand side is nonvariational.

Our approach is based on a semi-implicit time discretization, where the cortical flow term is evaluated at the old time step. This allows to solve the discrete problem by energy minimization. A priori estimates for the discrete solution allow to pass to the continuous limit.

These estimates are also uniform in the penalization parameter δ for the potential, so we can also carry out the high penalization limit:

Theorem 2. *(Limit of hard channel walls) With the assumptions of Theorem 1, the family $\{X_\delta : \delta > 0\}$ of solutions of (11), (12) contains a sequence, converging (as $\delta \rightarrow 0$) uniformly on bounded time intervals to*

$$X \in H_{\text{loc}}^1(\mathbb{R}_+; \mathcal{L}) \cap L_{\text{loc}}^\infty(\mathbb{R}_+; \mathcal{H}),$$

which satisfies $X \in \overline{\Omega_c}$ on $\mathbb{R}_+ \times \mathbb{T}^1$, (12), and

$$\partial_t X + \partial_s(S_{\text{flow}}[X]) = \partial_s \left((|\partial_s X| - 1)_+ \frac{\partial_s X}{|\partial_s X|} \right) - p \partial_s X^\perp, \quad (13)$$

for all (s, t) such that $X(s, t) \in \Omega_c$.

The properties of X stated in the theorem are not a complete formulation of the obstacle problem. Information on the behavior at the edges of contact regions is missing. Since the equation is degenerately parabolic, this is not so obvious. Under the additional assumption of convexity of the permissible set Ω_c , one could expect that X solves the variational inequality

$$\left\langle \partial_t X + \partial_s(S_{\text{flow}}[X]) - \partial_s \left((|\partial_s X| - 1)_+ \frac{\partial_s X}{|\partial_s X|} \right) + p \partial_s X^\perp, Y - X \right\rangle_2 \geq 0,$$

for all $Y \in \mathcal{H}$ such that $Y(s) \in \overline{\Omega_c}$, $s \in \mathbb{T}^1$.

For the numerical experiments, the obstacle is not modeled by a potential. For any $X(s, t) \in \Gamma(t)$, the total resulting force is rather restricted to the tangent cone to Ω_c at $X(s, t)$. In other words, our numerical simulations will be based on the formulation

$$\partial_t X(s, t) = \mathbf{P}_c \left(\partial_s \left((|\partial_s X| - 1)_+ \frac{\partial_s X}{|\partial_s X|} \right) - p \partial_s X^\perp - \partial_s(S_{\text{flow}}[X]) \right), \quad (14)$$

where the potential W_δ does not appear. The \mathbf{P}_c is the projection on the tangent cone:

$$\mathbf{P}_c(F(s, t)) = \begin{cases} (F(s, t) \cdot \tau_c(s, t)) \tau_c(s, t) & \text{for } X \in \partial\Omega_c, F(s, t) \cdot n_c(s, t) > 0, \\ F(s, t) & \text{else,} \end{cases} \quad (15)$$

where τ_c and n_c are a normalized tangent vector and, respectively, the normalized outward normal along $\partial\Omega_c$.

4.2 The energy functional and its properties

We introduce the total energy functional

$$\begin{aligned} E_\delta(X) &:= E_{\text{el}}(X) + E_{\text{p}}(X) + E_{\text{obst}, \delta}(X) \\ &= \begin{cases} \int_{\mathbb{T}^1} \left(\frac{1}{2} (|\partial_s X| - 1)_+^2 + \frac{p}{2} X \cdot \partial_s X^\perp + W_\delta(X) \right) ds & \text{if } X \in \mathcal{H} \\ +\infty & \text{otherwise,} \end{cases} \end{aligned}$$

and prove its coercivity:

Lemma 1 (Coercivity). *For $p < 2\pi$ and for all $X \in \mathcal{L}$*

$$E_\delta(X) \geq \frac{1}{2} \left(1 - \frac{p}{2\pi} \right) \|\partial_s X\|_2^2 - \|\partial_s X\|_2 \geq -\frac{\pi}{2\pi - p}. \quad (16)$$

Proof. W.l.o.g. we assume $X \in \mathcal{H}$ and denote by Ω the enclosed domain and by L the length of its boundary. Then the isoperimetric and the Cauchy-Schwarz inequalities imply

$$E_{\text{p}}(X) = -p|\Omega| \geq -p \frac{L^2}{4\pi} = -\frac{p}{4\pi} \|\partial_s X\|_1^2 \geq -\frac{p}{4\pi} \|\partial_s X\|_2^2.$$

For the elastic energy again the Cauchy-Schwarz inequality is used:

$$\begin{aligned} E_{\text{el}}(X) &= \frac{1}{2} \int_{\mathbb{T}^1} (|\partial_s X| - 1)^2 ds - \frac{1}{2} \int_{|\partial_s X| < 1} (|\partial_s X| - 1)^2 ds \geq \frac{1}{2} (\|\partial_s X\|_2^2 - 2\|\partial_s X\|_1 + 1) - \frac{1}{2} \\ &\geq \frac{1}{2} \|\partial_s X\|_2^2 - \|\partial_s X\|_2, \end{aligned}$$

which completes the proof, since $E_{\text{obst},\delta} \geq 0$. \square

Remark 2. This result shows that the definition of E_δ has been appropriate, since finiteness of $E_\delta(X)$ implies membership of X in \mathcal{H} .

Lemma 2. With respect to the weak topology in \mathcal{H} , the functional E_{el} is convex and lower semicontinuous, and the functionals E_p and $E_{\text{obst},\delta}$ are continuous.

Proof. The semicontinuity of E_{el} follows from [Giaquinta, 1983, Theorem 2.5]. The other properties are straightforward. \square

4.3 Time discretization

Choose $\tau > 0$ and $X^{n-1} \in \mathcal{H}$. Then, by the results of the preceding section, the functional

$$\Phi(\tau, X^{n-1}; Y) := \frac{\|Y - X^{n-1}\|_2^2}{2\tau} + E_\delta(Y) + \langle Y, \partial_s(S_{\text{flow}}[X^{n-1}]) \rangle_2$$

is weakly lower semicontinuous on \mathcal{H} . It is also bounded from below since E_δ is, and since

$$\begin{aligned} \frac{\|Y - X^{n-1}\|_2^2}{2\tau} + \langle Y, \partial_s(S_{\text{flow}}[X^{n-1}]) \rangle_2 &\geq -\frac{\tau}{2} \|\partial_s(S_{\text{flow}}[X^{n-1}])\|_2^2 - \|X^{n-1}\|_2 \|\partial_s(S_{\text{flow}}[X^{n-1}])\|_2 \\ &\geq -\left(\frac{\tau}{2} c_{\text{flow}}^2 + c_{\text{flow}}\right) \|X^{n-1}\|_{1,2}^2. \end{aligned}$$

Furthermore, sublevel sets are bounded in \mathcal{H} . This is sufficient for the minimum of $\Phi(\tau, X^{n-1}; \cdot)$ to be assumed in \mathcal{H} , and we choose

$$X^n \in \operatorname{argmin}_{Y \in \mathcal{H}} \Phi(\tau, X^{n-1}; Y), \quad n \geq 1.$$

We define $X_\tau \in C_{\text{loc}}^{0,1}(\mathbb{R}_+, \mathcal{H})$ as the piecewise linear interpolation of the X^n , $n \geq 0$. More precisely, we have

$$X_\tau(t) := X^{k_t} + \frac{(t - k_t\tau)}{\tau} (X^{k_t+1} - X^{k_t}), \quad (17)$$

where $k_t \in \mathbb{N}$ is such that $k_t\tau \leq t < (k_t + 1)\tau$.

Lemma 3 (Uniform estimates for the interpolation). *Let Assumptions (A1)–(A4) hold. Then*

$$X_\tau \in H_{\text{loc}}^1(\mathbb{R}_+, \mathcal{L}) \cap L_{\text{loc}}^\infty(\mathbb{R}_+, \mathcal{H}),$$

uniformly with respect to small enough τ and δ .

Proof. Denoting the duality bracket with $\langle \cdot, \cdot \rangle$, the variation of Φ in the direction $Y \in \mathcal{H}$ gives

$$\frac{\langle X^n - X^{n-1}, Y \rangle_2}{\tau} + \langle \partial E_\delta(X^n), Y \rangle - \langle \partial_s Y, S_{\text{flow}}[X^{n-1}] \rangle_2 = 0, \quad (18)$$

with the formal gradient of the energy determined from

$$\langle \partial E_\delta(X^n), Y \rangle_2 = \int_{\mathbb{T}^1} \left((|\partial_s X^n| - 1)_+ \frac{\partial_s X^n \cdot \partial_s Y}{|\partial_s X^n|} + Y \cdot (p \partial_s(X^n)^\perp + \nabla W_\delta(X^n)) \right) ds$$

With $Y = X^n$ and with Assumptions (A1) and (A3) we get, similarly to the proof of Lemma 1,

$$\langle \partial E_\delta(X^n), X^n \rangle_2 \geq \int_{\mathbb{T}^1} (|\partial_s X^n| - 1)_+ |\partial_s X^n| ds - 2p|\Omega^n| \geq \left(1 - \frac{p}{2\pi}\right) \|\partial_s X^n\|_2^2 - \|\partial_s X^n\|_2 - \frac{1}{4},$$

and we observe

$$\langle X^n - X^{n-1}, X^n \rangle_2 \geq \frac{1}{2} (\|X^n\|_2^2 - \|X^{n-1}\|_2^2).$$

Finally, we use

$$|\langle \partial_s X^n, S_{\text{flow}}[X^{n-1}] \rangle_2| \leq \gamma \|\partial_s X^n\|_2^2 + \frac{c_{\text{flow}}^2}{4\gamma} \|X^{n-1}\|_2^2$$

with $0 < \gamma < 1 - p/(2\pi)$. This implies the existence of positive constants A_1, A_2, A_3 (independent from n, τ and δ) such that

$$\|X^n\|_2^2 + \tau A_1 \|\partial_s X^n\|_2^2 \leq (1 + \tau A_2) \|X^{n-1}\|_2^2 + \tau A_3.$$

A discrete Gronwall estimate now gives

$$\|X^n\|_2^2 \leq \|X^0\|_2^2 e^{A_2 n \tau} + \frac{A_3}{A_2} (e^{A_2 n \tau} - 1),$$

and, as a consequence,

$$\tau \sum_{k=1}^n \|\partial_s X^k\|_2^2 \leq C(n\tau). \quad (19)$$

From the last two bounds we get for any $t_1, t_2 > 0$

$$\sup_{t \in [t_1, t_2]} \|X_\tau(t)\|_2^2 \leq \|X^0\|_2^2 e^{A_2 k_2 \tau} + \frac{A_3}{A_2} (e^{A_2 k_2 \tau} - 1),$$

and

$$\|\partial_s X_\tau\|_{L^2([t_1, t_2])}^2 \leq \sum_{k=k_1}^{k_2} \tau \left(\|\partial_s X^k\|_2^2 + \|\partial_s X^{k+1}\|_2^2 \right) \leq 2C(k_2 \tau),$$

where $k_2 \tau \leq t_2 < (k_2 + 1)\tau$. In other words, we have shown $X_\tau \in L_{\text{loc}}^\infty(\mathbb{R}_+, \mathcal{L}) \cap L_{\text{loc}}^2(\mathbb{R}_+, \mathcal{H})$ uniformly in τ and δ .

From the minimization we get

$$\frac{\|X^n - X^{n-1}\|_2^2}{2\tau} + E_\delta(X^n) + \langle X^n, \partial_s(S_{\text{flow}}[X^{n-1}]) \rangle_2 \leq E_\delta(X^{n-1}) + \langle X^{n-1}, \partial_s(S_{\text{flow}}[X^{n-1}]) \rangle_2.$$

With the velocity $\partial_t X_\tau^n = \frac{X^n - X^{n-1}}{\tau}$ of the linear interpolant this reads

$$\frac{\tau}{2} \|\partial_t X_\tau^n\|_2^2 + E_\delta(X^n) \leq E_\delta(X^{n-1}) - \tau \langle \partial_t X_\tau^n, \partial_s(S_{\text{flow}}[X^{n-1}]) \rangle_2$$

With the Young inequality and Assumption (A2) we obtain

$$\frac{\tau}{4} \|\partial_t X_\tau^n\|_2^2 + E_\delta(X^n) \leq E_\delta(X^{n-1}) + \tau c_{\text{flow}}^2 \|\partial_s X^{n-1}\|_2^2$$

Using (19), summation over n and an application of Lemma 1 complete the proof. \square

4.4 The continuous and high penalization limits

Considering the time discrete solution $(X^n)_{n \geq 0}$, we recall the definition (17) of X_τ and define two additional piecewise constant continuous-time approximations:

$$\begin{cases} X_\tau^{\text{old}}(t, s) & := X^{k_t}(s), \\ X_\tau^{\text{new}}(t, s) & := X^{k_t+1}(s), \end{cases} \quad \text{where } k_t \tau < t < (k_t + 1)\tau,$$

Then (18) implies

$$\partial_t X_\tau + \partial E_{\text{el}}(X_\tau^{\text{new}}) + p \partial_s (X_\tau^{\text{new}})^\perp + \nabla W_\delta(X_\tau^{\text{new}}) + \partial_s S_{\text{flow}}[X_\tau^{\text{old}}] = 0, \quad (20)$$

where $\partial E_{\text{el}}(X)$ is the subdifferential of the elastic energy, given by the distributional derivative with respect to s of $-(|\partial_s X| - 1)_+ \frac{\partial_s X}{|\partial_s X|}$. The equality only holds in the dual space of \mathcal{H} *a priori*, but thanks to Lemma 3, it also holds in \mathcal{L} . It is our goal to pass to the limit $\tau \rightarrow 0$ in this equation. By the results of the previous section, there exists a sequence $\tau_k \rightarrow 0$ such that

$$\lim_{k \rightarrow \infty} X_{\tau_k} = \lim_{k \rightarrow \infty} X_{\tau_k}^{\text{new}} = \lim_{k \rightarrow \infty} X_{\tau_k}^{\text{old}} = X \quad \text{in } L_{\text{loc}}^2(\mathbb{R}_+, \mathcal{L}).$$

We also have that all the terms in (20) except $\partial E_{\text{el}}(X_{\tau_k}^{\text{new}})$ are bounded in $L_{\text{loc}}^2(\mathbb{R}_+, \mathcal{L})$ uniformly in τ and δ . In all these terms we can pass to the limit in the sense of distributions, but also weakly in $L_{\text{loc}}^2(\mathbb{R}_+, \mathcal{L})$.

This also implies weak convergence of $\partial E_{\text{el}}(X_{\tau_k}^{\text{new}})$ to some $\eta \in L_{\text{loc}}^2(\mathbb{R}_+, \mathcal{L})$, and we obtain

$$\partial_t X + \eta + p \partial_s X^\perp + \nabla W_\delta(X) + \partial_s S_{\text{flow}}[X] = 0.$$

It remains to identify η .

Since the subdifferential is a maximal monotone operator, we shall apply the Minty trick [Hungerbühler, 2000, Theorem 2.2], [Minty, 1963]. Testing (20) against $X_{\tau_k}^{\text{new}}$, its strong convergence immediately implies

$$\lim_{k \rightarrow \infty} \int_0^T \langle \partial E_{\text{el}}(X_{\tau_k}^{\text{new}}), X_{\tau_k}^{\text{new}} \rangle_2 dt = \int_0^T \langle \eta, X \rangle_2 dt,$$

which is sufficient for $\eta \in \partial E_{\text{el}}(X)$ and, thus,

$$\eta = -\partial_s \left((|\partial_s X| - 1)_+ \frac{\partial_s X}{|\partial_s X|} \right),$$

to be understood as the weak derivative of an L^2 -function, since $X \in L_{\text{loc}}^\infty(\mathbb{R}_+, \mathcal{H})$. This completes the proof of Theorem 1.

For the proof of Theorem 2, we note that by the results of Theorem 1 and following Remark 1 the uniform convergence of a subsequence to X is immediate. The uniform boundedness of $E_{\text{obst}, \delta}$ implies $X \in \overline{\Omega_c}$ by its continuity and by the uniform convergence, since $W_\delta(x) \rightarrow \infty$ for $x \notin \overline{\Omega_c}$.

Similarly, since $\nabla W_\delta(x) \rightarrow 0$ for $x \in \Omega_c$, the weak formulation of (13) can be derived by using test functions vanishing away from $\{(s, t) : X(s, t) \in \Omega_c\}$. The convergence of the various terms is handled exactly as for the continuous limit.

5 Numerical results

5.1 Discretization

We start with the situation without obstacles and introduce $N \in \mathbb{N}$ grid-points for the discretization of s such that $s_i = i\Delta s$, $\Delta s = \frac{1}{N}$ for i considered on a discrete torus, meaning that i is identified with $i + N$. The time step is $\Delta t > 0$ and $t^n = n\Delta t$, $n \geq 0$. The numerical approximation for $X(s_i, t^n)$ is denoted by X_i^n .

We assume a cell which is polarized in the fixed direction $\omega \in \mathbb{R}^2$, $|\omega| = 1$ and define i_0^n and i_1^n such that

$$\omega \cdot X_{i_0^n}^n = \max_i \omega \cdot X_i^n, \quad \omega \cdot X_{i_1^n}^n = \min_i \omega \cdot X_i^n. \quad (21)$$

Actin is added to the cortex at the leading end $X_{i_0^n}^n$ and removed at the trailing end $X_{i_1^n}^n$, corresponding to $s_0(t^n) = i_0^n \Delta s$, $s_1(t^n) = i_1^n \Delta s$ in the notation of Section 3.

We use the explicit Euler scheme for the time discretization and symmetric finite differences for the discretization in the s -direction:

$$\frac{X_i^{n+1} - X_i^n}{\Delta t} = \mathbf{F}[X^n]_i := v \mathbb{1}_{i_1 \leq i \leq i_0} \frac{X_{i+1}^n - X_{i-1}^n}{2\Delta s} + \frac{G_{i+1/2}^n - G_{i-1/2}^n}{\Delta s} - p \frac{(X_{i+1}^n - X_{i-1}^n)^\perp}{2\Delta s} + F_{\text{comp},i}^n,$$

where the spring forces are given by

$$G_{i+1/2} = \left(\frac{|X_{i+1} - X_i|}{\Delta s} - 1 \right)_+ \frac{X_{i+1} - X_i}{|X_{i+1} - X_i|}, \quad (22)$$

and the compensating force by

$$F_{\text{comp},i}^n = \frac{v}{2} \left(\frac{X_{i_1^n-1}^n + X_{i_1^n}^n}{2} - \frac{X_{i_0^n+1}^n + X_{i_0^n}^n}{2} \right) (\tilde{\delta}_a(i - i_0^n) + \tilde{\delta}_a(i - i_1^n)).$$

Here $\tilde{\delta}_a(i)$ is a mollification of the Dirac Delta in s_i , with smoothing parameter a such that $\sum_i \tilde{\delta}_a(i) \Delta s = 1$. The approximation for the compensating force has been chosen such that the discrete center of mass $\sum_i X_i^n \Delta s$ is independent of time (in the absence of obstacles).

The restriction to the admissible domain Ω_c is enforced by an approximation of the formulation (14), (15):

$$\frac{X_i^{n+1} - X_i^n}{\Delta t} = \mathbf{P}_{c,\epsilon}(\mathbf{F}[X^n]_i),$$

where

$$\mathbf{P}_{c,\epsilon}(\mathbf{F}[X]_i) = \begin{cases} (\mathbf{F}[X]_i \cdot \vec{\mathbf{t}}) \vec{\mathbf{t}} & \text{for } \text{dist}(X_i, \partial\Omega_c) < \epsilon, \mathbf{F}[X]_i \cdot \vec{\mathbf{n}} > 0, \\ \mathbf{F}[X]_i & \text{else,} \end{cases}$$

with the tangent and normal vectors evaluated at the orthogonal projection of X_i to $\partial\Omega_c$ (see Fig. 4). The width ϵ of the tube, where the projection is applied, is coordinated with the choice of the time step such that the admissible domain is not left.




Figure 4: Sketch of the projection of the forces acting on the cell cortex close to the walls.

5.2 Dimensionalisation of the model parameters

We denote by κ_S the spring force and by μ the internal friction coefficient in front of the time derivative. Note that all these quantities have been set to 1 so far in the model, without loss of generality. We recover them here for the sake of computing their dimensions, Eq. (14) reads:

$$\mu \partial_t X = \mathbf{P}_c \left(\kappa_S \partial_s \left((|\partial_s X| - 1)_+ \frac{\partial_s X}{|\partial_s X|} \right) - p \partial_s X^\perp - \frac{1}{\mu} \partial_s (S_{\text{flow}}[X]) \right).$$

We denote by x_0 and t_0 the space and time units of the model. The space unit is chosen to be half the cortex length of leucocytes $L = 2x_0$, i.e $x_0 \approx 47.6 \mu\text{m}$. The polymerization speed in leucocytes is around $12 \mu\text{m} \cdot \text{min}^{-1}$ and the dimensionless polymerization speed writes $v = 2 \frac{x_0}{t_0}$ therefore 1 time unit of the model corresponds to $t_0 \approx 8 \text{min}$. The elastic properties of the human red blood cell have been studied via micropipette aspiration experiments in [Waugh and Evans, 1979] where the authors show that the stiffness constant of leucocytes membrane is of order $k_S = 7 \text{pN} \cdot \mu\text{m}^{-1}$. In our model $\frac{k_S}{\mu} = \frac{1}{t_0} = 0.1265 \text{min}^{-1}$ therefore we can deduce that the internal friction coefficient $\mu \approx 55 \text{pN} \cdot \mu\text{m}^{-1} \cdot \text{min}$. Finally, we chose the pressure constant $p = 3.2$. We deduce $\frac{p}{\mu} = 3.2 \frac{1}{t_0} = 0.4 \text{min}^{-1}$ and therefore $p \approx 22 \text{pN} \cdot \mu\text{m}^{-1}$ which is in biological range. All the model parameters are summarized in Table 1.

Parameters			
Symbol	Numerical Value	Biological value	Description
Numerical parameters			
Δt	$4 \cdot 10^{-2}$	N/A	Time step
Δs	$5 \cdot 10^{-3}$	N/A	Discretization step
N	200	N/A	Number of nodes
ϵ	0.1	N/A	Distance to the obstacle for the projection
a	adapted	N/A	Length of the polymerization zone
Model parameters			
L	1	$48 \mu m$	Reference membrane length
L_{eff}	2	$94 \mu m$	Effective membrane length -In biological range
κ_S	1	$7 \mu pN \cdot \mu m^{-1}$	Stiffness constant -[Waugh and Evans, 1979]
p	3.2	$22 pN \cdot \mu m^{-1}$	Pressure force -In biological range
v	2	$12 \mu m \cdot min^{-1}$	Polymerization speed -In biological range
μ	1	$55 pN \cdot \mu m^{-1} \cdot min$	Internal friction coefficient In biological range

Table 1: Numerical and model parameters for the simulations of the paper.

5.3 Simulations without Obstacle

In Fig. 5 we show computed equilibrium shapes of the cortex in the absence of obstacles. The polarization of the cell is always to the left (i.e. $\omega = (-1, 0)$ in (21)).

Fig. 5 (A) shows a simulation without compensating force. As expected, the cell migrates in the direction of the leading end with a speed slightly less than the polymerization/depolymerization speed. The equilibrium shape is a circle.

Figs. 5 (B,C,D) show the equilibrium shapes obtained for different values of $h = \frac{a}{L}$, where we recall that a measures the spread of the compensating force and L is the equilibrium circumference. The most important observation is that migration has been turned off successfully by including the compensating force. The deviation of the equilibrium shape from a circle is stronger for more concentrated compensating forces (smaller values of a therefore $\tilde{\sigma}$, compare Figs. (D) to (B)).





Figure 5: Different cell shapes obtained for different values of h : (A) Without compensating force. (B,C,D, E) with compensating force: (B) $h = 0.01$, (C) $h = 0.04$, (D) $h = 0.1$. Solid: cell cortex, dashed: circle with the same circumference.

5.4 Simulations of Migration in Channels

In order to reproduce the biological experiments, our protocol for simulations with channels is started by putting the cell with an initially circular shape close to the opening of a channel. Then we let it evolve to an equilibrium shape as in the preceding section, after which we turn on a force pushing it into the channel. This is realized by turning off the pressure force in a region around the trailing end. The pushing is removed as soon as the cell is entirely inside the channel.

(A) $\sigma = 0.04$,
Channel width = $2.16 \mu\text{m}$



(B) $\sigma = 0.04$,
Channel width = $3.7 \mu\text{m}$




Figure 6: Numerical simulations of cells pushed into channels with flat walls for $h = 0.04$ and channel width $2.16 \mu\text{m}$ (A) and for $h = 0.8$ and channel width $3.7 \mu\text{m}$ (B). For each simulation, we show in the top left corner the initial condition, and in the top right corner the final position.

In channels with flat walls with varying widths and spreads of the compensating force, no migration is observed in the simulations in spite of polarization and corresponding cortex flow (Fig.6). These results are in agreement with the experiments described in Section 2.

Ratchet channels are described by four parameters: (i) a wave length L_0 of the width variations, (ii) a minimal width $2w_0$, (iii) an amplitude of the width variations d_0 , and (iv) an asymmetry parameter α . For the length x along the channel, the walls of the channel are given by $\pm f(x)$, with the function

$$f(x) = d_0(g(x) - 1) - w_0,$$

where $g(x)$ solves the fixed point equation:

$$g(x) = \sin\left(\frac{2\pi x}{L_0} + \alpha g(x)\right).$$

In all our numerical simulations we choose $\alpha = 0.4$. Note that for $\alpha = 0$, the walls would be sinusoidal functions of x , i.e. symmetric with respect to the reflection $x \leftrightarrow -x$. As in the experiments, in each simulated channel we combine three wavelengths, $L_0 = 3.9, 7.6, 11.7 \mu\text{m}$, increasing in the migration direction. Fig. 7 (I) shows four examples with $2w_0 = 1.4 \mu\text{m}$ (A and C) and $2w_0 = 3.7 \mu\text{m}$ (B and D), with $d_0 = 0.78 \mu\text{m}$ (A and B) and $d_0 = 3.1 \mu\text{m}$ (C and D). As further illustration, Fig. 7 (II) shows a cell at equilibrium before being pushed into a ratchet with $L_0 = 7.6 \mu\text{m}$, $2w_0 = 3.7 \mu\text{m}$, and $d_0 = 1.54 \mu\text{m}$.




Figure 7: (I) Examples of the different obstacle geometries used in the simulations: we explore narrow obstacles of minimal width $1.4 \mu\text{m}$ (A and C) or wide obstacles of minimal width $3.7 \mu\text{m}$ (B and D), smooth ratchets of amplitude $0.78 \mu\text{m}$ (A and B) or larger ratchets of amplitude $3.1 \mu\text{m}$ (C and D). (II) Example of a stationary cell (at equilibrium), before being pushed into a ratchet channel of wavelength $7.6 \mu\text{m}$, minimal width $3.7 \mu\text{m}$ and amplitude $1.54 \mu\text{m}$.

Results of a typical simulation are shown in Fig. 8. In this situation the cell is able to migrate in all three different wavelengths. It seems that the average speed is highest in the part with the intermediate wavelength. Within each wavelength region, the behavior seems to be periodic related to the periodicity of the channel walls. The bigger the wavelength, the more we observe a jump-like migration pattern indicated by peaks in the cell speed.




Figure 8: Numerical simulation of a cell in a ratchet channel of minimal width $1.4 \mu m$, amplitude $2.7 \mu m$, and variable wave lengths (3.9 , 7.6 , and $11.7 \mu m$), for $h = 0.1$. Top left: initial condition. Top right: at time $2h$. Bottom left: circumference vs. time. Bottom right: Speed of the center of gravity vs. time.

A study of the dependence of the mean cell speed on the various parameters has been carried out (Fig.9). Not surprisingly, increasing the channel width decreases the cell speed and can prevent cell migration. For ratchets of the smallest wavelength $3.9 \mu m$, the cell is unable to migrate for large amplitudes. In this situation the cell is unable to enter the ratchets, thus reducing its contact to the wall. The channel acts in these cases as if it had flat walls. For period $7.6 \mu m$ ratchets, the optimal speed is obtained for large enough amplitude. This is expected, since for small amplitude the ratchet again acts as a flat walled channel. This fact is reinforced in channels of period $11.6 \mu m$, where only ratchets with large amplitude can induce cell migration.

To sum up, cell speed is directly linked to the cell compression induced by the channel geometry and to the possibility for the cell to push against parts of the channel wall facing towards the migration direction.




Figure 9: Average cell speeds ($\mu\text{m}\cdot\text{min}^{-1}$) in channels of different minimal widths w_0 : $1.4\mu\text{m}$ (A), $2.2\mu\text{m}$ (B), $2.9\mu\text{m}$ (C) and $3.7\mu\text{m}$ (D). In each case, the cell speed is represented in dependence of the spread $h \in [0.04, 0.8]$ of the compensating force (vertical direction) and of the amplitude $d_0 \in [0.78, 3.1]\mu\text{m}$ (horizontal direction). We consider channel walls with wave lengths $3.8\mu\text{m}$ (left figures), $7.6\mu\text{m}$ (middle figures) and $11.7\mu\text{m}$ (right figures).

6 Conclusions

This study has been motivated by the experimental results described in Section 2, where adhesion-free migration of leukocytes in artificial, structured micro-channels has been observed.

A mathematical model for this process has been formulated. Migration is assumed to be due to cortex flow, driven by a local imbalance of polymerization and depolymerization in a polarized cell. Cell shape is stabilized by cytoplasmic pressure and elastic behavior of the cortex. An undesired tangential friction, caused by the cortical flow, is balanced by a compensating force, which can be attributed to the internal transport of depolymerized actin.

The model has the form of an obstacle problem for a strongly nonlinear degenerate parabolic system. Global existence of solutions has been proven under natural assumptions on the data. The analysis relies on ideas from the theory of gradient flows, employing the structure of the dominating elastic and pressure terms. The results are complete for an approximate system with 'softened' obstacles. The limit for hard obstacles can be carried out, however with an incomplete characterization of the limiting problem.

For numerical simulations, a conservative, explicit-in-time discretization has been introduced. Under appropriate time step restrictions, simulations are stable and (at least qualitatively) reproduce the behavior observed in the experiments. In particular, migration needs both confinement and sufficiently structured channel walls. A parametric study shows the expected dependencies on geometric properties of the channel.

From a modeling point of view, this study has to be seen as a first step. Reliable experimental information on cell cortex structure and dynamics is still scarce. The fact that in our model migration strongly depends on the force compensating excess polymerization and depolymerization, is rather questionable. In ongoing work, the model is extended by a viscous resistance against cortex bending. This effect seems to be a reasonable alternative providing the necessary pushing force against the channel walls. However, its inclusion poses formidable new challenges both from an analytic and from a numerical point of view. The extended model will be parametrized with the goal of quantitative agreement with experimental data.

A Appendix: A modified Morrey inequality

Lemma 4. *Let $u \in L^\infty((0, T); H^1(0, 1)) \cap H^1((0, T); L^2(0, 1)) =: L_t^\infty H_s^1 \cap H_t^1 L_s^2$. Then*

$$|u(s_1, t_1) - u(s_0, t_0)| \leq 8 \left(\|\partial_s u\|_{L_t^\infty L_s^2} + \|\partial_t u\|_{L_{s,t}^2} \right) (|s_1 - s_0|^{1/2} + |t_1 - t_0|^{1/4}), \quad (s_0, t_0), (s_1, t_1) \in (0, 1) \times (0, T)$$

Proof. We introduce $\Delta s := |s_1 - s_0|$ and $\Delta t := |t_1 - t_0|$ and consider a rectangle $W \subset (0, 1) \times (0, T)$, containing the points $(s_0, t_0), (s_1, t_1)$ with sides parallel to the t -axis of lengths Δt and parallel to the s -axis of lengths $\Delta s + \sqrt{\Delta t}$, such that $|W| = \Delta t(\Delta s + \sqrt{\Delta t})$. We have

$$u(s_1, t_1) - u(s_0, t_0) = \frac{1}{|W|} \int_W (u(s_1, t_1) - u(\sigma, \tau)) d(\sigma, \tau) + \frac{1}{|W|} \int_W (u(\sigma, \tau) - u(s_0, t_0)) d(\sigma, \tau). \quad (23)$$

For estimating the second term, we introduce the curve $\{(s_0 + (\sigma - s_0)\sqrt{p}, t_0 + (\tau - t_0)p) : 0 \leq p \leq 1\}$, whence it can be estimated by

$$\frac{1}{|W|} \int_0^1 \frac{1}{2\sqrt{p}} \int_W |\sigma - s_0| |\partial_s u| d(\sigma, \tau) dp + \frac{1}{|W|} \int_0^1 \int_W |\tau - t_0| |\partial_t u| d(\sigma, \tau) dp,$$

where the derivatives of u are evaluated along the curve. Employing the Cauchy-Schwarz inequality, this can be estimated further by

$$\frac{\Delta s + \sqrt{\Delta t}}{\sqrt{|W|}} \int_0^1 \frac{1}{2\sqrt{p}} \sqrt{\int_W |\partial_s u|^2 d(\sigma, \tau)} dp + \frac{\Delta t}{\sqrt{|W|}} \int_0^1 \sqrt{\int_W |\partial_t u|^2 d(\sigma, \tau)} dp.$$

In the second integral over W we introduce the new coordinates $(s, t) = (s_0 + (\sigma - s_0)\sqrt{p}, t_0 + (\tau - t_0)p)$; in the first one we first estimate the integrand by its supremum with respect to t

and then make the coordinate transformation only in s :

$$\begin{aligned} & \sqrt{\Delta s + \sqrt{\Delta t}} \int_0^1 \frac{dp}{2p^{3/4}} \|\partial_s u\|_{L_t^\infty L_s^2} + \sqrt{\frac{\Delta t}{\Delta s + \sqrt{\Delta t}}} \int_0^1 \frac{dp}{p^{3/4}} \|\partial_t u\|_{L_{s,t}^2} \\ & \leq \left(2\|\partial_s u\|_{L_t^\infty L_s^2} + 4\|\partial_t u\|_{L_{s,t}^2} \right) (\Delta s^{1/2} + \Delta t^{1/4}). \end{aligned}$$

An analogous treatment of the first term on the right hand side of (23) completes the proof. \square

Acknowledgments This work has been supported by the Vienna Science and Technology Fund, Grant no. LS13-029. G.J. and C.S. also acknowledge support by the Austrian Science Fund, Grants no. W1245, F 65, and W1261, as well as by the Fondation Science Mathématiques de Paris, and by Paris Sciences et Lettre.

References

- [Abercrombie et al., 1970] Abercrombie, M., Heaysman, J. E., and Pegrum, S. M. (1970). The locomotion of fibroblasts in culture. 3. Movements of particles on the dorsal surface of the leading lamella. *Experimental Cell Research*, 62(2):389–398.
- [Bergert et al., 2015] Bergert, M., Erzberger, A., Desai, R. A., Aspalter, I. M., Oates, A. C., Charras, G., Salbreux, G., and Paluch, E. K. (2015). Force transmission during adhesion-independent migration. *Nature Cell Biology*, 17(4):524–529.
- [Blaser et al., 2006] Blaser, H., Reichman-Fried, M., Castanon, I., Dumstrei, K., Marlow, F. L., Kawakami, K., Solnica-Krezel, L., Heisenberg, C.-P., and Raz, E. (2006). Migration of Zebrafish Primordial Germ Cells: A Role for Myosin Contraction and Cytoplasmic Flow. *Developmental Cell*, 11(5):613–627.
- [Charras and Paluch, 2008] Charras, G. and Paluch, E. (2008). Blebs lead the way: how to migrate without lamellipodia. *Nature Reviews Molecular Cell Biology*, 9(9):730–736.
- [Friedl et al., 2001] Friedl, P., Borgmann, S., and Bröcker, E. B. (2001). Amoeboid leukocyte crawling through extracellular matrix: Lessons from the Dictyostelium paradigm of cell movement. *Journal of Leukocyte Biology*, 70(4):491–509.
- [Giaquinta, 1983] Giaquinta, M. (1983). *Multiple integrals in the calculus of variations and nonlinear elliptic systems*, volume 105 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ.
- [Hawkins et al., 2011] Hawkins, R. J., Poincloux, R., Bénichou, O., Piel, M., Chavrier, P., and Voituriez, R. (2011). Spontaneous Contractility-Mediated Cortical Flow Generates Cell Migration in Three-Dimensional Environments. *Biophysical Journal*, 101(5):1041–1045.
- [Hungerbühler, 2000] Hungerbühler, N. (2000). *Young measures and nonlinear PDEs*. Habilitationsschrift, ETH Zürich.
- [Lämmermann et al., 2008] Lämmermann, T., Bader, B., Monkley, S., Worbs, T., Wedlich-Söldner, R., Hirsch, K., and et al. (2008). Rapid leukocyte migration by integrin-independent flowing and squeezing. *Nature*, 453:51–55.

- [Malawista et al., 2000] Malawista, S. E., de Boisfleury Chevance, A., and Boxer, L. (2000). Random locomotion and chemotaxis of human blood polymorphonuclear leukocytes from a patient with Leukocyte Adhesion Deficiency-1: Normal displacement in close quarters via chimneying. *Cell Motility and the Cytoskeleton*, 46:183–189.
- [Minty, 1963] Minty, G. J. (1963). on a “monotonicity” method for the solution of non-linear equations in Banach spaces. *Proceedings of the National Academy of Sciences of the United States of America*, 50:1038–1041.
- [Paluch et al., 2016] Paluch, E. K., Aspalter, I. M., and Sixt, M. (2016). Focal Adhesion–Independent Cell Migration. *The Annual Review of Cell and Developmental Biology*, 32:469–490.
- [Phillipson et al., 2006] Phillipson, M., Heit, B., Colarusso, P., Liu, L., Ballantyne, C. M., and Kubes, P. (2006). Intraluminal crawling of neutrophils to emigration sites: a molecularly distinct process from adhesion in the recruitment cascade. *Journal of Experimental Medicine*, 203(12):2569–2575.
- [Rafelski and Theriot, 2004] Rafelski, S. M. and Theriot, J. A. (2004). Crawling toward a unified model of cell mobility: spatial and temporal regulation of actin dynamics. *Annual review of biochemistry*, 73:209–39.
- [Reversat et al., 2018] Reversat, A., Merrin, J., Vries, I., Hauschild, R., Stopp, J., Hons, M., Piel, M., Callan-Jones, A., Voituriez, R., and Sixt, M. (2018). Adhesion-free cell migration by topography-based force transmission. *submitted*.
- [Shalem et al., 2014] Shalem, O., Sanjana, N., Hartenian, E., Shi, X., Scott, D., Mikkelsen, T., and et al. (2014). Genome-scale crispr-cas9 knockout screening in human cells. *Science*, 343(6166):84–87.
- [Tinevez et al., 2009] Tinevez, J.-Y., Schulze, U., Salbreux, G., Roensch, J., Joanny, J.-F., and Paluch, E. (2009). Role of cortical tension in bleb growth. *Proceedings of the National Academy of Sciences*, 106(44):18581–18586.
- [Vargas et al., 2014] Vargas, P., Terriac, E., Lennon-Duménil, A.-M., and Piel, M. (2014). Study of cell migration in microfabricated channels. *J Vis Exp*, 84.
- [Vicente-Manzanares et al., 2009] Vicente-Manzanares, M., Choi, C. K., and Horwitz, A. R. (2009). Integrins in cell migration – the actin connection. *Journal of Cell Science*, 122(2):199–206.
- [Waugh and Evans, 1979] Waugh, R. and Evans, E. (1979). Thermoelasticity of red blood cell membrane. *Biophysical Journal*, pages 115–131.
- [Wolf and Friedl, 2006] Wolf, K. and Friedl, P. (2006). Molecular mechanisms of cancer cell invasion and plasticity. *British Journal of Dermatology*, 154:11–15.
- [Wolf et al., 2013] Wolf, K., Lindert, M. t., Krause, M., Alexander, S., Riet, J. t., Willis, A. L., Hoffman, R. M., Figdor, C. G., Weiss, S. J., and Friedl, P. (2013). Physical limits of cell migration: Control by ECM space and nuclear deformation and tuning by proteolysis and traction force. *J Cell Biol*, 201(7):1069–1084.
- [Yin Lim et al., 2013] Yin Lim, F., Ling Koon, Y., and Chiam, K.-H. (2013). A computational model of amoeboid cell migration. *Computer methods in biomechanics and biomedical engineering*, 16.