



Open cluster kinematics with Gaia DR2★ (Corrigendum)

C. Soubiran, T. Cantat-Gaudin, M. Romero-Gomez, L. Casamiquela, C. Jordi, A. Vallenari, T. Antoja, L. Balaguer-Núñez, D. Bossini, A. Bragaglia, et al.

► To cite this version:

C. Soubiran, T. Cantat-Gaudin, M. Romero-Gomez, L. Casamiquela, C. Jordi, et al.. Open cluster kinematics with Gaia DR2★ (Corrigendum). *Astronomy and Astrophysics - A&A*, 2019, 623, pp.C2. 10.1051/0004-6361/201834020e . hal-02079514

HAL Id: hal-02079514

<https://hal.science/hal-02079514>

Submitted on 21 Aug 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Open cluster kinematics with *Gaia* DR2[★] *(Corrigendum)*

C. Soubiran¹, T. Cantat-Gaudin², M. Romero-Gómez², L. Casamiquela¹, C. Jordi², A. Vallenari³, T. Antoja², L. Balaguer-Núñez², D. Bossini³, A. Bragaglia⁴, R. Carrera³, A. Castro-Ginard², F. Figueras², U. Heiter⁶, D. Katz⁷, A. Krone-Martins⁵, J.-F. Le Campion¹, A. Moitinho⁵, and R. Sordo³

¹ Laboratoire d’Astrophysique de Bordeaux, Univ. Bordeaux, CNRS, B18N, allée Geoffroy Saint-Hilaire, 33615 Pessac, France
 e-mail: caroline.soubiran@u-bordeaux.fr

² Institut de Ciències del Cosmos, Universitat de Barcelona (IEEC-UB), Martí i Franquès 1, 08028 Barcelona, Spain

³ INAF-Osservatorio Astronomico di Padova, vicolo Osservatorio 5, 35122 Padova, Italy

⁴ INAF-Osservatorio di Astrofisica e Scienza dello Spazio, via Gobetti 93/3, 40129 Bologna, Italy

⁵ CENTRA, Faculdade de Ciências, Universidade de Lisboa, Ed. C8, Campo Grande, 1749-016 Lisboa, Portugal

⁶ Department of Physics and Astronomy, Uppsala University, Box 516, 75120 Uppsala, Sweden

⁷ GEPI, Observatoire de Paris, Université PSL, CNRS, 5 Place Jules Janssen, 92190 Meudon, France

A&A, 619, A155 (2018), <https://doi.org/10.1051/0004-6361/201834020>

Key words. stars: kinematics and dynamics – Galaxy: kinematics and dynamics – open clusters and associations: general – errata, addenda

Due to an unfortunate error in the computation of distance between clusters, Tables 4 and 5, and Fig. 12 in Sect. 3.3 of the original paper contain incorrect values. Here we present the updated tables and figure.

Table 4 gives the list of cluster pairs that differ by less than 100 pc in distance and 5 km s⁻¹ in velocity in our high-quality sample. The closest pair includes ASCC 16 and ASCC 21, which are separated by ∼13 pc, with a velocity difference of 4.5 km s⁻¹. RSG 7 and RSG 8, as well as ASCC 16 and ASCC 19, are also close pairs separated by ∼24 pc, with velocity differences of 2.3 km s⁻¹ and 3.6 km s⁻¹ respectively, thus good candidates to be physically related. Several possibly larger complexes may be found in that table, formed by clusters that appear more than one time. Table 5 gives a list of candidates binaries from the literature. The most famous binary cluster formed by η and χ Persei (NGC 869 and NGC 884, Messow & Schorr 1913) appears to have a separation of 19.5 pc. An excellent candidate binary is the pair Collinder 394 and NGC 6716, which lie at a distance of ∼11 pc from each other. IC 2602 and Platais 8 are also close from each other (∼35 pc) and they are possibly physically related, owing to their common age in DAML and similar velocity (also reported in Table 4). The other pairs in Table 5 have much larger separation and they are more likely to result from a chance alignment than that they are a physical binary system.

Table 4. Pairs of OCs differing by less than 100 pc in their Galactic position and 5 km s⁻¹ in velocity in the high-quality sample.

Cluster 1	Cluster 2	Δ pos (pc)	Δ V (km s ⁻¹)
ASCC 105	Roslund 5	85.1	3.7
ASCC 127	RSG 7	61.8	3.3
ASCC 127	RSG 8	81.9	2.8
ASCC 16	ASCC 19	24.6	3.6
ASCC 16	ASCC 21	13.2	4.5
ASCC 19	NGC 2232	96.0	3.9
ASCC 58	Alessi 5	99.4	3.9
ASCC 58	BH 99	54.8	3.3
ASCC 97	IC 4725	84.0	3.8
Alessi 20	Stock 12	59.3	2.2
Alessi 5	BH 99	50.5	3.6
Collinder 135	Collinder 140	84.6	3.6
Collinder 135	NGC 2451B	69.2	2.4
Collinder 140	NGC 2451B	50.8	1.8
Collinder 69	Gulliver 6	83.4	3.5
Gulliver 20	IC 4665	96.3	2.7
IC 2391	Platais 9	43.9	1.6
IC 2602	Platais 8	35.9	4.3
NGC 2451B	NGC 2547	82.7	3.4
RSG 7	RSG 8	24.3	2.3
Stock 23	Trumpler 3	70.1	2.7

* The table with clusters velocities is only available at the CDS via anonymous ftp to <cdsarc.u-strasbg.fr> (130.79.128.5) or via <http://cdsarc.u-strasbg.fr/viz-bin/qcat?J/A+A/623/C2>




Fig. 12. Histogram of the distance between nearest neighbours in the high-quality sample (*upper panel*) and the same in velocity (*lower panel*).

Table 5. Separation in space and velocity of binary candidates from the literature, computed from our mean parameters for 861 OCs.

Cluster 1	Cluster 2	ref	Δpos (pc)	ΔV (km s ⁻¹)
Alessi 13	Mamajek 1	1	92.5	5.1
Alessi 21	NGC 2422	1	120.8	9.3
Platais 8	IC 2602	1	35.9	4.3
Turner 9	ASCC 110	1	275.6	9.0
Collinder 394	NGC 6716	1	11.4	13.8
IC 1396	NGC 7160	1	87.6	13.9
NGC 869	NGC 884	2	19.5	19.9
NGC 5617	Trumpler 22	3	79.8	10.4
IC 4756	NGC 6633	4	82.9	8.0

References. (1) [Conrad et al. \(2017\)](#), (2) [Messow & Schorr \(1913\)](#), (3) [De Silva et al. \(2015\)](#), (4) [Casamiquela et al. \(2016\)](#).

References

- Casamiquela, L., Carrera, R., Jordi, C., et al. 2016, [MNRAS](#), **458**, 3150
 Conrad, C., Scholz, R.-D., Kharchenko, N. V., et al. 2017, [A&A](#), **600**, A106
 De Silva, G. M., Carraro, G., D’Orazi, V., et al. 2015, [MNRAS](#), **453**, 106
 Messow, B., & Schorr, R. R. E. 1913, [Astronomische Abhandlungen der Hamburger Sternwarte](#), **2**, 1