



HAL
open science

Do maternal pushing techniques during labour affect obstetric or neonatal outcomes?

Chloé Barasinski, Didier Lémery, Françoise Vendittelli

► **To cite this version:**

Chloé Barasinski, Didier Lémery, Françoise Vendittelli. Do maternal pushing techniques during labour affect obstetric or neonatal outcomes?. *Gynécologie Obstétrique & Fertilité*, 2016, 44 (10), pp.578-583. 10.1016/j.gyobfe.2016.07.004 . hal-02081273

HAL Id: hal-02081273

<https://hal.science/hal-02081273>

Submitted on 27 Mar 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Do maternal pushing techniques during labour affect obstetric or neonatal outcomes?

Le type de poussée maternelle pendant le travail a-t-il un impact sur les issues obstétricales ou néonatales?

Short title: Maternal pushing during labour.

Chloé Barasinski, RM, MPH^{1,2}, Didier Lemery, professor, MD, PhD^{1,2,3}, Françoise Vendittelli, MD, PhD^{1,2,3}

¹The Clermont-Ferrand University Hospital, Site Estaing, 1 place Lucie et Raymond Aubrac, 63003 Clermont-Ferrand Cedex 1, France.

²Clermont Université, Université d'Auvergne, EA 4681, PEPRADE (Perinatology, pregnancy, Environment, medical care PRACTICES and DEVELOPMENT), 28 place Henri-Dunant BP 38, 63001 Clermont-Ferrand, France.

³AUDIPOG (Association des Utilisateurs de Dossiers informatisés en Pédiatrie, Obstétrique et Gynécologie), RTH Laennec Medical University, 7 rue Guillaume Paradin, 69372 Lyon Cedex 08, France.

CORRESPONDING AUTHOR: Chloé Barasinski, CHU de Clermont-Ferrand, Hôpital Estaing, Pôle de Gynécologie-Obstétrique, Recherche Clinique en Périnatalité, 1 place Lucie et Raymond Aubrac, 63003 Clermont-Ferrand Cedex 1, France. Tel: +33.(0)4.73.75.50.89, Fax: +33.(0)4.73.75.05.65. cbarasinski@chu-clermontferrand.fr

Abstract:

Objectives. — To assess, through a literature review, the maternal and neonatal morbidity associated with the type of pushing used during the second stage of labour.

Methods. — We searched the Cochrane Library and the Medline database for randomised controlled trials from 1980 to 2015, using the following key words: “ delivery”, “birth”, “birthing”, “bearing down, coached, uncoached, pushing”, “second and stage and labour” and “randomised controlled trials”, “meta-analysis”.

Results. — Seven randomised controlled trials were found. Interventions varied between the studies. In the intervention groups, open-glottis pushing was spontaneous or coached. The groups did not differ for perineal injuries, episiotomies or type of birth. Impact on pelvic floor structure varied between the studies. Only one study found a better 5-minute Apgar score and a better umbilical artery pH in the “open glottis” group.

Conclusion. — The low methodological quality of the studies and the differences between the protocols do not justify a recommendation of a particular pushing technique. Further studies appear necessary to study outcomes with each of these techniques.

Résumé:

Objectifs. — Évaluer, par une revue de la littérature, la morbidité maternelle et néonatale liée aux différents types de poussées lors du 2^{ème} stade du travail (glotte ouverte ou fermée).

Méthodes. — Nous avons recherché les essais randomisés publiés entre 1980 à 2015, en utilisant la banque de données du Medline, avec les mots clés suivants : “ delivery”, “birth”, “birthing”, “bearing down, coached, uncoached, pushing”, “second and stage and labour” et “randomised controlled trials”, “meta-analysis”.

Résultats. — Sept essais cliniques randomisés ont été retrouvés. Le type de poussée différait selon les études. La poussée en expiration pouvait être soit spontanée soit dirigée. Il n’y avait pas de différence entre les deux groupes concernant la survenue de déchirures, d’une

épisiotomie ou le mode d'accouchement. L'impact sur la statique pelvienne variait selon les auteurs. Seule une étude a retrouvé un meilleur Apgar à 5 minutes et un meilleur pH artériel dans le groupe « glotte ouverte ».

Conclusion. — La faible qualité méthodologique des études et leurs différences ne permettent pas de recommander un type particulier de poussée. D'autres études semblent nécessaires afin d'étudier les issues maternelles et néonatales avec chaque type de poussée.

Introduction

Practices concerning childbirth have evolved considerably in recent centuries. Until the 17th century, births traditionally took place at home with an attendant who had little or no training [1]. Later, childbirth slowly became more and more medicalised; practices for the management of parturients during labour and childbirth today are very different than they were. Since the end of the 20th century, however, the medicalisation of childbirth has come under increasing challenge. A school of thought promoting a return to so-called "natural childbirth", which is less medicalised, has grown. In some countries, even some professionals doubt, among other things, the dogma of the dorsal decubitus position and the appropriate type of pushing during the second stage of labour.

Closed-glottis pushing, also called "Valsalva", is widely used in Western countries, but the origin of this obstetric practice is obscure. Pushing while exhaling was nonetheless described in obstetrical treatises until the end of the 19th century [2,3]. Valsalva pushing probably developed with the dissemination of forceps births, but without any convincing scientific data to support it.

There are currently no French guidelines for the management of normal childbirth. These depend on the training and beliefs of each obstetrician, general practitioner and midwife as well as on the policies in each obstetrics department. Today, the customers, so to speak, of obstetrics care demand less medicalisation and more involvement in the decision making concerning their children's births. Professionals must therefore provide their patients with the best possible care according to up-to-date scientific data as well as inform them of the

different alternatives for care. In this context, an analysis of the risks and benefits of the type of pushing during the second stage of labour is useful.

The principal objective of this work was to assess, through a critical analysis of the literature, maternal morbidity according to type of pushing (open or closed glottis) during the second stage of labour. Similarly, the secondary objective was to assess the neonatal morbidity associated with these two types of pushing.

Methods

Definition of types of pushing during the second stage of labour

In closed-glottis pushing, the woman is asked to inhale completely and fill her lungs completely with air, to hold her breath, and to push downward very strongly during contractions, for as long as possible, normally three times for each contraction. In French practice, this type of pushing is often directed, or coached. Many French professionals consider it the reference technique for pushing. Anyone who has been in a French delivery room has heard this advice: "Inhale deeply, hold your breath, and push for as long as you can!" There is nonetheless another type of pushing used less often by professionals: called open-glottis pushing, it occurs while exhaling (and can be coached or spontaneous). In France, open-glottis pushing is not frequently used, although it is a spontaneous, physiological, and sometimes overwhelmingly urgent way of pushing among parturients, especially those without epidural analgesia (so called Ferguson's reflex).

Databases searched

We searched the Cochrane Library, which includes 6 databases (Pregnancy and Childbirth Group's Trials Register, Cochrane Database of Systematic Reviews, etc.), and the Medline database to identify all randomised trials on this topic. This research was completed manually

by studying the references of articles and book chapters published on this subject. The following keywords were used: "delivery", "birth", "birthing", "bearing down", "coached", "uncoached", "pushing", "second and stage and labour", and "randomised controlled trials" or "meta-analyses". We looked at English and French articles published between 1980 and the end of February, 2016.

Articles were retained for more detailed assessment when they described randomised clinical trials comparing the two types of pushing (open- or closed-glottis), including those comparing pushing coached by a professional to pushing without professional coaching during the second stage of labour.

The outcome measures considered were the following:

- for the mother: episiotomy, perineal lacerations, especially third and fourth degree, type of birth (spontaneous or not), and delayed damage to pelvic floor function.
- for the child: 5-minute Apgar score, cord artery pH, and need for resuscitation in the delivery room or admission to the neonatal intensive care unit (NICU).

Methodological quality assessment

The methodological quality of the studies was assessed independently by two of the authors, according to the recommendations of Chalmers et al. [4]. In particular, we looked for four factors creating a risk of bias: selection bias (inadequate random sequence generation and allocation concealment, subjects excluded after randomisation, or large numbers of women lost to follow-up), performance, attrition and detection bias. We also researched differences in co interventions, apart from pushing.

Results

Two meta-analyses [5,6] and seven single-centre, randomised trials reported in nine articles were identified [7-15]. Neither participants nor personnel can be blinded to the intervention we consider here.

Neither of the two meta-analyses compared exclusively directed open-glottis with directed closed-glottis breathing, although directed pushing is the usual practice in France. The meta-analysis of Prins et al., published in 2011, compared instructed closed-glottis pushing only with spontaneous pushing, although as we have underlined, in view of the lack of spontaneous practices in France, this was not the focus of our work. Moreover, the total number of cases was quite low for a meta-analysis (three studies, n = 425) [9,11,12,16], and it was limited to primiparous women without epidural analgesia [5]. The authors concluded that Valsalva type pushing should not be routinely recommended because it could have a deleterious effect on pelvic floor function. They also agreed that the studies on this topic are rare, heterogeneous, and often biased [5]. Lemos et al. published a second meta-analysis, in 2015, comparing spontaneous (in practice, undirected open-glottis) vs. directed (in practice, closed-glottis) pushing [6]. The authors included seven studies [9-17], all of which involved potential bias. Their principal analysis (five studies, n= 598) concerned the duration of the second stage, which was not an outcome we chose to consider. Except for the spontaneous vaginal deliveries (five studies, n=688), their sub-analyses concerned maternal and fetal comorbidities (perineal lacerations, etc.) and were based on only one to three studies. They found no differences for any of the criteria studied, except for the duration of expulsive efforts, which appeared shorter in the spontaneous pushing (mean difference -5.20 minutes [95%CI; -7.78 to -2.62], one study, n=100). They also concluded that randomised trials of good quality were

necessary for the reliable determination of the potentially beneficial and harmful effects of these different techniques.

To meet our objectives we review the published randomised controlled trials. The intervention studied varied between them: although the control groups always used coached Valsalva pushing, the open-glottis breathing in the intervention group was sometimes spontaneous (not coached) [9-12,14,15] and sometimes coached [7,8,13] (Table 1). Study inclusion criteria included a gestational age of at least 36 weeks' gestation or more, cephalic presentation, and a singleton pregnancy. All but two trials included only nulliparas [8,14]. Co-interventions varied between studies (Table 1). The mother's position during the second stage of labour differed between the studies; it was not specified in two [11,12,15], was left to the woman's choice in two others [8-10], and was mandated in two more: dorsal decubitus position in one [13] and semi-recumbent in the other [7]. In another study, position differed according to the study group (a vertical position for the intervention group and dorsal decubitus position for the control group) [14]. Cervical dilatation was specified in only two studies [9-12] (Table 1).

Globally, the quality of randomisation was not good in the randomised studies, especially in the study of Barnett et al., which does not report its randomisation method and indeed mentions randomisation only in the abstract [7]. Four studies reported using randomisation tables without providing further information [9-12,14,15] (Table 2). Two studies used sealed envelopes [8-10], another opaque envelopes [11,12], and the others did not specify the kind of envelope used [13,14] (Table 2). The timing of the randomisation was either not specified [7] or varied according to the study (Table 2).

Compliance with the allocated intervention was not described in three studies [11-14] and varied in the others [7-10,15] (Table 2). In one study, compliance was assessed by a self-administered questionnaire (from which we could not extract data because the trial tested perineal massage as a co-intervention and did not separate out those results) [15].

Most studies did not use an intention-to-treat analysis [8,11-15]. Only one study described the orientation of the fetal head at the start of pushing, and it did not consider it in its statistical analysis [9,10]. One study did not adjust its results for the children's weight, although it was higher in the closed-glottis group [8]. Post-randomisation exclusions were observed in some studies [8,11,12,14,15] (Table 2). Two studies had a relatively high number of women lost to follow-up (randomisation of women during pregnancy, and follow-up at six and twelve months post partum [15] or follow-up at three months post partum [11,12]).

The heterogeneity of the interventions studied, the other care and co-interventions provided, and the timing of the onset of pushing, as well as the relatively poor methodological quality of the studies (Table 2), made it impossible for us to perform a meta-analysis. Moreover, the outcomes measured varied according to the study.

Nonetheless, some results can be underlined. No woman in the open-glottis pushing group began pushing when uterine contractions began [9,10]. The duration of the second stage of labour varied according to the study. For Thomson [9,10] and for Bloom et al., it was shorter among the women in the coached Valsalva pushing group than among those in the open-glottis pushing group (58 minutes \pm 42 versus 121.4 minutes \pm 58.4; $P=0.002$) (second stage: 46.3 minutes \pm 41.5 versus 59.1 minutes \pm 49.1; $P=0.014$)[11,12]. On the other hand, for both Barnett et al. and Parnell et al, this did not differ between the two groups [7,8]. For Yildirim

et al., the duration of the second stage of labour was shorter for the open-glottis pushing group than for the coached Valsalva group (40.8 minutes \pm 19.1 versus 50.1 minutes \pm 23.3 minutes; $P=0.045$) [13]; this was also true for Jahdi et al., with both primiparous and multiparous patients (respectively 47.38 minutes \pm 36.75 versus 57.12 minutes \pm 33.1, $P < 0.0001$ and 26.12 minutes \pm 23.43 versus 33.20 minutes \pm 22.76, $P < 0.0001$) [14]. Only Yildirim et al. found that the duration of the expulsive (pushing) phase was shorter in the open-glottis pushing group than in the coached Valsalva group (9.6 minutes \pm 5.5 versus 14.8 minutes \pm 7.5; $P=0.001$)[13], while Parnell et al. found no difference between the two groups [8].

There was no statistically significant difference for the mode of birth (Table 3) for Barnett et al., Thomson, Schaffer et al., Low et al. or Jahdi et al.[7,9-11,14,15]. Similarly, Schaffer et al. and Yildirim et al. found no difference for operative vaginal births [11-13] (Table 3). The maternal outcomes did not differ statistically for perineal tears or lacerations, for Thomson, Schaffer et al., or Yildirim et al. [9-13]; nor did Schaffer et al. or Yildirim et al. find any difference for episiotomies [11-13] (Table 3). Few authors have looked at immediate postpartum haemorrhages [9,11,13]. The methods for estimating blood loss have varied substantially between studies.

For neonatal outcomes, Yildirim et al. found that the pH of the umbilical artery was significantly higher in the open-glottis compared with the Valsalva group [13] (Table 3), although Barnett et al., Parnell et al. and Schaffer et al. found no statistically significant differences [7,8,11,12] (Table 3). The Apgar score was also significantly higher in the open-glottis than in the Valsalva group, at both 1 minute and 5 minutes for Yildirim et al. (1 minute: 7.9 \pm 0.6 versus 7.3 \pm 0.8; $P =0,001$)[13] (Table 3), although Parnell et al. found no significant difference between the two groups for either of these Apgar scores [8] (Table 3).

Finally, Thomson and Schaffer et al. found no difference between the two groups for resuscitation in the delivery room or for NICU admission [9-12].

Two studies have looked at the consequences on pelvic floor function [11,15]. For Schaffer et al., urodynamic testing at 3 months post partum showed no reduction in bladder capacity (427 mL versus 482 mL, $P=0.51$), but did observe a reduction in the first urge to void (160 mL versus 202 mL, $P=0.025$) in the closed-glottis pushing group [9]. Detrusor overactivity increased in the closed-glottis group, but not significantly (16% versus 8%; $P=0.17$); the same was true for urodynamic stress incontinence ($P=0.42$) [11]. Low et al. found no difference between the two groups when they used the “leakage index” one year after birth [15].

Discussion

It is difficult to draw convincing scientific conclusions from the meta-analyses and randomised studies about the optimal type of pushing for reasons beyond their poor methodological quality and their frequent lack of power. The studies often do not consider either prognostic or confounding factors (direction of the fetal head, birth weight, maternal position during the second stage of labour, station when pushing begins, policy of early coaching and active management of labour, etc.). Moreover, most of these studies excluded patients with epidural analgesia, which is widely used in some western countries. In France, almost 80% of parturients receive epidural analgesia [18].

Questions have been raised about Valsalva-type pushing for many years. The exertion of pushing may increase intrathoracic and intra-abdominal pressure and thus lead to maternal haemodynamic modifications, including a drop in venous return followed by reduced oxygenation of the maternal blood and therefore also of the uterus, placenta, and fetus

[19,20]. Fetal cerebral oxygenation then drops, while cerebral blood volume increases [21]. These modifications of the oxygenation of maternal blood may not be observed during spontaneous pushing since women use their residual respiratory volume without exaggerated forced inspiration and with shorter pushing efforts in general (less than 6 seconds) [22]. The lithotomy position, principally used for patients with epidural analgesia in most western countries, may also reduce fetal oxygenation through maternal aortocaval compression by the gravid uterus [23].

Moreover, it is not clear that professionals are well trained in the different types of pushing. Open-glottis pushing is accompanied by noise made by the woman (grunts and groans). These noises are often interpreted by professionals as the expression of severe pain or loss of control by the woman. Generally, professionals respond by giving instructions and coaching the woman, instead of reassuring and encouraging her [24]. It is also unlikely that women in labour have been well trained in any of these different types of pushing. Labour is probably not the best moment to learn different types of pushing methods. However, this was the case in all randomised publications, except one [15].

One randomised study explored the influence of prenatal education on the type of pushing and on maternal outcomes, but found no difference between the two groups [25]. A meta-analysis about prenatal childbirth preparation, either individual or in groups, did not find any consistent results [26]. We also note that compliance with the specific type of pushing assigned is never 100%. Moreover, open-glottis pushing can be envisioned either by coaching women to exhale during contractions, or by encouraging them to push spontaneously [27,28]. Women's spontaneous physiological instinct might well be to push instinctively, to listen to

their bodies. In other words, at any given moment, they may be pushing with glottis intermittently open and closed.

After flexion, descent and intrapelvic rotation, the leading part of the fetus reaches the pelvic floor. The spontaneous reflex to push (Ferguson's reflex) can occur earlier, once the fetal head reaches station 0 to + 1 in the plane through the ischial spines [29]. Moreover, we know that a policy of late pushing compared to early pushing once full dilatation is reached does not appear to be harmful [30]. In a meta-analysis of 12 trials (1,584 women with immediate and 1,531 with delayed pushing), there were more spontaneous births (pooled RR = 1.09; 95% CI: 0.98-1.26) and a shorter duration of expulsive efforts in the delayed pushing group [30]. The meta-analysis of Lemos et al. reported (12 trials, n= 3,114) that there were more spontaneous births (pooled RR = 1.07; 95% CI: 1.03-1.11) and a 20-minute decrease in the duration of pushing in the delayed pushing group of women with epidural analgesia [6]. It is important to bear in mind that epidural analgesia may attenuate the physiological augmentation of oxytocin that takes place during the second stage of labour and therefore the reflex of spontaneous pushing [31]. Moreover, an older study reported that women who push spontaneously do not start until intrauterine pressure reaches at least 30 mm Hg [19]. Pushing therefore does not occur simultaneously with the beginning of a contraction.

A consequence of Valsalva-type pushing is increased pressure on perineal tissue and the prevention of venous return to the inferior vena cava. Although the study by Schaffer et al. lacks power [11], it is possible that this type of pushing is more harmful to pelvic floor function some time after childbirth. Low et al. found no difference between the two types of pushing, but the utility of the "leakage index" is questionable, in view of the rarity of its use in the literature [15]. Theoretically vulvar oedema may also increase in the closed-glottis

pushing group, which could increase perineal injuries, particularly if the mother's position magnifies this phenomenon, as the seated position does. The randomised trials about the type of pushing nonetheless do not support this risk [8,11-13]. The same is true for the question of fetal oxygenation. Nonetheless, one randomised study found that the Apgar score and arterial pH at birth were best in the open-glottis pushing group [13]. Buhimschi et al. report that Valsalva-type pushing during a contraction during the second stage of labour is associated with a 62% increase in intrauterine pressure, compared with baseline tone. This increased pressure is correlated with body mass index but inversely correlated with myometrial thickness, ultrasound-estimated fetal weight, and the need for oxytocin during labour [32].

Conclusion

Because of the low methodological quality of the studies, the lack of study power, the variability of their selection and outcome criteria, and of the co-interventions, these studies do not allow us to recommend a particular type of pushing. Moreover, most of the studies and meta-analyses compared spontaneous, non-coached open-glottis pushing to coached closed-glottis pushing. There is a lack of studies that compare coached open-glottis exhalation pushing to coached Valsalva-type pushing.

This assessment thus makes clear that a better understanding of the different modes of pushing might result in improving maternal satisfaction and even perhaps maternal and foetal morbidity during childbirth.

While awaiting such a study, the choice about the type of pushing during the active stage of birth should be given to the woman. Management must evolve by giving them more autonomy at the same time as we provide them the quality of care expected in light of current scientific data [27,33].

Acknowledgments and funding

This study was supported by a grant from the French Ministry of Health (PHRC 2015 N°05.09). The translation was supported by the research team PEPRADE EA 4681, University of Auvergne.

Author's contribution

All the authors have reviewed the contents of this article before its submission.

Disclosures

The authors have no conflicts of interest to declare.

References

- [1] Morel MF. Naître en France du XVII au XX^{ème} siècles. *Contact Santé* 2010 ; 230 : 47-49.
- [2] Playfair W.S : *Traité theorique et pratique de l'art des accouchements* (2nd edition; traduit par Dr Vermeil). Paris : Octave Doin ; 1879. p. 327-39.
- [3] Ribemont-Dessaignes A, Lepage G : *Précis d'Obstétrique*. 2nd edition Paris : G Masson ; 1896. p.317-18.
- [4] Chalmers TC, Smith H, Blackburn B, Silverman B, Schroeder B et al. A method for assessing the quality of a randomized control trial. *Control Clin Trials* 1981 ; 2 : 31-49.
- [5] Prins M, Boxem J, Lucas C, Hutton E. Effect of spontaneous pushing versus Valsalva pushing in the second stage of labour on mother and fetus: a systematic review of randomised trials. *BJOG* 2011 ; 118 : 662-70.
- [6] Lemos A, Amorim MMR, Dornelas de Andrade A, de Souza AI, Cabral Filho JE et al. Pushing/bearing down methods for the second stage of labour. *Cochrane Database Syst Rev*. 2015;10:CD009124.
- [7] Barnett MM, Humenick SS. Infant outcome in relation to second stage labor pushing method. *Birth* 1982 ; 9 : 221-229.
- [8] Parnell C, Langhoff-Roos J, Iversen R, Damgaard P. Pushing method in the expulsive phase of labor. A randomized trial. *Acta Obstet Gynecol Scand* 1993; 72 : 31-5.
- [9] Thomson AM. Pushing techniques in the second stage of labour. *J Adv Nurs* 1993 ; 18 : 171-7.
- [10] Thomson AM. Maternal behavior during spontaneous and directed pushing in the second stage of labour. *J Adv Nurs* 1995 ; 22 : 1027-34.
- [11] Schaffer JI, Bloom SL, Casey BM, McIntire DD, Nihira MA et al. A randomized trial of the effects of coached vs. uncoached maternal pushing during the second stage of labor on postpartum pelvic floor structure and function. *Am J Obstet Gynecol* 2005 ; 192 : 1692-6.

- [12] Bloom SL, Casey BM, Schaffer JI, McIntire DD, Leveno KJ. A randomized trial of coached versus uncoached maternal pushing during the second stage of labor. *Am J Obstet Gynecol* 2006 ; 194 : 10-3.
- [13] Yildirim G, Kizilkaya N. Effects of pushing techniques in birth on mother and fetus : a randomized study. *Birth* 2008 ; 35 : 25-30.
- [14] Jahdi F, Shahnazari M, Kashanian M, Farahani MA, Haghani H. A randomized controlled trial comparing the physiological and directed pushing on the duration of the second stage of labor, the mode of delivery and Apgar score. *Int J Nurs Midwifery* 2011 ; 3 : 55-9.
- [15] Low LK, Miller JM, Guo Y, Ashton-Miller JA, DeLancey JOL et al. Spontaneous pushing to prevent postpartum urinary incontinence: a randomized, controlled trial. *Int Urogynecol J*. 2013 ; 24 : 453-60.
- [16] Lam COC. Comparison of Two Pushing Techniques Used in the Second Stage of Labour for Their Effect on Maternal Perception of Fatigue in the Early Postpartum Period: a Randomised Controlled Trial [thesis]. Victoria: La Trobe University, 2006.
- [17] Knauth DG, Haloburdo EP. Effect of pushing techniques in birthing chair on length of second stage of labor. *Nursing Research* 1986;35:49–51.
- [18] Direction générale de la Santé et INSERM : Blondel B., Kermarrec M. Les naissances en 2010 et leur évolution depuis 2003. Available at : http://www.sante.gouv.fr/IMG/pdf/Les_naissances_en_2010_et_leur_evolution_depuis_2003.pdf. Accessed June 29, 2015.
- [19] Lemos A, Dean E, de Andrade AD. The Valsalva maneuver duration during labor expulsive stage: repercussions on the maternal and neonatal birth condition. *Braz J Phys Ther* 2011 ; 15 : 66-72.

- [20] Caldeyro-Barcia R, Giussi G, Storch E, Poseiro JJ, Lafaurie N, et al. The bearing-down efforts and their effects on fetal heart rate, oxygenation and acid base balance. *J Perinat Med* 1981 ; 9(s1) : 63-7.
- [21] Aldrich CJ, D'Antona D, Spencer JAD, Wyatt JS, Peebles DM et al. The effect of maternal pushing on fetal cerebral oxygenation and blood volume during the second stage of labour. *BJOG* 1995 ; 102:448-53.
- [22] Yeates DA, Roberts JE. A comparison of two bearing-down techniques during the second stage of labor. *J Nurse Midwifery* 1984 ; 29 : 3-11.
- [23] Carbonne B, Benachi A, Lévèque ML, Cabrol D, Papiernik E. Maternal Position During Labor : Effects on Fetal Oxygen Saturation Measured by Pulse Oximetry. *Obstet Gynecol* 1996 ; 88:797-800.
- [24] McKay S, Barrows T, Roberts J. Women's views of second stage as assessed by interviews and videotapes. *Birth* 1990 ; 17:192-8.
- [25] Phipps H, Charlton S, Dietz HP. Can antenatal education influence how women push in labour. *Aust N Z J Obstet Gynaecol* 2009 ; 49 : 274-8.
- [26] Gagnon AJ, Sandall J. Individual or group antenatal education for childbirth or parenthood, or both. *Cochrane Database Syst Rev* 2007 ; 3 : CD002869.
- [27] Roberts J, Hanson L. Best practices in second stage labor care: maternal bearing down and positioning. *J Midwifery Womens Health* 2007; 52 : 238-45.
- [28] Sampelle CM, Miller JM, Luecha Y, Fischer K, Rosten L. Provider support of spontaneous pushing during second stage of labor. *J Obstet Gynecol Neonatal Nurs* 2005 ; 34 : 695-702.
- [29] Roberts J, Goldstein S, Gruener J, Maggio M, Mendez-Bauer C. A descriptive analysis of involuntary bearing-down efforts during the expulsive phase of labor. *J Obstet Gynecol Neonatal Nurs* 1987 ; 16 : 48-55.

[30] Tuuli MG, Frey HA, Odibo AO, Macones GA, Cahill AG. Immediate compared with delayed pushing in the second stage of labor: a systematic review and meta-analysis. *Obstet Gynecol* 2012 ; 120 : 660-8.

[31] Goodfellow CF, Hull MGR, Swaab DF, Dogterom J, Buijs RM. Oxytocin deficiency at delivery with epidural analgesia. *Br J Obstet Gynaecol* 1983 ;90 : 214-9.

[32] Buhimschi CS, Buhimschi IA, Malinow AM, Kopelman JN, Weiner CP. Pushing in labor: performance and not endurance. *Am J Obstet Gynecol* 2002 ; 186 : 1339-44.

[33] Greene MF. Two Hundred Years of Progress in the Practice of Midwifery. *N Engl J Med* 2012 ; 367 : 1732-40.

Table 1. Methodological characteristics of the studies

Study	Year of study	Description of the study population	Parity	Eligible	Interventions studied	
					IG	CG
Barnett et al.[7]	?	USA, uncomplicated pregnancies ♀ between 16-30 years and between 38-42 weeks, EFW 2500-4000 g, no FHR abnormalities, no EA	Nulliparas	?	Coached open-glottis pushing	Coached Valsalva pushing
Parnell et al.[8]	1990-91	Denmark, ♀, > 18 years, ≥37 weeks, singleton in CP, planned vaginal birth, 1 previous caesarean possible	Nulliparas or primiparas	1413	Coached open-glottis pushing	Coached Valsalva pushing
Thomson [9,11]	?	England, spontaneous pregnancies, ♀ aged 18-40 years, between 37-42 weeks, singleton in CP, uncomplicated, no EA, dilatation ≥ 6 cm	Primiparas	?	Spontaneous open-glottis pushing	Coached Valsalva pushing
Schaffer et al.; Bloom et al.[11,12]	2000-02	USA, uncomplicated pregnancy, spontaneous labour (RG, dilatation ≥ 4 cm) between 36-41 weeks, singleton in CP, EFW < 4000 g, no pelvic prolapse or urogenital incontinence	Nulliparas	1534	Spontaneous open-glottis pushing	Coached Valsalva pushing

Yildirim et al.[13]	2003-04	Turkey, ♀ at low risk, spontaneous labour, planned VB, between 38-42 weeks, singleton, no disease in CP and EFW 2500-3999 g, without EA	Nulliparas	174	Coached open-glottis pushing	Coached Valsalva pushing
Jahdi et al.[14]	2009	Iran, ♀ between 18- 40 years, pregnancy at low risk and planned VB, spontaneous labour or induction for SRM or post-term and between 37-42 weeks, singleton, alive, in CP with EFW 2500-4000 g	Parity ranging from 0 to 5	258	Spontaneous open-glottis pushing	Coached Valsalva pushing
Low et al.[15]	2000-06	USA, ♀ >18 years, no urogenital disease (negative stress test) and continent before 20 weeks	Nulliparas	345	Spontaneous open-glottis pushing	Coached Valsalva pushing

Abbreviations: weeks (weeks of gestation); EFW (estimated fetal weight); FHR (foetal hart rate); EA (epidural analgesia); CP (cephalic presentation); RG (regular contractions); VB (vaginal birth); SRM (spontaneous rupture of the membranes); IG (interventional group); CG (control group); USA (United States of America).

Table 2. Description of the studies' other methodological criteria

Studies	Mode of randomisation	Time of randomisation	Women randomised	Exclusion post-randomisation	Score of studies	Compliance with the intervention	
						IG %	CG %
Barnett et al.[7]	?	?	10?	?	C	100	100
Parnell et al.[8]	Drawing by 1/3 women Sealed envelopes	Admission for childbirth	350	44 exclusions: 24 IG and 20 CG	C	34.4	84.5
Thomson[9,10]	Sealed envelopes Randomisation table	At 6 cm dilatation?	32	0	B	86.7	88.2
Schaffer et al. Bloom et al. [11,12]	Randomisation table Opaque envelopes	Beginning second stage	325	5 without data?	B	?	?
Yildirim et al. [13]	Envelopes	During first stage	100?	?	C	?	?
Jahdi et al. [14]	Drawing 1 envelope of 10. Randomisation blocks?	At full dilatation	258	67	C	?	?
Low et al. [15]	Randomisation table / computer	First appointment at 20 weeks	249 ^a	55 ^b ?	C	?	?

Abbreviations: weeks (weeks of gestation); IG (interventional group); CG (control group).

^aThis is the total number for the four study groups (control group, spontaneous open-glottis pushing group, perineal massage group, and perineal massage + spontaneous open-glottis pushing group).

^bTotal exclusions for the four study groups described in the note above.

Table 3. Maternal and neonatal outcomes

Studies	Episiotomies		Spontaneous birth		5-minute Apgar score		Umbilical artery pH	
	IG n/N (%)	CG n/N (%)	IG n/N (%)	CG n/N (%)	IG Mean [\pm SD]	CG Mean [\pm SD]	IG Mean [\pm SD]	CG Mean [\pm SD]
Barnett et al.[7]	?	?	5/5 (100%)	5/5 (100%)	?	?	7.32 ^c	7 ^c
Parnell et al.[8]	54/151 (36%)	46/155 (30%)	?	?	9.9 [\pm 0.4]	9.9 [\pm 0.3]	7.22 [\pm 0.08]	7.24 [\pm 0.08]
Thomson [9,10]	?	?	12/15 (80%)	15/17 (88.2%)	?	?	?	?
Schaffer et al. Bloom et al. [11,12]	32/157 (20%)	42/163 (26%)	149/157 (95%)	152/163 (93%)	?	?	7.2 [\pm 0.7]	7.2 [\pm 0.8]
Yildirim et al.[13]	45/50 (90%)	43/50 (86%)	-	-	9.9 [\pm 0.3]	9.5 [\pm 0.6] ^b	7.3 [\pm 0.1]	7.3 [\pm 0.1] ^d

Jahdi et al. [14]	?	?	97/99 (98%)	90/91 (98.9%)	?	?	?	?
Low et al. [15]	?	?	24/34 (68.6%) ^a	31/39 (86.1%) ^a	?	?	?	?

Abbreviations: IG (interventional group); CG (control group).

^aNote that for Low et al. there is a discordance in the tables between total number by categories and the addition of different numbers of individuals in the table.

^b $P=0.001$ for the authors (9.9 vs. 9.5).

^cStandard deviation found in the original article but value inconsistent with the umbilical artery pH values normally found at birth.

^d $P=0.004$ for the authors (7.3 vs. 7.3).