



A uniqueness theorem for Sturm–Liouville equations with a spectral parameter linearly contained in the boundary conditions

Namig Guliyev

► To cite this version:

Namig Guliyev. A uniqueness theorem for Sturm–Liouville equations with a spectral parameter linearly contained in the boundary conditions. Proceedings of the Institute of Mathematics and Mechanics, 2006. hal-02084845

HAL Id: hal-02084845

<https://hal.science/hal-02084845>

Submitted on 8 Apr 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Namig J. GULIYEV

A UNIQUENESS THEOREM FOR STURM-LIOUVILLE EQUATIONS WITH A SPECTRAL PARAMETER LINEARLY CONTAINED IN THE BOUNDARY CONDITIONS

Abstract

A uniqueness theorem for the Sturm–Liouville equation with a spectral parameter linearly contained in both boundary conditions is proved.

Consider the Sturm–Liouville equation

$$\ell y := -y''(x) + q(x)y(x) = \lambda y(x), \quad x \in [0, \pi] \quad (1)$$

with the boundary conditions

$$\lambda(y'(0) - hy(0)) = h_1y'(0) - h_2y(0), \quad (2)$$

$$\lambda(y'(\pi) + Hy(\pi)) = H_1y'(\pi) + H_2y(\pi), \quad (3)$$

where $q(x) \in \mathcal{L}_2(0, \pi)$ is a real-valued function, $h, h_1, h_2, H, H_1, H_2 \in \mathbb{R}$ and

$$\delta := hh_1 - h_2 > 0, \quad \rho := HH_1 - H_2 > 0.$$

We denote this problem by $\mathcal{P}(q, h, h_1, h_2, H, H_1, H_2)$.

Let $\varphi(x, \lambda)$ and $\psi(x, \lambda)$ be the solutions of (1) satisfying the initial conditions

$$\begin{aligned} \varphi(0, \lambda) &= -\lambda + h_1, & \varphi'(0, \lambda) &= -\lambda h + h_2, \\ \psi(\pi, \lambda) &= -\lambda + H_1, & \psi'(\pi, \lambda) &= \lambda H - H_2. \end{aligned} \quad (4)$$

We define

$$\chi(\lambda) := \varphi(x, \lambda)\psi'(x, \lambda) - \varphi'(x, \lambda)\psi(x, \lambda),$$

which is independent of $x \in [0, \pi]$. The function $\chi(\lambda)$ is entire and has zeros at the eigenvalues of the problem (1)–(3). The set of eigenvalues is countable, consists of real numbers $([1, 4])$ and for each eigenvalue λ_n there exists such a number k_n that

$$\psi(x, \lambda_n) = k_n \varphi(x, \lambda_n), \quad k_n \neq 0. \quad (5)$$

In the Hilbert space $\mathcal{H} = \mathcal{L}_2(0, \pi) \oplus \mathbb{C}^2$ let an inner product be defined by

$$(F, G) := \int_0^\pi F_1(x)\overline{G_1(x)}dx + \frac{1}{\delta}F_2\overline{G_2} + \frac{1}{\rho}F_3\overline{G_3}$$

for

$$F = \begin{pmatrix} F_1(x) \\ F_2 \\ F_3 \end{pmatrix}, \quad G = \begin{pmatrix} G_1(x) \\ G_2 \\ G_3 \end{pmatrix} \in \mathcal{H}.$$

We define operator

$$A(F) := \begin{pmatrix} -F_1''(x) + q(x)F_1(x) \\ h_1 F_1'(0) - h_2 F_1(0) \\ H_1 F_1'(\pi) + H_2 F_1(\pi) \end{pmatrix}$$

with

$$D(A) = \{F \in \mathcal{H} | F_1(x), F_1'(x) \in \mathcal{AC}[0, \pi], \ell F_1 \in \mathcal{L}_2(0, \pi), \\ F_2 = F_1'(0) - hF_1(0), F_3 = F_1'(\pi) + HF_1(\pi)\}.$$

Then

$$\Phi_n := \begin{pmatrix} \varphi(x, \lambda_n) \\ \varphi'(0, \lambda_n) - h\varphi(0, \lambda_n) \\ \varphi'(\pi, \lambda_n) + H\varphi(\pi, \lambda_n) \end{pmatrix} = \begin{pmatrix} \varphi(x, \lambda_n) \\ -\delta \\ \rho/k_n \end{pmatrix}$$

are orthogonal eigenlements of A :

$$(\Phi_m, \Phi_n) = \int_0^\pi \varphi(x, \lambda_m) \varphi(x, \lambda_n) dx + \delta + \frac{\rho}{k_m k_n} = 0, \quad m \neq n.$$

We also define *norming constants* by

$$\gamma_n := \|\Phi_n\|^2 = \int_0^\pi \varphi^2(x, \lambda_n) dx + \delta + \frac{\rho}{k_n^2}.$$

The numbers $\{\lambda_n, \gamma_n\}_{n \geq 0}$ are called the *spectral data* of the problem (1)–(3).

Lemma 1. *The following equality holds:*

$$\dot{\chi}(\lambda_n) = k_n \gamma_n, \quad n \geq 0,$$

where $\dot{\chi}(\lambda) = \frac{d}{d\lambda} \chi(\lambda)$.

Analogously to [3, Theorem 1(i)] the following theorem can be proved.

Theorem 1. *For $F \in \mathcal{H}$*

$$\|F\|^2 = \sum_{n=0}^{\infty} \frac{1}{\gamma_n} |(F, \Phi_n)|^2.$$

By taking $F = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}$ and $F = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}$ in this theorem we obtain

$$\frac{1}{\delta} = \sum_{n=0}^{\infty} \frac{1}{\gamma_n}, \quad \frac{1}{\rho} = \sum_{n=0}^{\infty} \frac{1}{k_n^2 \gamma_n}. \quad (6)$$

Theorem 2. *Following asymptotics hold:*

$$\sqrt{\lambda_n} = n - 2 + \frac{\omega}{n\pi} + \frac{\zeta_n}{n}, \quad \{\zeta_n\} \in l_2, \quad (7)$$

$$\gamma_n = (n - 2)^4 \left(\frac{\pi}{2} + \frac{\zeta'_n}{n} \right), \quad \{\zeta'_n\} \in l_2, \quad (8)$$

where

$$\omega = h + H + \frac{1}{2} \int_0^\pi q(x) dx.$$

Proof. We denote $s := \sqrt{\lambda}$. Then from

$$\begin{aligned} \varphi(x, \lambda) = & -s^2 \cos sx - \left(h + \frac{1}{2} \int_0^x q(t) dt \right) s \sin sx - \\ & - \frac{1}{2} \int_0^x q(t) s \sin s(x - 2t) dt + O\left(e^{|Im sx|}\right), \end{aligned}$$

$$\begin{aligned} \varphi'(x, \lambda) = & s^3 \sin sx - \left(h + \frac{1}{2} \int_0^x q(t) dt \right) s^2 \cos sx - \\ & - \frac{1}{2} \int_0^x q(t) s^2 \cos s(x - 2t) dt + O\left(|s| e^{|Im s x|}\right) \end{aligned}$$

using (4) we obtain

$$\chi(\lambda) = s^5 \sin s\pi - \left(h + H + \frac{1}{2} \int_0^\pi q(x) dx \right) s^4 \cos s\pi + I(s) s^4, \quad (9)$$

where

$$I(s) = -\frac{1}{2} \int_0^\pi q(t) \cos s(\pi - 2t) dt + O\left(\frac{e^{|Im s \pi|}}{|s|}\right).$$

Now using Bessel's inequality it's easy to obtain (7) and (8).

Lemma 2. *Let $v(t, \lambda)$ be any function entire on λ with*

$$v(t, \lambda) = O\left(e^{|Im \sqrt{\lambda}|t}\right), \quad t \in [0, \pi].$$

Then

$$\sum_{n=0}^{\infty} \frac{\varphi(x, \lambda_n) v(t, \lambda_n)}{\gamma_n} = 0, \quad (10)$$

uniformly on $0 \leq t \leq x \leq \pi$.

Proof. Using (5), Lemma 1. and the residue theorem we get

$$\begin{aligned} \sum_{n=0}^N \frac{\varphi(x, \lambda_n) v(t, \lambda_n)}{\gamma_n} &= \sum_{n=0}^N \frac{\psi(x, \lambda_n) v(t, \lambda_n)}{k_n \gamma_n} = \sum_{n=0}^N \frac{\psi(x, \lambda_n) v(t, \lambda_n)}{\dot{\chi}(\lambda_n)} = \\ &= \sum_{n=0}^N \operatorname{Res}_{\lambda=\lambda_n} \frac{\psi(x, \lambda) v(t, \lambda)}{\chi(\lambda)} = \frac{1}{2\pi i} \int_{C_N} \frac{\psi(x, \lambda) v(t, \lambda)}{\chi(\lambda)} d\lambda, \end{aligned}$$

where $C_N = \{\lambda : |\lambda| = (N - 3/2)^2\}$. Equality (9) implies that

$$\chi(\lambda) = \lambda^2 \sqrt{\lambda} \sin \sqrt{\lambda} \pi + O\left(|\lambda|^2 e^{Im\sqrt{\lambda}|\pi}\right).$$

We denote $G_\delta = \{\lambda : |\sqrt{\lambda} - n| \geq \delta, n = 0, \pm 1, \pm 2, \dots\}$ for some small fixed $\delta > 0$ and recall that (see e.g. [6, p. 15])

$$|\sin \sqrt{\lambda} \pi| \geq 2C_\delta e^{Im\sqrt{\lambda}|\pi}, \quad \lambda \in G_\delta,$$

where C_δ does not depend on λ . Therefore we obtain

$$|\chi(\lambda)| \geq C_\delta |\lambda|^2 \sqrt{\lambda} e^{Im\sqrt{\lambda}|\pi}, \quad \lambda \in G_\delta, |\lambda| \geq \lambda_\delta$$

for sufficiently large λ_δ . Since

$$|\psi(x, \lambda)| = O\left(|\lambda| e^{Im\sqrt{\lambda}(\pi-x)}\right), \quad x \in [0, \pi],$$

we obtain

$$|\psi(x, \lambda)v(t, \lambda)| = O\left(|\lambda| e^{Im\sqrt{\lambda}(\pi-x+t)}\right) = O\left(|\lambda| e^{Im\sqrt{\lambda}|\pi}\right), \quad 0 \leq t \leq x \leq \pi.$$

Thus the equality (10) holds. Uniformity of this convergence can be obtained from (7) and (8) by applying Weierstraß' theorem.

Putting $v(t, \lambda) \equiv 1$ and $x = 0$ in (10) we obtain

$$\sum_{n=0}^{\infty} \frac{h_1 - \lambda_n}{\gamma_n} = 0.$$

Similarly, we have

$$\sum_{n=0}^{\infty} \frac{H_1 - \lambda_n}{k_n^2 \gamma_n} = 0.$$

Using (6) in these equalities we get the representation of h_1 and H_1 by the spectral data:

$$h_1 = \delta \sum_{n=0}^{\infty} \frac{\lambda_n}{\gamma_n}, \quad H_1 = \rho \sum_{n=0}^{\infty} \frac{\lambda_n}{k_n^2 \gamma_n}. \quad (11)$$

Using the transformation operators (see [5]), we can write

$$\varphi(x, \lambda) = \dot{\varphi}(x, \lambda) + \int_{-x}^x G(x, \tau) \dot{\varphi}(\tau, \lambda) d\tau, \quad (12)$$

where

$$\dot{\varphi}(x, \lambda) := (h_1 - \lambda) \cos \sqrt{\lambda} x + (h_2 - \lambda h) \frac{\sin \sqrt{\lambda} x}{\sqrt{\lambda}}$$

is the solution of the equation $-y''(x) = \lambda y(x)$ satisfying the initial conditions

$$\dot{\varphi}(0, \lambda) = -\lambda + h_1, \quad \dot{\varphi}'(0, \lambda) = -\lambda h + h_2,$$

$G(x, \tau)$ is a real-valued continuous function and

$$G(x, x) = \frac{1}{2} \int_0^x q(t) dt.$$

Theorem 3. For each fixed $x \in (0, \pi]$ the kernel $G(x, \tau)$ satisfies the following equation:

$$A(x, t) + \int_{-x}^x G(x, \tau) A(\tau, t) d\tau = 0, \quad 0 \leq t \leq x \leq \pi, \quad (13)$$

where

$$A(x, t) := \sum_{n=0}^{\infty} \frac{\dot{\varphi}(x, \lambda_n) \cos \sqrt{\lambda_n} t}{\gamma_n}.$$

Proof. Using equality (12) we obtain

$$\begin{aligned} \sum_{n=0}^N \frac{\varphi(x, \lambda_n) \cos \sqrt{\lambda_n} t}{\gamma_n} &= \sum_{n=0}^N \frac{\dot{\varphi}(x, \lambda_n) \cos \sqrt{\lambda_n} t}{\gamma_n} + \\ &+ \sum_{n=0}^N \frac{\cos \sqrt{\lambda_n} t}{\gamma_n} \int_{-x}^x G(x, \tau) \dot{\varphi}(\tau, \lambda_n) d\tau. \end{aligned}$$

Applying Lemma 2. to $v(t, \lambda) = \cos \sqrt{\lambda} t$ we get the statement of the theorem.

We denote

$$\begin{aligned} F(x, t) &:= \frac{(h_1 - \lambda_0) \dot{\varphi}(x, \lambda_0) \cos \sqrt{\lambda_0} t}{\gamma_0} + \frac{(h_1 - \lambda_1) \dot{\varphi}(x, \lambda_1) \cos \sqrt{\lambda_1} t}{\gamma_1} + \\ &+ \sum_{n=0}^{\infty} \left(\frac{(h_1 - \lambda_{n+2}) \dot{\varphi}(x, \lambda_{n+2}) \cos \sqrt{\lambda_{n+2}} t}{\gamma_{n+2}} - \frac{(\cos nx + h \frac{\sin nx}{n}) \cos nt}{\alpha_n^0} \right), \end{aligned}$$

where

$$\alpha_n^0 = \begin{cases} \frac{\pi}{2}, & n \geq 1 \\ \pi, & n = 0. \end{cases}$$

By differentiating the equation (13) twice with respect to t we find

$$\begin{aligned} F(x, t) + h + G(x, t) + G(x, -t) + \\ + h \int_t^x (G(x, \tau) - G(x, -\tau)) d\tau + \int_{-x}^x G(x, \tau) F(\tau, t) d\tau = 0. \end{aligned}$$

Putting $x = t = 0$ we obtain

$$h = -\frac{(h_1 - \lambda_0)^2}{\gamma_0} - \frac{(h_1 - \lambda_1)^2}{\gamma_1} - \sum_{n=2}^{\infty} \left(\frac{(h_1 - \lambda_n)^2}{\gamma_n} - \frac{1}{\alpha_{n-2}^0} \right). \quad (14)$$

Analogously, we have

$$H = -\frac{(H_1 - \lambda_0)^2}{k_0^2 \gamma_0} - \frac{(H_1 - \lambda_1)^2}{k_1^2 \gamma_1} - \sum_{n=2}^{\infty} \left(\frac{(H_1 - \lambda_n)^2}{k_n^2 \gamma_n} - \frac{1}{\alpha_{n-2}^0} \right). \quad (15)$$

Theorem 4. Suppose that $\{\lambda_n, \gamma_n\}_{n \geq 0}$ and $\{\tilde{\lambda}_n, \tilde{\gamma}_n\}_{n \geq 0}$ are the spectral data of the problems $\mathcal{P}(q, h, h_1, h_2, H, H_1, H_2)$ and $\mathcal{P}(\tilde{q}, \tilde{h}, \tilde{h}_1, \tilde{h}_2, \tilde{H}, \tilde{H}_1, \tilde{H}_2)$, respectively. If

$$\lambda_n = \tilde{\lambda}_n, \quad \gamma_n = \tilde{\gamma}_n, \quad n \geq 0,$$

then

$$q(x) = \tilde{q}(x) \text{ a.e. on } (0, \pi),$$

$$h = \tilde{h}, \quad h_1 = \tilde{h}_1, \quad h_2 = \tilde{h}_2, \quad H = \tilde{H}, \quad H_1 = \tilde{H}_1, \quad H_2 = \tilde{H}_2.$$

Proof. Using (6), (11), (14) and (15) we obtain

$$h = \tilde{h}, \quad h_1 = \tilde{h}_1, \quad h_2 = \tilde{h}_2, \quad H = \tilde{H}, \quad H_1 = \tilde{H}_1, \quad H_2 = \tilde{H}_2.$$

The result now follows from [2, Theorem 5.2].

The author is grateful to Professor H. M. Huseynov for the problem statement and for useful discussions.

References

- [1]. Binding P.A., Browne P.J. and Seddighi K. *Sturm–Liouville problems with eigenparameter dependent boundary conditions*. Proc. Edinburgh Math. Soc. (2), 1993, v.37, No.1, pp.57–72.
- [2]. Binding P.A., Browne P.J. and Watson B.A. *Inverse spectral problems for Sturm–Liouville equations with eigenparameter dependent boundary conditions*. J. London Math. Soc. (2), 2000, v.62, No.1, pp.161–182.
- [3]. Fulton C.T. *Two-point boundary value problems with eigenvalue parameter contained in the boundary conditions*. Proc. Roy. Soc. Edinburgh Sect. A, 1977, v.77, No.3-4, pp.293–308.
- [4]. Kerimov N.B. and Allakhverdiev T.I. *On a boundary value problem. I*. Differ. Uravn., 1993, v.29, No.1, pp.54–60.
- [5]. Marchenko V.A. *The Sturm–Liouville operators and their applications*. Kiev: Naukova Dumka, 1977. (Russian) (*Engl. transl.*: (Basel: Birkhäuser Verlag, 1986.))
- [6]. Yurko V.A. *Inverse spectral problems and their applications*. Saratov: Izd-vo Saratovskogo pedinstituta, 2001. (Russian)

Namig J. Guliyev

Institute of Mathematics and Mechanics of NAS of Azerbaijan

9 F.Agayev str., AZ1141, Baku, Azerbaijan.

E-mail: njguliyev@yahoo.com

Received June 29, 2006; Revised September 28, 2006.

Translated by author