



HAL
open science

On numbers satisfying Robin's inequality, properties of the next counterexample and improved specific bounds

Robert Vojak

► **To cite this version:**

Robert Vojak. On numbers satisfying Robin's inequality, properties of the next counterexample and improved specific bounds. 2019. hal-02085868v1

HAL Id: hal-02085868

<https://hal.science/hal-02085868v1>

Preprint submitted on 31 Mar 2019 (v1), last revised 18 May 2020 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

On numbers satisfying Robin's inequality, properties of the next counterexample and improved specific bounds

Robert VOJAK

Croatia & France

Abstract

Define $s(n) := n^{-1}\sigma(n)$ ($\sigma(n) := \sum_{d|n} d$) and $\omega(n)$ is the number of prime divisors of n . One of the properties of s plays a central role: $s(p^a) > s(q^b)$ if $p < q$ are prime numbers, with no special condition on a, b other than $a, b \geq 1$. This result, combined with the Multiplicity Permutation theorem, will help us establish properties of the next counterexample (say c) to Robin's inequality $s(n) < e^\gamma \log \log n$. The number c is superabundant, and $\omega(c)$ must be greater than a number close to one billion. In addition, the ratio $p_{\omega(c)}/\log c$ has a lower and upper bound. At most $\omega(c)/14$ multiplicity parameters are greater than 1. Last but not least we apply simple methods to sharpen Robin's inequality for various categories of numbers.

1. Introduction

The following notations will be used throughout the article :

- γ : Euler's constant,
- $\lfloor x \rfloor$ denotes the largest integer not exceeding x
- p_i : i -th prime number,
- $\sigma(n) = \sum_{d|n} d$,
- $\varphi(n)$ counts the positive integers up to a given integer n that are relatively prime to n
- $\omega(n)$ is the number of distinct primes dividing n ,
- $N_i = \prod_{j \leq i} p_j$ (primorial),
- $\text{rad}(n) = \prod_{\substack{p|n \\ p \text{ prime}}} p$, radical of n .

Email address: vojakrob@gmail.com (Robert VOJAK).

Define $s(n) = \frac{\sigma(n)}{n}$, $f(n) = \frac{n}{\varphi(n)}$.

A number n satisfies *Robin's inequality*[3] if and only if for all $n \geq 5041$.

$$\forall n \geq 5041 \quad s(n) < e^\gamma \log \log n$$

Définition 1.1 A number n is *superabundant (SA)* [1] if for all $m < n$, we have $s(m) < s(n)$.

Définition 1.2 A number n is a *Hardy-Ramanujan number (HR)* [6] if $n = \prod_{i \leq \omega(n)} p_i^{a_i}$ with $a_i \geq a_{i+1} \geq 0$ for all $i \leq \omega(n) - 1$.

All the results presented hereafter concern Robin's inequality, which is known to be equivalent to Riemann's Hypothesis[3]. We start by establishing an important result about the multiplicities in the prime factorization of a number n , and how switching these multiplicities affects n and $s(n)$.

Next, we give some categories of numbers satisfying Robin's inequality, and for those who do not, we exhibit some of their properties (theorem 1.4).

The last part is devoted to sharpening Robin's inequality in specific cases.

In the sequel, the notation q_1, q_2, \dots will be used to define a finite increasing sequence of prime numbers. The following theorems represent our main results (the proofs will be given later on) :

Theorem 1.1 Let $n = \prod_{i=1}^{\omega(n)} q_i^{a_i}$ be a positive integer. If

- $\omega(n) \geq 18$ and $\#\{i \leq \omega(n) ; a_i \neq 1\} \geq \frac{\omega(n)}{2}$, or
- $\omega(n) \geq 39$ and $\#\{i \leq \omega(n) ; a_i \neq 1\} \geq \frac{\omega(n)}{3}$, or
- $\omega(n) \geq 969672728$ and $\#\{i \leq \omega(n) ; a_i \neq 1\} \geq \frac{\omega(n)}{14}$

then n satisfies Robin's inequality.

Theorem 1.2 Let $n = \prod_{i \leq k} q_i^{a_i}$. For all positive integers q , define

$$M(k) = e^{e^{-\gamma} f(N_k)} - \log N_k \tag{1}$$

$$M_k(q) = 1 + \frac{M(k)}{\log q}$$

If for some $i \leq \omega(n)$, we have $a_i \geq M_{\omega(n)}(q_i)$, then the integer n satisfies Robin's inequality, regardless of the multiplicities of other prime divisors.

For instance, if $n = \prod_{i \leq 5} p_i^{a_i}$, $M_5(2) = 11.3367 \dots$. All integers of the form $2^{a_1} \prod_{i=2}^5 p_i^{a_i}$ satisfy Robin's inequality if $a_1 \geq 12$, and for all $a_2, a_3, a_4, a_5 \geq 1$.

Corollary 1.3 Let n be a positive integer. If $f(N_{\omega(n)}) - e^\gamma \log \log N_{\omega(n)} \leq 0$, then n satisfies Robin's inequality.

Proof: If $f(N_{\omega(n)}) - e^\gamma \log \log N_{\omega(n)} \leq 0$, we have $M(k) \leq 0$, and for all $i \leq \omega(n)$ and all prime numbers p ,

$$a_i \geq 1 \geq M_{\omega(n)}(p)$$

We conclude using theorem 1.2. □

Theorem 1.4 Let $c = \prod_{i=1}^{\omega(n)} q_i^{c_i}$ ($c \geq 5041$) be the least number (if it exists) not satisfying Robin's inequality.

The following properties hold :

1. c is superabundant,
2. $\omega(c) \geq 969\,672\,728$,
3. $\#\{i \leq \omega(c); c_i \neq 1\} < \frac{\omega(c)}{14}$,
4. $e^{-\frac{1}{\log p_{\omega(c)}}} < \frac{p_{\omega(c)}}{\log c} < 1$,
5. for all $1 < i \leq \omega(c)$,

$$p_i^{c_i} < \min\left(2^{c_i+2}, p_i e^{M(k)}\right)$$

2. Preliminary results

We will need some properties about the functions s and f to prove the above-mentioned theorems.

Lemma 2.1 Let p, q be two prime numbers, and k, a and b positive integers.

1. the functions s and f are multiplicative
2. $f(n) = \prod_{\substack{p|n \\ p \text{ prime}}} \frac{p}{p-1}$ and $f(n) = f(\text{rad}(n))$.
3. $1 + \frac{1}{p} \leq s(p^k) < \frac{p}{p-1}$ and $\lim_{k \rightarrow \infty} s(p^k) = \frac{p}{p-1} = f(p)$
4. If $p < q$, $s(p^a) > s(q^b)$ for all positive integers a, b
5. $s(p^k)$ increases as $k > 0$ increases, decreases as p increases,
6. $\frac{s(p^{k+1})}{s(p^k)}$ decreases as k increases,
7. If $a > b > 0$, $\frac{s(p^a)}{s(p^b)}$ decreases as p increases,
8. Let p be a prime number, and q any prime number, $q < p(p+1)$. Then, for all $a, b \geq 1$

$$\frac{s(p^{a+1})}{s(p^a)} < s(q^b)$$

For instance : $s(p_n^{a+1}) < s(p_n^a) s(p_{n+1}^b)$ for all $a, b \geq 1$.

9. If $m, n \geq 2$, $s(mn) < s(m)s(n)$.

IMPORTANT : note that $s(p^a) > s(q^b)$ requires only to have $p < q$, and no condition on positive integers a, b .

Proof:

1. The multiplicativity of s and f is a consequence of the multiplicativity of σ and φ ,
2. The proof is straightforward : recall that $\text{rad}(p^k) = p$, $\varphi(p^k) = p^k - p^{k-1}$, and hence $f(p^k) = \frac{p}{p-1} = f(\text{rad}(p^k))$, and use the multiplicativity of f and rad to conclude.
3. It is deduced from the properties $\sigma(p^k) = \frac{p^{k+1}-1}{p-1}$ and $\varphi(p^k) = p^k - p^{k-1}$. To prove the asymptotic result, use $s(p^k) = \frac{p - \frac{1}{p^k}}{p-1}$,

4. We have

$$1 + \frac{1}{p} \geq \frac{q}{q-1}$$

when $q > p$, and we deduce

$$s(p^a) \geq 1 + \frac{1}{p} \geq \frac{q}{q-1} > s(q^b)$$

5. Use $s(p^k) = \frac{p - \frac{1}{p^k}}{p-1}$, and we can conclude that $s(p^k)$ increases as k increases. Use (2.1) with $a = b = k$ to prove that if $p < q$, $s(p^k) > s(q^k)$.
6. We have

$$\frac{s(p^{k+1})}{s(p^k)} = \frac{p^{k+2} - 1}{p^{k+2} - p}$$

which decreases as p^k increases.

Let us show that it also decreases as p increases. The sign of the first derivative of $p \mapsto \frac{s(p^{k+1})}{s(p^k)}$ is the same as the sign of $p^{k+1}(k+2) - p^{k+2}(k+1) - 1 = p^{k+1}(k+2 - p(k+1)) - 1 \leq p^{k+1}(k+2 - 2(k+1)) - 1 = -p^{k+1}(k-1) < 0$.

7. The equality

$$\frac{s(p^a)}{s(p^b)} = \prod_{k=b}^{a-1} \frac{s(p^{k+1})}{s(p^k)}$$

along with the monotonicity of each fraction of the product yields the desired result.

8. The function $a \mapsto \frac{s(p^{a+1})}{s(p^a)}$ is decreasing, and we have

$$\frac{s(p^{a+1})}{s(p^a)} - s(q) < \frac{s(p^2)}{s(p)} - 1 - \frac{1}{q} = \frac{1}{p(p+1)} - \frac{1}{q}$$

Hence

$$\frac{s(p^{a+1})}{s(p^a)} < s(q) \leq s(q^b)$$

9. Let $n = \prod_{p|n} p^{a_p}$ and $m = \prod_{p|m} p^{b_p}$. We have

$$m n = \prod_{\substack{p|n \\ p \nmid m}} p^{a_p} \prod_{\substack{p \nmid n \\ p|m}} p^{b_p} \prod_{\substack{p|n \\ p|m}} p^{a_p + b_p}$$

yielding

$$s(mn) = \prod_{\substack{p|n \\ p \nmid m}} s(p^{a_p}) \prod_{\substack{p|n \\ p \nmid m}} s(p^{b_p}) \prod_{\substack{p|n \\ p \nmid m}} s(p^{a_p+b_p})$$

If p is a prime number, and a, b two positive integers, then

$$s(p^{a+b}) - s(p^a)s(p^b) = -\frac{(p^a-1)(p^b-1)}{p^{a+b-1}(p-1)^2} < 0$$

Now we have

$$s(mn) \leq \prod_{\substack{p|n \\ p \nmid m}} s(p^{a_p}) \prod_{\substack{p|n \\ p \nmid m}} s(p^{b_p}) \prod_{\substack{p|n \\ p \nmid m}} (s(p^{a_p})s(p^{b_p})) = s(m)s(n)$$

□

Lemma 2.2 For all positive integers $n \geq 2$, we have $s(n) < f(N_{\omega(n)})$

Proof: Let $n = \prod_{i=1}^k q_i^{a_i}$ where q_1, q_2, q_3, \dots is a finite increasing sequence of prime numbers. Therefore, for all i , $p_i \leq q_i$. Set $m = \prod_{i=1}^k p_i^{a_i}$. Clearly, $m \leq n$ and

$$\frac{s(p_i^{a_i})}{s(q_i^{a_i})} \geq 1$$

yielding

$$s(n) \leq s(m) < f(m) = f(N_{\omega(n)})$$

□

Theorem 2.3 (Multiplicity Permutation) Let n be a positive integer, p and q be two prime divisors of n ($p < q$), and a and b their multiplicity respectively.

Consider the integer n^* obtained by switching a and b . If $a < b$

$$n^* < n \quad \text{and} \quad s(n^*) > s(n)$$

and if $a > b$

$$n^* > n \quad \text{and} \quad s(n^*) < s(n)$$

Proof: Switching a and b means that $n^* = n \left(\frac{p}{q}\right)^{b-a}$. Hence, if $a < b$, we have $n^* < n$ and

$$\frac{s(n)}{s(n^*)} = \frac{s(p^a)s(q^b)}{s(p^b)s(q^a)} < \frac{s(p^a)s(p^b)}{s(p^b)s(p^a)} = 1$$

Indeed, since $a < b$, we know from lemma 2.1 that $\frac{s(q^b)}{s(q^a)}$ decreases as q increases, and since $q > p$, we have

$$\frac{s(q^b)}{s(q^a)} < \frac{s(p^b)}{s(p^a)}$$

The proof for the case $a > b$ follows the same logic.

□

Let $n = \prod_{i=1}^k q_i^{a_i}$, and b_1, b_2, \dots, b_k be a reordering of a_1, a_2, \dots, a_k such that $b_i \geq b_{i+1}$. Define the function A and H as follows :

$$A \left(\prod_{i \leq k} q_i^{a_i} \right) = \prod_{i \leq k} q_i^{b_i} \quad (2)$$

$$H \left(\prod_{i \leq k} q_i^{a_i} \right) = \prod_{i \leq k} p_i^{b_i} \quad (3)$$

We claim that $A(n) \leq n$ and $s(n) \leq s(A(n))$: let n_1 be the integer obtained by switching a_1 and b_1 . Using theorem 2.3, we have $n_1 \leq n$ and $s(n_1) \geq s(n)$. Continue by switching b_2 and the multiplicity of p_2 in the prime factorization of n_1 . The result is an integer n_2 such that $n_2 \leq n_1 \leq n$ and $s(n_2) \geq s(n_1) \geq s(n)$. Repeat the process a total of $k - 1$ times, and you obtain a positive integer n_{k-1} such that

$$n_{k-1} = \prod_{i \leq k} q_i^{b_i} = A(n) \quad \text{with } b_i \geq b_{i+1}$$

$$n_{k-1} \leq n \quad \text{and} \quad s(n_{k-1}) \geq s(n)$$

This proves that $A(n) \leq n$ and $s(A(n)) \geq s(n)$. The same inequalities hold for the function H . The function H replaces all the prime divisors by the first prime divisors, and reorganize the multiplicities in a decreasing order. It is easy to check that $H(n) \leq n$ and $s(H(n)) \geq s(n)$. Indeed, note that $H(n) \leq A(n) \leq n$ and $s(H(n)) \geq s(A(n)) \geq s(n)$. If $H(n)$ satisfies Robin's inequality, so does n . Indeed,

$$s(n) \leq s(H(n)) < e^\gamma \log \log H(n) < e^\gamma \log \log n$$

thus proving :

Theorem 2.4 *If Robin's inequality is valid for all Hardy-Ramanujan numbers greater than 5040, then it is valid for all positive integers greater than 5040.*

Note : compare this result to proposition 5.1 in [6] : "if Robins inequality holds for all Hardy-Ramanujan integers $5041 \leq n \leq x$, then it holds for all integers $5041 \leq n \leq x$ ".

3. Numbers satisfying Robin's inequality

Some categories of numbers satisfy Robin's inequality. Most of them have already been identified in previous works[6], and we report these results below. Note that the proofs are not the original ones. Instead, we used simple methods to provide shorter and simpler proofs. To do so, we will need the following inequalities ([3], th. 2).

Theorem 3.1 *For all $n \geq 3$,*

$$s(n) \leq e^\gamma \log \log n + \frac{0.6483}{\log \log n}$$

and [2](th. 15)

Theorem 3.2 For all $n \geq 3$,

$$f(n) \leq e^\gamma \log \log n + \frac{2.51}{\log \log n}$$

We will start with primorials. We will then investigate odd positive integers, and integers n such that $\omega(n) \leq 4$ (all known integers not satisfying Robin's inequality are such that $\omega(n) \leq 4$), and we will conclude with square-free and square-full integers.

3.1. Primorial numbers

Theorem 3.3 All primorials N_k ($k \geq 4$) satisfy Robin's inequality :

$$s(N_k) = \prod_{i=1}^k \left(1 + \frac{1}{p_i}\right) < e^\gamma \log \log N_k$$

The following inequality is a bit sharper (for $k \geq 2$) :

$$\prod_{i=1}^k \left(1 + \frac{1}{p_i}\right) \leq \frac{3}{4} \left(e^\gamma \log \log N_k + \frac{2.51}{\log \log N_k} \right)$$

Proof: Let $k \geq 2$ be an integer. We have

$$s(N_k) = s(2)s\left(\frac{N_k}{2}\right) \leq s(2)f\left(\frac{N_k}{2}\right) = \frac{s(2)}{f(2)}f(N_k) = \frac{3}{4}f(N_k)$$

Using theorem 3.2, we have

$$\begin{aligned} s(N_k) - e^\gamma \log \log N_k &\leq \frac{3}{4} \left(e^\gamma \log \log N_k + \frac{2.51}{\log \log N_k} \right) - e^\gamma \log \log N_k \\ &\leq -0.25e^\gamma \log \log N_k + \frac{1.8825}{\log \log N_k} \\ &< 0 \end{aligned}$$

if $k \geq 6$. For $k \leq 5$, only the numbers $N_1 = 2$, $N_2 = 6$ and $N_3 = 30$ do not satisfy Robin's inequality. \square

3.2. Odd integers

The following theorem can be found in [6], but the proof is not the original one. A very simple proof is given instead.

Theorem 3.4 Any odd positive integer n distinct from 3, 5 and 9 satisfies Robin's inequality.

Proof: Let $n \geq 3$ be a positive odd integer. Using theorem 3.1, we have

$$s(n) = \frac{s(2n)}{s(2)} < \frac{2}{3} \left(e^\gamma \log \log(2n) + \frac{0.6483}{\log \log(2n)} \right) < e^\gamma \log \log n$$

for all $n \geq 17$. Indeed, set

$$g(x) = \frac{2}{3} \left(e^\gamma \log \log(2x) + \frac{0.6483}{\log \log(2x)} \right) - e^\gamma \log \log x$$

We have, for $x \geq 2$,

$$g'(x) < -\frac{1.23(\log \log(2x))^2 + (0.43 + 0.6(\log \log(2x))^2) \log x}{x(\log x)(\log(2x))(\log \log(2x))^2} < 0$$

When $x \geq 17$, $g(x) \leq g(17) < 0$. Therefore, all odd integers greater than 15 satisfy Robin's inequality. For odd integers up to 15, only 3, 5 and 9 do not satisfy Robin's inequality. \square

3.3. Integers n such that $\omega(n) \leq 4$

Theorem 3.5 *Let n such that $\omega(n) \leq 4$. Then n satisfies Robin's inequality if and only if it does not belong to $\mathcal{C} = \mathcal{C}_1 \cup \mathcal{C}_2 \cup \mathcal{C}_3 \cup \mathcal{C}_4$ with*

$$\mathcal{C}_1 = \{2, 3, 4, 5, 8, 9, 16\}$$

$$\mathcal{C}_2 = \{6, 10, 12, 18, 20, 24, 36, 48, 72\}$$

$$\mathcal{C}_3 = \{30, 60, 84, 120, 180, 240, 360, 720\}$$

$$\mathcal{C}_4 = \{840, 2520, 5040\}$$

For each $k = 1, 2, 3$ or 4 , \mathcal{C}_k is the set of counterexamples n to Robin's inequality such that $\omega(n) = k$.

Proof: Numbers n under 5041 that do not satisfy Robin's inequality are well known, and the number of prime divisors $\omega(n)$ of these counterexamples does not exceed 4. We will now show that if $\omega(n) \leq 4$, the only counterexamples are the elements of \mathcal{C} .

Let n be an integer such that $\omega(n) \leq 4$. We have (lemma 2.2) using the monotonicity of $k \mapsto f(N_k)$

$$s(n) < f(N_{\omega(n)}) \leq f(N_4) = 4.375$$

Therefore, if $n > 116144$, $\log \log n \geq e^{-\gamma} 4.375$,

$$e^\gamma \log \log n \geq 4.375 > s(n)$$

and Robin's inequality is satisfied if $n > 116144$ and $\omega(n) \leq 4$. Numerical computations confirm that there are no counterexamples to Robin's inequality in the range $5041 < n \leq 116144$. \square

3.4. Square-free integers

A positive integer n is called *square-free* if for every prime number p , p^2 is not a factor of n . Hence n has the form

$$n = \prod_{i \leq k} q_i$$

The following two theorems can be found in [6], but not the proofs (we provide simpler proofs).

Theorem 3.6 *Any square-free positive integer distinct from 2, 3, 5, 6, 10, 30 satisfy Robin's inequality.*

Proof: If n is square-free and even, the prime number 2 has multiplicity $a_1 = 1$. This implies

$$\frac{s(n)}{s(2n)} = \frac{s(2)}{s(4)} = \frac{6}{7}$$

and using theorem 3.1

$$s(n) < \frac{6}{7} \left(e^\gamma \log \log(2n) + \frac{0.6483}{\log \log(2n)} \right) < e^\gamma \log \log n$$

if $n > 418$. If $n \leq 418$, numerical computations show that only the square-free numbers 2, 6, 10, 30 do not satisfy Robin's inequality.

If n is odd, see theorem 3.4. □

3.5. Square-full integers

Let n be an integer. If for every prime divisor p of n , we have $p^2|n$, the integer n is said to be *square-full*. This result can also be found in [6]. The proof we give here is shorter and simpler.

Theorem 3.7 *The only square-full integers not satisfying Robin's inequality are 4, 8, 9, 16 and 36.*

Proof:

Case 1 : $\omega(n) \leq 4$. The answer is given in theorem 3.5. All square-full counterexamples are 4, 8, 9, 16, 36.

Case 2 : $\omega(n) \geq 5$. Let $n = \prod_{i=1}^k q_i^{a_i}$. Since n is square-full, we have $n \geq N_k^2$ and $s(n) < f(N_k)$ (lemma 2.2).

We can now write, using theorem 3.2,

$$s(n) - e^\gamma \log \log n < f(N_k) - e^\gamma \log \log N_k^2 < \frac{2.51}{\log \log N_k} - 1.2345 < -0.0083$$

□

4. Proof of theorem 1.1

Lemma 4.1 *Let $M(k) = e^{e^{-\gamma} f(N_k)} - \log N_k$*

1. *For all $k \geq 39$, we have $M(k) \leq \log N_{\lfloor \frac{k}{3} \rfloor}$*
2. *For all $k \geq 18$, we have $M(k) \leq \log N_{\lfloor \frac{k}{2} \rfloor}$*
3. *For all $k \geq 969672728$, we have $M(k) \leq \log N_{\lfloor \frac{k}{14} \rfloor}$*

Proof: Let us prove property 1.

We claim that for all $k > 38$, we have $M(3k+j) \leq \log N(k)$ for all $0 \leq j < 3$. Using theorem 3.2, we have

$$M(k) \leq \left(e^{\frac{1.41}{\log \log N_k}} - 1 \right) \log N_k$$

Define

$$\epsilon_k = e^{\frac{1.41}{\log(3k \log(3k))}} - 1 - \frac{k \log k}{(3k+2)(\log(3k+2) + \log \log(3k+2))}$$

Using the inequalities [4]

$$k \log k < \log N_k < k (\log k + \log \log k) \quad \text{for } k \geq 13$$

we see that

$$\epsilon_k \geq e^{\frac{1.41}{\log \log N_{3k}}} - 1 - \frac{\log N_k}{\log N_{3k+2}}$$

We claim that ϵ_k is decreasing, and that $\epsilon_k < 0$ for $k > 109$. Indeed, rewrite the term

$$\frac{k \log k}{(3k+2)(\log(3k+2) + \log \log(3k+2))} = \frac{k}{3k+2} \times \frac{\log k}{\log(3k+2)} \times \frac{\log(3k+2)}{\log(3k+2) + \log \log(3k+2)}$$

as a product of three increasing functions : to prove they are increasing, you can use the monotonicity of $x \mapsto \frac{\log x}{x}$ ($x \geq e$), and the monotonicity of $x \mapsto \frac{\log x}{\log(x+a)}$, where a is a positive real number .

We have $\epsilon_{109} < -0.0003$ implying for all $0 \leq i < 3$

$$e^{\frac{1.41}{\log \log N_{3k+i}}} - 1 < e^{\frac{1.41}{\log \log N_{3k}}} - 1 \leq \frac{\log N_k}{\log N_{3k+2}} \leq \frac{\log N_k}{\log N_{3k+i}}$$

yielding

$$M(3k+i) \leq \left(e^{\frac{1.41}{\log \log N_{3k+i}}} - 1 \right) \log N_{3k+i} \leq \log N_k$$

or simply

$$M(k) \leq \log N_{\lfloor \frac{k}{3} \rfloor}$$

for $k \geq 109$, and was confirmed for $39 \leq k < 109$ with the help of numerical computations.

• **Proof of property 2 :** This result is deduced the same way : consider the function

$$\epsilon_k = e^{\frac{1.41}{\log(2k \log(2k))}} - 1 - \frac{k \log k}{(2k+1)(\log(2k+1) + \log \log(2k+1))}$$

We have

$$\epsilon_k \geq e^{\frac{1.41}{\log \log N_{2k}}} - 1 - \frac{\log N_k}{\log N_{2k+1}}$$

and ϵ_k is a decreasing function with $\epsilon_{28} < -0.003$ yielding

$$M(k) \leq \log N_{\lfloor \frac{k}{2} \rfloor}$$

for $k > 27$, and was confirmed for $18 \leq k < 28$ with the help of numerical computations.

• **Proof of property 3 :** we proceed the same way. Consider the function

$$\epsilon_k = e^{\frac{1.41}{\log(14k \log(14k))}} - 1 - \frac{k \log k}{(14k+13)(\log(14k+13) + \log \log(14k+13))}$$

We have

$$\epsilon_k \geq e^{\frac{1.41}{\log \log N_{14k}}} - 1 - \frac{\log N_k}{\log N_{14k+13}}$$

and ϵ_k is a decreasing function with $\epsilon_{969672728} < -0.001$ yielding

$$M(k) \leq \log N_{\lfloor \frac{k}{14} \rfloor}$$

for $k \geq 969672728$. □

Proof of theorem 1.1

We will need the function H defined in (3).

$$H(n) \leq n \quad \text{and} \quad s(H(n)) \geq s(n)$$

The integer $H(n)$ is a Hardy-Ramanujan number. In addition, if $H(n)$ satisfies Robin's inequality, so does n . Indeed,

$$s(n) \leq s(H(n)) < e^\gamma \log \log H(n) \leq e^\gamma \log \log n$$

Note that the value of $\#\{i \leq \omega(n); a_i \neq 1\}$ is the same if we replace n with $H(n)$, and that $\omega(H(n)) = \omega(n)$. So without loss of generality, we will prove the theorem for Hardy-Ramanujan numbers only.

Let n be a Hardy-Ramanujan number. Define $j(n) := \min\{1 \leq j \leq \omega(n); M(\omega(n)) \leq \log N_j\}$ and $i(n) := \#\{i \leq \omega(n); a_i \neq 1\}$. A sufficient condition for n to satisfy Robin's inequality is $i(n) \geq j(n)$. Since n is a Hardy-Ramanujan number, we have $a_i \geq 2$ for $i \leq i(n)$, and $a_i = 1$ otherwise, implying that $n \geq N_{i(n)}N_{\omega(n)}$ and yielding

$$s(n) - e^\gamma \log \log n < f(N_{\omega(n)}) - e^\gamma \log \log(N_{i(n)}N_{\omega(n)}) < 0$$

if $i(n) \geq j(n)$. Hence, n satisfies Robin's inequality.

If $\omega(n) \geq 18$, and if $\#\{i \leq \omega(n); a_i \neq 1\} \geq \frac{\omega(n)}{2}$, n satisfies Robin's inequality. Indeed, if $\omega(n) \geq 18$, lemma 4.1 tells us that $j(n) \leq \frac{\omega(n)}{2}$. Using the assumption $\#\{i \leq \omega(n); a_i \neq 1\} \geq \frac{\omega(n)}{2}$ yields $i(n) \geq j(n)$.

Proceed the same way with the two cases : $\omega(n) \geq 39$ and $\omega(n) \geq 969672728$ □

5. Proof of theorem 1.2

According to lemma 2.2, we have $s(n) < f(N_{\omega(n)})$ and

$$n = \prod_{j=1}^{\omega(n)} q_j^{a_j} \geq q_i^{a_i-1} \left(q_i \prod_{j \neq i} q_j^{a_j} \right) \geq q_i^{a_i-1} \left(p_i \prod_{j \neq i} p_j \right) = q_i^{a_i-1} N_{\omega(n)}$$

implying

$$s(n) - e^\gamma \log \log n < f(N_{\omega(n)}) - e^\gamma \log \log(q_i^{a_i-1} N_{\omega(n)})$$

If $a_i \geq M_{\omega(n)}(q_i)$ with

$$M_k(q) := 1 + \frac{e^{e^{-\gamma} f(N_k)} - \log N_k}{\log q}$$

then

$$f(N_{\omega(n)}) - e^\gamma \log \log(q_i^{a_i-1} N_{\omega(n)}) \leq 0$$

and therefore

$$s(n) < e^\gamma \log \log n$$

6. Proof of theorem 1.4

Proof of 1. The superabundance of the number c is proved in [7].

Proof of 2. In [5], numerical computations have confirmed that $c > 10^{10^{10}}$.

Using lemma 2.2, we have

$$e^\gamma \log \log c \leq s(c) < f(N_{\omega(c)}) < e^\gamma \left(\log p_{\omega(c)} + \frac{1}{\log p_{\omega(c)}} \right)$$

yielding

$$\log p_{\omega(c)} > \frac{1}{2} \left(\log \log c + \sqrt{\log \log c - 4} \right)$$

Since $\log \log c = 10 \log 10 + \log \log 10 > 23.85988$, we obtain

$$\log p_{\omega(c)} > 23.81789$$

yielding

$$\omega(c) \geq 969\,672\,728$$

Proof of 3. \star *Part 1* : $p_{\omega(c)} < \log c$

By definition, the counterexample c does not satisfy Robin's inequality :

$$s(c) \geq e^\gamma \log \log c$$

and since $c > 5040$ is the least one, the integer $c' = \frac{c}{p_{\omega(c)}}$ satisfy Robin's inequality, because $\omega(c') = \omega(c) - 1 > 4$. Therefore, we have

$$s\left(\frac{c}{p_{\omega(c)}}\right) < e^\gamma \log \log \frac{c}{p_{\omega(c)}}$$

yielding

$$\frac{s(c)}{s\left(\frac{c}{p_{\omega(c)}}\right)} > \frac{\log \log c}{\log \log \frac{c}{p_{\omega(c)}}} \quad (4)$$

Now use $\frac{s(c)}{s\left(\frac{c}{p_{\omega(c)}}\right)} = s(p_{\omega(c)}) = 1 + \frac{1}{p_{\omega(c)}}$, and using inequality $\log(1-x) < x$ ($x \neq 0$), we establish that

$$\log \log \frac{c}{p_{\omega(c)}} = \log \log c + \log \left(1 - \frac{\log p_{\omega(c)}}{\log c} \right) < \log \log c - \frac{\log p_{\omega(c)}}{\log c} \quad (5)$$

Inequality (4) becomes

$$\frac{1}{p_{\omega(c)}} > \frac{\log \log c}{\log \log \frac{c}{p_{\omega(c)}}} - 1 > \frac{\log \log c}{\log \log c - \frac{\log p_{\omega(c)}}{\log c}} - 1 = \frac{\log p_{\omega(c)}}{\log c \log \log c - \log p_{\omega(c)}}$$

and

$$p_{\omega(c)} \log p_{\omega(c)} < \log c \log \log c - \log p_{\omega(c)} < \log c \log \log c$$

implying $p_{\omega(c)} < \log c$.

★ *Part 2* : $\log c < p_{\omega(c)} e^{\frac{1}{\log p_{\omega(c)}}}$

The counterexample inequality $s(c) \geq e^\gamma \log \log c$ yields

$$e^\gamma \log \log c \leq s(c) < f(N_{\omega(c)}) < e^\gamma \left(\log p_{\omega(c)} + \frac{1}{\log p_{\omega(c)}} \right)$$

implying $\log c < p_{\omega(c)} e^{\frac{1}{\log p_{\omega(c)}}}$.

Proof of 4. See theorem 1.1.

Proof of 5. The proof of the inequality $p_i^{a_i} < 2^{a_i+2}$ can be found in [1] (lemma 1). The other inequality

$$p_i^{c_i} < p_i e^{M(\omega(c))}$$

is deduced from theorem 1.2.

7. Sharper bounds

We can use some of the ideas of this article to improve upper bounds for the arithmetic functions s and f .

7.1. Primorials

For instance, when dealing with primorials N_k , we can write

$$s(N_k) = \frac{s(2)}{s(4)} s(2N_k)$$

and then use an upper bound for $s(2N_k)$ (see theorem 3.1) to sharpen Robin's bound for $s(N_k)$. Or we can use another method :

Theorem 7.1 *For all $i \geq 2$, for all $n \geq i$,*

$$\prod_{k=1}^n \left(1 + \frac{1}{p_k} \right) < \alpha_i \left(e^\gamma \log \log N_n + \frac{2.51}{\log \log N_n} \right)$$

with

$$\alpha_i = \prod_{j=1}^i \left(1 - \frac{1}{p_j^2} \right)$$

Ex : for all $n \geq 4$,

$$\prod_{k=1}^n \left(1 + \frac{1}{p_k} \right) < 0.627 \left(e^\gamma \log \log N_n + \frac{2.51}{\log \log N_n} \right) < e^\gamma \log \log N_n$$

Proof: Let $i \geq 2, n \geq i$. We have

$$s(N_n) = s(N_i) s\left(\frac{N_n}{N_i}\right) < s(N_i) f\left(\frac{N_n}{N_i}\right) = \frac{s(N_i)}{f(N_i)} f(N_n) = \alpha_i f(N_n)$$

Use theorem 3.2 to conclude. \square

7.2. Odd numbers

Lemma 7.2 *If $n \geq 17$ is odd,*

$$\sigma(2n) < 2e^\gamma n \log \log(2n)$$

Proof: Write

$$s(2n) = \frac{s(2)}{s(4)} s(4n) \leq \frac{6}{7} \left(e^\gamma \log \log(4n) + \frac{0.6483}{\log \log(4n)} \right) < e^\gamma \log \log(2n)$$

To prove the last inequality, define

$$g(x) = \frac{6}{7} \left(e^\gamma \log \log(4x) + \frac{0.6483}{\log \log(4x)} \right) - e^\gamma \log \log(2x)$$

We have

$$g'(x) < -\frac{0.38 + 1.41(\log \log(4x))^2 + (0.55 + 0.25(\log \log(4x))^2) \log x}{x \log(2x) \log(4x) (\log \log(4x))^2} < 0$$

and $g(210) < 0 < g(209)$. The inequality $s(2n) < e^\gamma \log \log(2n)$ is valid for $n > 209$, and numerical computations show that this is also true when $17 \leq n \leq 209$. \square

Corollary 7.3 *If $n \geq 17$ is odd,*

$$\sigma(n) < \frac{2}{3} e^\gamma n \log \log(2n)$$

The following result is an improvement of theorem 1.2 in [6].

Proof: Use the previous result to get

$$s(n) = \frac{s(2n)}{s(2)} < \frac{2}{3} e^\gamma \log \log(2n)$$

The following result is an improvement of theorem 2.1 in [6]. \square

Theorem 7.4 *If $n \geq 3$ is odd,*

$$\frac{n}{\varphi(n)} \leq \frac{1}{2} \left(e^\gamma \log \log(2n) + \frac{2.51}{\log \log(2n)} \right) < e^\gamma \log \log n$$

Proof: Write

$$f(n) = \frac{f(2n)}{f(2)}$$

and use theorem 3.2 :

$$f(n) \leq \frac{1}{2} \left(e^\gamma \log \log(2n) + \frac{2.51}{\log \log(2n)} \right)$$

\square

7.3. The general case

In previous sections, we have mentioned superabundant numbers and pointed out some of their properties, thus confirming that all superabundant numbers are Hardy-Ramanujan numbers.

Robin's inequality can be enhanced for non superabundant numbers. Indeed, if n is not a superabundant number, there exists $m < n$ such that $s(m) \geq s(n)$. Let $I(n) = \{m < n; s(n) \leq s(m)\}$. The set $I(n)$ is not empty, and we can define $B(n) = \min I(n)$. We have

$$B(n) < n \quad \text{and} \quad s(n) \leq s(B(n))$$

yielding

$$s(n) \leq s(B(n)) < e^\gamma \log \log B(n)$$

if $B(n)$ satisfies Robin's inequality. Hence, the upper bound is better because $B(n) < n$.

But this method to sharpen Robin's bound does not say much about $B(n)$. In order to find an upper bound with exploitable informations, we can use the function H (see (3)). This method concerns non Hardy-Ramanujan numbers, whereas the previous method concerns the larger set of non superabundant numbers.

In (3), we defined the function H which transforms any number into a Hardy-Ramanujan number, leaves the multiplicities unchanged (but not the way they are ordered) and is such that $H(n) \leq n$ and $s(n) \leq s(H(n))$. If $H(n)$ satisfies Robin's inequality, then

$$s(n) \leq s(H(n)) < e^\gamma \log \log H(n)$$

The property $H(n) \leq n$ improves Robin's inequality, and unlike the function B , there is an explicit method to determine its value for all positive integers.

Références

- [1] Alaoglu, L. and Erdos, P. On highly composite and similar numbers. Trans. Amer. Math. Soc., 56 :448469, 1944.
- [2] Rosser, J. Barkley and Schoenfeld, Lowell. Approximate formulas for some functions of prime numbers. Illinois J. Math. 6 (1) : 6494, 1962.
- [3] Robin, Guy. Grandes valeurs de la fonction somme des diviseurs et hypothèse de Riemann, Journal de Mathématiques Pures et Appliquées, Neuvième Série, 63 (2) : 187213, ISSN 0021-7824, MR 0774171, 1984.
- [4] Massias, Jean-Pierre and Robin, Guy. Bornes effectives pour certaines fonctions concernant les nombres premiers. Journal de théorie des nombres de Bordeaux, Tome 8, no. 1, p. 215-242, 1996.
- [5] Briggs, K. Abundant numbers and the Riemann hypothesis. Experiment. Math., 15(2) :251256, 2006.
- [6] Choie, YoungJu and Lichiardopol, Nicolas and Moree, Pieter and Sole, Patrick. On Robin's criterion for the Riemann hypothesis, Journal de Théorie des Nombres de Bordeaux 19, 357372, 2007.
- [7] Akbary, Amir and Friggstad, Zachary. Superabundant numbers and the Riemann hypothesis. Amer. Math. Monthly, 116(3) :273275, 2009.