



HAL
open science

Deciphering multi-stage ore-forming processes in metasedimentary-rock-hosted orogenic gold deposit settings using LA ICP-MS sulphide analysis

Blandine Gourcerol, D J Kontak, P C Thurston, J A Petrus

► To cite this version:

Blandine Gourcerol, D J Kontak, P C Thurston, J A Petrus. Deciphering multi-stage ore-forming processes in metasedimentary-rock-hosted orogenic gold deposit settings using LA ICP-MS sulphide analysis. GAC-MAC-IAH / CNC Conference, May 2019, Québec, Canada. hal-02086616

HAL Id: hal-02086616

<https://hal.science/hal-02086616>

Submitted on 1 Apr 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Deciphering multi-stage ore-forming processes in metasedimentary-rock-hosted orogenic gold deposit settings using LA ICP-MS sulphide analysis

B. Gourcerol^{1*}, D.J. Kontak², P.C. Thurston² and J.A. Petrus²

¹ BRGM, French Geological Survey, 3 avenue C. Guillemin BP 36009, 45 060 Orléans Cedex 2, France

² Mineral Exploration Research Centre, Harquail School of Earth Sciences, Laurentian University, Sudbury, Ontario P3E 2C6, Canada

*Corresponding author: gourcerol.blandine@gmail.com; b.gourcerol@brgm.fr

Orogenic gold deposits span a spectrum in regards to setting and style of mineralization, nature of gold (e.g., invisible, coarse) and favourable host rock. In addition, deposit formation is often attributed to protracted multi-stage hydrothermal processes. That gold mineralization also depends on a variety of features, such as metal-source reservoirs, metal-transport processes and wall-rock stratigraphy, which add to the challenge of investigating origin of paleo-mineralized settings. Recent work using quantitative laser ablation inductively coupled plasma-mass spectrometry (LA ICP-MS) element distribution maps/profiles and their corresponding time slice datasets (TSD) provides new insight into identifying and assessing elemental paragenesis, multi-dimensional element coupling/decoupling processes, and corresponding mineralizing events. To further assess complexities of mineralization, application of geostatistical tools (e.g., multidimensional scaling, principal component analysis, linear discriminant analysis) and various innovative multi-element binary plots (e.g., Ag versus Au, Ni versus Co) is advised.

To illustrate the application of this methodology, the results LA ICP-MS mapping and data processing for Fe sulphides from several metasedimentary-rock-hosted Canadian gold systems are presented: three Archean Algoma-type BIF-hosted gold deposits (~ 4 Moz Au Meadowbank, ≥ 2.8 Moz Au Meliadine district, ~ 6 Moz Au Musselwhite) and eight slate-belt style vein gold deposits from the Paleozoic Meguma terrane (Nova Scotia). The maps and derived elemental plots generated from the various settings demonstrate that: 1) the gold mineralization present is the product of multi-stage processes; 2) elemental associations vary as mineralization progresses, such as the early growth history of sulphides versus later coupled dissolution-precipitation reactions; and 3) different metal-source reservoirs and stratigraphy influence the fluid signature. These results contribute to better deciphering the complex processes involved in the protracted evolution of these and other orogenic-type gold systems (e.g., remobilization of invisible gold from early sulphide, precipitation of visible gold in later sites, increase of Au fineness).